



More Efficient Production of Polymer Chemical from Biomass Glucose

INVENTORS • James Dumesic, Yomaira Pagan-Torres, Brent Shanks, Tianfu Wang

WARF: P120054US01

[View U.S. Patent No. 8,642,791 in PDF format.](#)

The Wisconsin Alumni Research Foundation (WARF) is seeking commercial partners interested in developing a cost-effective method of deriving the platform chemical 5-hydroxymethylfurfural from glucose by utilizing Lewis acid and Brønsted-Lowry acid catalysts.

OVERVIEW

Beyond its fuel potential, biomass is poised to supplant nonrenewable sources in the production of important building block chemicals. One such intermediary is 5-hydroxymethylfurfural (HMF), a primary component used to make plastic materials analogous to those derived from the petroleum polymer industry.

Efforts to render HMF from biomass traditionally have focused on dehydrating fructose in high yields using ionic liquids, which are expensive, susceptible to impurities and difficult to separate from the product stream. Glucose, another renewably derived carbohydrate, contains only one-twentieth of the appropriate tautomers in solution and because of low yield has remained an underutilized resource.

A new approach is needed that exploits glucose and glucose-containing saccharides and polymers at yields comparable to fructose, without the drawbacks of ionic acids.

THE INVENTION

UW-Madison researchers have developed a method to produce HMF from biomass-derived glucose in a two-phase reaction system using Lewis acid and Brønsted-Lowry acid catalysts.

Conducted in a two-phase reaction vessel, the method isomerizes the glucose feedstock material, chemically transforming it into fructose while simultaneously converting that fructose via dehydration into HMF. The aqueous component of the medium comprises the glucose and both types of homogenous acid catalysts. The solution may be saturated with sodium chloride. The organic extraction layer preferably contains one alkylphenol, and into this layer the HMF spontaneously separates.

The two sugar reactions—*isomerization and dehydration into HMF*—occur in tandem, result

THE WARF ADVANTAGE

Since its founding in 1925 as the patenting and licensing organization for the University of Wisconsin-Madison, WARF has been working with business and industry to transform university research into products that benefit society. WARF intellectual property managers and licensing staff members are leaders in the field of university-based technology transfer. They are familiar with the intricacies of patenting, have worked with researchers in relevant disciplines, understand industries and markets, and have negotiated innovative licensing strategies to meet the individual needs of business clients.



in high yields, increase separation efficiency and react in a system that can be conducted continuously or in batch fashion.

APPLICATIONS

- Production of furanic polyesters, polyamides and polyurethanes that can be used in the manufacture of plastics from renewable resources

KEY BENEFITS

- Combined, efficient reaction
- High yield from glucose
- No use of ionic liquids
- Amenable to one- or two-phase reaction method

STAGE OF DEVELOPMENT

The researchers have shown HMF yields of 67 percent.

ADDITIONAL INFORMATION

Related Technologies

[WARF reference number P06305US describes a cost-effective procedure to synthesize furan derivatives, including HMF, from fructose.](#)

Tech Fields

Clean Technology - Bio-based & renewable chemicals

Materials & Chemicals - Polymers

CONTACT INFORMATION

For current licensing status, please contact Mark Staudt at mstaudt@warf.org or 608-960-9845.

