



VeA, a Global Regulator of Secondary Metabolism, Can Increase Production of Secondary Metabolites

[View U.S. Patent No. 8,871,494 in PDF format.](#)

WARF: P09056US02

Inventors: Nancy Keller, Saori Campen

The Wisconsin Alumni Research Foundation (WARF) is seeking commercial partners interested in developing methods of using VeA, a newly identified global regulator of secondary metabolism, to increase or decrease production of secondary metabolites in fungi.

Overview

Microorganisms, such as fungi, produce a variety of secondary metabolites. These secondary metabolites display a broad range of activities, including antibiotic, immunosuppressant, phytotoxic and mycotoxic activities, and are useful for drug or technological development. For example, the antibiotic penicillin and the cholesterol-lowering drug lovastatin are secondary metabolites.

However, producing large amounts of secondary metabolites is difficult, and available techniques often provide unpredictable results. Because they are formed from a relatively small number of metabolic pathways, identifying the genes that control these pathways may provide an alternative method of generating secondary metabolites.

The inventors previously identified a global regulator of secondary metabolism, called *LaeA*, in fungi (see WARF reference number P02379US). Overexpression of the *laeA* gene upregulates production of secondary metabolites, greatly increasing penicillin production in *Aspergillus nidulans* and lovastatin production in *A. terreus*. On the other hand, deletion of *laeA* in *A. fumigatus* eliminates the production of gliotoxin and other secondary metabolites, decreasing the virulence of this human pathogen.

The Invention

UW-Madison researchers now have identified another global regulator of secondary metabolism, called VeA. VeA is a conserved protein that interacts with *LaeA* in an as yet unknown mechanism. Overexpression of *veA* upregulates secondary metabolism in *A. flavus* to a greater degree than overexpression of *laeA*. This gene could be used to increase the production of important natural products, including novel products with medicinal value.

Applications

- Increasing production of useful secondary metabolites, such as penicillin or lovastatin
- Decreasing production of toxic secondary metabolites, such as aflatoxin

Key Benefits

- Provides a simple method of increasing or decreasing secondary metabolite production

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete

- May enable new technologies, you agree to the storing of cookies and related technologies on your device. [See our privacy policy.](#)

- May be used to identify new secondary metabolite biosynthesis gene clusters



Additional Information

For More Information About the Inventors

- [Nancy Keller](#)

Related Technologies

- [See WARF reference number P02379US for more information on LaeA, another global regulator of secondary metabolism.](#)

Related Intellectual Property

- [View Continuation Patent in PDF format.](#)

Publications

- Bayram O., Krappmann S., Ni M., Bok J.W., Helmstaedt K., Valerius O., Braus-Stromeyer S., Kwon N.J., Keller N.P., Yu J.H. & Braus G.H. 2008. VelB/VeA/LaeA Coordinated Light Information, Fungal Development and Secondary Metabolism. *Science* 320, 1504-1506.
- Amaike S. & Keller N.P. 2009. Distinct Roles for VeA and LaeA in Development and Pathogenesis of *Aspergillus flavus*. *Eukary. Cell* 8, 1051-1060.

Tech Fields

- [Drug Discovery & Development : Drug production & design](#)

For current licensing status, please contact Rafael Diaz at rdiaz@warf.org or 608-960-9847

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete cookies, you agree to the storing of cookies and related technologies on your device. [See our privacy policy.](#)

OK

