



Elastographic Imaging of the Cervix and Uterine Wall

[View U.S. Patent No. 7,297,116 in PDF format.](#)

WARF: P03009US

Inventors: Tomy Varghese, Mark Kliewer, James Zagzebski

The Wisconsin Alumni Research Foundation (WARF) is seeking commercial partners interested in developing an elastographic method and device for producing diagnostic images of the cervix, uterus and pelvic floor.

Overview

Elastography is a new ultrasound imaging technique that detects and images the local stiffness properties of tissues under compression. Elastography promises to improve the diagnosis of several uterine and cervical disorders. For example, it could provide a more direct measure of cervical incompetence – a disorder believed to be the principal cause of an estimated 25 percent of premature deliveries – because the stiffness properties of the cervix relate more closely to cervical competence than does the current measure, cervical length. In addition, unlike conventional ultrasound, elastography can potentially distinguish uterine fibroids from adenomyosis foci, due to the differences in stiffness between these tissue types.

The Invention

A group of UW-Madison researchers has now developed an elastographic method and device for producing diagnostic images of the cervix, uterus and pelvic floor. Their invention includes a number of ways to achieve the controlled tissue compression needed for imaging, such as using the ultrasound probe itself to compress the uterus or cervix, or inflating a balloon (similar to the balloons used in angioplasty) inside these organs to compress them.

Applications

- Elastography imaging of the cervix, uterus and pelvic floor
- Diagnosis of incompetent cervix, endometrial cancer and other disorders

Key Benefits

- Provides a simple, low-cost alternative to magnetic resonance imaging (MRI) for the diagnosis of uterine, cervical and pelvic disorders
- Could provide a more reliable measure of cervical incompetence and the risk of premature delivery than the current measure, cervical length
- Unlike conventional ultrasound, holds potential for distinguishing between fibroids and adenomyosis in the uterus
- Could be used to distinguish endometrial cancer from benign uterine disorders such as hyperplasia

Additional Information

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete cookies, you agree to the storing of cookies and related technologies on your device. [See our privacy policy.](#)

• [Tomy Varghese](#)

OK

Tech Fields



WARF
Wisconsin Alumni Research Foundation

| info@warf.org | 608.960.9850

- [Medical Imaging : Ultrasound](#)

For current licensing status, please contact Jeanine Burmania at jeanine@warf.org or 608-960-9846

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete cookies, you agree to the storing of cookies and related technologies on your device. [See our privacy policy.](#)

OK



WARF
Wisconsin Alumni Research Foundation

| info@warf.org | 608.960.9850