

Tracking Tumors for Real-Time Radiation Therapy by Automatic Segmentation

View U.S. Patent No. 8,992,405 in PDF format.

WARF: P120018US01

Inventors: Bhudatt Paliwal, Venkata Chebrolu

The Wisconsin Alumni Research Foundation (WARF) is seeking commercial partners interested in developing a method that reduces clinician labor and error by automatically defining volumes of healthy and malignant tissue, and guiding therapy despite changes in tumor location or form.

Overview

External beam radiation therapy degrades the rapidly dividing cells of a tumor by directing high-energy radiation into the target of interest to eradicate it. The efficacy of the treatment is strongly impacted by dosage, which is constrained by the need to spare surrounding tissue and organs. Directing beams to intersect across a tumor, while reducing the dose to areas outside the intersection, can reduce damage to healthy tissue.

Imaging obtained by CT scan or MRI prior to therapy provides a computerized treatment plan designed to 'segment' tissue types, defining volumes of healthy versus afflicted tissue and assigning to each a minimum or maximum dose. But current automated segmentation is inaccurate and laborious when performed manually, requiring a health worker to draw boundary lines on the multiple data slices composing an image. Further confounding ideal treatment is the tendency of tumors to shift and change size during and between sessions—in response to breathing, regression or even bladder filling.

The latest generation of imaging equipment is capable of snapshotting a patient every quarter of a second. To exploit this fully, an automated and equally rapid method to deliver radiation into a tumor despite its deviations is called for.

The Invention

UW-Madison researchers have developed an extremely fast algorithm-based segmentation technique to guide radiation at a rate commensurate with real-time tissue imaging.


Novel Morphological Processing and Successive Localization (MPSL) can be applied to auto-contour the volume, shape and position of a target. The method utilizes predetermined knowledge of the general location and size range of the tumor and based on similarities within value and positioning data, isolates healthy and diseased regions for radiation. More efficient than manual segmentation and more accurate than existing algorithms, the method enables flexible, medical image-guided radiotherapy.

Applications

- Adaptive treatment planning
- Image-guided radiotherapy in real time

We use cookies of this site to entine by the storing of cookies and related technologies on your device. See our privacy policy

Key Benefits



- · Efficient automated tissue segmentation
- No manual contouring
- · Applied to quantify changes in tumor volume, shape and position

Stage of Development

Researchers used MPSL to segment CT and MRI images of lung cancer and lung disease patients. The method successfully defined regions of interest from images with both high and low signal-to-noise ratio. The average computational time for a volume was two seconds.

Additional Information

For More Information About the Inventors

<u>Bhudatt Paliwal</u>

Publications

• Tewatia D.K., Tolakanahalli R.P., Paliwal B.R. and Tomé W.A. 2011. Time Series Analyses of Breathing Patterns of Lung Cancer Patients Using Nonlinear Dynamical System Theory. Phys. Med. Biol. 56, 2161-2181.

Tech Fields

- Information Technology : Computing methods, software & machine learning
- Radiation Therapy : Treatment planning

For current licensing status, please contact Jeanine Burmania at jeanine@warf.org or 608-960-9846

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete cookies, you agree to the storing of cookies and related technologies on your device. See our privacy policy

