

Green Method for Producing 1,5-Pentanediol Slashes Catalyst Cost 10,000-fold

View U.S. Patent No. 10,183,904 in PDF format.

WARF: P160103US01

Inventors: George Huber, James Dumesic, Kevin Barnett, Zach Brentzel

The Wisconsin Alumni Research Foundation (WARF) is seeking commercial partners interested in developing a new route for producing the high value chemical 1,5-PD via upgrading of biomass-derived tetrahydrofurfuryl alcohol.

Overview

Commercial interest in the production of commodity chemicals from renewable sources continues to surge. Among these chemicals, a, wdiols are particularly attractive because of the high market prices they command (\$2,700 - 6,000/MT in 2015 U.S. dollars).

1,5-Pentanediol (I,5-PD), which is used as a plasticizer and also as a precursor in the manufacture of polyurethanes, is especially lucrative because it is currently produced in small quantities from petroleum feedstocks. Research has shown that 1,5-PD can be made from the conversion of tetrahydrofurfuryl alcohol (THFA) using noble metal catalysts. However, subprime yields, high temperature and reliance on high cost catalysts render the process economically infeasible.

The Invention

Seeking a commercially viable alternative, UW-Madison researchers have developed a new route for producing 1,5-PD from biomassderived THFA. Their three-step process is orders of magnitude cheaper than competing methods, green and exceeds 90 percent overall yields.

More specifically the new method includes hydration of THFA to dihydropyran, conversion to 2-hydroxy-tetrahydropyran (no need for a mineral acid catalyst) and subsequent production of 1,5-PD. The entire method can be conducted entirely in the absence of noble metal catalysts.

Applications

• Production of 1,5-PD for use in coatings, polymer resins, plasticizers, acrylates, adhesives and more

Key Benefits

- New route is cheaper, faster and simpler than competing methods.
- · Competes on price with petroleum-derived sources
- · Uses inexpensive metal-oxide and base metal catalysts
- · Relatively mild reaction conditions
- · Higher reactant concentrations cut distillation costs.

Eliminates difficult and costly acid neutralization step
We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete
Incurs far lower separation costs
cookies, you agree to the storing of cookies and related technologies on your device. See our privacy policy

Stage of Development



The new method results in >90 percent overall yield for conversion of furfural into 1,5 pentanediol.

The development of this technology was supported by WARF Accelerator. WARF Accelerator selects WARF's most commercially promising technologies and provides expert assistance and funding to enable achievement of commercially significant milestones. WARF believes that these technologies are especially attractive opportunities for licensing.

Additional Information

For More Information About the Inventors

<u>George Huber</u>

Publications

- Brentzel Z.J., Barnett K.J., Huang K., Maravelias C.T., Dumesic J.A. and Huber G.W. 2017. Chemicals from Biomass: Combining Ring-Opening Tautomerization and Hydrogenation Reactions to Produce 1,5-Pentanediol from Furfural. ChemSusChem. 10, 1351-1355.
- Read a news story about this technology.

Tech Fields

<u>Clean Technology : Biobased & renewable chemicals & fuels</u>

For current licensing status, please contact Jennifer Gottwald at jennifer@warf.org or 608-960-9854

We use cookies on this site to enhance your experience and improve our marketing efforts. By continuing to browse without changing your browser settings to block or delete cookies, you agree to the storing of cookies and related technologies on your device. See our privacy policy

