



US007910565B2

(12) **United States Patent**
McNeel et al.

(10) **Patent No.:** **US 7,910,565 B2**
(45) **Date of Patent:** **Mar. 22, 2011**

(54) **PROSTATE CANCER VACCINE**

(75) Inventors: **Douglas G. McNeel**, Madison, WI (US);
Brian M. Olson, Madison, WI (US)

(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 647 days.

(21) Appl. No.: **11/848,607**

(22) Filed: **Aug. 31, 2007**

(65) **Prior Publication Data**

US 2008/0063654 A1 Mar. 13, 2008

Related U.S. Application Data

(60) Provisional application No. 60/841,769, filed on Sep. 1, 2006.

(51) **Int. Cl.**
A61K 48/00 (2006.01)

(52) **U.S. Cl.** **514/44**; 435/320.1; 435/455

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,548,064	B1	4/2003	Tureci et al.
6,800,730	B1	10/2004	Tureci et al.
7,179,797	B2	2/2007	McNeel et al.
2004/0142890	A1	7/2004	McNeel et al.

OTHER PUBLICATIONS

Eder et al. *Future Oncol* 2005;1:93-101.*
Scher et al. *Endocrine-Related Cancer* 2004;11:459-76.*
Chen et al. *Nat Med* 2004;10:33-9.*
Olson et al. *Prostate* 2007;1729-39.*
Suarez-Quian et al. *J Clin Endocrinol Metab* 1999;84:350-8.*
Chichet et al. *Int J Morphol* 2007;25:579-85.*
Meidenbauer et al., Direct visualization of antigen-specific T cells using peptide-MHC-class I tetrameric complexes, *Methods* 31 (2003): 160-171.
Palmowski, et al., The use of HLA class I tetramers to design a vaccination strategy for melanoma patients, *Immunological Reviews* 188 (2002): 155-163.
Mincheff M, Tchakarov S, Zoubak S, et al, Naked DNA and adenoviral immunizations for immunotherapy of prostate cancer; a phase I/II clinical trial. *Eur Urol*. 2000;38:208-217.
Irvine and Restifo, 1995, *Seminars in Canc. Biol.* 6:337-347.
Iwasaki A, Torres C A, Ohashi P S, Robinson H L, Barber B H. *J Immunol*. 1997;159:11-4.
Chen et al., 1998, Induction of CD8+ T Cell Responses to Dominant and Subdominant Epitopes and Protective Immunity to Sendai Virus Infection by DNA Vaccination. *J. Immunol.* 160:2425-2432.
Thomson et al., 1998, Delivery of Multiple CD8 Cytotoxic T Cell Epitopes by DNA Vaccination. *J. Immunol.* 160:1717-1723.
Cho et al., 2000, *Nat. Biotechnol.* 18:509-14.
Wolff et al., 1990, Direct Gene Transfer into Mouse Muscle In Vivo, *Science* 247-1465-8.

Tighe et al., 1998, Gene vaccination: plasmid DNA is more than just a blueprint, *Immunol. Today* 19:89-97.

Raz E, Carson DA, Parker SE, Parr TB, Abai AM, Aichinger G, Gromkowski SH, Singh M, Lew D, Yankauckas MA: Intradermal gene immunization: The possible role of DNA uptake in the induction of cellular immunity to viruses. *Proc Natl Acad Sci USA* 1994, 91:9519-23.

Lee, A. H., Y. S. Suh, J. H. Sung, S. H. Yang, and Y. C. Sung 1997. Comparison of various expression plasmids for the induction of immune response by DNA immunization. *Mol. Cell* 7:495-501.

Hartmann G, Weeratna RD, Belles ZK et al. Delineation of a CpG phosphorothioate oligodeoxynucleotide for activating primate immune responses in vitro and in vivo. *J Immunol* 2000, 164: 1617-24.

Acsadi et al., 1991, Direct gene transfer and expression into rat heart in vivo, *New Biol.* 3:71-81.

Wolff et. al., 1991, Conditions Affecting Direct Gene Transfer into Rodent Muscle in Vivo, *Biotechniques* 11(4):474-85.

Budker et. al., 1996, *Nat. Biotechnol.* 14:760-4.

Davis HL, Whalen RG, Demeneix BA. Direct gene transfer into skeletal muscle in vivo: factors affecting efficiency of transfer and stability of expression. *Hum Gene Ther* 1993;4:151-9.

Danko et al., 1994, *Gene Ther.* 1:114-21.

Manthorpe et al., Gene Therapy by Intramuscular Injection of Plasmid DNA: Studies on Firefly Luciferase Gene Expression in Mice, *Hum. Gene Ther.* 4:419-31 (1993).

Budker et al., 1996, Naked DNA delivered intraportally expresses efficiently in hepatocytes, *Gene Ther.* 3:593-8.

Budker et al., 1998, The efficient expression intravascularly delivered DNA in rat muscle. *Gene Ther.* 5:272-276.

Zhang G et al. Efficient expression of naked DNA delivered intraarterially to limb muscles of nonhuman primates. *Hum Gene Ther* 2001; 12: 427-438.

Wagner et al., 2003, *Cancer Immun* 3:18-32.

Maha Ayyoub, Charles S. Hesdorffer, Genevieve Matthez, Stefan Stevanovic, Gerd Ritter, Yao-Tseng Chen, Lloyd J. Old, Daniel Speiser, Jean-Charles Cerottini, Danila Valmori. (2004) identification of an SSX-2 Epitope Presented by Dendritic Cells to circulating Autologous CD4+ T Cells, *J. of Immunol.* 172:7206-11.

Tureci O, Sahin U, Zwick C, Koslowski M, Seitz G, Pfreundschuh M. Identification of a meiosis-specific protein as a member of the class of cancer/testis antigens. *Proc Natl Acad Sci U S A* 1998; 95: 5211-6.

Scanlan M.J. et al., The cancer/testis genes: review, standardization, and commentary (2004) *Cancer Immun.* 4:1. Abstract, Prikler L. et al., 2004, Adaptive Immunotherapy of the Advanced Prostate Cancer—Cancer Testis Antigen (CTA) as Possible Target Antigens, *Klinische Originalarbeit* 35:326-330.

Kaplan-Lefko P. J., Chen T. M., Ittmann M. M., Barrios R. J., Ayala G. E., Huss W. J., Maddison L. A., Foster B. A., and Greenberg N. M. (2003) "Pathobiology of autochthonous prostate cancer in a pre-clinical transgenic mouse model." *Prostate* 55:219-237.

* cited by examiner

Primary Examiner — Q. Janice Li

(74) *Attorney, Agent, or Firm* — Quarles & Brady LLP

(57) **ABSTRACT**

Androgen receptor-based vaccines for eliciting an immune reaction in vivo against cells expressing androgen receptor are disclosed. The vaccines are useful in the treatment of prostate cancer. Also disclosed are methods for inducing immune reaction to androgen receptor or treating prostate cancer in a mammal, using the vaccines and pharmaceutical compositions comprising the vaccines.

10 Claims, 15 Drawing Sheets




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A





FIG. 2B




FIG. 2C






FIG. 3

**FIG. 4**

**FIG. 5**





FIG. 6






FIG. 8




FIG. 9




FIG. 10




FIG. 11

1

PROSTATE CANCER VACCINE**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims the benefit of U.S. provisional application Ser. No. 60/841,769, filed on Sep. 1, 2006, which is herein incorporated by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was made with United States government support awarded by the following agency: NIH RR016489. The United States has certain rights in this invention.

BACKGROUND OF THE INVENTION

Prostate cancer is a significant health risk for men over the age of 50, with about 200,000 newly diagnosed cases each year in the United States (Jemal A. et al., Cancer Statistics, 2005 (2005) CA Cancer J Clin, 55:10-30). It is the most common tumor diagnosed among men and the second leading cause of male cancer-related death in the United States (Jemal et al., Cancer Statistics, 2003 (2003) CA Cancer J Clin, 53:5-26). Despite advances in screening and early detection, approximately 30% of patients undergoing definitive prostatectomy or ablative radiation therapy will have recurrent disease at 10 years (Oefelein et al., 1997, J Urol, 158:1460-1465). At present, there is no accepted adjuvant treatment for patients undergoing radical prostatectomy or ablative radiation therapy that has been shown to prevent the progression to metastatic disease. In addition to new treatments for metastatic disease, new strategies are needed to eradicate microscopic disease to prevent the progression to clinically apparent metastasis.

In patients who have undergone definitive ablative therapy for prostate cancer, the presence of detectable serum levels of prostate-specific antigen (PSA) has provided a valuable indicator of microscopic metastatic disease. In a retrospective review of 1,997 men treated with radical prostatectomy, 15% were found to have evidence of a PSA-only recurrence over a median 5-year follow up, so-called stage D0 disease (Pound et al., 1999, JAMA 281:1591-7). Of these, 34% developed radiographically apparent metastatic disease, with a median time to development of metastatic disease of 8 years. In all patients with metastatic disease, the median time to death was 5 years (Pound et al., 1999, JAMA 281:1591-7). These findings suggest that patients with stage D0 disease are at high risk for progressive disease, however with a long window of time to test adjuvant therapies. Similarly, many patients are found to have microscopic pelvic lymph node metastases at the time of radical prostatectomy, so-called stage D1 disease. At present, the best treatment for these patients is controversial, with most treated with androgen deprivation, and others are expectantly observed without specific treatment. In retrospective studies, 10-year disease-specific recurrence and mortality is on the order of 50 to 66% for patients with stage D1 disease (Sgrignoli et al., 1994, J Urol, 152:1077-81; and Cadeddu et al., 1997, Urology, 50:251-5). This high-risk stage of minimal residual disease also provides an opportunity to test novel adjuvant therapies.

Immunological therapies, and vaccines in particular, are appealing as possible treatment options for prostate cancer for several reasons. Such therapies may be relatively safe and inexpensive treatments compared with chemotherapies for a disease for which no standard adjuvant treatments exist (Kent

2

et al., Immunity of prostate specific antigens in the clinical expression of prostatic carcinoma (1976) In: Crispin R G, ed. Neoplasm immunity: mechanisms. Chicago, ITR, pp. 85-95; Guinan et al., 1984, Prostate, 5:221-230; and McNeel et al., 2000, Arch. Immunol. Ther. Exp., 48:85-93). Moreover, prostate cancer is a slow-growing disease, with typically over five years from the time of diagnosis of organ-confined disease to the development of clinically apparent metastatic disease. Such a slow-growing disease might be more amenable to vaccine-based treatments than a rapidly growing tumor, assuming that microscopic amounts of disease would be easier to treat than bulky or rapidly growing disease by vaccines. In fact, vaccines have already entered clinical trials for prostate cancer targeting a variety of prostate-specific proteins, with at least two dendritic cell-based vaccines suggesting clinical benefit in patients with low-volume metastatic disease (Murphy et al., 1999, Prostate, 39:54-59; and Small et al., 2000, J. Clin. Oncol. 18:3894-3903).

The use of plasmid DNA alone as a means of in vivo gene delivery by direct injection into muscle tissue was first described by Wolff et al. (Wolff et al., 1990, Science, 247: 1465-1468). It was subsequently found that intramuscular or intradermal administration of plasmids expressing foreign genes elicited immune responses (Tang, et al., 1992, Nature, 356:152-154; Wang et al., 1993, Proc Natl. Acad. Sci. USA, 90:4156-4160; and Raz et al., 1994, Proc Natl. Acad. Sci. USA, 91:9519-9523). This has quickly led to numerous investigations into the use of plasmid DNA as a means of vaccine antigen delivery, both in animal and human models. DNA vaccines, like peptide-based vaccines, are relatively easy and inexpensive to manufacture, and are not individualized for patients as are dendritic cell-based vaccines. With recombinant protein vaccines, the antigen is taken up by antigen presenting cells and expressed predominantly in the context of MHC class II. DNA in nucleic acid vaccines is taken up and expressed by antigen-presenting cells directly, leading to antigen presentation through naturally processed MHC class I and II epitopes (Iwasaki, et al. 1997, J Immunol, 159:11-14). This direct expression by host cells, including MHC class I expressing bystander cells, has been demonstrated to lead to vigorous CD8+CTL responses specific for the targeted antigen (Iwasaki et al., 1997, J. Immunol. 159: 11-14; Chen et al., 1998, J. Immunol., 160:2425-2432; Thomson et al., 1998, J. Immunol., 160:1717-1723; and Cho et al., 2000, Nat. Biotechnol. 18:509-514).

Clinical trials have suggested that plasmid DNA vaccines are safe and immunologically effective in humans. Boyer and colleagues reported that doses of 300 µg of plasmid DNA encoding HIV rev and env proteins administered intramuscularly were capable of eliciting antigen-specific, IFN γ -secreting T cell responses in HIV-seronegative patients (Boyer et al., 2000, J. Infect. Dis. 181:476-83). In addition, results of a clinical trial targeting prostate-specific membrane antigen (PSMA) in patients with prostate cancer by means of plasmid DNA and adenovirus have been reported (Mincheff et al., 2000, Eur. Urol., 38:208-217). In this study, 26 patients were immunized either in a prime/boost strategy with an adenoviral vector expressing PSMA followed by immunization with plasmid DNA expressing PSMA, or with plasmid DNA alone. The authors report no significant toxicity with doses of 100-800 µg of plasmid DNA administered intradermally, and suggest that patients receiving plasmid DNA expressing PSMA and CD86 with soluble GM-CSF as an adjuvant were all successfully immunized.

A DNA vaccine for the treatment of prostate cancer based on prostatic acid phosphatase (PAP) has also been described (US 2004/0142890).

BRIEF SUMMARY OF THE INVENTION

The present invention is based, in part, on the inventors' discovery that patients with prostate cancer have antibodies specific for the androgen receptor, that androgen receptor ligand-binding domain as well as four fragments thereof (SEQ ID NO:9-12) can elicit immune responses in vivo, and that animals vaccinated with a DNA vaccine encoding the androgen receptor (AR) ligand-binding domain (LBD) inhibited prostate tumor growth in vivo.

In one aspect, the invention relates a method for inducing an immune reaction to androgen receptor in a mammal in need thereof, the method comprising administering to the mammal an effective amount of a recombinant DNA construct comprising a polynucleotide operatively linked to a transcriptional regulatory element (e.g., a promoter such as a heterologous promoter) wherein the polynucleotide encodes a member selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, whereby the mammal develops immune reaction against the androgen receptor. In one form, the polynucleotide employed in the method encodes the ligand-binding domain of a mammalian androgen receptor. In another form, multiple DNA constructs with each comprising a polynucleotide that encodes a different fragment selected from (iii)-(vi) are administered. For example, two DNA constructs covering fragments (iii) and (iv) can be administered together. As another example, four DNA constructs covering all four fragments (iii)-(vi) can be administered together. The method disclosed can be practiced with a mammal, preferably a human, who either currently has or previously had prostate cancer.

In one embodiment, the polynucleotide encodes a human androgen receptor or a fragment of the human androgen receptor that comprises the ligand-binding domain. The polynucleotide is preferably a nucleotide sequence of the human androgen receptor gene. In one form of this embodiment, the polynucleotide encodes the ligand-binding domain of a human androgen receptor.

The above method employing the DNA construct induces cytotoxic immune reaction against cells expressing androgen receptor. Preferably, both humoral and cellular immune reactions against androgen receptor are induced.

In another aspect, the present invention relates to a method for inducing an immune reaction to androgen receptor in a mammal in need thereof, the method comprising administering to the mammal an effective amount of a polypeptide selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, whereby the mammal develops immune reaction against the androgen receptor. In one form, the polypeptide employed is the ligand-binding domain of a mammalian androgen receptor. In another form, multiple fragments of the ligand-binding domain (e.g. SEQ ID NO:9 and SEQ ID NO:10, and optionally SEQ ID NO:11 and SEQ ID NO:12) are administered.

The method disclosed can be practiced with a mammal, preferably a human, who either currently has or previously had prostate cancer.

In one embodiment, the human androgen receptor or a fragment of the human androgen receptor that comprises the ligand-binding domain is administered. In one form of this embodiment, the ligand-binding domain of the human androgen receptor is administered.

The above method employing the polypeptide induces cellular or humoral immune reaction against cells expressing androgen receptor. Preferably, both humoral and cellular immune reactions against androgen receptor are induced.

According to one embodiment of the invention, the recombinant DNA construct or the polypeptide is administered to the mammal intradermally, intramuscularly, subcutaneously, or intravascularly, including intravenously and intraarterially. Preferably, the recombinant DNA construct is administered intradermally, intramuscularly, or intravascularly and the polypeptide is administered subcutaneously.

In another aspect, the present invention relates to an isolated polypeptide selected from SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12. In another aspect, the present invention relates to a composition that comprises one or more of the above polypeptides and a pharmaceutically acceptable carrier.

According to another aspect of the present invention, a DNA vaccine is contemplated which comprises a plasmid vector comprising a polynucleotide operatively linked to a transcriptional regulatory element (e.g., a promoter such as a heterologous promoter) wherein the polynucleotide encodes a member selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, wherein upon administration of said vaccine to a mammal a cytotoxic immune reaction against cells expressing androgen receptor is induced. The vaccine of the present invention preferably is suitable for intradermal, intramuscular, subcutaneous, or intravascular (including intravenous and intraarterial) administration to a mammal such as a human. According to a preferred embodiment, the plasmid vector comprises (a) a backbone of pNGVL3, (b) a polynucleotide operably inserted therein wherein the polynucleotide encodes a member selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, and, optionally, (c) one or a plurality of an immunostimulatory sequence (ISS) motif.

Preferably, the DNA vaccine according to the invention comprises a plasmid vector that comprises (a) a polynucleotide operatively linked to a CMV promoter wherein the polynucleotide encodes a member selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID

5

NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, (b) a CMV intron A operatively linked to the polynucleotide for enhancing expression of the polynucleotide, and, optionally, (c) at least one copy of an immunostimulatory fragment comprising 5'-GTCGTT-3'. In one embodiment, the plasmid construct does not express in eukaryotic cells any gene other than a member selected from (i) a mammalian androgen receptor, (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12. The plasmid vector pTVG4 is particularly preferred.

According to another aspect of the present invention, a peptide vaccine is contemplated which comprises a member selected from (i) a mammalian androgen receptor (e.g., a human androgen receptor), (ii) a fragment of the androgen receptor that comprises the ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12. The peptide vaccine also comprises a pharmaceutically acceptable carrier. The peptide vaccine preferably is suitable for intradermal, intramuscular, subcutaneous, or intravascular (including intravenous and intraarterial) administration to a mammal such as a human.

Also disclosed are pharmaceutical compositions comprising a DNA or peptide vaccine of the invention (the polypeptides or recombinant plasmid vectors described above), and a pharmaceutically acceptable carrier. Preferably, the pharmaceutical composition further comprises a suitable amount of immuno-stimulant such as GM-CSF.

A kit containing the DNA or peptide vaccine of the invention and an instruction manual directing administering the vaccine to a mammal that has or previously had prostate cancer (e.g., a human prostate cancer patient) is also within the scope of the invention.

In another aspect, the present invention relates to a method for determining the effectiveness of a treatment for prostate cancer. The method includes the steps of (a) measuring the frequency or amount of cytotoxic T lymphocytes (CTLs) specific for a peptide selected from SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12 prior to providing at least a portion of the treatment to a mammal (e.g., a human) having prostate cancer, (b) measuring the frequency or amount of CTLs specific for the peptide after said portion of the treatment is provided to the mammal, and (c) comparing the frequency or amount of CTLs of (a) and that of (b) wherein the frequency or amount of CTLs of (b) being higher than that of (a) indicates that the treatment is effective.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

FIG. 1 shows that patients with prostate cancer have antibodies specific for the androgen receptor. Panel A: Sera from patients with various stages of prostate cancer were analyzed for the presence of AR-specific antibodies by screening titrated sera samples using ELISA. Panel B: The presence or absence of AR-specific antibodies was confirmed using Western blotting against thioredoxin-tagged AR LBD or thioredoxin (trx) alone, followed by incubation with patient's sera.

6

Panel C: ELISA was used to evaluate sera samples for the presence of AR-specific antibodies. Samples were analyzed from healthy male blood donors (n=41), patients with prostatitis (n=38), or patients with prostate cancer (n=105), and relative antibody concentrations were calculated by referencing delta absorbance values to titrated Ig protein standards. Positive antibody responses were defined by values higher than three standard deviations above the mean of the healthy donor group (greater than 0.22 µg/mL, indicated by the line). Statistically significant differences were calculated using the Chi square test.

FIG. 2 shows that patients with AR LBD-specific IgG antibodies have concurrent AR-specific cellular immune responses. PBMC were analyzed for the presence of AR-specific T cells and IFN γ -secreting cells among patients with antibody responses specific for the AR (n=6) or patients with no detectable antibody responses (n=9). PBMC were stimulated with thioredoxin-tagged AR LBD, thioredoxin alone, media only, or PHA. After a 96-hour stimulation, these cells were analyzed for the presence of CD4+ and CD8+ T cell proliferation in response to antigen stimulation. Example data is shown in panel A from a subject with a strong IgG₂ response. The numbers in the upper-right corner of each panel indicate the percentage of CD4+ or CD8+ T cells that co-stained with BrdU. Proliferation indexes (PI) were calculated by normalizing experimental values to values obtained from PBMCs stimulated with media alone, and compiled PI values are shown for CD4+ (panel B, left) and CD8+ (panel B, right) T-cells. Supernatants from cultured PBMC were also analyzed for the presence of IFN γ secretion by capture ELISA (panel C). A comparison of results among different antigen-stimulation conditions was performed using the Student's T test.

FIG. 3 shows that patients with AR LBD-specific IgG antibodies have a mixed Th1/Th2-type immune response. PBMC from prostate cancer patients with (n=5) or without (n=5) antibody responses specific for the AR were analyzed for the presence of antigen-specific IFN γ or IL-10-secretion. PBMC were stimulated with thioredoxin-tagged AR LBD, thioredoxin alone, media only, or PHA. After a 96-hour stimulation, supernatants were analyzed for the presence of IFN γ (panel A) and IL-10 (panel B) secretion by quantitative ELISA. A comparison of results among different antigen-stimulation conditions was performed using a Student's T test. The results shown are representative of duplicate experiments with the same peripheral blood samples.

FIG. 4 shows the plasmid map of pTVG4.

FIG. 5 shows the plasmid map of pTVG-ARLBD (pTVG-androgen receptor ligand-binding domain).

FIG. 6 shows that AR LBD peptide-specific T-cells can lyse peptide-pulsed and prostate cancer cell line target cells. Peptide-specific T-cell lines were cultured from the peripheral blood of HLA-A2-expressing patients with prostate cancer using peptides AR811 (LLLFSIIPV, SEQ ID NO:9, panel A) or AR859 (QLTKLLDSV, SEQ ID NO:12, panel B). After several in vitro restimulations, cultures were tested for cytolytic activity to T2 cells loaded with AR811 peptide (diamond), AR859 (square), or the LNCaP HLA-A2 expressing prostate cancer cell line (triangle). Shown is a representative graph results obtained from the cytotoxicity assay of T-cell cultures performed in triplicate at three different effector-to-target ratios as indicated.

FIG. 7 shows that a T-cell clone specific for the AR811 peptide secretes IFN γ and TNF α in response to peptide stimulation. A clonal T-cell line specific for the AR811 peptide was derived by limited dilution following multiple restimulations in vitro. This line was restimulated for 5 hours in vitro with T2

7

cells and media only (panel A), an irrelevant peptide (panel B), AR811 peptide (panel C), or PMA/Ionomycin (panel D). IFN γ and TNF α cytokine accumulation were assessed by intracellular flow cytometric analysis (Cytofix/cytoperm kit, BD Pharmingen). Cells were first stained for CD3 and CD8 expression and CD3+/CD8+ cells were analyzed for IFN γ and TNF α expression. Shown are the plots gated on CD3+/CD8+ cells.

FIG. 8 shows that the AR811 epitope is a naturally processed HLA-A2-restricted T-cell epitope. The AR811 T-cell clone was tested for its ability to lyse prostate cancer cell lines (expressing the AR) that express HLA-A2 (LNCaP, diamond and MDA-Pca-2B, triangle), or do not express HLA-A2 (DU145, X). In addition, lysis was evaluated following pretreatment of LNCaP with an HLA-A2 blocking monoclonal antibody (starburst). Shown is a cytotoxicity assay of an AR811CD8+ cell clone. Cells were restimulated for seven days using peptide-pulsed antigen presenting cells, followed on day seven by a rest period of three days before they were analyzed for their cytotoxic activity.

FIG. 9 shows that HHD-II mice immunized with AR811 peptide develop peptide-specific CTL (contain splenocytes that can specifically lyse AR811 target cells). Male HHD-II mice (n=3 per group) were immunized once with 100 μ g AR811 peptide in CFA or with CFA alone. One week after immunization, splenocytes were collected and stimulated in vitro with 10 μ g/mL peptide for two hours, and on the second day, recombinant murine IL-2 and IL-7 (Fitzgerald Industries) were added to 10 U/mL and 30 U/mL, respectively. The cultures were then allowed to incubate an additional six days before analysis. Cultured cells from AR811-immunized animals (panel A) or control immunized animals (panel B) were then tested for cytolytic activity to T2 cells pulsed with the AR811 peptide (square) or T2 cells pulsed with an influenza matrix peptide (diamond). Shown are the mean and standard deviation of triplicate wells at three effector-to-target ratios as indicated, each from a single animal per group, and representative of the other animals per group.

FIG. 10 shows that rats immunized with a plasmid vector encoding AR LBD develop AR LBD-specific antibody responses. Copenhagen rats were immunized 4 times at 14-day intervals with pTVG4 control vector or pTVG-AR-LBD. Two weeks after the fourth immunization, blood was collected and assessed for antibody responses to the AR-LBD protein by ELISA. Shown are the mean and standard error for 10 animals per experimental group.

FIG. 11 shows tumor growth in pTVG4 control vector or pTVG-ARLBD immunized rats (Kaplan-Meier analysis of animal endpoint survival following treatment with the pTVG4 control vector or pTVG-ARLBD). Male Copenhagen rats were immunized four times every other week with pTVG4 control vector or pTVG-ARLBD. Two weeks after immunization, the rats were challenged with 1×10^4 syngeneic Mat-LyLu prostate cancer cells given with Matrigel. Tumors were measured every other day, and rats were sacrificed when tumors grew larger than 10 cm³ (Kaplan-Meier survival endpoint being tumor volume over 10 cm³).

DETAILED DESCRIPTION OF THE INVENTION

This invention provides pharmaceutical compositions and methods that relate to the use of plasmid DNA and peptide vaccines for the treatment of prostate cancer. Specifically, this invention provides polypeptides such as the ligand-binding domain of an androgen receptor or certain fragments thereof and recombinant plasmid vectors comprising genes or polynucleotide molecules encoding the polypeptides for prevent-

8

ing or treating prostate cancer, including metastatic tumors thereof. In a preferred embodiment, the polypeptides or recombinant plasmid vectors are administered to prostate cancer patients to treat prostate cancer. In another preferred embodiment, the polypeptides or recombinant plasmid vectors are administered to stage D0 or D1 prostate cancer patients to prevent recurrence or metastasis of prostate cancer.

A polypeptide vaccine of the present invention, which comprises a pharmaceutically acceptable carrier and an effective amount of a mammalian androgen receptor, a fragment of the mammalian androgen receptor that comprises the ligand-binding domain, or certain fragments of the ligand-binding domain, can be administered into a mammal such as a human being to elicit an immune response against androgen receptor in the mammal. An "effective amount" or an "immunologically effective amount" means that the administration of that amount to a subject, either in a single dose or as part of a series, is effective for inducing an immune reaction and preferably for treating or preventing prostate cancer. Pharmaceutically acceptable carriers are well known to those of ordinary skill in the art (Amon, R. (Ed.) *Synthetic Vaccines* 1:83-92, CRC Press, Inc., Boca Raton, Fla., 1987). They include liquid media suitable for use as vehicles to introduce the peptide into a patient but should not in themselves induce the production of antibodies harmful to the individual receiving the composition. An example of such liquid media is saline solution. Moreover, the vaccine formulation may also contain an adjuvant for stimulating the immune response and thereby enhancing the effect of the vaccine.

The plasmid DNA vaccines of the present invention, when directly introduced into mammals such as humans in vivo, induce the expression of encoded polypeptides within the mammals, and cause the mammals' immune system to become reactive against the polypeptides. The vaccines may be any polynucleotides that are capable of eliciting immune responses to an encoded polypeptide.

The instant invention also provides a method of using a polynucleotide which, upon introduction into a mammal, induces the expression, in vivo, of the polynucleotide thereby producing the encoded polypeptide, and causes the mammal to become immune reactive against the polypeptide so produced.

DNA vaccines, like peptide-based vaccines, are relatively easy and inexpensive to manufacture, and are not individualized for patients, as are dendritic cell-based vaccines. With recombinant protein vaccines, the antigen is taken up by antigen presenting cells and expressed predominantly in the context of MHC class II. DNA in nucleic acid vaccines is taken up and expressed by antigen-presenting cells directly, leading to antigen presentation through naturally processed MHC class I and II epitopes (Iwasaki, et al. 1997, *J Immunol*, 159:11-14).

Given their potential ability to elicit antigen-specific cytotoxic T lymphocytes (CTL) immunity in an MHC class I diverse population, DNA-based vaccines for various diseases have recently entered human clinical trials (Mincheff et al., 2000, *Eur. Urol.*, 38:208-217). This method of immunization is similar to the use of viral immunization vectors, but without the additional foreign antigens introduced with a viral vector and therefore with less risk of an overwhelming immune response to the vector itself (Irvine and Restifo, 1995, *Seminars in Canc. Biol.* 6:337-347). Direct expression by host cells, including MHC class I-expressing bystander cells, has been demonstrated to lead to vigorous CD8+ CTL responses specific for the targeted antigen (Iwasaki et al., 1997, *J. Immunol.* 159:11-4; Chen et al., 1998, *J. Immunol.* 160:2425-2432;

Thomson et al., 1998, J. Immunol. 160:1717-1723; and Cho et al., 2000, Nat. Biotechnol. 18:509-14). In addition, plasmid DNA used for immunization may remain within cells at the site of immunization, providing a constant source of antigenic stimulation. Persistent antigen expression may lead to long-lived immunity (Raz et al., 1994, Proc. Natl. Acad. Sci. USA 91:9519-23).

The present invention provides DNA-based vaccines that express a polypeptide antigen, the ligand-binding domain of a mammalian androgen receptor or certain fragments thereof, and methods for treating prostate cancers in a human or non-human animal using the vaccines. In addition to the reasons explained above, plasmid vaccines are advantageous over viral vaccines. For example, viral vaccines are not amenable to repeated immunizations. With viral vectors, one is trying to elicit an immune response against a "self" protein encoded by a foreign virus. The immune system preferentially recognizes the foreign proteins, sometimes hundreds of proteins, encoded by the virus. For example, the inventors have found in rats that repeated immunizations with a vaccinia virus encoding human prostatic acid phosphatase (hPAP) elicits a strong vaccinia response but no hPAP-specific response (Johnson et al., 2007, Canc. Immunol. Immunoth. 56:885). That same finding was also shown in humans, in a trial in which repeated immunization with the vaccinia virus encoding human prostate-specific antigen (PSA) elicited weak PSA-specific immunity, but potent vaccinia immunity (Sanda et al., 1999, Urology 53:260). The direction in the field of viral-based vaccines is to "prime" with a virus encoding the antigen, and then "boost" with a different virus (like adenovirus or fowl pox) encoding the same antigen. The advantage of plasmid DNA vaccines is that they encode a defined, often small, number of proteins. Therefore, one can repetitively immunize the animal or patient. Furthermore, a virus may kill cells, incorporate into the genome, or potentially induce other unwanted immune responses. All these are disadvantages that are likely avoided by DNA plasmid vaccines.

It is readily recognizable that the ligand-binding domain of an androgen receptor of any origin, or any of the ligand-binding domain's derivatives, equivalents, variants, mutants etc., is suitable for the instant invention, as long as the ligand-binding domain or derivatives, equivalents, variants, or mutants thereof is able to induce an immune reaction in the host human or non-human animal substantially similar to that induced by an autoantigenic or xenoantigenic ligand-binding domain of the androgen receptor in the animal. Similarly, a polynucleotide sequence of an androgen receptor gene of any origin that encodes the ligand-binding domain of the receptor, or any of the polynucleotide's derivatives, equivalents, variants, mutants etc., is suitable for the instant invention, as long as the polynucleotide sequence and the polypeptide or protein encoded by the polynucleotide sequence, or derivatives, equivalents, variants, or mutants thereof is able to induce an immune reaction in the host human or non-human animal substantially similar to that induced by an autoantigenic or xenoantigenic ligand-binding domain of the androgen receptor in the animal.

Androgen receptor genes are known and have been cloned from many species. For example, the human, mouse, rat, dog, chimpanzee, macaque, and lemur androgen receptor cDNA along with amino acid sequences can be found at GenBank Accession Nos. NM_000044 (cDNA-SEQ ID NO:1 and amino acid sequence-SEQ ID NO:2), NM_013476 (cDNA-SEQ ID NO:3 and amino acid sequence-SEQ ID NO:4), NM_012502 (cDNA-SEQ ID NO:5 and amino acid sequence-SEQ ID NO:6), NM_001003053, NM_001009012, U94179, and U94178, respectively. Androgen receptor genes from other species are also known.

These species include but are not limited to *Sus scrofa*, *Astatotilapia burtoni*, *Gallus gallus*, *Kryptolebias marmoratus*, *Alligator mississippiensis*, *Leucoraja erinacea*, *Haplochromis burtoni*, *Pimephales promelas*, *Dicentrarchus labrax*, *Gambusia affinis*, *Micropogonias undulatus*, *Oryzias latipes*, *Acanthopagrus schlegelii*, *Rana catesbeiana*, *Crocota crocuta*, *Eulemur fulvus collaris*, and *Anguilla japonica* (see GenBank Accession Nos. NM_214314 (or AF161717), AY082342, NM_001040090, DQ339105, AB186356, DQ382340, AF121257, AY727529, AY647256, AB099303, AY701761, AB076399, AY219702, AY324231, AY128705, U94178, and AB023960, respectively). The ligand-binding domains of androgen receptors are well known in the art. For the purpose of the present invention, the ligand-binding domain of the human androgen receptor refers to a polypeptide that starts at any amino acid from amino acid positions 651 to 681 and ends at any amino acid from amino acid positions 900 to 920. For example, human androgen receptor or a fragment of the human androgen receptor that comprises amino acids 681-900 as well as DNA vaccines containing a polynucleotide encoding the above are suitable vaccines. The corresponding ligand-binding domains of androgen receptors from other species can be readily determined by sequence alignment (to the human sequence) (e.g., by the methods described below in connection with sequence identity or homology). In a preferred embodiment, a polypeptide from the human androgen receptor that starts at any amino acid from amino acid positions 661 to 671 and ends at any amino acid from amino acid positions 910 to 920 is used in the present invention. In a more preferred embodiment, a polypeptide containing amino acids 661 to 920 or 664 to 920 of the human androgen receptor is used in the present invention. To help determine the corresponding fragments of the androgen receptors from other species, it is noted here that the amino acid positions on rat, dog, chimpanzee, macaque, and lemur androgen receptors that correspond to amino acid positions 661 to 920 of the human androgen receptor are 640 to 899, 643 to 902, 648 to 907, 652 to 910, 636 to 895, and 625 to 884, respectively. It is noted that the above fragments of the human, mouse, rat, dog, chimpanzee, macaque, and lemur androgen receptors have the same amino acid sequence. The ligand-binding domains of the androgen receptors of other species are also known or can be readily identified through sequence alignment. As will be readily recognized by one of ordinary skill in the art, any DNA sequence that encodes a ligand-binding domain or a larger fragment of an androgen receptor including the full-length receptor from one of the above species as well as other animals is suitable for the present invention.

As is well-known to those skilled in the art, polypeptides having substantial sequence similarities cause identical or very similar immune reaction in a host animal. As discussed below, this phenomenon is the basis for using a xenoantigen for inducing autoreactive reaction to an otherwise tolerated autoantigen. Accordingly, a derivative, equivalent, variant, fragment, or mutant of the ligand-binding domain of any of the known or to-be-identified androgen receptors or any DNA sequence encoding the above is also suitable for the present invention. The polypeptides encoded by these DNA sequences are useful as long as the polypeptides encoded by the DNA sequences are structurally similar to the ligand-binding domain of the autologous androgen receptor, and are sufficiently immunogenic.

It is readily apparent to those ordinarily skilled in the art that variations or derivatives of the nucleotide sequence

encoding the polypeptide or protein antigen can be produced which alter the amino acid sequence of the encoded polypeptide or protein. The altered polypeptide or protein may have an altered amino acid sequence, for example by conservative substitution, yet still elicits immune responses which react with the unaltered protein antigen, and are considered functional equivalents. According to a preferred embodiment, the derivative, equivalents, variants, or mutants of the ligand-binding domain of an androgen receptor are polypeptides that are at least 85% homologous to the ligand-binding domain of a human androgen receptor. More preferably, the homology is at least 88%, at least 90%, at least 95%, or at least 98%.

As used in this application, "percent identity" between amino acid or nucleotide sequences is synonymous with "percent homology," which can be determined using the algorithm of Karlin and Altschul (Proc. Natl. Acad. Sci. USA 87, 2264-2268, 1990), modified by Karlin and Altschul (Proc. Natl. Acad. Sci. USA 90, 5873-5877, 1993). The noted algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (*J. Mol. Biol.* 215, 403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, score=100, wordlength=12, to obtain nucleotide sequences homologous to a polynucleotide of the invention. BLAST protein searches are performed with the XBLAST program, score=50, wordlength=3, to obtain amino acid sequences homologous to a reference polypeptide. To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized as described in Altschul et al. (*Nucleic Acids Res.* 25, 3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used.

As used herein, the term "conservative substitution" denotes the replacement of an amino acid residue by another, biologically similar residue. It is well known in the art that the amino acids within the same conservative group can typically substitute for one another without substantially affecting the function of a protein. For the purpose of the present invention, such conservative groups are set forth in Table 1 based on shared properties.

TABLE 1

Conservative substitution.	
Original Residue	Conservative Substitution
Ala (A)	Val, Leu, Ile
Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg
Ile (I)	Leu, Val, Met, Ala, Phe
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	Gly
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr, Phe
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

In addition, fragments of a ligand binding domain of an androgen receptor such as those that can bind to HLA-A2 are also useful antigens which elicit cytotoxic responses against cells expressing the androgen receptor or its ligand binding domain. Polynucleotides that encode these fragments are

considered functional equivalents. Examples of these fragments are provided in the examples below. In particular, the use of the following four fragments are contemplated: SEQ ID NO:9 (amino acids 811-819 of SEQ ID NO:2), SEQ ID NO:10 (amino acids 761-770 of SEQ ID NO:2), SEQ ID NO:11 (amino acids 805-813 of SEQ ID NO:2), and SEQ ID NO:12 (amino acids 859-867 of SEQ ID NO:2).

A polynucleotide useful in the present invention is preferably ligated into an expression vector which has been specifically optimized for polynucleotide vaccinations. Elements include a transcriptional promoter, immunogenic epitopes, and additional cistrons encoding immunoenhancing or immunomodulatory genes, with their own promoters, transcriptional terminator, bacterial origin of replication and antibiotic resistance gene, as well known to those skilled in the art. Optionally, the vector may contain internal ribosome entry sites (IRES) for the expression of polycistronic mRNA.

In one embodiment of this invention, a polynucleotide useful in the present invention is directly linked to a transcriptional promoter. The use of tissue-specific promoters or enhancers, for example the muscle creatine kinase (MCK) enhancer element, may be desirable to limit expression of the polynucleotide to a particular tissue type. For example, myocytes are terminally differentiated cells which do not divide. Integration of foreign DNA into chromosomes appears to require both cell division and protein synthesis. Thus, limiting protein expression to non-dividing cells such as myocytes may be preferable. In addition, a PSA promoter may be used to limit expression of the protein to prostate tissue. In one embodiment, tissue- or cell-specific promoters may be used to target the expression of the protein to antigen-presenting cells. For example, an α -fetoprotein (AFP) promoter (see e.g., Peyton et al. 2000, Proc. Natl. Acad. Sci., USA, 97:10890-10894) may be used to limit expression to liver tissues. However, use of the CMV promoter is adequate for achieving expression in many tissues into which the plasmid DNA vaccine is introduced.

Suitable vectors include any plasmid DNA construct encoding an androgen receptor, a fragment of the androgen receptor that comprises the ligand-binding domain, a suitable fragment of the ligand-binding domain, or a functional equivalent or derivative thereof, operatively linked to a suitable promoter. Examples of such vectors include the pCMV series of expression vectors, commercially available from Stratagene (La Jolla, Calif.); or the pcDNA or pREP series of expression vectors by Invitrogen Corporation (Carlsbad, Calif.).

A preferred vector is pNGVL3 available from the National Gene Vector Laboratory at the University of Michigan. This vector, similar to the pcDNA3.1 eukaryotic expression vector of Invitrogen Corp. (Carlsbad, Calif.), drives transcription from the CMV promoter, but also includes the CMV intron A sequence to enhance protein expression (Lee et al., 1997, Mol. Cells. 7:495-501). The vector also contains a multi-cloning site, and does not express a eukaryotic antibiotic resistance gene, such that the only protein expression expected in a eukaryotic system is the one driven from the CMV promoter, unlike the pcDNA vector. Another preferred vector is the pTVG4 vector described in US 2004/0142890, which is herein incorporated by reference in its entirety. The pTVG4 vector can be made by incorporating 2 copies of a 36-bp immunostimulatory (ISS) fragment containing the 5'-GTCGTT-3' motif previously identified (Hartmann et al., 2000, J. Immunol. 164:1617-24) into pNGVL3.

There are many embodiments of the instant invention which those skilled in the art can appreciate from the specification. Thus, different transcriptional promoters, terminators, and other transcriptional regulatory elements may be used successfully. Examples of other eukaryotic transcription promoters include the Rous sarcoma virus (RSV) promoter, the simian virus 40 (SV40) promoter, the human elongation factor-1 α (EF-1 α) promoter, and the human ubiquitin C (UbC) promoter.

A Kozak sequence can be provided upstream of the polynucleotide useful in the present invention to enhance the translation of the corresponding mRNA from the polynucleotide. For vertebrates, the Kozak sequence is (GCC)NC-CATGG (SEQ ID NO:7) wherein N is A or G and GCC is less conserved. For example, ACCATGG can be used. See Kozak, M. *Nucleic Acids Res.* 1987, 15:8125-48.

The vectors of the present invention may be delivered intradermally, intramuscularly, subcutaneously, or intravascularly (including intravenously and intraarterially). In preferred embodiments, delivery may be a combination of two or more of the various delivery methods.

"Naked" plasmid DNA expressing a transgene could be directly injected intradermally or intramuscularly, taken up, and expressed (see e.g., Wolff et al., 1990, *Science* 247:1465-8). The efficiency of this approach may be low, with only a small percentage of myocytes being directly transformed in vivo, and within only a limited area of muscle tissue targeted by this directed delivery. Various alternative approaches yielding a higher gene delivery efficiency are known (see e.g., Acsadi et al., 1991, *New Biol.* 3:71-81). Subsequent work on strategies that increase uptake of plasmid DNA by muscle tissue focused on various carrier solutions and molecules (Wolff et al., 1991, *Biotechniques* 11:474-85; and Budker et al., 1996, *Nat. Biotechnol.* 14:760-4), the use of myotoxic agents to enhance DNA uptake (Davis et al., 1993, *Hum. Gene Ther.* 4:151-9; and Danko et al., 1994, *Gene Ther.* 1:114-21), and the use of various transcriptional promoters and plasmid DNA backbones (Manthorpe et al., 1993, *Hum. Gene Ther.* 4:419-31).

In a preferred embodiment, plasmid vectors of the present invention may be delivered to the patient in need thereof intravascularly. Plasmid DNA delivered intravascularly resulted in 100-fold higher uptake in downstream tissues in rodent studies (Budker et al., 1996, *Gene Ther.* 3:593-8). Intravascular delivery may be intravenous, e.g. by direct injection of plasmid DNA into the portal vein of rodents with uptake and expression demonstrated in hepatocytes (Budker et al., 1996, *Gene Ther.* 3:593-8; and Zhang et al., 1997, *Hum. Gene Ther.* 8:1763-72). Intravascular delivery may also be performed more directly by intraarterial delivery. For example, initial studies in rodents demonstrated that high levels of gene expression in hind limb muscle could be obtained by rapid injection of plasmid DNA into the femoral artery (Budker et al., 1998, *Gene Ther.* 5:272-276). This approach is efficient and safe in non-human primates as well, with an average of 7% of downstream myofibers expressing a β -galactosidase reporter construct two weeks after intraarterial DNA administration (Zhang et al., 2001, *Hum. Gene Ther.* 12:427-438). Parallel studies in T cell immuno-suppressed rats showed that gene expression was stable for at least 10 weeks (Zhang et al., 2001, *Hum. Gene Ther.* 12:427-438).

Accordingly, delivery of plasmid DNA vaccines of the present invention can be done by direct intraarterial administration. This method provides more effective delivery to

MHC class I expressing cells. Administrations of plasmid DNA vaccines intravascularly may result in increased antigen expression and subsequently lead to enhanced immune responses, and increased antigen expression in MHC class I expressing cells by means of intraarterial delivery of DNA plasmid may lead to a more robust immune response with androgen receptor-specific CTL. An intraarterial method of DNA delivery has been shown to be at least as effective as or more effective than traditional intradermal administration of DNA in eliciting prostatic acid phosphatase-specific immunity.

In another embodiment, intravenous delivery may also be used, employing methods well known to those skilled in the art (See e.g., Budker et al., 1998, *Gene Ther.* 5:272-276; and Budker et al., 1996, *Gene Ther.* 3:593-598). This delivery method may lead to a high level of antigen expression in hepatocytes. Expression of the antigen in liver, a tissue more rich with antigen-presenting cells, may lead to a more pronounced Th1/CTL response than expression in muscle tissue.

The DNA or peptide vaccines of the present invention can be used in a prime-boost strategy to induce robust and long-lasting immune response to androgen receptor. Priming and boosting vaccination protocols based on repeated injections of the same antigenic construct are well known and result in strong CTL responses. In general, the first dose may not produce protective immunity, but only "primes" the immune system. A protective immune response develops after the second or third dose.

In one embodiment, the DNA or peptide vaccines of the present invention may be used in a conventional prime-boost strategy, in which the same antigen is administered to the animal in multiple doses. In a preferred embodiment, the DNA or peptide vaccine is used in one or more inoculations. These boosts are performed according to conventional techniques, and can be further optimized empirically in terms of schedule of administration, route of administration, choice of adjuvant, dose, and potential sequence when administered with another vaccine, therapy or homologous vaccine.

The peptide or DNA vaccines of the present invention may be used in a prime-boost strategy using an alternative administration of xenoantigen and autoantigen or xenoantigen- and autoantigen-encoding vectors. Specifically, according to the present invention, the animal is first treated, or "primed," with a peptide antigen of foreign origin (a "xenoantigen") or DNA vaccine encoding the antigen of foreign origin. The animal is then treated with another peptide antigen which corresponds to the xenoantigen but is of self origin ("autoantigen") or another DNA vaccine encoding the autoantigen. This way, the immune reaction to the antigen is boosted. The boosting step may be repeated one or more times.

A xenoantigen, as compared to a self-antigen or an autoantigen, is an antigen originated in or derived from a species different from the species that generates an immune reaction against the antigen. Xenoantigens usually are highly homologous molecules to a corresponding autoantigen. Xenoantigens have been shown to be able to elicit auto-reactive immunity. For example, molecular mimicry by highly homologous viral antigens has been one theory to explain the occurrence of several autoimmune diseases (von Herrat and Oldstone, 1996, *Curr. Opin. Immunol.* 8:878-885; and Oldstone, 1998, *Faseb J.* 12:1255-1265). That is, the induction of immune responses following infection by viral antigens that closely resemble normal autologous proteins may then lead to an autoimmune reaction to the autologous protein.

The use of highly homologous foreign antigens or xenoantigens as vaccine antigens to elicit autoreactive immunity has been explored in animal models. For example, xenoantigens derived from *zona pellucida* of foreign species can elicit autoreactive T cell responses and disrupt ovarian function in a variety of animal species studied (Mahi-Brown et al., 1992, J. Reprod. Immunol. 21:29-46; and Mahi-Brown, 1996, J. Reprod. Fertil. Suppl. 50:165-74). While not wishing to be bound by any theory on mechanism, it is believed that because T cells involved in autoimmune processes recognize peptide epitopes presented in the context of MHC molecules, the uptake and MHC presentation of a homologous foreign antigen presumably exposes T cell epitopes with enhanced MHC binding or unmasks cryptic epitopes of the native antigen not normally recognized.

While the prime-boost strategy is known to work with antigens of different origins, it is readily apparent to those ordinarily skilled in the art that variants, derivatives or equivalents, as discussed above, of the nucleotide sequence encoding a self-antigen can also be used to achieve the same results as xenoantigens.

The peptide or DNA vaccines of the present invention may be used together with prostate cancer vaccines based on other antigens such as prostatic acid phosphatase-based antigens. The androgen receptor-based vaccines and vaccines based on other antigens can be used simultaneously or at different times. Each may be used in a prime-boost strategy.

The present invention also provides a method for determining the effectiveness of a treatment for prostate cancer. The method includes the steps of (a) measuring the frequency or amount of cytotoxic T lymphocytes (CTLs) specific for a peptide selected from SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12 prior to providing at least a portion of the treatment to a mammal having prostate cancer, (b) measuring the frequency or amount of CTLs specific for the peptide after said portion of the treatment is provided to the mammal, and (c) comparing the frequency or amount of CTLs of (a) and that of (b) wherein the frequency or amount of CTLs of (b) being higher than that of (a) indicates that the treatment is effective. For example, a biological sample containing CTLs such as a blood sample or a sample of peripheral blood mononuclear cells (PBMC) can be taken from the mammal and the frequency or amount of CTLs in the blood sample can be measured. In one embodiment, the method is used to determine the effectiveness of a treatment provided to a human prostate cancer patient.

One of ordinary skill in the art is familiar with the techniques for functional and quantitative measurements of antigen-specific T cells. Examples include but are not limited to limited dilution assays (LDA), enzyme linked immunosorbent assay on a single cell level (ELISPOT), intracellular staining, and MHC/HLA multimer (e.g., dimer, tetramer, and pentamer) staining. Description on the MHC/HLA multimer staining technique can be found, for example, in Arnold H Bakker and Ton N M Schumacher (Current Opinion in Immunology, 2005, 17:428-433), Meidenbauer N et al. (Methods, 2003, 31:160-171), and U.S. patent application publication 200723036812.

In one embodiment, a biological sample (e.g., a blood sample or PBMC sample) containing CTLs from a patient is obtained and the sample is brought into contacted with an HLA multimer (e.g., an HLA tetramer). The frequency or amount of CTLs specific for a peptide antigen bound the HLA tetramer can then be measured by known techniques such as flow cytometry.

The invention will be more fully understood upon consideration of the following non-limiting examples.

Example 1

Immune Responses to Androgen Receptor Ligand-Binding Domain can Exist in Patients with Prostate Cancer

We evaluated whether or not patients with prostate cancer have existing immune responses to the androgen receptor (AR). We focused these studies on responses specific for the AR ligand-binding domain (LBD). As shown in FIG. 1, we found that prostate cancer patients, but not healthy male blood donors, have antibodies that are specific for the AR LBD. These antibodies were predominantly of the IgG isotype and IgG₂ sub-isotype (data not shown). Moreover, we identified that patients with antibody responses to the AR LBD have CD4⁺ and CD8⁺ T cells that proliferate, as well as cells that secrete interferon-gamma (IFN γ), in response to stimulation with the AR LBD (FIG. 2). Antigen-specific IL-10 secretion was observed in many patients who did not have evidence of antibody responses to AR LBD (FIG. 3). Taken together, these results demonstrate that some patients with prostate cancer can have pre-existing cellular immune responses specific for the AR. These findings further suggest that tolerance against the AR, which may be prevalent, is not absolute and can be overcome in some patients with prostate cancer.

Example 2

Materials and Methods

Subject Population: Peripheral blood mononuclear cells (PBMC) were obtained from eleven patients with prostate cancer at the University of Wisconsin Hospital and Clinics between 2001 and 2007. All subjects gave Institutional Review Board-approved written informed consent for their blood products to be used for immunological research. PBMC were prepared from heparinized blood by gradient centrifugation.

T2 Binding Assay: After passing T2 cells into fresh media the day before the assay, these cells were pulsed with 50 μ g/mL peptide overnight at 37° C./5% CO₂. The next day, the levels of HLA-A2 expression on the surface of these cells were measured using a fluorescently-labeled HLA-A2 antibody (Clone BB7.2, BD Biosciences, San Jose, Calif.) followed by flow cytometry using FACSCaliber system (BD Biosciences). The reported "fold change" in fluorescence intensity was calculated by averaging the mean fluorescent intensity results from triplicate samples and dividing these values by the average mean fluorescent intensity of the negative control (media only) samples.

T-Cell Culturing: Human prostate cancer patients underwent leukapheresis, and PBMCs were isolated using a Ficoll-Paque gradient (Pharmacia, Kirkland, Quebec). Immature dendritic cells (iDCs) were generated by incubating flask-adherent PBMCs with 20 ng/mL granulocyte-macrophage colony stimulating factor and 10 ng/mL interleukin four (IL-4) for six days at 37° C./5% CO₂ (cytokines from: Fitzgerald Industries, Concord, Mass.). These iDCs were then treated with 150 ng/mL IL-6, 10 ng/mL IL-1 β , 10 ng/mL tumor

necrosis factor alpha (TNF- α), and 1 mg/mL prostaglandin E2 (Fitzgerald Industries) to generate mature dendritic cells (mDCs). These mDCs were then pulsed with 20 μ g/mL AR LBD-derived peptide, and after being irradiated were co-incubated with CD4+ and CD8+ T-cells negatively isolated from autologous PBMCs (Invitrogen, Carlsbad, Calif.). After incubating for 24 hours, these cultures were then treated with 10 U/mL IL-2 and 30 U/mL IL-7 (Fitzgerald Industries), and incubated for another 6 days. These cultured T-cells were re-stimulated weekly using peptide-pulsed antigen presenting cells (either mDCs or the TK6 lymphoma cell line).

Lactate Dehydrogenase Cytotoxicity Assay: T-cell cultures that underwent at least two stimulations (or splenocyte cultures that had undergone one stimulation) were collected and incubated for four hours with target cells (either T2 cells pulsed for two hours with a specific or non-specific peptide, the LNCaP prostate cancer cell line, or media alone) at various effector-to-target ratios. After incubation, supernatants were collected and levels of lactate dehydrogenase were measured using the CytoTox 96 Non-Radioactive Assay (Promega). The relative percentage of peptide-specific lysis was quantitated using the following equation:

% Cytotoxicity =

$$\frac{\text{Experimental} - \text{Effector Spontaneous} - \text{Target Spontaneous}}{\text{Target Maximum} - \text{Target Spontaneous}}$$

To conduct a mini-cytotoxicity assay of limited-dilution clones, 50 μ l (1/4) of the cultured clones were incubated for four hours with either T2 cells pulsed with the specific peptide, a non-specific peptide, or media alone. For the subsequent characterization of CTL clone cytotoxicity against prostate cancer cells, cryopreserved CTL clones were thawed, washed 3 times, and restimulated by incubating the clones with peptide-pulsed irradiated TK6 cells for 7 days. These clones were then resuspended in 10 U/mL IL-2, and allowed to incubate for three days at 37° C./5% CO₂. They were then analyzed for cytotoxicity (as described above) against four prostate cancer cell lines: LNCaP, DU145, LAPC4, and MDAPCa-2b cell lines. To characterize HLA-A2-restriction of observed cytotoxicity, a HLA-A2 antibody was added to the reaction (Clone BB7.2, BD Biosciences, 1 μ g/mL).

Limited-Dilution T-Cell Cloning and T-cell expansion: After T-cell cultures containing AR LBD peptide-specific T-cells were identified, peptide-specific T-cell clones were isolated using limited-dilution cloning. Briefly, cultured T-cells were diluted to 400 cells/mL, and were diluted at a 1:1 ratio down the rows of a 96-well plate. These T-cells were mixed with 2 $\times$ 10⁵ autologous PBMCs, as well as an anti-CD3 antibody (Clone UCHT1, BD Biosciences, 120 ng/mL) and IL-2 (Fitzgerald Industries, 200 U/mL), and were incubated at 37° C./5% CO₂ for 12-14 days. Cultures generated from a single cell were identified, and their peptide-specificity was analyzed using a mini-cytotoxicity assay (as above). Peptide-specific T-cell clones were then expanded in the presence of autologous PBMCs, TK6 cells, and an anti-CD3 antibody (30 ng/mL). The next day, 50 U/mL IL-2 was added to the cultures. Six days later, CTL clones were resuspended in media with 80 U/mL IL2, and three days later were again resuspended in 20 U/mL IL-2. After three additional days,

expanded T-cell clones were analyzed for cytotoxicity against peptide-specific or non-specific T2-pulsed cells (as described above).

Surface Molecule Staining: T-cell clones were thawed, washed 3 times, and resuspended in media supplemented with 10 U/mL IL-2 for 18 hours at 37° C./5% CO₂. These recovered clones were then resuspended in staining buffer (phosphate-buffered saline+5% fetal bovine serum) and incubated with fluorescently-labeled antibodies specific for CD3, CD4, CD8, and CD56 (clones SK7, RPA-T4, RPA-T8, or NCAM16.2, respectively; BD Biosciences) or the appropriate controls, for 30 minutes on ice. Cells were subsequently analyzed using an LSR II flow cytometer (BD Biosciences), counting 100,000 events. Cells were gated based on CD3+/CD56+ expression and CD4+/CD8+ expression.

Intracellular Cytokine Staining: Recovered CTL clones were restimulated for one hour with media alone, the specific peptide, a non-specific peptide (peptides both at 2 μ g/mL), or Phorbol Myristate Acetate (Sigma-Aldrich, St. Louis, Mo.; 10 μ g/mL) and Ionomycin (MP Biomedicals, Solon, Ohio; 1 μ g/mL). Cells were then treated with monensin (BD Biosciences; 1 μ l per 1.5 mL cell culture) for four hours at 37° C./5% CO₂, followed by a brief blocking treatment with mouse IgG. Cells were then resuspended in staining buffer (phosphate-buffered saline+5% fetal bovine serum) and incubated with fluorescently-labeled CD3- and CD8-specific antibodies (BD Biosciences), or the appropriate controls, for 30 minutes on ice. After fixation and permeabilization, intracellular staining was conducted using fluorescently-labeled IFN γ and TNF α antibodies (Clones 4S.B3 and MA611, respectively; BD Biosciences), or the appropriate isotype controls. Cells were subsequently analyzed using an LSR II flow cytometer, counting 100,000 events. IFN γ and TNF α -positive events were determined by gating CD3+/CD8+ cells and analyzing this population for co-expression of IFN γ and TNF α .

Immunization of HLA-A2/HLA-DR1 mice: Groups of four 6-10 week old HLA-A2/DR1 transgenic male mice (Charles River Laboratory—France with the permission of Dr. François Lemmonier) were immunized subcutaneously with 100 μ g AR811 peptide with Complete Freund's Adjuvant (CFA) or with CFA alone (Sigma-Aldrich), and seven days later, the mice were euthanized. Spleens were collected, and splenocytes were isolated by gradient centrifugation (Histopaque 3130, Sigma-Aldrich). Splenocytes were stimulated with 10 μ g/mL peptide for two hours, and on the second day, recombinant murine IL-2 and IL-7 (Fitzgerald Industries) were added to 10 U/mL and 30 U/mL, respectively. The cultures were then allowed to incubate an additional six days before analysis.

Construction of pTVG4 and pTVG-ARLBD: Plasmid DNA expression vectors have been developed for use in human vaccines. Shown in FIG. 4 is a plasmid map of the pTVG4 vector as constructed for animal (e.g., rat and mouse) and human immunization. The coding sequence for the ligand-binding domain of the human androgen receptor gene has been inserted into the pTVG4 vector to create the immunization vector pTVG-ARLBD (see below).

The plasmid vector pNGVL3 was obtained from the National Gene Vector Laboratory at the University of Michigan (courtesy, Dr. Robert Gerard). This vector, similar to the

pcDNA3.1 expression vector from Invitrogen Corp. (Carlsbad, Calif.), drives transcription from the CMV promoter, but also includes the CMV intron A sequence to enhance transcription (Lee et al., 1997, Mol. Cells. 7:495-501). The vector also contains a multi-cloning site, and does not express a eukaryotic antibiotic resistance gene, such that the only protein expression expected in a eukaryotic system is the one driven from the CMV promoter, unlike the pcDNA vector. To this vector has been added 2 copies of a 36-bp immunostimulatory (ISS) fragment containing the 5'-GTCGTT-3' motif previously identified (Hartmann et al., 2000, J. Immunol. 164:1617-24), to create the vector pTVG4 (FIG. 4). ColE1, the DNA sequence for Colicin E1 which can be used for cloning purpose, is provided in the vector. Kan, the DNA sequence encoding a kanamycin resistance gene which can also be used for cloning purpose, is also provided in the vector. The coding sequence for the ligand-binding domain of the human androgen receptor gene has been cloned into this-vector and a Kozak sequence has been provided directly upstream of the coding sequence to enhance the translation of the corresponding mRNA (FIG. 5). Expression of the ligand-binding domain has been confirmed by in vitro expression studies (not shown). This construct, named pTVG-ARLBD, is used for the immunization of animals and humans.

Immunization and tumor protection of Copenhagen rats: Groups of ten 9-11 week old Copenhagen rats (Harlan) were immunized intradermally with 100 µg pTVG-ARLBD or 100 µg pTVG4 alone with 1 µg rat GM-CSF. Rats received three booster immunizations (100 µg) every 14 days, and 14 days after the last immunization, rats were challenged with 10,000 syngeneic Mat-LyLu prostate tumor cells, given along

Matrigel Matrix (BD Biosciences). Tumors (long and short diameters) were measured every two days, and volumes were calculated using the following equation:

$$\text{Tumor volume} = (\pi/6) \times (d_{\text{short}})^2 \times (d_{\text{long}})$$

Results

AR LBD-specific CD8+ T-cells from prostate cancer patients can lyse prostate cancer cells: To characterize CD8+ T-cell responses to the AR LBD, the amino acid sequence of the AR LBD was evaluated for potential HLA-A2-binding epitopes that fit the consensus peptide binding sequence of X-L/M-X-X-X-V-X-X-V/L (SEQ ID NO:8), using the algorithm of Parker and colleagues (Parker K C et al., 1994, J. Immunol. 152:163-175). As demonstrated in Table 1, ten unique peptides were identified. These peptides were synthesized and then characterized for their affinity for HLA-A2 in vitro using T2 binding assays. The results from these binding studies are also shown in Table 1. As demonstrated, several potential epitopes were predicted and found experimentally to bind strongly to HLA-A2. Peptide-specific T-cell lines were then cultured from the peripheral blood of 11 HLA-A2+ patients with prostate cancer, using each of these peptides, and then tested for their cytolytic activity against both peptide-pulsed HLA-A2-expressing target cells and the HLA-A2+ LNCaP prostate cancer cell line. Specifically, naive T cells were isolated by magnetic negative selection (Dyna) from the peripheral blood mononuclear cells (PBMCs) of HLA-A2-expressing prostate cancer patients, and cultured in the presence of autologous cytokine-matured irradiated dendritic cells (mDC) that had been loaded with individual peptides. These cultures received interleukin 2 (IL-2) and IL-7 the day after the culture, and were re-stimulated weekly with peptide-pulsed mDCs. Beginning after two weeks of stimulation, the T-cell cultures were tested weekly for their cytolytic activity using peptide-loaded T2 cells as target cells. Peptide-specific T-cells could be cultured from the majority of patients to at least one of these peptides, and peptide-specific T-cell lines could be cultured from the majority (7/11) of HLA-A2-expressing patients against the AR811 peptide in particular (Table 2).

TABLE 1

Prediction of AR LBD-derived HLA-A2-specific peptide epitopes.			
Peptide	Sequence	Predicted Binding Affinity (t _{1/2} (min) of Dissociation)	In Vitro HLA-A2 Expression (Relative Mean Fluorescent Intensity)
AR677	VLEAIEPGV (SEQ ID NO:13)	7.6	1.37 ± 0.14
AR700	ALLSSLNEL (SEQ ID NO:14)	182	2.66 ± 0.28
AR708	LGERQLVHV (SEQ ID NO:15)	0.114	1.16 ± 0.06
AR742	WMGLMVFAM (SEQ ID NO:16)	220	1.32 ± 0.06
AR761	RMLYFAPDLV (SEQ ID NO:10)	217	2.15 ± 0.13
AR805	FLCMKALLL (SEQ ID NO:11)	98	2.19 ± 0.14
AR811	LLLFSTIPV (SEQ ID NO:9)	1006	2.54 ± 0.25

TABLE 1-continued

Prediction of AR LBD-derived HLA-A2-specific peptide epitopes.			
Peptide	Sequence	Predicted Binding Affinity (t _{1/2} (min) of Dissociation)	In Vitro HLA-A2 Expression (Relative Mean Fluorescent Intensity)
AR814	FSIIPVDGL (SEQ ID NO:17)	111	1.01 ± 0.12
AR859	QLTKLLDSV (SEQ ID NO:12)	78	1.33 ± 0.15
AR862	KLLDSVQPI (SEQ ID NO:18)	1274	1.65 ± 0.19
Influenza	GILGFVFTL (SEQ ID NO:19)	30	1.88 ± 0.24
Negative Control		0	1.00 ± 0.07

AR LBD peptide epitopes were identified by scanning the protein sequence of the AR LBD for 9-mer or 10-mer peptides that fit the HLA-A2 consensus binding sequence X-L/M-X-X-X-V-X-X-V/L (SEQ ID NO:8) and by their predicted binding affinity to HLA-A2 (Bioinformatics and Molecular Analysis Section). These peptides were synthesized and then analyzed for their affinity for HLA-A2 using a T2 binding assay. Shown is the ratio of the mean fluorescent intensity, calculated from triplicatesamples, of peptide-loaded T cells normalized against unloaded T cells. Influenza: positive control influenza matrix protein peptide; negative control: vehicle.

TABLE 2

Peptide	Peptide-specific lysis (Number of patients)	Prostate cancer cell lysis
AR677	2/11	-
AR700	1/11	-
AR708	1/11	-
AR742	2/11	-
AR761	6/11	+/-
AR805	3/11	+/-
AR811	7/11	+
AR814	3/11	-
AR859	2/11	+/-
AR862	5/11	-

As shown in FIG. 6, AR811 peptide-specific T-cells were found to lyse T2 cells in a peptide-specific fashion, and could lyse the LNCaP cell line. Several other peptides demonstrated peptide-specific lysis and variable amounts of lysis against the LNCaP cell line (Table 2). In contrast, T-cells specific for the AR859 peptide could be cultured, and while these showed peptide-specific lysis they did not lyse the LNCaP cell line (FIG. 6B). AR811-specific T-cells were cloned by limited dilution and found to be CD8+, and to secrete both IFN γ and TNF α in response to peptide stimulation (FIG. 7). Moreover, these cells lysed prostate cancer cells in an MHC class I-restricted fashion (FIG. 8). These findings confirm that the AR811 peptide is a naturally processed and presented HLA-A2 epitope from the AR.

HLA-A2 transgenic mice immunized with the AR811 peptide developed peptide-specific CTL: Human HLA transgenic mice have become a valuable tool for the identification and study of human MHC class I-specific epitopes and CTL responses. In work published by others, transgenic mice expressing human HLA-A201 have been immunized directly with peptides, or with DNA encoding antigens, or protein antigens, to identify HLA-A2-specific epitopes (Carralot J P et al., 2005, Int Immunol 17:591-7; Gallez-Hawkins G et al., 2003, J Virol 77:4457-62; and Loirat D et al., 2000, J Immunol 165:4748-55). Unfortunately, many of the early studies

with these transgenic strains were complicated by the preference to develop H-2-restricted murine responses rather than HLA-A2-restricted CTL responses, thus limiting the usefulness of these strains. This led to the development of the HHD strains by Dr. François Lemonnier and colleagues at the Institut Pasteur, in which the mouse MHC class I H-2 Db was knocked out, and mice express human β 2-microglobulin and HLA-A201 monochains fused to the α 3, transmembrane and cytoplasmic domains of the mouse MHC class I molecule (Pascolo S et al., 1997, J Exp Med 185:2043-51). These strains and derivatives have been particularly useful as these mice are forced to use a diverse repertoire of CD8+ T-cells specific for HLA-A2 (Pascolo S et al., 1997, J Exp Med 185:2043-51), and have been demonstrated to be superior in eliciting HLA-A2-restricted CTL (Ramage J M et al., 2004, Vaccine 22:1728-31). The HHD-II transgenic mouse strain developed by Dr. Lemonnier expresses both human HLA-A201 and HLA-DR1, and has both the murine H-2 class I and MHC class II knocked out (Pajot A et al., 2004, Eur J Immunol 34:3060-9). This particular strain has been used for the identification of HLA-DR1-restricted CD4+ T-cell epitopes as well as HLA-A2-restricted epitopes (Pajot A et al., 2006, Microbes Infect 8:2783-90). HHD-II mice were immunized once with 100 μ g of the AR811 peptide in complete Freund's adjuvant (CFA) or with CFA alone. As shown in FIG. 9, AR811 peptide-specific CTL could be identified after immunization.

DNA vaccine encoding AR LBD can elicit antigen-specific responses and retard prostate cancer cell growth in vivo: cDNA was prepared from a prostate cancer cell line, and AR LBD (amino acids 664-920) was cloned into the pTVG4 vector as described above (similar to cloning prostatic acid phosphatase into the pTVG4 vector described in Johnson L E, et al., 2007, Canc Immunol Immunoth 56:885-895, which is herein incorporated by reference in its entirety). CHO cells transiently transfected with this pTVG-ARLBD construct produced AR LBD mRNA and protein that could be detected by RT-PCR and by Western blot analysis (data not shown).

Male Copenhagen rats, 2-3 months of age, were then immunized with 100 µg of pTVG-ARLBD four times at 14-day intervals, intradermally with 5 µg rat GM-CSF given as a vaccine adjuvant. Two weeks after the final immunization, blood was collected for immunological analysis. As shown in FIG. 10, animals immunized with pTVG-ARLBD, but not the pTVG4 vector, developed AR LBD-specific IgG antibody responses. To assess anti-tumor efficacy, Copenhagen rats that had been immunized four times at 14-day intervals were then challenged with 1×10^4 syngeneic Mat-LyLu prostate tumor cells implanted subcutaneously. As shown in FIG. 11, immunization with pTVG-ARLBD, but not the pTVG4 vector, retarded the growth of these Mat-LyLu prostate tumors.

Example 3

Prophetic

Prostate Cancer Therapy with pTVG-ARLBD DNA Vaccine

Groups of ten suitable rats or mice such as Copenhagen rats are challenged with a suitable number of prostate cancer cells

(e.g., 1×10^4 Mat-LyLu prostate cancer cells). These rats or mice are then immunized with either the pTVG4 (negative control) or pTVG-ARLBD constructs. One example is to immunize the rats or mice at days 2, 9, and 16 after the tumor challenge with 100 µg injected intradermally along with 5 µg rat or mouse GM-CSF as an adjuvant. Suitable schemes with fewer or additional immunizations may be used as alternatives. Optionally, boosts (e.g., on weekly basis) can be provided. Other suitable amounts of DNA or adjuvant can be used, as can different adjuvants (such as Freund's adjuvant) or additional vaccines (such as those targeting prostatic acid phosphatase or the synovial sarcoma X chromosome family of proteins). In addition, other suitable routes of administration may be used (such as intravenously). Tumor growth is monitored daily using bi-dimensional measurements. Sera from these rats or mice may be obtained and used to evaluate the presence of AR LBD antibodies. It is expected that immunization with pTVG-ARLBD DNA vaccine will elicit therapeutic anti-tumor response.

The present invention is not intended to be limited to the foregoing examples, but encompasses all such modifications and variations as come within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 19

<210> SEQ ID NO 1

<211> LENGTH: 4314

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1116) .. (3878)

<400> SEQUENCE: 1

```

cgagatcccg gggagccagc ttgctgggag agcgggacgg tccggagcaa gccagagggc      60
agaggaggcg acagaggga aaagggccga gctagccgct ccagtgtgt acaggagccg      120
aagggaacga ccacgccagc cccagcccgg ctccagcgac agccaacgcc tcttgccagc      180
cggcggttc gaagccgcc cccggagctg ccctttcttc ttcgggtgaag tttttaaaag      240
ctgctaaaga ctggaggaa gcaaggaaag tgctggttag gactgacggc tgctttgtc      300
ctctctctct ccacccgcgc tccccccacc ctgccttccc cccctcccc gtcttctctc      360
ccgcagctgc ctacgtggc tactctcagc caacccccct caccaccctt ctccccacc      420
gccccccgc ccccgctggc ccagcgtgc cagcccgagt ttgcagagag gtaactcct      480
ttggtgcga gcgggcgagc tagctgcaca ttgcaaagaa ggctcttagg agccaggcga      540
ctggggagcg gcttcagcac tgcagccacg acccgctgg ttaggctgca cgcggagaga      600
acctctgtt tccccccact ctctctccac ctctctctgc ctccccacc ccgagtgcgg      660
agccagagat caaaagatga aaaggcagtc aggtcttcag tagccaaaa acaaaacaaa      720
caaaaacaaa aaagccgaaa taaaagaaa agataataac tcagttctta ttgcaccta      780
cttcagtga cactgaattt ggaagggtga ggattttgtt tttttctttt aagatctggg      840
catcttttga atctaccctt caagtattaa gagacagact gtgagcctag cagggcagat      900
cttgctccacc gtgtgtcttc ttctgcacga gactttgagg gtgtcagagc gctttttgag      960
tggttgctcc cgcaagtctt ctctctctga gcttcccga ggtgggcagc tagctgcagc      1020
gactaccgca tcatcacagc ctgttgaact ctctgagca agagaagggg aggcggggta      1080

```

-continued

aggggaagtag gtggaagatt cagccaagct caagg atg gaa gtg cag tta ggg															1133
Met Glu Val Gln Leu Gly															
1 5															
ctg gga agg gtc tac cct cgg ccg ccg tcc aag acc tac cga gga gct	1181														
Leu Gly Arg Val Tyr Pro Arg Pro Ser Lys Thr Tyr Arg Gly Ala															
10 15 20															
ttc cag aat ctg ttc cag agc gtg cgc gaa gtg atc cag aac ccg ggc	1229														
Phe Gln Asn Leu Phe Gln Ser Val Arg Glu Val Ile Gln Asn Pro Gly															
25 30 35															
ccc agg cac cca gag gcc gcg agc gca gca cct ccc ggc gcc agt ttg	1277														
Pro Arg His Pro Glu Ala Ala Ser Ala Ala Pro Pro Gly Ala Ser Leu															
40 45 50															
ctg ctg ctg cag cag cag cag cag cag cag cag cag cag cag cag	1325														
Leu Leu Leu Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln															
55 60 65 70															
cag cag cag cag cag cag cag cag cag caa gag act agc ccc agg cag	1373														
Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Glu Thr Ser Pro Arg Gln															
75 80 85															
cag cag cag cag cag ggt gag gat ggt tct ccc caa gcc cat cgt aga	1421														
Gln Gln Gln Gln Gln Gly Glu Asp Gly Ser Pro Gln Ala His Arg Arg															
90 95 100															
ggc ccc aca ggc tac ctg gtc ctg gat gag gaa cag caa cct tca cag	1469														
Gly Pro Thr Gly Tyr Leu Val Leu Asp Glu Glu Gln Gln Pro Ser Gln															
105 110 115															
ccg cag tcg gcc ctg gag tgc cac ccc gag aga ggt tgc gtc cca gag	1517														
Pro Gln Ser Ala Leu Glu Cys His Pro Glu Arg Gly Cys Val Pro Glu															
120 125 130															
cct gga gcc gcc gtg gcc gcc agc aag ggg ctg ccg cag cag ctg cca	1565														
Pro Gly Ala Ala Val Ala Ala Ser Lys Gly Leu Pro Gln Gln Leu Pro															
135 140 145 150															
gca cct ccg gac gag gat gac tca gct gcc cca tcc acg ttg tcc ctg	1613														
Ala Pro Pro Asp Glu Asp Asp Ser Ala Ala Pro Ser Thr Leu Ser Leu															
155 160 165															
ctg ggc ccc act ttc ccc ggc tta agc agc tgc tcc gct gac ctt aaa	1661														
Leu Gly Pro Thr Phe Pro Gly Leu Ser Ser Cys Ser Ala Asp Leu Lys															
170 175 180															
gac atc ctg agc gag gcc agc acc atg caa ctc ctt cag caa cag cag	1709														
Asp Ile Leu Ser Glu Ala Ser Thr Met Gln Leu Leu Gln Gln Gln Gln															
185 190 195															
cag gaa gca gta tcc gaa ggc agc agc agc ggg aga gcg agg gag gcc	1757														
Gln Glu Ala Val Ser Glu Gly Ser Ser Ser Gly Arg Ala Arg Glu Ala															
200 205 210															
tcg ggg gct ccc act tcc tcc aag gac aat tac tta ggg ggc act tcg	1805														
Ser Gly Ala Pro Thr Ser Ser Lys Asp Asn Tyr Leu Gly Gly Thr Ser															
215 220 225 230															
acc att tct gac aac gcc aag gag ttg tgt aag gca gtg tcg gtg tcc	1853														
Thr Ile Ser Asp Asn Ala Lys Glu Leu Cys Lys Ala Val Ser Val Ser															
235 240 245															
atg ggc ctg ggt gtg gag gcg ttg gag cat ctg agt cca ggg gaa cag	1901														
Met Gly Leu Gly Val Glu Ala Leu Glu His Leu Ser Pro Gly Glu Gln															
250 255 260															
ctt cgg ggg gat tgc atg tac gcc cca ctt ttg gga gtt cca ccc gct	1949														
Leu Arg Gly Asp Cys Met Tyr Ala Pro Leu Leu Gly Val Pro Pro Ala															
265 270 275															
gtg cgt ccc act cct tgt gcc cca ttg gcc gaa tgc aaa ggt tct ctg	1997														
Val Arg Pro Thr Pro Cys Ala Pro Leu Ala Glu Cys Lys Gly Ser Leu															
280 285 290															
cta gac gac agc gca ggc aag agc act gaa gat act gct gag tat tcc	2045														
Leu Asp Asp Ser Ala Gly Lys Ser Thr Glu Asp Thr Ala Glu Tyr Ser															
295 300 305 310															

-continued

cct ttc aag gga ggt tac acc aaa ggg cta gaa ggc gag agc cta ggc Pro Phe Lys Gly Gly Tyr Thr Lys Gly Leu Glu Gly Glu Ser Leu Gly 315 320 325	2093
tgc tct ggc agc gct gca gca ggg agc tcc ggg aca ctt gaa ctg ccg Cys Ser Gly Ser Ala Ala Ala Gly Ser Ser Gly Thr Leu Glu Leu Pro 330 335 340	2141
tct acc ctg tct ctc tac aag tcc gga gca ctg gac gag gca gct gcg Ser Thr Leu Ser Leu Tyr Lys Ser Gly Ala Leu Asp Glu Ala Ala Ala 345 350 355	2189
tac cag agt cgc gac tac tac aac ttt cca ctg gct ctg gcc gga ccg Tyr Gln Ser Arg Asp Tyr Tyr Asn Phe Pro Leu Ala Leu Ala Gly Pro 360 365 370	2237
ccg ccc cct ccg ccg cct ccc cat ccc cac gct cgc atc aag ctg gag Pro Pro Pro Pro Pro Pro Pro His Pro His Ala Arg Ile Lys Leu Glu 375 380 385 390	2285
aac ccg ctg gac tac ggc agc gcc tgg gcg gct gcg gcg gcg cag tgc Asn Pro Leu Asp Tyr Gly Ser Ala Trp Ala Ala Ala Ala Ala Gln Cys 395 400 405	2333
cgc tat ggg gac ctg gcg agc ctg cat ggc gcg ggt gca gcg gga ccc Arg Tyr Gly Asp Leu Ala Ser Leu His Gly Ala Gly Ala Ala Gly Pro 410 415 420	2381
ggc tct ggg tca ccc tca gcc gcc gct tcc tca tcc tgg cac act ctc Gly Ser Gly Ser Pro Ser Ala Ala Ala Ser Ser Ser Trp His Thr Leu 425 430 435	2429
ttc aca gcc gaa gaa ggc cag ttg tat gga ccg tgt ggt ggt ggt ggg Phe Thr Ala Glu Glu Gly Gln Leu Tyr Gly Pro Cys Gly Gly Gly Gly 440 445 450	2477
ggc ggt ggc ggc ggc ggc ggc ggc ggc ggc ggc ggc ggc ggc ggc Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly 455 460 465 470	2525
ggc ggc ggc gag gcg gga gct gta gcc ccc tac ggc tac act ccg ccc Gly Gly Gly Glu Ala Gly Ala Val Ala Pro Tyr Gly Tyr Thr Arg Pro 475 480 485	2573
cct cag ggg ctg gcg ggc cag gaa agc gac ttc acc gca cct gat gtg Pro Gln Gly Leu Ala Gly Gln Glu Ser Asp Phe Thr Ala Pro Asp Val 490 495 500	2621
tgg tac cct ggc ggc atg gtg agc aga gtg ccc tat ccc agt ccc act Trp Tyr Pro Gly Gly Met Val Ser Arg Val Pro Tyr Pro Ser Pro Thr 505 510 515	2669
tgt gtc aaa agc gaa atg ggc ccc tgg atg gat agc tac tcc gga cct Cys Val Lys Ser Glu Met Gly Pro Trp Met Asp Ser Tyr Ser Gly Pro 520 525 530	2717
tac ggg gac atg cgt ttg gag act gcc agg gac cat gtt ttg ccc att Tyr Gly Asp Met Arg Leu Glu Thr Ala Arg Asp His Val Leu Pro Ile 535 540 545 550	2765
gac tat tac ttt cca ccc cag aag acc tgc ctg atc tgt gga gat gaa Asp Tyr Tyr Phe Pro Pro Gln Lys Thr Cys Leu Ile Cys Gly Asp Glu 555 560 565	2813
gct tct ggg tgt cac tat gga gct ctc aca tgt gga agc tgc aag gtc Ala Ser Gly Cys His Tyr Gly Ala Leu Thr Cys Gly Ser Cys Lys Val 570 575 580	2861
ttc ttc aaa aga gcc gct gaa ggg aaa cag aag tac ctg tgc gcc agc Phe Phe Lys Arg Ala Ala Glu Gly Lys Gln Lys Tyr Leu Cys Ala Ser 585 590 595	2909
aga aat gat tgc act att gat aaa ttc cga agg aaa aat tgt cca tct Arg Asn Asp Cys Thr Ile Asp Lys Phe Arg Arg Lys Asn Cys Pro Ser 600 605 610	2957
tgt cgt ctt ccg aaa tgt tat gaa gca ggg atg act ctg gga gcc ccg Cys Arg Leu Arg Lys Cys Tyr Glu Ala Gly Met Thr Leu Gly Ala Arg 615 620 625 630	3005

-continued

aag ctg aag aaa ctt ggt aat ctg aaa cta cag gag gaa gga gag gct Lys Leu Lys Lys Leu Gly Asn Leu Lys Leu Gln Glu Glu Gly Glu Ala 635 640 645	3053
tcc agc acc acc agc ccc act gag gag aca acc cag aag ctg aca gtg Ser Ser Thr Thr Ser Pro Thr Glu Glu Thr Thr Gln Lys Leu Thr Val 650 655 660	3101
tca cac att gaa ggc tat gaa tgt cag ccc atc ttt ctg aat gtc ctg Ser His Ile Glu Gly Tyr Glu Cys Gln Pro Ile Phe Leu Asn Val Leu 665 670 675	3149
gaa gcc att gag cca ggt gta gtg tgt gct gga cac gac aac aac cag Glu Ala Ile Glu Pro Gly Val Val Cys Ala Gly His Asp Asn Asn Gln 680 685 690	3197
ccc gac tcc ttt gca gcc ttg ctc tct agc ctc aat gaa ctg gga gag Pro Asp Ser Phe Ala Ala Leu Leu Ser Ser Leu Asn Glu Leu Gly Glu 695 700 705 710	3245
aga cag ctt gta cac gtg gtc aag tgg gcc aag gcc ttg cct ggc ttc Arg Gln Leu Val His Val Val Lys Trp Ala Lys Ala Leu Pro Gly Phe 715 720 725	3293
cgc aac tta cac gtg gac gac cag atg gct gtc att cag tac tcc tgg Arg Asn Leu His Val Asp Asp Gln Met Ala Val Ile Gln Tyr Ser Trp 730 735 740	3341
atg ggg ctc atg gtg ttt gcc atg ggc tgg cga tcc ttc acc aat gtc Met Gly Leu Met Val Phe Ala Met Gly Trp Arg Ser Phe Thr Asn Val 745 750 755	3389
aac tcc agg atg ctc tac ttc gcc cct gat ctg gtt ttc aat gag tac Asn Ser Arg Met Leu Tyr Phe Ala Pro Asp Leu Val Phe Asn Glu Tyr 760 765 770	3437
cgc atg cac aag tcc cgg atg tac agc cag tgt gtc cga atg agg cac Arg Met His Lys Ser Arg Met Tyr Ser Gln Cys Val Arg Met Arg His 775 780 785 790	3485
ctc tct caa gag ttt gga tgg ctc caa atc acc ccc cag gaa ttc ctg Leu Ser Gln Glu Phe Gly Trp Leu Gln Ile Thr Pro Gln Glu Phe Leu 795 800 805	3533
tgc atg aaa gca ctg cta ctc ttc agc att att cca gtg gat ggg ctg Cys Met Lys Ala Leu Leu Leu Phe Ser Ile Ile Pro Val Asp Gly Leu 810 815 820	3581
aaa aat caa aaa ttc ttt gat gaa ctt cga atg aac tac atc aag gaa Lys Asn Gln Lys Phe Phe Asp Glu Leu Arg Met Asn Tyr Ile Lys Glu 825 830 835	3629
ctc gat cgt atc att gca tgc aaa aga aaa aat ccc aca tcc tgc tca Leu Asp Arg Ile Ile Ala Cys Lys Arg Lys Asn Pro Thr Ser Cys Ser 840 845 850	3677
aga cgc ttc tac cag ctc acc aag ctc ctg gac tcc gtg cag cct att Arg Arg Phe Tyr Gln Leu Thr Lys Leu Leu Asp Ser Val Gln Pro Ile 855 860 865 870	3725
gcg aga gag ctg cat cag ttc act ttt gac ctg cta atc aag tca cac Ala Arg Glu Leu His Gln Phe Thr Phe Asp Leu Leu Ile Lys Ser His 875 880 885	3773
atg gtg agc gtg gac ttt ccg gaa atg atg gca gag atc atc tct gtg Met Val Ser Val Asp Phe Pro Glu Met Met Ala Glu Ile Ile Ser Val 890 895 900	3821
caa gtg ccc aag atc ctt tct ggg aaa gtc aag ccc atc tat ttc cac Gln Val Pro Lys Ile Leu Ser Gly Lys Val Lys Pro Ile Tyr Phe His 905 910 915	3869
acc cag tga agcattggaa accctatttc cccacccag ctcatgcccc Thr Gln 920	3918
ctttcagatg tcttctgcct gttataactc tgcactactc ctctgcagtg ccttggggaa	3978
tttctcttat tgatgtacag tctgtcatga acatgttctt gaattctatt tgctgggctt	4038

-continued

```

tttttttctc tttctctcct ttctttttct tcttccctcc ctatctaacc ctcccatggc 4098
accttcagac tttgcttccc attgtggctc ctatctgtgt tttgaatggt gttgtatgcc 4158
tttaaattcg tgatgatect catatggccc agtgtcaagt tgtgcttggt tacagcacta 4218
ctctgtgccca gccacacaaa cgtttactta tcttatgccca cggaagtgtt agagagctaa 4278
gattatctgg ggaaatcaaa acaaaaacaa gcaaac 4314

```

```

<210> SEQ ID NO 2
<211> LENGTH: 920
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Glu Val Gln Leu Gly Leu Gly Arg Val Tyr Pro Arg Pro Pro Ser
1           5           10          15

Lys Thr Tyr Arg Gly Ala Phe Gln Asn Leu Phe Gln Ser Val Arg Glu
20          25          30

Val Ile Gln Asn Pro Gly Pro Arg His Pro Glu Ala Ala Ser Ala Ala
35          40          45

Pro Pro Gly Ala Ser Leu Leu Leu Leu Gln Gln Gln Gln Gln Gln
50          55          60

Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln
65          70          75          80

Glu Thr Ser Pro Arg Gln Gln Gln Gln Gln Gln Gly Glu Asp Gly Ser
85          90          95

Pro Gln Ala His Arg Arg Gly Pro Thr Gly Tyr Leu Val Leu Asp Glu
100         105         110

Glu Gln Gln Pro Ser Gln Pro Gln Ser Ala Leu Glu Cys His Pro Glu
115         120         125

Arg Gly Cys Val Pro Glu Pro Gly Ala Ala Val Ala Ala Ser Lys Gly
130         135         140

Leu Pro Gln Gln Leu Pro Ala Pro Pro Asp Glu Asp Asp Ser Ala Ala
145         150         155         160

Pro Ser Thr Leu Ser Leu Leu Gly Pro Thr Phe Pro Gly Leu Ser Ser
165         170         175

Cys Ser Ala Asp Leu Lys Asp Ile Leu Ser Glu Ala Ser Thr Met Gln
180         185         190

Leu Leu Gln Gln Gln Gln Gln Glu Ala Val Ser Glu Gly Ser Ser Ser
195         200         205

Gly Arg Ala Arg Glu Ala Ser Gly Ala Pro Thr Ser Ser Lys Asp Asn
210         215         220

Tyr Leu Gly Gly Thr Ser Thr Ile Ser Asp Asn Ala Lys Glu Leu Cys
225         230         235         240

Lys Ala Val Ser Val Ser Met Gly Leu Gly Val Glu Ala Leu Glu His
245         250         255

Leu Ser Pro Gly Glu Gln Leu Arg Gly Asp Cys Met Tyr Ala Pro Leu
260         265         270

Leu Gly Val Pro Pro Ala Val Arg Pro Thr Pro Cys Ala Pro Leu Ala
275         280         285

Glu Cys Lys Gly Ser Leu Leu Asp Asp Ser Ala Gly Lys Ser Thr Glu
290         295         300

Asp Thr Ala Glu Tyr Ser Pro Phe Lys Gly Gly Tyr Thr Lys Gly Leu
305         310         315         320

Glu Gly Glu Ser Leu Gly Cys Ser Gly Ser Ala Ala Ala Gly Ser Ser

```

-continued

325								330					335				
Gly	Thr	Leu	340			Pro	Ser	Thr	Leu	Ser	Leu	Tyr	Lys	Ser	Gly	Ala	
Leu	Asp	Glu	Ala	Ala	Ala	Tyr	Gln	Ser	Arg	Asp	Tyr	Tyr	Asn	Phe	Pro		
Leu	Ala	Leu	Ala	Gly	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	His	Pro	His		
Ala	Arg	Ile	Lys	Leu	Glu	Asn	Pro	Leu	Asp	Tyr	Gly	Ser	Ala	Trp	Ala		
Ala	Ala	Ala	Ala	Gln	Cys	Arg	Tyr	Gly	Asp	Leu	Ala	Ser	Leu	His	Gly		
Ala	Gly	Ala	Ala	Gly	Pro	Gly	Ser	Gly	Ser	Pro	Ser	Ala	Ala	Ala	Ser		
Ser	Ser	Trp	His	Thr	Leu	Phe	Thr	Ala	Glu	Glu	Gly	Gln	Leu	Tyr	Gly		
Pro	Cys	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly		
Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Glu	Ala	Gly	Ala	Val	Ala	Pro		
Tyr	Gly	Tyr	Thr	Arg	Pro	Pro	Gln	Gly	Leu	Ala	Gly	Gln	Glu	Ser	Asp		
Phe	Thr	Ala	Pro	Asp	Val	Trp	Tyr	Pro	Gly	Gly	Met	Val	Ser	Arg	Val		
Pro	Tyr	Pro	Ser	Pro	Thr	Cys	Val	Lys	Ser	Glu	Met	Gly	Pro	Trp	Met		
Asp	Ser	Tyr	Ser	Gly	Pro	Tyr	Gly	Asp	Met	Arg	Leu	Glu	Thr	Ala	Arg		
Asp	His	Val	Leu	Pro	Ile	Asp	Tyr	Tyr	Phe	Pro	Pro	Gln	Lys	Thr	Cys		
Leu	Ile	Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Ala	Leu	Thr		
Cys	Gly	Ser	Cys	Lys	Val	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln		
Lys	Tyr	Leu	Cys	Ala	Ser	Arg	Asn	Asp	Cys	Thr	Ile	Asp	Lys	Phe	Arg		
Arg	Lys	Asn	Cys	Pro	Ser	Cys	Arg	Leu	Arg	Lys	Cys	Tyr	Glu	Ala	Gly		
Met	Thr	Leu	Gly	Ala	Arg	Lys	Leu	Lys	Lys	Leu	Gly	Asn	Leu	Lys	Leu		
Gln	Glu	Glu	Gly	Glu	Ala	Ser	Ser	Thr	Thr	Ser	Pro	Thr	Glu	Glu	Thr		
Thr	Gln	Lys	Leu	Thr	Val	Ser	His	Ile	Glu	Gly	Tyr	Glu	Cys	Gln	Pro		
Ile	Phe	Leu	Asn	Val	Leu	Glu	Ala	Ile	Glu	Pro	Gly	Val	Val	Cys	Ala		
Gly	His	Asp	Asn	Asn	Gln	Pro	Asp	Ser	Phe	Ala	Ala	Leu	Leu	Ser	Ser		
Leu	Asn	Glu	Leu	Gly	Glu	Arg	Gln	Leu	Val	His	Val	Val	Lys	Trp	Ala		
Lys	Ala	Leu	Pro	Gly	Phe	Arg	Asn	Leu	His	Val	Asp	Asp	Gln	Met	Ala		
Val	Ile	Gln	Tyr	Ser	Trp	Met	Gly	Leu	Met	Val	Phe	Ala	Met	Gly	Trp		

-continued

Arg	Ser	Phe	Thr	Asn	Val	Asn	Ser	Arg	Met	Leu	Tyr	Phe	Ala	Pro	Asp
		755					760					765			
Leu	Val	Phe	Asn	Glu	Tyr	Arg	Met	His	Lys	Ser	Arg	Met	Tyr	Ser	Gln
	770					775					780				
Cys	Val	Arg	Met	Arg	His	Leu	Ser	Gln	Glu	Phe	Gly	Trp	Leu	Gln	Ile
785					790					795					800
Thr	Pro	Gln	Glu	Phe	Leu	Cys	Met	Lys	Ala	Leu	Leu	Leu	Phe	Ser	Ile
				805					810					815	
Ile	Pro	Val	Asp	Gly	Leu	Lys	Asn	Gln	Lys	Phe	Phe	Asp	Glu	Leu	Arg
			820					825					830		
Met	Asn	Tyr	Ile	Lys	Glu	Leu	Asp	Arg	Ile	Ile	Ala	Cys	Lys	Arg	Lys
		835					840					845			
Asn	Pro	Thr	Ser	Cys	Ser	Arg	Arg	Phe	Tyr	Gln	Leu	Thr	Lys	Leu	Leu
	850					855					860				
Asp	Ser	Val	Gln	Pro	Ile	Ala	Arg	Glu	Leu	His	Gln	Phe	Thr	Phe	Asp
865					870					875					880
Leu	Leu	Ile	Lys	Ser	His	Met	Val	Ser	Val	Asp	Phe	Pro	Glu	Met	Met
			885						890					895	
Ala	Glu	Ile	Ile	Ser	Val	Gln	Val	Pro	Lys	Ile	Leu	Ser	Gly	Lys	Val
			900					905					910		
Lys	Pro	Ile	Tyr	Phe	His	Thr	Gln								
	915						920								

<210> SEQ ID NO 3
 <211> LENGTH: 2998
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (33)..(2732)

<400> SEQUENCE: 3

gaattcgggtg gaagctacag acaagctcaa gg atg gag gtg cag tta ggg ctg	53
Met Glu Val Gln Leu Gly Leu	
1 5	
gga agg gtc tac cca cgg ccc cca tcc aag acc tat cga gga gcg ttc	101
Gly Arg Val Tyr Pro Arg Pro Pro Ser Lys Thr Tyr Arg Gly Ala Phe	
10 15 20	
cag aat ctg ttc cag agc gtg cgc gaa gcg atc cag aac ccg ggc ccc	149
Gln Asn Leu Phe Gln Ser Val Arg Glu Ala Ile Gln Asn Pro Gly Pro	
25 30 35	
agg cac cct gag gcc gct aac ata gca cct ccc ggc gcc tgt tta cag	197
Arg His Pro Glu Ala Ala Asn Ile Ala Pro Pro Gly Ala Cys Leu Gln	
40 45 50 55	
cag agg cag gag act agc ccc cgg cgg cgg cgg cag cag cac act	245
Gln Arg Gln Glu Thr Ser Pro Arg Arg Arg Arg Gln Gln His Thr	
60 65 70	
gag gat ggt tct cct caa gcc cac atc aga ggc ccc aca ggc tac ctg	293
Glu Asp Gly Ser Pro Gln Ala His Ile Arg Gly Pro Thr Gly Tyr Leu	
75 80 85	
gcc ctg gag gag gaa cag cag cct tca cag cag cag gca gcc tcc gag	341
Ala Leu Glu Glu Glu Gln Gln Pro Ser Gln Gln Gln Ala Ala Ser Glu	
90 95 100	
ggc cac cct gag agc agc tgc ctc ccc gag cct ggg gcg gcc acc gct	389
Gly His Pro Glu Ser Ser Cys Leu Pro Glu Pro Gly Ala Ala Thr Ala	
105 110 115	
cct ggc aag ggg ctg ccg cag cag cca cca gct cct cca gat cag gat	437
Pro Gly Lys Gly Leu Pro Gln Gln Pro Pro Ala Pro Pro Asp Gln Asp	
120 125 130 135	

-continued

gac tca gct gcc cca tcc acg ttg tcc ctg ctg ggc ccc act ttc cca	485
Asp Ser Ala Ala Pro Ser Thr Leu Ser Leu Leu Gly Pro Thr Phe Pro	
140 145 150	
ggc tta agc agc tgc tcc gcc gac att aaa gac att ttg aac gag gcc	533
Gly Leu Ser Ser Cys Ser Ala Asp Ile Lys Asp Ile Leu Asn Glu Ala	
155 160 165	
ggc acc atg caa ctt ctt cag cag cag caa caa cag cag cag cac caa	581
Gly Thr Met Gln Leu Leu Gln Gln Gln Gln Gln Gln Gln His Gln	
170 175 180	
cag cag cac caa cag cac caa cag cag cag gta atc tcc gaa ggc	629
Gln Gln His Gln Gln His Gln Gln Gln Gln Glu Val Ile Ser Glu Gly	
185 190 195	
agc agc gca aga gcc agg gag gcc acg ggg gct ccc tct tcc tcc aag	677
Ser Ser Ala Arg Ala Arg Glu Ala Thr Gly Ala Pro Ser Ser Ser Lys	
200 205 210 215	
gat agt tac cta ggg ggc aat tca acc ata tct gac agt gcc aag gag	725
Asp Ser Tyr Leu Gly Gly Asn Ser Thr Ile Ser Asp Ser Ala Lys Glu	
220 225 230	
ttg tgt aaa gca gtg tct gtg tcc atg gga ttg ggt gtg gaa gca ttg	773
Leu Cys Lys Ala Val Ser Val Ser Met Gly Leu Gly Val Glu Ala Leu	
235 240 245	
gaa cat ctg agt cca ggg gaa cag ctt cgg gga gac tgc atg tac gcg	821
Glu His Leu Ser Pro Gly Glu Gln Leu Arg Gly Asp Cys Met Tyr Ala	
250 255 260	
tcg ctc ctg gga ggt cca ccc gcg gtg cgt ccc act cct tgt gcg ccg	869
Ser Leu Leu Gly Gly Pro Pro Ala Val Arg Pro Thr Pro Cys Ala Pro	
265 270 275	
ctg ccc gaa tgc aaa ggt ctt ccc ctg gac gaa ggc cca ggc aaa agc	917
Leu Pro Glu Cys Lys Gly Leu Pro Leu Asp Glu Gly Pro Gly Lys Ser	
280 285 290 295	
act gaa gag act gct gag tat tcc tct ttc aag gga ggt tac gcc aaa	965
Thr Glu Glu Thr Ala Glu Tyr Ser Ser Phe Lys Gly Gly Tyr Ala Lys	
300 305 310	
gga ttg gaa ggt gag agc ttg ggg tgc tct ggc agc agt gaa gca ggt	1013
Gly Leu Glu Gly Glu Ser Leu Gly Cys Ser Gly Ser Ser Glu Ala Gly	
315 320 325	
agc tct ggg aca ctt gag atc ccg tcc tct ctg tct ctg tat aaa tct	1061
Ser Ser Gly Thr Leu Glu Ile Pro Ser Ser Leu Ser Leu Tyr Lys Ser	
330 335 340	
gga gca cta gac gag gca gca gca tac cag aat cgc gac tac tac aac	1109
Gly Ala Leu Asp Glu Ala Ala Ala Tyr Gln Asn Arg Asp Tyr Tyr Asn	
345 350 355	
ttt ccg ctg gct ctg tcc ggg ccg ccg cac ccc ccg ccc cct acc cat	1157
Phe Pro Leu Ala Leu Ser Gly Pro Pro His Pro Pro Pro Pro Thr His	
360 365 370 375	
cca cac gcc cgt atc aag ctg gag aac cca ttg gac tac ggc agc gcc	1205
Pro His Ala Arg Ile Lys Leu Glu Asn Pro Leu Asp Tyr Gly Ser Ala	
380 385 390	
tgg gct gcg gcg gca gcg caa tgc cgc tat ggg gac ttg ggt agt cta	1253
Trp Ala Ala Ala Ala Ala Gln Cys Arg Tyr Gly Asp Leu Gly Ser Leu	
395 400 405	
cat gga ggg agt gta gcc ggg ccc agc act gga tcg ccc cca gcc acc	1301
His Gly Gly Ser Val Ala Gly Pro Ser Thr Gly Ser Pro Pro Ala Thr	
410 415 420	
acc tct tct tcc tgg cat act ctc ttc aca gct gaa gaa ggc caa tta	1349
Thr Ser Ser Ser Trp His Thr Leu Phe Thr Ala Glu Glu Gly Gln Leu	
425 430 435	
tat ggg cca gga ggc ggg gcc gcc agc agc agc cca agc gat gcc ggg	1397
Tyr Gly Pro Gly Gly Gly Gly Gly Ser Ser Ser Pro Ser Asp Ala Gly	
440 445 450 455	

-continued

cct gta gcc ccc tat ggc tac act cgg ccc cct cag ggg ctg aca agc	1445
Pro Val Ala Pro Tyr Gly Tyr Thr Arg Pro Pro Gln Gly Leu Thr Ser	
460 465 470	
cag gag agt gac tac tct gcc tcc gaa gtg tgg tat cct ggt gga gtt	1493
Gln Glu Ser Asp Tyr Ser Ala Ser Glu Val Trp Tyr Pro Gly Gly Val	
475 480 485	
gtg aac aga gta ccc tat ccc agt ccc aat tgt gtc aaa agt gaa atg	1541
Val Asn Arg Val Pro Tyr Pro Ser Pro Asn Cys Val Lys Ser Glu Met	
490 495 500	
gga cct tgg atg gag aac tac tcc gga cct tat ggg gac atg cgt ttg	1589
Gly Pro Trp Met Glu Asn Tyr Ser Gly Pro Tyr Gly Asp Met Arg Leu	
505 510 515	
gac agt acc agg gac cat gtt tta ccc atc gac tat tac ttt cca ccc	1637
Asp Ser Thr Arg Asp His Val Leu Pro Ile Asp Tyr Tyr Phe Pro Pro	
520 525 530 535	
cag aag acc tgc ctg atc tgt gga gat gaa gct tct ggc tgt cac tac	1685
Gln Lys Thr Cys Leu Ile Cys Gly Asp Glu Ala Ser Gly Cys His Tyr	
540 545 550	
gga gct ctc act tgt ggc agc tgc aag gtc ttc ttc aaa aga gcc gct	1733
Gly Ala Leu Thr Cys Gly Ser Cys Lys Val Phe Phe Lys Arg Ala Ala	
555 560 565	
gaa ggg aaa cag aag tat cta tgt gcc agc aga aac gat tgt acc att	1781
Glu Gly Lys Gln Lys Tyr Leu Cys Ala Ser Arg Asn Asp Cys Thr Ile	
570 575 580	
gat aaa ttt cgg agg aaa aat tgc cca tct tgt cgt ctc cgg aaa tgt	1829
Asp Lys Phe Arg Arg Lys Asn Cys Pro Ser Cys Arg Leu Arg Lys Cys	
585 590 595	
tat gaa gca ggg atg act ctg gga gct cgt aag ctg aag aaa ctt gga	1877
Tyr Glu Ala Gly Met Thr Leu Gly Ala Arg Lys Leu Lys Lys Leu Gly	
600 605 610 615	
aat cta aaa cta cag gag gaa gga gaa aac tcc aat gct ggc agc ccc	1925
Asn Leu Lys Leu Gln Glu Glu Gly Glu Asn Ser Asn Ala Gly Ser Pro	
620 625 630	
act gag gac cca tcc cag aag atg act gta tca cac att gaa ggc tat	1973
Thr Glu Asp Pro Ser Gln Lys Met Thr Val Ser His Ile Glu Gly Tyr	
635 640 645	
gaa tgt cag cct atc ttt ctt aac gtc ctg gaa gcc att gag cca gga	2021
Glu Cys Gln Pro Ile Phe Leu Asn Val Leu Glu Ala Ile Glu Pro Gly	
650 655 660	
gtg gtg tgt gcc gga cat gac aac aac caa cca gat tcc ttt gct gcc	2069
Val Val Cys Ala Gly His Asp Asn Asn Gln Pro Asp Ser Phe Ala Ala	
665 670 675	
ttg tta tct agc ctc aat gag ctt gga gag agg cag ctt gtg cat gtg	2117
Leu Leu Ser Ser Leu Asn Glu Leu Gly Glu Arg Gln Leu Val His Val	
680 685 690 695	
gtc aag tgg gcc aag gcc ttg cct ggc ttc cgc aac ttg cat gtg gat	2165
Val Lys Trp Ala Lys Ala Leu Pro Gly Phe Arg Asn Leu His Val Asp	
700 705 710	
gac cag atg gcg gtc att cag tat tcc tgg atg gga ctg atg gta ttt	2213
Asp Gln Met Ala Val Ile Gln Tyr Ser Trp Met Gly Leu Met Val Phe	
715 720 725	
gcc atg ggt tgg cgg tcc ttc act aat gtc aac tcc agg atg ctc tac	2261
Ala Met Gly Trp Arg Ser Phe Thr Asn Val Asn Ser Arg Met Leu Tyr	
730 735 740	
ttt gca cct gac ttg gtt ttc aat gag tac cgc atg cac aag tct cgg	2309
Phe Ala Pro Asp Leu Val Phe Asn Glu Tyr Arg Met His Lys Ser Arg	
745 750 755	
atg tac agc cag tgt gtg agg atg agg cac ctg tct caa gag ttt gga	2357
Met Tyr Ser Gln Cys Val Arg Met Arg His Leu Ser Gln Glu Phe Gly	
760 765 770 775	

-continued

tgg ctc caa ata acc ccc cag gaa ttc ctg tgc atg aaa gca ctg ctg	2405
Trp Leu Gln Ile Thr Pro Gln Glu Phe Leu Cys Met Lys Ala Leu Leu	
780 785 790	
ctc ttc agc att att cca gtg gat ggg ctg aaa aat caa aaa ttc ttt	2453
Leu Phe Ser Ile Ile Pro Val Asp Gly Leu Lys Asn Gln Lys Phe Phe	
795 800 805	
gat gaa ctt cga atg aac tac atc aag gaa ctc gat cgc atc att gca	2501
Asp Glu Leu Arg Met Asn Tyr Ile Lys Glu Leu Asp Arg Ile Ile Ala	
810 815 820	
tgc aaa aga aag aat ccc aca tcc tgc tca agg cgc ttc tac cag ctc	2549
Cys Lys Arg Lys Asn Pro Thr Ser Cys Ser Arg Arg Phe Tyr Gln Leu	
825 830 835	
acc aag ctc ctg gat tct gtg cag cct att gca aga gag ctg cat cag	2597
Thr Lys Leu Leu Asp Ser Val Gln Pro Ile Ala Arg Glu Leu His Gln	
840 845 850 855	
ttc act ttt gac ctg cta atc aag tcc cat atg gtg agc gtg gac ttt	2645
Phe Thr Phe Asp Leu Leu Ile Lys Ser His Met Val Ser Val Asp Phe	
860 865 870	
cct gaa atg atg gca gag atc atc tct gtg caa gtg ccc aag atc ctt	2693
Pro Glu Met Met Ala Glu Ile Ile Ser Val Gln Val Pro Lys Ile Leu	
875 880 885	
tct ggg aaa gtc aag ccc atc tat ttc cac aca cag tga agatttgga	2742
Ser Gly Lys Val Lys Pro Ile Tyr Phe His Thr Gln	
890 895	
accctaatac ccaaaaccca ccttggtccc tttccagatg tcttctgcct gttatataac	2802
tctgcactac ttctctgcag tgccttgggg gaaattcctc tactgatgta cagtctgtcg	2862
tgaacagggt cctcagttct atttccctggg cttctccttc ttttttttcc ttcttccctc	2922
cctcttttcac cctcccatgg cacattttga atctgtgtgt gattgtgggt ctgcctttgt	2982
tttgatttct gttgta	2998

<210> SEQ ID NO 4

<211> LENGTH: 899

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 4

Met Glu Val Gln Leu Gly Leu Gly Arg Val Tyr Pro Arg Pro Pro Ser	
1 5 10 15	
Lys Thr Tyr Arg Gly Ala Phe Gln Asn Leu Phe Gln Ser Val Arg Glu	
20 25 30	
Ala Ile Gln Asn Pro Gly Pro Arg His Pro Glu Ala Ala Asn Ile Ala	
35 40 45	
Pro Pro Gly Ala Cys Leu Gln Gln Arg Gln Glu Thr Ser Pro Arg Arg	
50 55 60	
Arg Arg Arg Gln Gln His Thr Glu Asp Gly Ser Pro Gln Ala His Ile	
65 70 75 80	
Arg Gly Pro Thr Gly Tyr Leu Ala Leu Glu Glu Glu Gln Gln Pro Ser	
85 90 95	
Gln Gln Gln Ala Ala Ser Glu Gly His Pro Glu Ser Ser Cys Leu Pro	
100 105 110	
Glu Pro Gly Ala Ala Thr Ala Pro Gly Lys Gly Leu Pro Gln Gln Pro	
115 120 125	
Pro Ala Pro Pro Asp Gln Asp Asp Ser Ala Ala Pro Ser Thr Leu Ser	
130 135 140	
Leu Leu Gly Pro Thr Phe Pro Gly Leu Ser Ser Cys Ser Ala Asp Ile	
145 150 155 160	

Lys	Asp	Ile	Leu	Asn	Glu	Ala	Gly	Thr	Met	Gln	Leu	Leu	Gln	Gln	Gln
			165						170			175			
Gln	Gln	Gln	Gln	Gln	His	Gln	Gln	Gln	His	Gln	Gln	His	Gln	Gln	Gln
			180						185			190			
Gln	Glu	Val	Ile	Ser	Glu	Gly	Ser	Ser	Ala	Arg	Ala	Arg	Glu	Ala	Thr
			195			200			205						
Gly	Ala	Pro	Ser	Ser	Ser	Lys	Asp	Ser	Tyr	Leu	Gly	Gly	Asn	Ser	Thr
			210			215			220						
Ile	Ser	Asp	Ser	Ala	Lys	Glu	Leu	Cys	Lys	Ala	Val	Ser	Val	Ser	Met
225			230						235			240			
Gly	Leu	Gly	Val	Glu	Ala	Leu	Glu	His	Leu	Ser	Pro	Gly	Glu	Gln	Leu
			245						250			255			
Arg	Gly	Asp	Cys	Met	Tyr	Ala	Ser	Leu	Leu	Gly	Gly	Pro	Pro	Ala	Val
			260			265						270			
Arg	Pro	Thr	Pro	Cys	Ala	Pro	Leu	Pro	Glu	Cys	Lys	Gly	Leu	Pro	Leu
			275			280						285			
Asp	Glu	Gly	Pro	Gly	Lys	Ser	Thr	Glu	Glu	Thr	Ala	Glu	Tyr	Ser	Ser
			290			295			300						
Phe	Lys	Gly	Gly	Tyr	Ala	Lys	Gly	Leu	Glu	Gly	Glu	Ser	Leu	Gly	Cys
305			310						315			320			
Ser	Gly	Ser	Ser	Glu	Ala	Gly	Ser	Ser	Gly	Thr	Leu	Glu	Ile	Pro	Ser
			325						330			335			
Ser	Leu	Ser	Leu	Tyr	Lys	Ser	Gly	Ala	Leu	Asp	Glu	Ala	Ala	Ala	Tyr
			340			345						350			
Gln	Asn	Arg	Asp	Tyr	Tyr	Asn	Phe	Pro	Leu	Ala	Leu	Ser	Gly	Pro	Pro
			355			360						365			
His	Pro	Pro	Pro	Pro	Thr	His	Pro	His	Ala	Arg	Ile	Lys	Leu	Glu	Asn
			370			375			380						
Pro	Leu	Asp	Tyr	Gly	Ser	Ala	Trp	Ala	Ala	Ala	Ala	Ala	Gln	Cys	Arg
385			390						395			400			
Tyr	Gly	Asp	Leu	Gly	Ser	Leu	His	Gly	Gly	Ser	Val	Ala	Gly	Pro	Ser
			405						410			415			
Thr	Gly	Ser	Pro	Pro	Ala	Thr	Thr	Ser	Ser	Ser	Trp	His	Thr	Leu	Phe
			420			425						430			
Thr	Ala	Glu	Glu	Gly	Gln	Leu	Tyr	Gly	Pro	Gly	Gly	Gly	Gly	Gly	Ser
			435			440			445						
Ser	Ser	Pro	Ser	Asp	Ala	Gly	Pro	Val	Ala	Pro	Tyr	Gly	Tyr	Thr	Arg
			450			455			460						
Pro	Pro	Gln	Gly	Leu	Thr	Ser	Gln	Glu	Ser	Asp	Tyr	Ser	Ala	Ser	Glu
465			470						475			480			
Val	Trp	Tyr	Pro	Gly	Gly	Val	Val	Asn	Arg	Val	Pro	Tyr	Pro	Ser	Pro
			485						490			495			
Asn	Cys	Val	Lys	Ser	Glu	Met	Gly	Pro	Trp	Met	Glu	Asn	Tyr	Ser	Gly
			500			505						510			
Pro	Tyr	Gly	Asp	Met	Arg	Leu	Asp	Ser	Thr	Arg	Asp	His	Val	Leu	Pro
			515			520						525			
Ile	Asp	Tyr	Tyr	Phe	Pro	Pro	Gln	Lys	Thr	Cys	Leu	Ile	Cys	Gly	Asp
			530			535			540						
Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Ala	Leu	Thr	Cys	Gly	Ser	Cys	Lys
545			550						555			560			
Val	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr	Leu	Cys	Ala
			565						570			575			
Ser	Arg	Asn	Asp	Cys	Thr	Ile	As								

-continued

Ser Cys Arg Leu Arg Lys Cys Tyr Glu Ala Gly Met Thr Leu Gly Ala
 595 600 605
 Arg Lys Leu Lys Lys Leu Gly Asn Leu Lys Leu Gln Glu Glu Gly Glu
 610 615 620
 Asn Ser Asn Ala Gly Ser Pro Thr Glu Asp Pro Ser Gln Lys Met Thr
 625 630 635 640
 Val Ser His Ile Glu Gly Tyr Glu Cys Gln Pro Ile Phe Leu Asn Val
 645 650 655
 Leu Glu Ala Ile Glu Pro Gly Val Val Cys Ala Gly His Asp Asn Asn
 660 665 670
 Gln Pro Asp Ser Phe Ala Ala Leu Leu Ser Ser Leu Asn Glu Leu Gly
 675 680 685
 Glu Arg Gln Leu Val His Val Val Lys Trp Ala Lys Ala Leu Pro Gly
 690 695 700
 Phe Arg Asn Leu His Val Asp Asp Gln Met Ala Val Ile Gln Tyr Ser
 705 710 715 720
 Trp Met Gly Leu Met Val Phe Ala Met Gly Trp Arg Ser Phe Thr Asn
 725 730 735
 Val Asn Ser Arg Met Leu Tyr Phe Ala Pro Asp Leu Val Phe Asn Glu
 740 745 750
 Tyr Arg Met His Lys Ser Arg Met Tyr Ser Gln Cys Val Arg Met Arg
 755 760 765
 His Leu Ser Gln Glu Phe Gly Trp Leu Gln Ile Thr Pro Gln Glu Phe
 770 775 780
 Leu Cys Met Lys Ala Leu Leu Leu Phe Ser Ile Ile Pro Val Asp Gly
 785 790 795 800
 Leu Lys Asn Gln Lys Phe Phe Asp Glu Leu Arg Met Asn Tyr Ile Lys
 805 810 815
 Glu Leu Asp Arg Ile Ile Ala Cys Lys Arg Lys Asn Pro Thr Ser Cys
 820 825 830
 Ser Arg Arg Phe Tyr Gln Leu Thr Lys Leu Leu Asp Ser Val Gln Pro
 835 840 845
 Ile Ala Arg Glu Leu His Gln Phe Thr Phe Asp Leu Leu Ile Lys Ser
 850 855 860
 His Met Val Ser Val Asp Phe Pro Glu Met Met Ala Glu Ile Ile Ser
 865 870 875 880
 Val Gln Val Pro Lys Ile Leu Ser Gly Lys Val Lys Pro Ile Tyr Phe
 885 890 895
 His Thr Gln

<210> SEQ ID NO 5
 <211> LENGTH: 4137
 <212> TYPE: DNA
 <213> ORGANISM: Rattus norvegicus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (994)..(3702)

<400> SEQUENCE: 5

atccctagga gccagcctgc tgggagaacc agagggtccg gagcaaacct ggaggctgag	60
agggcatcag aggggaaaag actgagttag ccactccagt gccatacaga agcttaaggg	120
acataccacg ccagccccag cccagcgaca gccaacgcct gttgcagagc ggcggttcg	180
aagccgccgc ccagaagctg ccttttcctc ttcggtgaag tttctaaaag ctgcgggaga	240
ctcggaggaa gcgaagaaag tgtccggtag gactacgact gcctttgtcc tcctccctcc	300

-continued

taccctace cctctgggt cccctctccc tgagcggact aggcaggctt cctggccage	360
cctctccct acaccaccag ctctgccagc cagtttgac agaggttaact ccctttggct	420
gaaagcagac gagctgttg cccattggaa gggaggcttt tgggagccca gagactgagg	480
agcaacagca cgctggagag tcctgtatc caggttctcc cccctgcacc tcctactgcc	540
cgccccac cctgtgtgtg cagctagaat tgaaaagatg aaaagacagt tggggcttca	600
gtagtcgaaa gcaaaacaaa agcaaaaaga aaacaaaag aaaatagccc agttcttatt	660
tgcacctgct tcagtggaca ttgactttgg aaggcagaga atttctctc cccccagtca	720
agctttgagc atcttttaat ctgttcttca agtatttagg gacaaactgt gaaactagca	780
gggcagatcc tgtctagegc gtgccttcc ttacaggaga ctttgaggct atctgggcgc	840
tccccccct cctgcaagt tttcttccct ggagcttccc gcaggtgggc agctagctgc	900
agatactaca tcactagtc gtagaactct tcagagcaag agacgaggag gcaggataag	960
ggaattcggg ggaagctaga gacaagctaa agg atg gag gtg cag tta ggg ctg	1014
Met Glu Val Gln Leu Gly Leu	
1 5	
gga agg gtc tac cca cgg ccc cgg tcc aag acc tat cga gga gcg ttc	1062
Gly Arg Val Tyr Pro Arg Pro Pro Ser Lys Thr Tyr Arg Gly Ala Phe	
10 15 20	
cag aat ctg ttc cag agc gtg cgc gaa gcg atc cag aac ccg ggc ccc	1110
Gln Asn Leu Phe Gln Ser Val Arg Glu Ala Ile Gln Asn Pro Gly Pro	
25 30 35	
agg cac cct gag gcc gct agc ata gca cct ccc ggt gcc tgt tta cag	1158
Arg His Pro Glu Ala Ala Ser Ile Ala Pro Pro Gly Ala Cys Leu Gln	
40 45 50 55	
cag cgg cag gag act agc ccc cgg cgg cgg cgg cgg cag cag cac cct	1206
Gln Arg Gln Glu Thr Ser Pro Arg Arg Arg Arg Arg Gln Gln His Pro	
60 65 70	
gag gat ggc tct cct caa gcc cac atc aga ggc acc aca ggc tac ctg	1254
Glu Asp Gly Ser Pro Gln Ala His Ile Arg Gly Thr Thr Gly Tyr Leu	
75 80 85	
gcc ctg gag gag gaa cag cag cct tca cag cag cag tca gcc tcc gag	1302
Ala Leu Glu Glu Glu Gln Gln Pro Ser Gln Gln Gln Ser Ala Ser Glu	
90 95 100	
ggc cac cct gag agc ggc tgc ctc ccg gag cct gga gct gcc acg gct	1350
Gly His Pro Glu Ser Gly Cys Leu Pro Glu Pro Gly Ala Ala Thr Ala	
105 110 115	
cct ggc aag ggg ctg ccg cag cag cca cca gct cct cca gat cag gat	1398
Pro Gly Lys Gly Leu Pro Gln Gln Pro Pro Ala Pro Pro Asp Gln Asp	
120 125 130 135	
gac tca gct gcc cca tcc acg ttg tcc cta ctg ggc ccc act ttc cca	1446
Asp Ser Ala Ala Pro Ser Thr Leu Ser Leu Leu Gly Pro Thr Phe Pro	
140 145 150	
ggc tta agc agc tgc tcc gca gac att aaa gac atc ctg agc gag gcc	1494
Gly Leu Ser Ser Cys Ser Ala Asp Ile Lys Asp Ile Leu Ser Glu Ala	
155 160 165	
ggc acc atg caa ctt ctt cag cag cag cag caa cag caa cag cag cag	1542
Gly Thr Met Gln Leu Leu Gln Gln Gln Gln Gln Gln Gln Gln Gln	
170 175 180	
cag cag cag cag cag cag cag cag caa cag cag cag gta ata tcc	1590
Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Glu Val Ile Ser	
185 190 195	
gaa ggc agc agc agc gtg aga gca agg gag gcc act ggg gct ccc tct	1638
Glu Gly Ser Ser Ser Val Arg Ala Arg Glu Ala Thr Gly Ala Pro Ser	
200 205 210 215	
tcc tcc aag gat agt tac cta ggg ggc aat tcg acc ata tct gac agt	1686

-continued

Ser	Ser	Lys	Asp	Ser	Tyr	Leu	Gly	Gly	Asn	Ser	Thr	Ile	Ser	Asp	Ser	
				220					225					230		
gcc	aag	gag	ttg	tgt	aaa	gca	gtg	tct	gtg	tcc	atg	ggg	ttg	ggt	gtg	1734
Ala	Lys	Glu	Leu	Cys	Lys	Ala	Val	Ser	Val	Ser	Met	Gly	Leu	Gly	Val	
			235					240					245			
gaa	gca	ctg	gaa	cat	ctg	agt	cca	ggg	gag	cag	ctt	cgg	ggc	gac	tgc	1782
Glu	Ala	Leu	Glu	His	Leu	Ser	Pro	Gly	Glu	Gln	Leu	Arg	Gly	Asp	Cys	
		250					255					260				
atg	tac	gcg	tcg	ctc	ctg	gga	ggt	cca	ccc	gcc	gtg	cgt	ccc	act	cct	1830
Met	Tyr	Ala	Ser	Leu	Leu	Gly	Gly	Pro	Pro	Ala	Val	Arg	Pro	Thr	Pro	
		265				270					275					
tgt	gcg	cct	ctg	gcc	gaa	tgc	aaa	ggt	ctt	tcc	ctg	gac	gaa	ggc	ccg	1878
Cys	Ala	Pro	Leu	Ala	Glu	Cys	Lys	Gly	Leu	Ser	Leu	Asp	Glu	Gly	Pro	
280					285					290					295	
ggc	aaa	ggc	act	gaa	gag	act	gct	gag	tat	tcc	tct	ttc	aag	gga	ggt	1926
Gly	Lys	Gly	Thr	Glu	Glu	Thr	Ala	Glu	Tyr	Ser	Ser	Phe	Lys	Gly	Gly	
				300					305					310		
tac	gcc	aaa	ggg	ttg	gaa	ggt	gag	agt	ctg	ggc	tgc	tct	ggc	agc	agt	1974
Tyr	Ala	Lys	Gly	Leu	Glu	Gly	Glu	Ser	Leu	Gly	Cys	Ser	Gly	Ser	Ser	
			315					320					325			
gaa	gca	ggt	agc	tct	ggg	aca	ctt	gag	atc	ccg	tcc	tca	ctg	tct	ctg	2022
Glu	Ala	Gly	Ser	Ser	Gly	Thr	Leu	Glu	Ile	Pro	Ser	Ser	Leu	Ser	Leu	
		330					335					340				
tat	aag	tct	gga	gca	gta	gac	gag	gca	gca	gca	tac	cag	aat	cgc	gac	2070
Tyr	Lys	Ser	Gly	Ala	Val	Asp	Glu	Ala	Ala	Ala	Tyr	Gln	Asn	Arg	Asp	
	345					350					355					
tac	tac	aac	ttt	ccg	ctc	gct	ctg	tcc	ggg	ccg	ccg	cac	ccc	ccg	ccc	2118
Tyr	Tyr	Asn	Phe	Pro	Leu	Ala	Leu	Ser	Gly	Pro	Pro	His	Pro	Pro	Pro	
360					365					370					375	
cct	acc	cat	cca	cac	gcc	cgc	atc	aag	ctg	gag	aac	ccg	tcg	gac	tac	2166
Pro	Thr	His	Pro	His	Ala	Arg	Ile	Lys	Leu	Glu	Asn	Pro	Ser	Asp	Tyr	
				380				385						390		
ggc	agc	gcc	tgg	gct	gcg	gcg	gca	gcg	caa	tgc	cgc	tat	ggg	gac	ttg	2214
Gly	Ser	Ala	Trp	Ala	Ala	Ala	Ala	Ala	Gln	Cys	Arg	Tyr	Gly	Asp	Leu	
			395					400					405			
gct	agc	cta	cat	gga	ggg	agt	gta	gcc	gga	ccc	agc	act	gga	tcg	ccc	2262
Ala	Ser	Leu	His	Gly	Gly	Ser	Val	Ala	Gly	Pro	Ser	Thr	Gly	Ser	Pro	
		410					415					420				
cca	gcc	acc	gcc	tct	tct	tcc	tgg	cat	act	ctc	ttc	aca	gct	gaa	gaa	2310
Pro	Ala	Thr	Ala	Ser	Ser	Ser	Trp	His	Thr	Leu	Phe	Thr	Ala	Glu	Glu	
		425				430					435					
ggc	caa	tta	tat	ggg	cca	gga	ggc	ggg	ggc	ggc	agc	agt	agc	cca	agc	2358
Gly	Gln	Leu	Tyr	Gly	Pro	Gly	Gly	Gly	Gly	Gly	Ser	Ser	Ser	Pro	Ser	
440					445					450				455		
gat	gct	ggg	cct	gta	gcc	ccc	tat	ggc	tac	act	cgg	ccc	cct	cag	ggg	2406
Asp	Ala	Gly	Pro	Val	Ala	Pro	Tyr	Gly	Tyr	Thr	Arg	Pro	Pro	Gln	Gly	
			460					465						470		
ctg	gca	agc	cag	gag	ggt	gac	ttc	tct	gcc	tct	gaa	gtg	tgg	tat	cct	2454
Leu	Ala	Ser	Gln	Glu	Gly	Asp	Phe	Ser	Ala	Ser	Glu	Val	Trp	Tyr	Pro	
			475				480						485			
ggt	gga	ggt	gtg	aac	aga	gtc	ccc	tat	ccc	agt	ccc	agt	tgt	ggt	aaa	2502
Gly	Gly	Val	Val	Asn	Arg	Val	Pro	Tyr	Pro	Ser	Pro	Ser	Cys	Val	Lys	
		490					495					500				
agt	gaa	atg	gga	cct	tgg	atg	gag	aac	tac	tcc	gga	cct	tat	ggg	gac	2550
Ser	Glu	Met	Gly	Pro	Trp	Met	Glu	Asn	Tyr	Ser	Gly	Pro	Tyr	Gly	Asp	
		505				510					515					
atg	cgt	ttg	gac	agt	acc	agg	gac	cac	gtt	tta	ccc	atc	gac	tat	tac	2598
Met	Arg	Leu	Asp	Ser	Thr	Arg	Asp	His	Val	Leu	Pro	Ile	Asp	Tyr	Tyr	
520					525					530					535	
ttc	cca	ccc	cag	aag	acc	tgc	ctg	atc	tgt	gga	gat	gaa	gct	tct	ggt	2646

-continued

Phe	Pro	Pro	Gln	Lys	Thr	Cys	Leu	Ile	Cys	Gly	Asp	Glu	Ala	Ser	Gly	
				540					545					550		
tgt	cac	tac	gga	gct	ctc	act	tgt	ggc	agc	tgc	aag	gtc	ttc	ttc	aaa	2694
Cys	His	Tyr	Gly	Ala	Leu	Thr	Cys	Gly	Ser	Cys	Lys	Val	Phe	Phe	Lys	
			555					560					565			
aga	gct	gcg	gaa	ggg	aaa	cag	aag	tat	cta	tgt	gcc	agc	aga	aat	gat	2742
Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr	Leu	Cys	Ala	Ser	Arg	Asn	Asp	
			570				575					580				
tgc	acc	att	gat	aaa	ttt	cgg	agg	aaa	aat	tgt	cca	tcg	tgt	cgt	ctc	2790
Cys	Thr	Ile	Asp	Lys	Phe	Arg	Arg	Lys	Asn	Cys	Pro	Ser	Cys	Arg	Leu	
	585					590					595					
cgg	aaa	tgt	tat	gaa	gca	ggg	atg	act	ctg	gga	gct	cgt	aag	ctg	aag	2838
Arg	Lys	Cys	Tyr	Glu	Ala	Gly	Met	Thr	Leu	Gly	Ala	Arg	Lys	Leu	Lys	
	600				605					610					615	
aaa	ctt	gga	aat	ctc	aaa	cta	cag	gaa	gaa	gga	gaa	aac	tcc	agt	gct	2886
Lys	Leu	Gly	Asn	Leu	Lys	Leu	Gln	Glu	Glu	Gly	Glu	Asn	Ser	Ser	Ala	
				620					625					630		
ggg	agc	ccc	act	gag	gac	cca	tcc	cag	aag	atg	act	gta	tca	cac	att	2934
Gly	Ser	Pro	Thr	Glu	Asp	Pro	Ser	Gln	Lys	Met	Thr	Val	Ser	His	Ile	
			635					640					645			
gaa	ggc	tat	gaa	tgt	caa	cct	atc	ttt	ctt	aat	gtc	ctg	gaa	gcc	att	2982
Glu	Gly	Tyr	Glu	Cys	Gln	Pro	Ile	Phe	Leu	Asn	Val	Leu	Glu	Ala	Ile	
		650					655					660				
gag	cca	gga	gtg	gtg	tgt	gcc	gga	cat	gac	aac	aac	cag	cct	gat	tcc	3030
Glu	Pro	Gly	Val	Val	Cys	Ala	Gly	His	Asp	Asn	Asn	Gln	Pro	Asp	Ser	
	665					670					675					
ttt	gct	gcc	ttg	tta	tct	agt	ctc	aac	gag	ctt	ggc	gag	aga	cag	ctt	3078
Phe	Ala	Ala	Leu	Leu	Ser	Ser	Leu	Asn	Glu	Leu	Gly	Glu	Arg	Gln	Leu	
					685					690				695		
gta	cat	gtg	gtc	aag	tgg	gcc	aag	gcc	ttg	cct	ggc	ttc	cgc	aac	ttg	3126
Val	His	Val	Val	Lys	Trp	Ala	Lys	Ala	Leu	Pro	Gly	Phe	Arg	Asn	Leu	
				700					705					710		
cat	gtg	gat	gac	cag	atg	gca	gtc	att	cag	tat	tcc	tgg	atg	gga	ctg	3174
His	Val	Asp	Asp	Gln	Met	Ala	Val	Ile	Gln	Tyr	Ser	Trp	Met	Gly	Leu	
			715					720					725			
atg	gta	ttt	gcc	atg	ggg	tgg	cgg	tcc	ttc	act	aat	gtc	aac	tct	agg	3222
Met	Val	Phe	Ala	Met	Gly	Trp	Arg	Ser	Phe	Thr	Asn	Val	Asn	Ser	Arg	
			730				735					740				
atg	ctc	tac	ttt	gca	cct	gac	ctg	gtt	ttc	aat	gag	tat	cgc	atg	cac	3270
Met	Leu	Tyr	Phe	Ala	Pro	Asp	Leu	Val	Phe	Asn	Glu	Tyr	Arg	Met	His	
			745			750					755					
aag	tct	cga	atg	tac	agc	cag	tgc	gtg	agg	atg	agg	cac	ctt	tct	caa	3318
Lys	Ser	Arg	Met	Tyr	Ser	Gln	Cys	Val	Arg	Met	Arg	His	Leu	Ser	Gln	
	760				765					770				775		
gag	ttt	gga	tgg	ctc	cag	ata	acc	ccc	cag	gaa	ttc	ctg	tgc	atg	aaa	3366
Glu	Phe	Gly	Trp	Leu	Gln	Ile	Thr	Pro	Gln	Glu	Phe	Leu	Cys	Met	Lys	
				780					785					790		
gca	ctg	cta	ctc	ttc	agc	att	att	cca	gtg	gat	ggg	ctg	aaa	aat	caa	3414
Ala	Leu	Leu	Leu	Phe	Ser	Ile	Ile	Pro	Val	Asp	Gly	Leu	Lys	Asn	Gln	
				795				800					805			
aaa	ttc	ttt	gat	gaa	ctt	cga	atg	aac	tac	atc	aag	gaa	ctt	gat	cgc	3462
Lys	Phe	Phe	Asp	Glu	Leu	Arg	Met	Asn	Tyr	Ile	Lys	Glu	Leu	Asp	Arg	
		810					815					820				
atc	att	gca	tgc	aaa	aga	aaa	aat	ccc	aca	tcc	tgc	tca	agg	cgc	ttc	3510
Ile	Ile	Ala	Cys	Lys	Arg	Lys	Asn	Pro	Thr	Ser	Cys	Ser	Arg	Arg	Phe	
		825				830					835					
tac	cag	ctc	acc	aag	ctc	ctg	gat	tct	gtg	cag	cct	att	gca	aga	gag	3558
Tyr	Gln	Leu	Thr	Lys	Leu	Leu	Asp	Ser	Val	Gln	Pro	Ile	Ala	Arg	Glu	
		840			845					850				855		
ctg	cat	caa	ttc	act	ttt	gac	ctg	cta	atc	aag	tcc	cat	atg	gtg	agc	3606

-continued

Leu	His	Gln	Phe	Thr	Phe	Asp	Leu	Leu	Ile	Lys	Ser	His	Met	Val	Ser		
				860					865					870			
gtg	gac	ttt	cct	gaa	atg	atg	gca	gag	atc	atc	tct	gtg	caa	gtg	ccc		3654
Val	Asp	Phe	Pro	Glu	Met	Met	Ala	Glu	Ile	Ile	Ser	Val	Gln	Val	Pro		
			875					880					885				
aag	atc	ctt	tct	ggg	aaa	gtc	aag	ccc	atc	tat	ttc	cac	aca	cag	tga		3702
Lys	Ile	Leu	Ser	Gly	Lys	Val	Lys	Pro	Ile	Tyr	Phe	His	Thr	Gln			
		890					895				900						
agatttgaa	accctaatac	ccaaaccac	cttggtccct	tttcagatgt	cttctgcctg												3762
ttatataact	ctgcactact	tctctggcat	gggccttggg	ggaaattcct	ctactgatgt												3822
acagtctgtc	atgaacatgt	tccccaagtt	ctatttcctg	ggcttttcct	tctttctttt												3882
tcttcttctc	tgccctctttt	accctcccat	ggcacatttt	gaatccgctg	cgtgttggtg												3942
ctctctgcctg	tgttttgagt	ttgtttgtat	ttcttcaagt	ctgtgatgat	cttcttggtg												4002
cccagtgta	actgtgcttg	tttatagcac	tgtgctgtgt	gccaaaccaag	caaatgttta												4062
ctcaccttat	gccatggcaa	gtttagagag	ctataagtat	cttggaaga	aacaaacaga												4122
gagagtaaaa	aaacc																4137

<210> SEQ ID NO 6

<211> LENGTH: 902

<212> TYPE: PRT

<213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 6

Met	Glu	Val	Gln	Leu	Gly	Leu	Gly	Arg	Val	Tyr	Pro	Arg	Pro	Pro	Ser		
1				5					10					15			
Lys	Thr	Tyr	Arg	Gly	Ala	Phe	Gln	Asn	Leu	Phe	Gln	Ser	Val	Arg	Glu		
			20				25					30					
Ala	Ile	Gln	Asn	Pro	Gly	Pro	Arg	His	Pro	Glu	Ala	Ala	Ser	Ile	Ala		
		35				40					45						
Pro	Pro	Gly	Ala	Cys	Leu	Gln	Gln	Arg	Gln	Glu	Thr	Ser	Pro	Arg	Arg		
		50			55					60							
Arg	Arg	Arg	Gln	Gln	His	Pro	Glu	Asp	Gly	Ser	Pro	Gln	Ala	His	Ile		
65				70				75						80			
Arg	Gly	Thr	Thr	Gly	Tyr	Leu	Ala	Leu	Glu	Glu	Gln	Gln	Pro	Ser			
			85					90					95				
Gln	Gln	Gln	Ser	Ala	Ser	Glu	Gly	His	Pro	Glu	Ser	Gly	Cys	Leu	Pro		
			100					105					110				
Glu	Pro	Gly	Ala	Ala	Thr	Ala	Pro	Gly	Lys	Gly	Leu	Pro	Gln	Gln	Pro		
		115				120						125					
Pro	Ala	Pro	Pro	Asp	Gln	Asp	Asp	Ser	Ala	Ala	Pro	Ser	Thr	Leu	Ser		
		130			135						140						
Leu	Leu	Gly	Pro	Thr	Phe	Pro	Gly	Leu	Ser	Ser	Cys	Ser	Ala	Asp	Ile		
145				150					155					160			
Lys	Asp	Ile	Leu	Ser	Glu	Ala	Gly	Thr	Met	Gln	Leu	Leu	Gln	Gln	Gln		
			165					170					175				
Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Gln		
			180				185						190				
Gln	Gln	Gln	Glu	Val	Ile	Ser	Glu	Gly	Ser	Ser	Ser	Val	Arg	Ala	Arg		
		195				200						205					
Glu	Ala	Thr	Gly	Ala	Pro	Ser	Ser	Ser	Lys	Asp	Ser	Tyr	Leu	Gly	Gly		
		210			215					220							
Asn	Ser	Thr	Ile	Ser	Asp	Ser	Ala	Lys	Glu	Leu	Cys	Lys	Ala	Val	Ser		
225				230					235					240			

Val	Ser	Met	Gly	Leu	Gly	Val	Glu	Ala	Leu	Glu	His	Leu	Ser	Pro	Gly
				245					250					255	
Glu	Gln	Leu	Arg	Gly	Asp	Cys	Met	Tyr	Ala	Ser	Leu	Leu	Gly	Gly	Pro
			260					265					270		
Pro	Ala	Val	Arg	Pro	Thr	Pro	Cys	Ala	Pro	Leu	Ala	Glu	Cys	Lys	Gly
		275					280					285			
Leu	Ser	Leu	Asp	Glu	Gly	Pro	Gly	Lys	Gly	Thr	Glu	Glu	Thr	Ala	Glu
	290					295					300				
Tyr	Ser	Ser	Phe	Lys	Gly	Gly	Tyr	Ala	Lys	Gly	Leu	Glu	Gly	Glu	Ser
305					310					315					320
Leu	Gly	Cys	Ser	Gly	Ser	Ser	Glu	Ala	Gly	Ser	Ser	Gly	Thr	Leu	Glu
				325					330					335	
Ile	Pro	Ser	Ser	Leu	Ser	Leu	Tyr	Lys	Ser	Gly	Ala	Val	Asp	Glu	Ala
				340				345					350		
Ala	Ala	Tyr	Gln	Asn	Arg	Asp	Tyr	Tyr	Asn	Phe	Pro	Leu	Ala	Leu	Ser
		355					360					365			
Gly	Pro	Pro	His	Pro	Pro	Pro	Pro	Thr	His	Pro	His	Ala	Arg	Ile	Lys
	370					375					380				
Leu	Glu	Asn	Pro	Ser	Asp	Tyr	Gly	Ser	Ala	Trp	Ala	Ala	Ala	Ala	Ala
385					390					395					400
Gln	Cys	Arg	Tyr	Gly	Asp	Leu	Ala	Ser	Leu	His	Gly	Gly	Ser	Val	Ala
				405					410					415	
Gly	Pro	Ser	Thr	Gly	Ser	Pro	Pro	Ala	Thr	Ala	Ser	Ser	Ser	Trp	His
			420					425					430		
Thr	Leu	Phe	Thr	Ala	Glu	Glu	Gly	Gln	Leu	Tyr	Gly	Pro	Gly	Gly	Gly
		435					440					445			
Gly	Gly	Ser	Ser	Ser	Pro	Ser	Asp	Ala	Gly	Pro	Val	Ala	Pro	Tyr	Gly
	450					455					460				
Tyr	Thr	Arg	Pro	Pro	Gln	Gly	Leu	Ala	Ser	Gln	Glu	Gly	Asp	Phe	Ser
465					470					475					480
Ala	Ser	Glu	Val	Trp	Tyr	Pro	Gly	Gly	Val	Val	Asn	Arg	Val	Pro	Tyr
				485					490					495	
Pro	Ser	Pro	Ser	Cys	Val	Lys	Ser	Glu	Met	Gly	Pro	Trp	Met	Glu	Asn
			500					505					510		
Tyr	Ser	Gly	Pro	Tyr	Gly	Asp	Met	Arg	Leu	Asp	Ser	Thr	Arg	Asp	His
		515					520					525			
Val	Leu	Pro	Ile	Asp	Tyr	Tyr	Phe	Pro	Pro	Gln	Lys	Thr	Cys	Leu	Ile
	530					535					540				
Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Ala	Leu	Thr	Cys	Gly
545					550					555					560
Ser	Cys	Lys	Val	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr
				565					570					575	
Leu	Cys	Ala	Ser	Arg	Asn	Asp	Cys	Thr	Ile	Asp	Lys	Phe	Arg	Arg	Lys
			580					585					590		
Asn	Cys	Pro	Ser	Cys											

-continued

Asp Asn Asn Gln Pro Asp Ser Phe Ala Ala Leu Leu Ser Ser Leu Asn
 675 680 685
 Glu Leu Gly Glu Arg Gln Leu Val His Val Val Lys Trp Ala Lys Ala
 690 695 700
 Leu Pro Gly Phe Arg Asn Leu His Val Asp Asp Gln Met Ala Val Ile
 705 710 715 720
 Gln Tyr Ser Trp Met Gly Leu Met Val Phe Ala Met Gly Trp Arg Ser
 725 730 735
 Phe Thr Asn Val Asn Ser Arg Met Leu Tyr Phe Ala Pro Asp Leu Val
 740 745 750
 Phe Asn Glu Tyr Arg Met His Lys Ser Arg Met Tyr Ser Gln Cys Val
 755 760 765
 Arg Met Arg His Leu Ser Gln Glu Phe Gly Trp Leu Gln Ile Thr Pro
 770 775 780
 Gln Glu Phe Leu Cys Met Lys Ala Leu Leu Leu Phe Ser Ile Ile Pro
 785 790 795 800
 Val Asp Gly Leu Lys Asn Gln Lys Phe Phe Asp Glu Leu Arg Met Asn
 805 810 815
 Tyr Ile Lys Glu Leu Asp Arg Ile Ile Ala Cys Lys Arg Lys Asn Pro
 820 825 830
 Thr Ser Cys Ser Arg Arg Phe Tyr Gln Leu Thr Lys Leu Leu Asp Ser
 835 840 845
 Val Gln Pro Ile Ala Arg Glu Leu His Gln Phe Thr Phe Asp Leu Leu
 850 855 860
 Ile Lys Ser His Met Val Ser Val Asp Phe Pro Glu Met Met Ala Glu
 865 870 875 880
 Ile Ile Ser Val Gln Val Pro Lys Ile Leu Ser Gly Lys Val Lys Pro
 885 890 895
 Ile Tyr Phe His Thr Gln
 900

<210> SEQ ID NO 7
 <211> LENGTH: 10
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic oligonucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (4)..(4)
 <223> OTHER INFORMATION: n is a or g

<400> SEQUENCE: 7

gccnccatgg

10

<210> SEQ ID NO 8
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic peptide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(1)
 <223> OTHER INFORMATION: X can be any amino acid
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (2)..(2)
 <223> OTHER INFORMATION: X is Leu or Met
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (3)..(5)

-continued

<223> OTHER INFORMATION: X can be any amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(8)
<223> OTHER INFORMATION: X can be any amino acid
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: X is Val or Leu

<400> SEQUENCE: 8

Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa
1 5

<210> SEQ ID NO 9
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 9

Leu Leu Leu Phe Ser Ile Ile Pro Val
1 5

<210> SEQ ID NO 10
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 10

Arg Met Leu Tyr Phe Ala Pro Asp Leu Val
1 5 10

<210> SEQ ID NO 11
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 11

Phe Leu Cys Met Lys Ala Leu Leu Leu
1 5

<210> SEQ ID NO 12
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 12

Gln Leu Thr Lys Leu Leu Asp Ser Val
1 5

<210> SEQ ID NO 13
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 13

Val Leu Glu Ala Ile Glu Pro Gly Val
1 5

-continued

<210> SEQ ID NO 14
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 14

Ala Leu Leu Ser Ser Leu Asn Glu Leu
1 5

<210> SEQ ID NO 15
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 15

Leu Gly Glu Arg Gln Leu Val His Val Val
1 5 10

<210> SEQ ID NO 16
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 16

Trp Met Gly Leu Met Val Phe Ala Met
1 5

<210> SEQ ID NO 17
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 17

Phe Ser Ile Ile Pro Val Asp Gly Leu
1 5

<210> SEQ ID NO 18
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 18

Lys Leu Leu Asp Ser Val Gln Pro Ile
1 5

<210> SEQ ID NO 19
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic peptide

<400> SEQUENCE: 19

Gly Ile Leu Gly Phe Val Phe Thr Leu
1 5

63

We claim:

1. A method for inducing an immune reaction to androgen receptor in a mammal having prostate cancer, comprising administering to the mammal an effective amount of a recombinant DNA construct comprising a polynucleotide operatively linked to a transcriptional regulatory element wherein the polynucleotide encodes a member selected from a group consisting of (i) a mammalian androgen receptor, (ii) a fragment of the androgen receptor that comprises a ligand-binding domain, (iii) a fragment of the ligand-binding domain defined by SEQ ID NO:9, (iv) a fragment of the ligand-binding domain defined by SEQ ID NO:10, (v) a fragment of the ligand-binding domain defined by SEQ ID NO:11, and (vi) a fragment of the ligand-binding domain defined by SEQ ID NO:12, whereby the mammal develops immune reaction against androgen receptor.

2. The method of claim 1, wherein the polynucleotide encodes the ligand-binding domain of the androgen receptor.

3. The method of claim 2, wherein the androgen receptor is a human androgen receptor.

4. The method of claim 3, wherein the polynucleotide encodes a fragment of SEQ ID NO:2 wherein the fragment

64

starts from a position ranging from amino acid 651 to amino acid 681 of SEQ ID NO:2 and ends at a position ranging from amino acid 900 to amino acid 920 of SEQ ID NO:2.

5. The method of claim 1, wherein the recombinant DNA construct is administered to the mammal intradermally, intramuscularly, intravascularly, or subcutaneously.

6. The method of claim 1, wherein the mammal is a human.

7. The method of claim 1, wherein cellular immune reaction against cells expressing androgen receptor is induced.

10. 8. The method of claim 7, wherein both humoral and cellular immune reactions against androgen receptor are induced.

15. 9. A method for inducing an immune reaction to androgen receptor in a mammal having prostate cancer, comprising administering to the mammal an effective amount of a recombinant DNA construct comprising a polynucleotide encoding amino acids 664-920 of SEQ ID NO:2 operatively linked to a transcriptional regulatory element whereby the mammal develops immune reaction against androgen receptor.

20. 10. The method of claim 9, wherein the mammal is a human being.

* * * * *