



US008114859B2

(12) **United States Patent**
DeLuca et al.

(10) **Patent No.:** **US 8,114,859 B2**
(45) **Date of Patent:** ***Feb. 14, 2012**

(54) **2-METHYLENE-(20S,25S)-19,27-DINOR-(22E)-VITAMIN D ANALOGS**

5,536,713 A 7/1996 Deluca et al.
5,843,928 A 12/1998 Deluca et al.
5,945,410 A 8/1999 DeLuca et al.
6,566,352 B1 5/2003 DeLuca et al.
6,579,861 B2 6/2003 DeLuca et al.

(Continued)

FOREIGN PATENT DOCUMENTS

WO 02/009427 11/2002

(Continued)

(75) Inventors: **Hector F. DeLuca**, Deerfield, WI (US);
Margaret Clagett-Dame, Deerfield, WI (US);
Lori A. Plum, Arena, WI (US);
Grazia Chiellini, Madison, WI (US);
Pawel Grzywacz, Madison, WI (US)

(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 116 days.

This patent is subject to a terminal disclaimer.

OTHER PUBLICATIONS

Shevde et al, "A Potent Analog of 1α -25-Dihydroxyvitamin D_3 Selectively Induces Bone Formation," PNAS, vol. 99 No. 21, pp. 13487-13491, (2002).

(Continued)

(21) Appl. No.: **11/860,863**

(22) Filed: **Sep. 25, 2007**

(65) **Prior Publication Data**

US 2008/0081799 A1 Apr. 3, 2008

Related U.S. Application Data

(60) Provisional application No. 60/848,805, filed on Sep. 28, 2006.

(51) **Int. Cl.**
A61K 31/593 (2006.01)
C07C 401/00 (2006.01)

(52) **U.S. Cl.** **514/167; 552/653**

(58) **Field of Classification Search** **514/167; 552/653**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,486,333 A * 12/1984 Sebba 516/14
4,666,634 A 5/1987 Miyamoto et al.
5,086,191 A 2/1992 DeLuca et al.


Primary Examiner — Sabiha Qazi

(74) *Attorney, Agent, or Firm* — Andrus, Scales, Starke & Sawall, LLP

(57) **ABSTRACT**

This invention discloses 2-methylene-(20S,25S)-19,27-dinor-(22E)-vitamin D analogs, and specifically 2-methylene-(20S,25S)-19,27-dinor-(22E)- 1α ,25-dihydroxyvitamin D_3 , and pharmaceutical uses therefor. This compound exhibits pronounced activity in arresting the proliferation of undifferentiated cells and inducing their differentiation to the monocyte thus evidencing use as an anti-cancer agent and for the treatment of skin diseases such as psoriasis as well as skin conditions such as wrinkles, slack skin, dry skin and insufficient sebum secretion. This compound also has little, if any, calcemic activity and therefore may be used to treat autoimmune disorders or inflammatory diseases in humans as well as renal osteodystrophy. This compound may also be used for the treatment or prevention of obesity.

79 Claims, 5 Drawing Sheets



U.S. PATENT DOCUMENTS

| | | | | |
|--------------|------|---------|------------------|---------|
| 6,627,622 | B2 | 9/2003 | DeLuca et al. | |
| 6,753,013 | B1 * | 6/2004 | Didriksen et al. | 424/484 |
| 6,887,860 | B2 | 5/2005 | DeLuca et al. | |
| 6,992,074 | B2 | 1/2006 | DeLuca et al. | |
| 7,241,749 | B2 | 7/2007 | DeLuca et al. | |
| 2007/0238702 | A1 | 10/2007 | DeLuca et al. | |

FOREIGN PATENT DOCUMENTS

| | | |
|----|----------------|--------|
| WO | 2005/018648 | 3/2005 |
| WO | 2005/051323 | 6/2005 |
| WO | WO 2007/092720 | 8/2007 |

OTHER PUBLICATIONS

Sicinski et al, "New $1\alpha,25$ -Dihydroxy-19-Norvitamin D_3 Compounds of High Biological Activity: Synthesis and Biological Evaluation of 2-Hydroxymethyl, 2-Methyl, and 2-Methylene Analogues," *Journal of Medical Chemistry*, 41, pp. 4662-4674, (1998).

Yamamoto et al, "Vitamin D Receptor: Ligand Recognition and Allosteric Network," *J. Med. Chem.*, 29, pp. 1313-1324, (2006).

Baggiolini et al., "Stereocontrolled Total Syntheses of $1\alpha,25$ -Dihydroxycholecalciferol and $1\alpha,25$ -Dihydroxyergocalciferol," *Journal of Organic Chemistry*, 51, pp. 3098-3108, (1986).

Lythgoe et al, "Calciferol and its Relatives. Part 22. A Direct Total Synthesis of Vitamin D_2 and Vitamin D_3 ," *J. Chem. Soc. Perkin Trans. 1*, p. 590, (1978).

Lythgoe, "Synthetic Approaches to Vitamin D and its Relatives," *Chem. Soc. Rev.* 9, p. 449, (1983).

Miyamoto et al, "Synthetic Studies of Vitamin D Analogues. XIV. Synthesis and Calcium Regulating Activity of Vitamin D_3 Analogues Bearing a Hydroxyalkoxy Group at the 2β -Position," *Chem. Pharm.*, vol. 41 No. 6, pp. 1111-1113, (1993).

Nishii et al, "The Development of Vitamin D_3 Analogues for the Treatment of Osteoporosis," *Osteoporosis Int.*, Suppl. 1, pp. S190-S193, (1993).

Okano et al, "Regulatory Activities of 2β -(3-Hydroxypropoxy)- $1\alpha,25$ -Dihydroxy-Vitamin D_3 , a Novel Synthetic Vitamin D_3 Derivative, on Calcium Metabolism," *Biochemical and Biophysical Research Communications*, vol. 163 No. 3, pp. 1444-1449, (1989).

Ostrem et al, "24- and 26-homo- $1,25$ -dihydroxyvitamin D_3 : Preferential activity in inducing differentiation of human leukemia cells HL-60 in vitro," *Proc. Natl. Acad. Sci. USA*, vol. 84, pp. 2610-2614, (1987).

Perlman et al, " $1\alpha,25$ -Dihydroxy-19-Nor-Vitamin D_3 . A Novel Vitamin D-Related Compound with Potential Therapeutic Activity," *Tetrahedron Letters*, vol. 31 No. 13, pp. 1823-1824, (1990).

Perlman et al, "Novel Synthesis of 19-Nor-Vitamin D Compounds," *Tetrahedron Letters*, vol. 32 No. 52, pp. 7663-7666, (1991).

Posner et al, "Stereocontrolled Total Synthesis of Calcitriol Derivatives: $1,25$ -Dihydroxy-2-(4'-hydroxybutyl)vitamin D_3 Analogs of an Osteoporosis Drug," *Journal Of Organic Chemistry*, vol. 59 No. 25, pp. 7855-7861, (1994).

Posner et al, "2-Fluoroalkyl A-Ring Analogs of $1,25$ -Dihydroxyvitamin D_3 . Stereocontrolled Total Synthesis via Intramolecular and Intermolecular Diels-Alder Cycloadditions. Preliminary Biological Testing," *Journal of Organic Chemistry*, vol. 60 No. 14, pp. 4617-4628, (1995).

Sardina et al, "Studies on the Synthesis of Side-Chain Hydroxylated Metabolites of Vitamin D. 2. Stereocontrolled Synthesis of 25 -Hydroxyvitamin D_2 ," *J. Org. Chem.*, 51, pp. 1264-1269, (1986).

Toh et al, "Studies on a Convergent Route to Side-Chain Analogues of Vitamin D: 25 -Hydroxy-23-Oxavitamin D_3 ," *J. Org. Chem.*, 48, 1414, (1983).

* cited by examiner






Figure 1

HL-60 Cell Differentiation



EC₅₀: 1,25(OH)₂D₃ = 2×10^{-9} M
SS-22 = 1×10^{-9} M

Figure 2



EC₅₀: 1,25(OH)₂D₃ = 2×10^{-10} M
SS-22 = 1×10^{-10} M

Figure 3




Figure 4




Figure 5

1

**2-METHYLENE-(20S,25S)-19,27-DINOR-(22E)-
VITAMIN D ANALOGS****CROSS REFERENCE TO RELATED
APPLICATIONS**

This application claims priority to and the benefit of U.S. Provisional Application No. 60/848,805, filed Sep. 28, 2006.

BACKGROUND OF THE INVENTION

This invention relates to vitamin D compounds, and more particularly to 2-methylene-(20S,25S)-19,27-dinor-(22E)-vitamin D analogs and their pharmaceutical uses.

The natural hormone, 1 α ,25-dihydroxyvitamin D₃ and its analog in ergosterol series, i.e. 1 α ,25-dihydroxyvitamin D₂ are known to be highly potent regulators of calcium homeostasis in animals and humans, and their activity in cellular differentiation has also been established, Ostrem et al., Proc. Natl. Acad. Sci. USA, 84, 2610 (1987). Many structural analogs of these metabolites have been prepared and tested, including 1 α -hydroxyvitamin D₃, 1 α -hydroxyvitamin D₂, various side chain homologated vitamins and fluorinated analogs. Some of these compounds exhibit an interesting separation of activities in cell differentiation and calcium regulation. This difference in activity may be useful in the treatment of a variety of diseases such as renal osteodystrophy, vitamin D-resistant rickets, osteoporosis, psoriasis, and certain malignancies.

Another class of vitamin D analogs, i.e. the so called 19-nor-vitamin D compounds, is characterized by the replacement of the A-ring exocyclic methylene group (carbon 19), typical of the vitamin D system, by two hydrogen atoms. Biological testing of such 19-nor-analogs (e.g., 1 α ,25-dihydroxy-19-nor-vitamin D₃) revealed a selective activity profile with high potency in inducing cellular differentiation, and very low calcium mobilizing activity. Thus, these compounds are potentially useful as therapeutic agents for the treatment of malignancies, or the treatment of various skin disorders. Two different methods of synthesis of such 19-nor-vitamin D analogs have been described (Perlman et al., Tetrahedron Lett. 31, 1823 (1990); Perlman et al., Tetrahedron Lett. 32, 7663 (1991), and DeLuca et al., U.S. Pat. No. 5,086,191).

In U.S. Pat. No. 4,666,634, 2 β -hydroxy and alkoxy (e.g., ED-71) analogs of 1 α ,25-dihydroxyvitamin D₃ have been described and examined by Chugai group as potential drugs for osteoporosis and as antitumor agents. See also Okano et al., Biochem. Biophys. Res. Commun. 163, 1444 (1989). Other 2-substituted (with hydroxyalkyl, e.g., ED-120, and fluoroalkyl groups) A-ring analogs of 1 α ,25-dihydroxyvitamin D₃ have also been prepared and tested (Miyamoto et al., Chem. Pharm. Bull. 41, 1111 (1993); Nishii et al., Osteoporosis Int. Suppl. 1, 190 (1993); Posner et al., J. Org. Chem. 59, 7855 (1994), and J. Org. Chem. 60, 4617 (1995)).

2-substituted analogs of 1 α ,25-dihydroxy-19-nor-vitamin D₃ have also been synthesized, i.e. compounds substituted at 2-position with hydroxy or alkoxy groups (DeLuca et al., U.S. Pat. No. 5,536,713), with 2-alkyl groups (DeLuca et al U.S. Pat. No. 5,945,410), and with 2-alkylidene groups (DeLuca et al U.S. Pat. No. 5,843,928), which exhibit interesting and selective activity profiles. All these studies indicate that binding sites in vitamin D receptors can accommodate different substituents at C-2 in the synthesized vitamin D analogs.


2

In a continuing effort to explore the 19-nor class of pharmacologically important vitamin D compounds, analogs which are characterized by the presence of a methylene substituent at carbon 2 (C-2), a hydroxyl group at carbon 1 (C-1), and a shortened side chain attached to carbon 20 (C-20) have also been synthesized and tested. 1 α -hydroxy-2-methylene-19-nor-pregnacalciferol is described in U.S. Pat. No. 6,566,352 while 1 α -hydroxy-2-methylene-19-nor-homopregnacalciferol is described in U.S. Pat. No. 6,579,861 and 1 α -hydroxy-2-methylene-19-nor-bishomopregnacalciferol is described in U.S. Pat. No. 6,627,622. All three of these compounds have relatively high binding activity to vitamin D receptors and relatively high cell differentiation activity, but little if any calcemic activity as compared to 1 α ,25-dihydroxyvitamin D₃. Their biological activities make these compounds excellent candidates for a variety of pharmaceutical uses, as set forth in the '352, '861 and '622 patents.

SUMMARY OF THE INVENTION


The present invention is directed toward 2-methylene-(20S,25S)-19,27-dinor-(22E)-vitamin D analogs, and more specifically toward 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃, their biological activity, and various pharmaceutical uses for these compounds. These new vitamin D compounds not known heretofore are the 19-nor-vitamin D analogs having a methylene group at the 2-position (C-2), a methyl group at the 20-position (C-20) in its S-configuration, a double bond between the 22 and 23 positions (C-22 and C-23) in the side chain attached at the 17-position (C-17), and the replacement of the methyl group typically located at the 27 position (C-27) in the side chain with a hydrogen atom.

Structurally these 2-methylene-(20S,25S)-19,27-dinor-(22E)-vitamin D analogs are characterized by the general formula I shown below:



where X₁, X₂ and X₃, which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group. The preferred analog is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ which has the following formula Ia:

3



The above compounds I, particularly Ia, exhibit a desired, and highly advantageous, pattern of biological activity. These compounds are characterized by relatively high binding to vitamin D receptors, but very low intestinal calcium transport activity, as compared to that of $1\alpha,25$ -dihydroxyvitamin D_3 , and have very low ability to mobilize calcium from bone, as compared to $1\alpha,25$ -dihydroxyvitamin D_3 . Hence, these compounds can be characterized as having little, if any, calcemic activity. It is undesirable to raise serum calcium to supra-physiologic levels when suppressing the preproparathyroid hormone gene (Darwish & DeLuca, Arch. Biochem. Biophys. 365, 123-130, 1999) and parathyroid gland proliferation. These analogs having little or no calcemic activity while very active on differentiation are expected to be useful as a therapy for suppression of secondary hyperparathyroidism of renal osteodystrophy.

The compounds I, particularly Ia, of the invention have also been discovered to be especially suited for treatment and prophylaxis of human disorders which are characterized by an imbalance in the immune system, e.g. in autoimmune diseases, including multiple sclerosis, lupus, diabetes mellitus, host versus graft rejection, and rejection of organ transplants; and additionally for the treatment of inflammatory diseases, such as rheumatoid arthritis, asthma, and inflammatory bowel diseases such as celiac disease, ulcerative colitis and Crohn's disease. Acne, alopecia and hypertension are other conditions which may be treated with the compounds of the invention.

The above compounds I, and particularly Ia, are also characterized by relatively high cell differentiation activity. Thus, these compounds also provide a therapeutic agent for the treatment of psoriasis, or as an anti-cancer agent, especially against leukemia, colon cancer, breast cancer, skin cancer and prostate cancer. In addition, due to their relatively high cell differentiation activity, these compounds provide a therapeutic agent for the treatment of various skin conditions including wrinkles, lack of adequate dermal hydration, i.e. dry skin, lack of adequate skin firmness, i.e. slack skin, and insufficient sebum secretion. Use of these compounds thus not only results in moisturizing of skin but also improves the barrier function of skin.

The compounds of the invention of formula I, and particularly formula Ia, are also useful in preventing or treating obesity, inhibiting adipocyte differentiation, inhibiting SCD-1 gene transcription, and/or reducing body fat in animal subjects. Therefore, in some embodiments, a method of preventing or treating obesity, inhibiting adipocyte differentiation, inhibiting SCD-1 gene transcription, and/or reducing

4

body fat in an animal subject includes administering to the animal subject, an effective amount of one or more of the compounds or a pharmaceutical composition that includes one or more of the compounds of formula I. Administration of one or more of the compounds or the pharmaceutical compositions to the subject inhibits adipocyte differentiation, inhibits gene transcription, and/or reduces body fat in the animal subject.

One or more of the compounds may be present in a composition to treat the above-noted diseases and disorders in an amount from about 0.01 $\mu\text{g/gm}$ to about 1000 $\mu\text{g/gm}$ of the composition, preferably from about 0.1 $\mu\text{g/gm}$ to about 500 $\mu\text{g/gm}$ of the composition, and may be administered topically, transdermally, orally, rectally, nasally, sublingually or parenterally in dosages of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$, preferably from about 0.1 $\mu\text{g/day}$ to about 500 $\mu\text{g/day}$.

BRIEF DESCRIPTION OF THE DRAWINGS

In the drawings:

FIGS. 1-5 illustrate various biological activities of 2-methylene-(20S,25S)-19,27-dinor-(22E)- $1\alpha,25$ -dihydroxyvitamin D_3 , hereinafter referred to as "SS-22," as compared to the native hormone $1\alpha,25$ -dihydroxyvitamin D_3 , hereinafter " $1,25(\text{OH})_2D_3$."

FIG. 1 is a graph illustrating the relative activity of SS-22 and $1,25(\text{OH})_2D_3$ to compete for binding with [^3H]- $1,25(\text{OH})_2D_3$ to the full-length recombinant rat vitamin D receptor;

FIG. 2 is a graph illustrating the percent HL-60 cell differentiation as a function of the concentration of SS-22 and $1,25(\text{OH})_2D_3$;

FIG. 3 is a graph illustrating the in vitro transcription activity of $1,25(\text{OH})_2D_3$ as compared to SS-22;


FIG. 4 is a bar graph illustrating the bone calcium mobilization activity of $1,25(\text{OH})_2D_3$ as compared to SS-22; and

FIG. 5 is a bar graph illustrating the intestinal calcium transport activity of $1,25(\text{OH})_2D_3$ as compared to SS-22.

DETAILED DESCRIPTION OF THE INVENTION

2-methylene-(20S,25S)-19,27-dinor-(22E)- $1\alpha,25$ -dihydroxyvitamin D_3 (referred to herein as "SS-22") a 19-nor vitamin D analog which is characterized by the presence of a methylene substituent at the carbon 2 (C-2), a methyl group at the 20-position (C-20) in its S-configuration, a double bond between the 22 and 23 positions (C-22 and C-23) in the side chain attached at the 17-position (C-17), and the replacement of the methyl group typically located at the 27-position (C-27) in the side chain with a hydrogen atom, was synthesized and tested. Such vitamin D analog seemed an interesting target because the relatively small methylene group at the C-2 position should not interfere with binding to the vitamin D receptor. Structurally, this 19-nor analog is characterized by the general formula Ia previously illustrated herein, and its pro-drug (in protected hydroxy form) is characterized by general formula I previously illustrated herein.

The preparation of 2-methylene-(20S,25S)-19,27-dinor-(22E)-vitamin D analogs having the structure I can be accomplished by a common general method, i.e. the condensation of a bicyclic Windaus-Grundmann type ketone II with the allylic phosphine oxide III to the corresponding 2-methylene-19,27-dinor-vitamin D analog IV followed by deprotection at C-1 and C-3 in the latter compound:



In the structures II, III and IV, groups X_1 , X_2 and X_3 are hydroxy-protecting groups, preferably *t*-butyldimethylsilyl, it being also understood that any functionalities that might be sensitive, or that interfere with the condensation reaction, be suitably protected as is well-known in the art. The process shown above represents an application of the convergent synthesis concept, which has been applied effectively for the preparation of vitamin D compounds [e.g. Lythgoe et al., *J. Chem. Soc. Perkin Trans. I*, 590 (1978); Lythgoe, *Chem. Soc. Rev.* 9, 449 (1983); Toh et al., *J. Org. Chem.* 48, 1414 (1983); Baggiolini et al., *J. Org. Chem.* 51, 3098 (1986); Sardina et al., *J. Org. Chem.* 51, 1264 (1986); *J. Org. Chem.* 51, 1269 (1986); DeLuca et al., U.S. Pat. No. 5,086,191; DeLuca et al., U.S. Pat. No. 5,536,713].

The hydrindanone of the general structure II is not known. It can be prepared by the method shown in Schemes 1, 2 and 3 herein (see the preparation of compound SS-22).

For the preparation of the required phosphine oxides of general structure III, a synthetic route has been developed starting from a methyl quinate derivative which is easily obtained from commercial (1*R*,3*R*,4*S*,5*R*)-(-)-quinic acid as described by Perlman et al., *Tetrahedron Lett.* 32, 7663 (1991) and DeLuca et al., U.S. Pat. No. 5,086,191.

The overall process of the synthesis of compounds I and Ia is illustrated and described more completely in U.S. Pat. No. 5,843,928 entitled "2-Alkylidene-19-Nor-Vitamin D Compounds" the specification of which is specifically incorporated herein by reference.

As used in the description and in the claims, the term "hydroxy-protecting group" signifies any group commonly used for the temporary protection of hydroxy functions, such as for example, alkoxy carbonyl, acyl, alkylsilyl or alkylaryl-silyl groups (hereinafter referred to simply as "silyl" groups), and alkoxyalkyl groups. Alkoxy carbonyl protecting groups are alkyl-O-CO- groupings such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, *tert*-butoxycarbonyl, benzyloxycarbonyl or allyloxycarbonyl. The term "acyl" signifies an alkanoyl group of 1 to 6 carbons, in all of its isomeric forms, or a carboxyalkanoyl group of 1 to 6 carbons, such as an oxalyl, malonyl, succinyl, glutaryl group, or an aromatic acyl group such as benzoyl, or a halo, nitro or alkyl substituted benzoyl group. The word "alkyl" as used in the description or of the claims, denotes a straight-chain or branched alkyl radical of 1 to 10 carbons, in all its isomeric forms. Alkoxyalkyl protecting groups are groupings such as methoxymethyl, ethoxymethyl, methoxyethoxymethyl, or tetrahydrofuranyl and tetrahydropyranyl. Preferred silyl-protecting groups are trimethylsilyl, triethylsilyl, *t*-butyldimethylsilyl, dibutylmethylsilyl, diphenylmethylsilyl, phenyldimethylsilyl, diphenyl-*t*-butylsilyl and analogous alkylated silyl radicals. The term "aryl" specifies a phenyl-, or an alkyl-, nitro- or halo-substituted phenyl group.

A "protected hydroxy" group is a hydroxy group derivatised or protected by any of the above groups commonly used for the temporary or permanent protection of hydroxy functions, e.g. the silyl, alkoxyalkyl, acyl or alkoxy carbonyl groups, as previously defined. The terms "hydroxyalkyl", "deuteroalkyl" and "fluoroalkyl" refer to an alkyl radical substituted by one or more hydroxy, deuterium or fluoro groups respectively.

More specifically, reference should be made to the following illustrative example and description as well as to Schemes 1-4 herein for a detailed illustration of the preparation of compound SS-22.

In this example specific products identified by Arabic numerals (1, 2, 3) refer to the specific structures so identified in the Schemes 1, 2, 3 and 4.

EXAMPLE

Chemistry. Ultraviolet (UV) absorption spectra were recorded with a Hitachi Model 60-100 UV-vis spectrometer in the solvent noted. ^1H nuclear magnetic resonance (NMR) spectra were recorded at 500 MHz with a Bruker AM-500 FT spectrometer in deuteriochloroform. Chemical shifts (δ) are reported downfield from internal Me_4Si (δ 0.00). Mass spectra were recorded at 70 eV on a Kratos DS-50 TC instrument equipped with a Kratos MS-55 data system. Samples were introduced into the ion source maintained at 120-250° C. via a direct insertion probe. High-performance liquid chromatography (HPLC) was performed on a Waters Associates liquid chromatograph equipped with a Model 6000A solvent delivery system, a Model 6 UK Universal injector, a Model 486 tunable absorbance detector, and a differential R 401 refractometer.

Example 1

Preparation of (3*S*)-1-*p*-Toluenesulfonyloxy-3-triethylsilyloxy-butane (2)

To a stirred solution of the (S)-(+)-1,3-butanediol 1 (1 g, 11.1 mmol), DMAP (30 mg, 0.25 mmol) and Et_3N (4.6 mL, 3.33 g, 33 mmol) in anhydrous methylene chloride (20 mL)

7

p-toluenesulfonyl chloride (2.54 g, 13.3 mmol) was added at 0° C. The reaction mixture was stirred at 4° C. for 22 h. Methylene chloride was added and the mixture was washed with water, dried (Na₂SO₄) and concentrated under reduced pressure. A residue was chromatographed on silica gel with hexane/ethyl acetate (8:2, then 1:1) to afford the tosylate (2.31 g, 85% yield) as a colorless oil.

To a stirred solution of the tosylate (2.31 g, 9.5 mmol) and 2,6-lutidine (1.2 mL, 1.12 g, 10.5 mmol) in anhydrous methylene chloride (15 mL) triethylsilyl trifluoromethanesulfonate (2.1 mL, 2.51 g, 9.5 mmol) was added at -50° C. The reaction mixture was allowed to warm to room temperature (4 h) and stirring was continued for additional 20 h. Methylene chloride was added and the mixture was washed with water, dried (Na₂SO₄) and concentrated under reduced pressure. A residue was chromatographed on silica gel with hexane/ethyl acetate (97:3) to afford the product 2 (2.71 g, 80% yield) as a colorless oil:

[α]_D+18.0 (c 2.38, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.77 (2H, d, J=8.2 Hz, o-H_{TS}), 7.33 (2H, d, J=8.2 Hz, m-H_{TS}), 4.10 (2H, t, J=6.1 Hz, 1-H₂), 3.90 (1H, m, 3-H), 2.43 (3H, s, Me_{TS}), 1.72 (2H, m, 2-H₂), 1.10 (3H, d, J=6.2 Hz, 4-H₃), 0.88 (9H, t, J=8.0 Hz, 3 \times SiCH₂CH₃), 0.50 (6H, q, J=8.0 Hz, 3 \times SiCH₂CH₃); ¹³C NMR (100 MHz) δ 144.62 (s, p-C_{TS}), 133.03 (s, i-C_{TS}), 129.72 (d, m-C_{TS}), 127.82 (d, o-C_{TS}), 67.78 (t, C-1), 64.46 (d, C-3), 38.47 (t, C-2), 23.82 (q, C-4), 21.52 (q, Me_{TS}), 6.71 (q, SiCH₂CH₃), 4.77 (t, SiCH₂CH₃); MS (EI) m/z 359 (5, MH⁺), 329 (87, M⁺-C₂H₅), 259 (100), 233 (54), 197 (50), 179 (74), 163 (40), 149 (48), 135 (38), 115 (53), 91 (71); exact mass calculated for C₁₅H₂₅O₄SSi (M⁺-C₂H₅) 329.1243, found 329.1239.

Preparation of (3S)-1-Iodo-3-triethylsilyloxy-butane (3)

To a stirred solution of the tosylate 2 (2.71 g, 7.6 mmol) in anhydrous acetone (50 mL) potassium iodide (8 g, 48 mmol) was added and the reaction mixture was refluxed for 10 h. Water (30 mL) was added and the solution was extracted with ethyl acetate. The combined organic phases were dried (Na₂SO₄) and concentrated under reduced pressure. The residue was chromatographed on silica gel with hexane/ethyl acetate (97:3) to give the alcohol 3 (2.26 g, 95% yield) as a colorless oil:

[α]_D+36.3 (c 2.12, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.89 (1H, m, 3-H), 3.22 (2H, t, J=7.0 Hz, 1-H₂), 1.91 (2H, m, 2-H₂), 1.16 (3H, d, J=6.1 Hz, 4-H₃), 0.96 (9H, t, J=7.9 Hz, 3 \times SiCH₂CH₃), 0.61 (6H, q, J=7.9 Hz, 3 \times SiCH₂CH₃); ¹³C NMR (100 MHz) δ 68.13 (d, C-3), 43.23 (d, C-2), 23.45 (q, C-4), 6.86 (q, SiCH₂CH₃), 4.99 (t, SiCH₂CH₃), 3.34 (t, C-1); MS (EI) m/z 314 (1, M⁺), 299 (1, M⁺-CH₃), 285 (100, M⁺-C₂H₅), 257 (97, M⁺-C₄H₉), 228 (51), 212 (98), 184 (58), 157 (62), 129 (33), 115 (31); exact mass calculated for C₈H₁₈OISi (M⁺-C₂H₅) 285.0172, found 285.0169.

Preparation of (3S)-Hydroxybutyl-triphenylphosphonium iodide (4)

To a stirred solution of the iodide 3 (1.67 g, 5.3 mmol) in acetonitrile (50 mL) triphenylphosphine (4.2 g, 16 mmol) was added and the reaction mixture was refluxed for 2 days. Acetonitrile was evaporated under reduced pressure, ethyl acetate (50 mL) was added and the mixture was stirred at room temperature for 4 h. After removal of the solvent by filtration the solid was washed with ethyl acetate, filtered off and dried. The pure phosphonium salt 4 (1.77 g, 87% yield) was obtained as white crystals:

8

¹H NMR (400 MHz, CD₃OD) δ 8.00-7.70 (15H, m, H_{Ph}), 3.89 (1H, m, 3-H), 3.48 (2H, m, 1-H₂), 1.73 (2H, m, 2-H₂), 1.19 (3H, d, J=6.2 Hz, 4-H₃); ¹³C NMR (100 MHz) δ 136.42 (d, p-C_{Ph}), 134.99 (d, J_{C-P}=10.1 Hz, m-C_{Ph}), 131.71 (d, J_{C-P}=13.1 Hz, o-C_{Ph}), 120.04 (s, J_{C-P}=86.5 Hz, i-C_{Ph}), 67.94 (d, J_{C-P}=16.2 Hz, C-3), 32.52 (t, J_{C-P}=4.1 Hz, C-2), 23.38 (q, C-4), 19.84 (t, J_{C-P}=53.7 Hz, C-1);

Preparation of (8S,20S)-de-A,B-20-(hydroxymethyl)pregnan-8-ol (5)

Ozone was passed through a solution of vitamin D₂ (3 g, 7.6 mmol) in methanol (250 mL) and pyridine (2.44 g, 2.5 mL, 31 mmol) for 50 min at -78° C. The reaction mixture was then flushed with an oxygen for 15 min to remove the residual ozone and the solution was treated with NaBH₄ (0.75 g, 20 mmol). After 20 min the second portion of NaBH₄ (0.75 g, 20 mmol) was added and the mixture was allowed to warm to room temperature. The third portion of NaBH₄ (0.75 g, 20 mmol) was then added and the reaction mixture was stirred for 18 h. The reaction was quenched with water (40 mL) and the solution was concentrated under reduced pressure. The residue was extracted with ethyl acetate and the combined organic phases were washed with 1M aq. HCl, saturated aq. NaHCO₃, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was chromatographed on silica gel with hexane/ethyl acetate (75:25) to give the diol 5 (1.21 g, 75% yield) as white crystals:

m.p. 106-108° C.; [α]_D+30.2° (c 1.46, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 4.08 (1H, d, J=2.0 Hz, 8 α -H), 3.63 (1H, dd, J=10.5, 3.1 Hz, 22-H), 3.38 (1H, dd, J=10.5, 6.8 Hz, 22-H), 1.99 (1H, br.d, J=13.2 Hz), 1.03 (3H, d, J=6.6 Hz, 21-H₃), 0.956 (3H, s, 18-H₃); ¹³C NMR (100 MHz) δ 69.16 (d, C-8), 67.74 (t, C-22), 52.90 (d), 52.33 (d), 41.83 (s, C-13), 40.19 (t), 38.20 (d), 33.53 (t), 26.62 (t), 22.54 (t), 17.36 (t), 16.59 (q, C-21), 13.54 (q, C-18); MS (EI) m/z 212 (2, M⁺), 194 (34, M⁺-H₂O), 179 (33, M⁺-H₂O-CH₃), 163 (18, M⁺-CH₂OH-H₂O), 135 (36), 125 (54), 111 (100), 95 (63), 81 (67); exact mass calculated for C₁₃H₂₂O (M⁺-H₂O) 194.1671, found 194.1665.

Preparation of (8S,20S)-de-A,B-8-benzoyloxy-20-(hydroxymethyl)pregnane (6)

Benzoyl chloride (2.4 g, 2 mL, 17 mmol) was added to a solution of the diol 5 (1.2 g, 5.7 mmol) and DMAP (30 mg, 0.2 mmol) in anhydrous pyridine (20 mL) at 0° C. The reaction mixture was stirred at 4° C. for 24 h, diluted with methylene chloride (100 mL), washed with 5% aq. HCl, water, saturated aq. NaHCO₃, dried (Na₂SO₄) and concentrated under reduced pressure. The residue (3.39 g) was treated with a solution of KOH (1 g, 15.5 mmol) in anhydrous ethanol (30 mL) at room temperature. After stirring of the reaction mixture for 3 h, ice and 5% aq. HCl were added until pH=6. The solution was extracted with ethyl acetate (3 \times 50 mL) and the combined organic phases were washed with saturated aq. NaHCO₃, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was chromatographed on silica gel with hexane/ethyl acetate (75:25) to give the alcohol 6 (1.67 g, 93% yield) as a colorless oil:

[α]_D+56.0 (c 0.48, CHCl₃); ¹H NMR (400 MHz, CDCl₃+TMS) δ 8.08-8.02 (2H, m, o-H_{Bz}), 7.59-7.53 (1H, m, p-H_{Bz}), 7.50-7.40 (2H, m, m-H_{Bz}), 5.42 (1H, d, J=2.4 Hz, 8 α -H), 3.65 (1H, dd, J=10.5, 3.2 Hz, 22-H), 3.39 (1H, dd, J=10.5, 6.8 Hz, 22-H), 1.08 (3H, d, J=5.3 Hz, 21-H₃), 1.07 (3H, s, 18-H₃); ¹³C NMR (125 MHz) δ 166.70 (s, C=O), 132.93 (d, p-C_{Bz}), 131.04 (s, i-C_{Bz}), 129.75 (d, o-C_{Bz}), 128.57 (d, m-C_{Bz}), 72.27

9

(d, C-8), 67.95 (t, C-22), 52.96 (d), 51.60 (d), 42.15 (s, C-13), 39.98 (t), 38.61 (d), 30.73 (t), 26.81 (t), 22.91 (t), 18.20 (t), 16.87 (q, C-21), 13.81 (q, C-18); MS (EI) m/z 316 (5, M^+), 301 (3, M^+ -Me), 299 (1, M^+ -OH), 298 (2, M^+ -H₂O), 285 (10, M^+ -CH₂OH), 257 (6), 230 (9), 194 (80), 135 (84), 105 (100); exact mass calculated for C₂₀H₂₈O₃ 316.2038, found 316.2019.

Preparation of
(8S,20S)-de-A,B-8-benzoyloxy-20-formylpregnane
(7)

Sulfur trioxide pyridine complex (1.94 g, 12.2 mmol) was added to a solution of the alcohol 6 (640 mg, 2.03 mmol), triethylamine (1.41 mL, 1.02 g, 10.1 mmol) in anhydrous methylene chloride (10 mL) and anhydrous DMSO (2 mL) at 0° C. The reaction mixture was stirred under argon at 0° C. for 1 h and then concentrated. The residue was diluted with ethyl acetate, washed with brine, dried (Na₂SO₄) and concentrated. The residue was purified by column chromatography on silica gel with hexane/ethyl acetate (95:5) to give the aldehyde 7 (529 mg, 83% yield) as an oil:

¹H NMR (400 MHz, CDCl₃+TMS) δ 9.60 (1H, d, J=3.1 Hz, CHO), 8.05 (2H, m, o-H_{Bz}), 7.57 (1H, m, p-H_{Bz}), 7.45 (2H, m, m-H_{Bz}), 5.44 (1H, s, 8α-H), 2.39 (1H, m, 20-H), 2.03 (2H, dm, J=11.5 Hz), 1.15 (3H, d, J=6.9 Hz, 21-H₃), 1.10 (3H, s, 18-H₃); ¹³C NMR (100 MHz) δ 204.78 (d, CHO), 132.78 (d, p-Bz), 130.69 (s, i-Bz), 129.50 (d, o-Bz), 128.38, (d, m-Bz), 71.66 (d, C-8), 51.30 (d), 50.95 (d), 49.20 (d), 42.38 (s, C-13), 39.62 (t), 30.47 (t), 25.99 (t), 22.92 (t), 17.92 (t), 13.90 (q), 13.35 (q); MS (EI) m/z 314 (1, M^+), 299 (0.5, M^+ -Me), 286 (1, M^+ -CO), 285 (5, M^+ -CHO), 257 (1, M^+ -C₃H₅O), 209 (10, M^+ -PhCO), 192 (38), 134 (60), 105 (100), 77 (50); exact mass calculated for C₂₀H₂₆O₃ 314.1882, found 314.1887.

Preparation of (8S,20R)-de-A,B-8-benzoyloxy-20-
(hydroxymethyl)pregnane (8)

The aldehyde 7 (1.36 g, 4.3 mmol) was dissolved in CH₂Cl₂ (15 mL) and a 40% aq. N-Bu₄NOH solution (5.6 mL, 5.57 g, 8.6 mmol) was added. The resulting mixture was stirred at room temperature for 16 h, diluted with methylene chloride (30 mL), washed with water, dried (Na₂SO₄) and concentrated under reduced pressure. A residue was chromatographed on silica gel with hexane/ethyl acetate (95:5) to afford a mixture of aldehyde 7 and its epimer (730 mg, 53% yield) in ca. 1:1.7 ratio (by ¹H NMR).

This mixture of aldehydes (730 mg, 2.3 mmol) was dissolved in THF (5 mL) and NaBH₄ (175 mg, 4.6 mmol) was added, followed by a drop wise addition of ethanol (5 mL). The reaction mixture was stirred at room temperature for 30 min and it was quenched with saturated aq. NH₄Cl solution. The mixture was extracted with ether (3×30 mL) and the combined organic phase was washed with water, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was chromatographed on silica gel with hexane/ethyl acetate (95:5 80:20) to give the desired, pure (20R)-alcohol 8 (366 mg, 52% yield) as an oil and a mixture of 8 and its 20-epimer 6 (325 mg, 45% yield) in ca. 1:4 ratio (by ¹H NMR).

8: [α]_D+43.0 (c 0.54, CHCl₃); ¹H NMR (500 MHz, CDCl₃+TMS) δ 8.10-8.00 (2H, m, o-H_{Bz}), 7.60-7.53 (1H, m, p-H_{Bz}), 7.48-7.41 (2H, m, m-H_{Bz}), 5.42 (1H, br s, 8α-H), 3.75 (1H, dd, J=10.6, 3.5 Hz, 22-H), 3.48 (1H, dd, J=10.6, 7.0 Hz, 22-H), 1.069 (3H, s, 18-H₃), 0.973 (3H, d, J=6.7 Hz, 21-H₃); ¹³C NMR (125 MHz) δ 166.70 (s, C=O), 132.94 (d, p-C_{Bz}),

10

131.05 (s, i-C_{Bz}), 129.76 (d, o-C_{Bz}), 128.59 (d, m-C_{Bz}), 72.28 (d, C-8), 66.95 (t, C-22), 52.94 (d), 51.77 (d), 41.96 (s, C-13), 39.56 (t), 37.78 (d), 30.75 (t), 26.67 (t), 22.71 (t), 18.25 (t), 16.76 (q, C-21), 14.14 (q, C-18); MS (EI) m/z 316 (16, M^+), 301 (5, M^+ -Me), 299 (2, M^+ -OH), 298 (3, M^+ -H₂O), 285 (9, M^+ -CH₂OH), 257 (5), 242 (11), 230 (8), 194 (60), 147 (71), 105 (100); exact mass calculated for C₂₀H₂₈O₃ 316.2038, found 316.2050.

Preparation of
(8S,20R)-de-A,B-8-benzoyloxy-20-formylpregnane
(9)

Sulfur trioxide pyridine complex (1.94 g, 12.2 mmol) was added to a solution of the alcohol 8 (640 mg, 2.03 mmol), triethylamine (1.41 mL, 1.02 g, 10.1 mmol) in anhydrous methylene chloride (10 mL) and anhydrous DMSO (2 mL) at 0° C. The reaction mixture was stirred under argon at 0° C. for 1 h and then concentrated. The residue was diluted with ethyl acetate, washed with brine, dried (Na₂SO₄) and concentrated. The residue was purified by column chromatography on silica gel with hexane/ethyl acetate (95:5) to give the aldehyde 9 (529 mg, 83% yield) as an oil:

¹H NMR (400 MHz, CDCl₃+TMS) δ 9.56 (1H, d, J=3.1 Hz, CHO), 8.03 (2H, m, o-H_{Bz}), 7.56 (1H, m, p-H_{Bz}), 7.45 (2H, m, m-H_{Bz}), 5.44 (1H, s, 8α-H), 2.36 (1H, m, 20-H), 1.06-1.04 (6H, m, 21-H₃ and 18-H₃); ¹³C NMR (100 MHz) δ 205.56, 132.79, 130.68, 129.52, 128.39, 71.59, 52.18, 51.17, 48.48, 41.64, 38.63, 30.49, 25.37, 22.16, 17.77, 14.58, 13.46; exact mass calculated for C₂₀H₂₆O₃ 314.1882, found 314.1887.

Preparation of (8S,20S)-de-A,B-8-benzoyloxy-20-
[(4S)-hydroxy-pent-(1E)-en-yl]pregnane (10)

To a stirred suspension of the phosphonium salt 4 (310 mg, 0.67 mmol) in anhydrous THF (5 mL) butyllithium (1.6 M, 840 μL, 1.34 mmol) was added at -20° C. The solution turned deep orange. After 1 h a precooled (-20° C.) solution of the aldehyde 9 (70 mg, 0.22 mmol) in anhydrous THF (2 mL) was added and the reaction mixture was stirred at -20° C. for 3 h and at room temperature for 18 h. The reaction was quenched with water and the mixture was extracted with ethyl acetate. Combined organic phases were washed with brine, dried (Na₂SO₄) and evaporated. The residue was chromatographed on silica gel with hexane/ethyl acetate (95:5) to give the product 10 (42 mg, 52% yield):

¹H NMR (400 MHz, CDCl₃+TMS) δ 8.05 (2H, m, o-H_{Bz}), 7.55 (1H, m, p-H_{Bz}), 7.44 (2H, m, m-H_{Bz}), 5.46-5.25 (3H, m, 8α-H, 22-H and 23-H), 3.82 (1H, m, 25-H), 1.20 (3H, d, J=6.2 Hz, 27-H₃), 1.03 (3H, s, 18-H₃), 0.94 (3H, d, J=6.7 Hz, 21-H₃); ¹³C NMR (100 MHz) δ 166.45 (s, C=O), 141.15 (d, C-22), 132.67 (d, p-C_{Bz}), 130.86 (s, i-C_{Bz}), 129.53 (d, o-C_{Bz}), 128.32 (d, m-C_{Bz}), 123.40 (d, C-23), 72.09 (d, C-8), 67.21 (d, C-25), 56.33 (d), 51.46 (d), 42.56 (t), 41.95 (s, C-13), 40.16 (d), 39.36 (t), 30.59 (t), 26.81 (t), 22.73 (t), 22.49 (q, C-27), 21.57 (q, C-21), 17.83 (t), 13.85 (q, C-18); exact mass calculated for C₂₄H₃₄O₃ (M⁺) 370.2508, found 370.2498.

Preparation of (8S,20S)-de-A,B-8-benzoyloxy-20-
[(4S)-tert-butylidimethylsilyloxy-pent-(1E)-enyl]
pregnane (11)

tert-Butyldimethylsilyl trifluoromethanesulfonate (33 μL, 37 mg, 0.14 mmol) was added to a solution of the alcohol 10 (30 mg, 0.08 mmol) and 2,6-lutidine (33 μL, 30 mg, 0.28 mmol) in anhydrous methylene chloride (3 mL) at -20° C.

11

The mixture was stirred under argon at 0° C. for 1 h. The reaction was quenched with water and extracted with methylene chloride. The combined organic phases were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was chromatographed on silica gel with hexane and hexane/ethyl acetate (97:3) to give the product 11 (35 mg, 89%); ¹H NMR (600 MHz, CDCl₃+TMS) δ 8.05 (2H, m, o-H_{Bz}), 7.56 (1H, m, p-H_{Bz}), 7.45 (2H, m, m-H_{Bz}), 5.45 (1H, s, 8α-H), 5.33-5.24 (2H, m, 22-H and 23-H), 3.80 (1H, m, 25-H), 1.18 (3H, d, J=6.0 Hz, 27-H₃), 1.05 (3H, s, 18-H₃), 0.95 (3H, d, J=6.6 Hz, 21-H₃), 0.88 (9H, s, Si-t-Bu), 0.051 (6H, s, SiMe₂); ¹³C NMR (600 MHz) δ 166.47, 139.55, 132.85, 131.11, 129.73, 128.52, 124.42 (d, C-23), 72.40 (d, C-8), 69.17 (d, C-25), 56.67, 51.26, 43.25, 42.15, 40.22, 39.40, 30.77, 26.81, 26.12, 23.45, 22.85, 19.42, 18.26, 18.04, 13.96, -4.32, -4.45.

Preparation of (8S,20S)-de-A,B-20-[(4S)-tert-butylidimethylsilyloxy-pent-(1E)-en-yl]pregnan-8-ol (12)

A solution of sodium hydroxide in ethanol (2.5M, 2 mL) was added to a stirred solution of the benzoate 11 (30 mg, 62 μmol) in anhydrous ethanol (10 mL) and the reaction mixture was refluxed for 18 h. The mixture was cooled to room temperature, neutralized with 5% aq. HCl and extracted with dichloromethane. Combined organic phases were washed with saturated aq. NaHCO₃, dried (Na₂SO₄) and evaporated. The residue was chromatographed on silica gel with hexane/ethyl acetate (95:5) to give the alcohol 12 (17 mg, 72% yield): ¹H NMR (400 MHz, CDCl₃+TMS) δ 5.36-5.23 (2H, m, 22-H and 23-H), 4.09 (1H, d, J=1.9 Hz, 8α-H), 3.78 (1H, m, 25-H), 1.15 (3H, d, J=6.1 Hz, 27-H₃), 1.02 (3H, s, 18-H₃), 0.93 (3H, d, J=6.6 Hz, C-21), 0.88 (9H, s, Si-t-Bu), 0.051 (6H, s, SiMe₂); ¹³C NMR (100 MHz) δ 139.19, 124.24, 69.45, 69.15, 56.57, 52.66, 42.95, 41.84, 40.28, 39.63, 33.54, 27.61, 25.90, 23.21, 22.47, 20.35, 18.19, 17.43, 13.68, -4.36, -4.58.

Preparation of (20S)-de-A,B-20-[(4S)-tert-butylidimethylsilyloxy-pent-(1E)-enyl]pregnan-8-one (13)

Pyridinium dichromate (77 mg, 207 μmol) was added to a solution of the alcohol 12 (15 mg, 39 μmol) and pyridinium p-toluenesulfonate (3 mg, 12 μmol) in anhydrous methylene chloride (5 mL). The resulting suspension was stirred at room temperature for 3 h. The reaction mixture was filtered through a Waters silica Sep-Pak cartridge (5 g) that was further washed with hexane/ethyl acetate (8:2). After removal of solvents the ketone 13 (10 mg, 67% yield) was obtained as a colorless oil:

¹H NMR (400 MHz, CDCl₃+TMS) δ 5.35-5.22 (2H, m, 22-H and 23-H), 3.76 (1H, m, 25-H), 2.41 (1H, dd, J=11.5, 7.6 Hz), 1.15 (3H, d, J=6.1 Hz, 27-H₃), 1.01 (3H, s, 18-H₃), 0.92 (3H, d, J=6.6 Hz), 0.89 (9H, s, Si-t-Bu), 0.052 (6H, s, SiMe₂); ¹³C NMR (100 MHz) δ 211.97, 139.10, 124.44, 68.72, 62.32, 56.46, 51.76, 42.89, 41.15, 40.19, 33.85, 27.73, 25.88, 24.05, 23.60, 20.61, 19.03, 18.27, 12.68, -4.54, -4.69; exact mass calculated for C₂₃H₄₂O₂Si Na (MNa)⁺ 401.2852, found 401.2848.

Preparation of (20S,25S)-2-Methylene-19,27-dinor-22-(E)-ene-1α,25-dihydroxyvitamin D₃ (15)

To a solution of phosphine oxide 12 (52 mg, 89 μmol) in anhydrous THF (400 μL) at -20° C. was slowly added PhLi (1.8 M in di-n-butylether, 61 μL, 110 μmol) under argon with stirring. The solution turned deep orange. After 30 min the

12

mixture was cooled to -78° C. and a precooled (-78° C.) solution of ketone 13 (9 mg, 24 μmol) in anhydrous THF (200+100 μL) was slowly added. The mixture was stirred under argon at -78° C. for 3 h and at 0° C. for 18 h. Ethyl acetate was added, and the organic phase was washed with brine, dried (Na₂SO₄) and evaporated. The residue was dissolved in hexane and applied on a Waters silica Sep-Pak cartridge (2 g). The cartridge was washed with hexane and hexane/ethyl acetate (99.5:0.5) to give 19-norvitamin derivative 14 (10 mg, 46% yield). Then the Sep-Pak was washed with ethyl acetate to recover diphenylphosphine oxide 12 (22 mg). For analytical purpose a sample of the protected vitamin 14 was further purified by HPLC (9.4×250 mm Zorbax Sil column, 4 mL/min, hexane/2-propanol (99.9:0.1) solvent system, R_f=3.80 min):


UV (in hexane) λ_{max} 263.1, 253.2, 244.3 nm; ¹H NMR (500 MHz, CDCl₃) δ 6.22 and 5.83 (each 1H, each d, J=11.2 Hz, 6- and 7-H), 5.38-5.27 (2H, m, 22-H and 23-H), 4.96 and 4.90 (each 1H, each s, =CH₂), 4.43 (2H, m, 1β- and 3α-H), 3.78 (1H, m, 25-H), 2.85 (1H, dm, J=11.8 Hz, 9β-H), 2.52 (1H, dd, J=13.1, 5.9 Hz, 10α-H), 2.47 (1H, dd, J=12.6, 4.3 Hz, 4α-H), 2.33 (1H, dd, J=13.1, 2.3 Hz, 10β-H), 1.11 (3H, d, J=6.0 Hz, 27-H₃), 1.02 (3H, d, J=6.4 Hz, 21-H₃), 0.898 (9H, s, Si-t-Bu), 0.895 (9H, s, Si-t-Bu), 0.867 (9H, s, Si-t-Bu), 0.52 (3H, s, 18-H₃), 0.082 (3H, s, SiMe), 0.067 (3H, s, SiMe), 0.055 (9H, s, 3×SiMe), 0.027 (3H, s, SiMe); ¹³C NMR (100 MHz) δ 152.96, 141.13, 138.98, 132.76, 124.74, 122.40, 116.09, 106.25, 72.54, 71.63, 68.73, 56.63, 56.29, 47.61, 45.67, 40.61, 40.24, 38.55, 36.13, 35.98, 28.76, 27.73, 25.93, 25.85, 25.78, 23.89, 23.45, 22.33, 22.22, 18.77, 18.25, 18.17, 12.06, -4.37, -4.66, -4.86, -5.09; exact mass calculated for C₄₄H₈₄O₃Si₃Na (MNa)⁺ 767.5626, found 767.5646.

The protected vitamin 14 (7 mg, 9 μmol) was dissolved in THF (2 mL) and acetonitrile (2 mL). A solution of aq. 48% HF in acetonitrile (1:9 ratio, 2 mL) was added at 0° C. and the resulting mixture was stirred at room temperature for 8 h. Saturated aq. NaHCO₃ solution was added and the reaction mixture was extracted with ethyl acetate. The combined organic phases were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was diluted with 2 mL of hexane/ethyl acetate (8:2) and applied on a Waters silica Sep-Pak cartridge (2 g). An elution with hexane/ethyl acetate (8:2) and later with ethyl acetate gave the crude product 15 (4 mg). The vitamin 15 was further purified by straight phase HPLC [9.4×250 mm Zorbax Sil column, 4 mL/min, hexane/2-propanol (85:15) solvent system, R_f=9.27 min.] and by reverse phase HPLC [9.4×250 mm Zorbax Eclipse XDB-C18 column, 2.5 mL/min, methanol/water (85:15) solvent system, R_f=11.14 min.] to give a colorless oil (1.3 mg, 36% yield):



UV (in EtOH) λ_{max} 262.1, 252.6, 244.1 nm; ¹H NMR (800 MHz, CDCl₃) δ 6.35 and 5.87 (1H and 1H, each d, J=11.2 Hz, 6- and 7-H), 5.45-5.42 (1H, m, 22-H), 5.34-5.30 (1H, m, 23-H), 5.11 and 5.09 (each 1H, each s, =CH₂), 4.48 (2H, m, 1β- and 3α-H), 3.78 (1H, m, 25-H), 2.84 (1H, dd, J=12.8, 4.8 Hz, 10β-H), 2.80 (1H, br d, J=12.8 Hz, 9β-H), 2.57 (1H, dd, J=12.8, 2.5 Hz, 4α-H), 2.32 (1H, dd, J=12.8, 6.4 Hz, 4β-H), 2.29 (1H, dd, J=12.8, 8.0 Hz, 10α-H), 1.19 (3H, d, J=6.4 Hz, 27-H₃), 0.95 (3H, d, J=6.4 Hz, 21-H₃), 0.52 (3H, s, 18-H₃); exact mass calculated for C₂₆H₄₀O₃Na⁺ (MNa)⁺ 423.2875, found 402.2870.

13

Scheme 1





Scheme 2







14

-continued




Scheme 3



15

Scheme 4



BIOLOGICAL ACTIVITY OF 2-METHYLENE-
(20S,25S)-19,27-DINOR-(22E)-1 α ,25-DIHY-
DROXYVITAMIN D₃

The introduction of a methylene group to the 2-position, a methyl group at the 20-position (C-20) in its S-configuration, a double bond between the 22 and 23 positions (C-22 and C-23) in the side chain attached at the 17-position (C-17), and the replacement of the methyl group typically located at the 27 position (C-27) in the side chain with a hydrogen atom had little effect on binding of SS-22 to the full length recombinant rat vitamin D receptor, as compared to 1 α ,25-dihydroxyvita-

16

min D₃. The compound SS-22 bound equally to the receptor as compared to the standard 1,25-(OH)₂D₃ (FIG. 1). It might be expected from these results that compound SS-22 would have equivalent biological activity. Surprisingly, however, compound SS-22 is a highly selective analog with unique biological activity.

FIG. 5 shows that SS-22 has very little activity as compared to that of 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃), the natural hormone, in stimulating intestinal calcium transport.

FIG. 4 demonstrates that SS-22 has very little bone calcium mobilization activity, as compared to 1,25(OH)₂D₃.

FIGS. 4 and 5 thus illustrate that SS-22 may be characterized as having little, if any, calcemic activity.

FIG. 2 illustrates that SS-22 is as potent as 1,25(OH)₂D₃ on HL-60 cell differentiation, making it an excellent candidate for the treatment of psoriasis and cancer, especially against leukemia, colon cancer, breast cancer, skin cancer and prostate cancer. In addition, due to its relatively high cell differentiation activity, this compound provides a therapeutic agent for the treatment of various skin conditions including wrinkles, lack of adequate dermal hydration, i.e. dry skin, lack of adequate skin firmness, i.e. slack skin, and insufficient sebum secretion. Use of this compound thus not only results in moisturizing of skin but also improves the barrier function of skin.

FIG. 3 illustrates that the compound SS-22 has about the same transcriptional activity as 1 α ,25-dihydroxyvitamin D₃ in bone cells. This result, together with the cell differentiation activity of FIG. 2, suggests that SS-22 will be very effective in psoriasis because it has direct cellular activity in causing cell differentiation, gene transcription, and in suppressing cell growth. These data also indicate that SS-22 may have significant activity as an anti-cancer agent, especially against leukemia, colon cancer, breast cancer, skin cancer and prostate cancer.

The strong activity of SS-22 on HL-60 differentiation suggests it will be active in suppressing growth of parathyroid glands and in the suppression of the preproparathyroid gene.

EXPERIMENTAL METHODS

Vitamin D Receptor Binding

Test Material

Protein Source

Full-length recombinant rat receptor was expressed in *E. coli* BL21 (DE3) Codon Plus RIL cells and purified to homogeneity using two different column chromatography systems. The first system was a nickel affinity resin that utilizes the C-terminal histidine tag on this protein. The protein that was eluted from this resin was further purified using ion exchange chromatography (S-Sepharose Fast Flow). Aliquots of the purified protein were quick frozen in liquid nitrogen and stored at -80° C. until use. For use in binding assays, the protein was diluted in TEDK₅₀ (50 mM Tris, 1.5 mM EDTA, pH7.4, 5 mM DTT, 150 mM KCl) with 0.1% Chaps detergent. The receptor protein and ligand concentration were optimized such that no more than 20% of the added radiolabeled ligand was bound to the receptor.

Study Drugs

Unlabeled ligands were dissolved in ethanol and the concentrations determined using UV spectrophotometry (1,25(OH)₂D₃: molar extinction coefficient=18,200 and λ_{max} =265 nm; Analogs: molar extinction coefficient=42,000 and λ_{max} =252 nm). Radiolabeled ligand (³H-1,25(OH)₂D₃, ~159 Ci/mmol) was added in ethanol at a final concentration of 1 nM.

Assay Conditions

Radiolabeled and unlabeled ligands were added to 100 μ l of the diluted protein at a final ethanol concentration of $\leq 10\%$, mixed and incubated overnight on ice to reach binding equilibrium. The following day, 100 μ l of hydroxylapatite slurry (50%) was added to each tube and mixed at 10-minute intervals for 30 minutes. The hydroxylapatite was collected by centrifugation and then washed three times with Tris-EDTA buffer (50 mM Tris, 1.5 mM EDTA, pH 7.4) containing 0.5% Titron X-100. After the final wash, the pellets were transferred to scintillation vials containing 4 ml of Biosafe II scintillation cocktail, mixed and placed in a scintillation counter. Total binding was determined from the tubes containing only radiolabeled ligand.

HL-60 Differentiation

Test Material

Study Drugs

The study drugs were dissolved in ethanol and the concentrations determined using UV spectrophotometry. Serial dilutions were prepared so that a range of drug concentrations could be tested without changing the final concentration of ethanol ($\leq 0.2\%$) present in the cell cultures.

Cells

Human promyelocytic leukemia (HL60) cells were grown in RPMI-1640 medium containing 10% fetal bovine serum. The cells were incubated at 37° C. in the presence of 5% CO₂.

Assay Conditions

HL60 cells were plated at 1.2×10^5 cells/ml. Eighteen hours after plating, cells in duplicate were treated with drug. Four days later, the cells were harvested and a nitro blue tetrazolium reduction assay was performed (Collins et al., 1979; J. Exp. Med. 149:969-974). The percentage of differentiated cells was determined by counting a total of 200 cells and recording the number that contained intracellular black-blue formazan deposits. Verification of differentiation to monocytic cells was determined by measuring phagocytic activity (data not shown).

In vitro Transcription Assay

Transcription activity was measured in ROS 17/2.8 (bone) cells that were stably transfected with a 24-hydroxylase (24Oase) gene promoter upstream of a luciferase reporter gene (Arbour et al., 1998). Cells were given a range of doses. Sixteen hours after dosing the cells were harvested and luciferase activities were measured using a luminometer.

RLU=Relative Luciferase Units.

Intestinal Calcium Transport and Bone Calcium Mobilization

Male, weanling Sprague-Dawley rats were placed on Diet 11 (0.47% Ca) diet+AEK for one week followed by Diet 11 (0.02% Ca)+AEK for 3 weeks. The rats were then switched to a diet containing 0.47% Ca for one week followed by two weeks on a diet containing 0.02% Ca. Dose administration began during the last week on 0.02% calcium diet. Four consecutive ip doses were given approximately 24 hours apart. Twenty-four hours after the last dose, blood was collected from the severed neck and the concentration of serum calcium determined as a measure of bone calcium mobilization. The first 10 cm of the intestine was also collected for intestinal calcium transport analysis using the everted gut sac method.

Interpretation of Data

VDR binding, HL60 cell differentiation, and transcription activity. SS-22 ($K_i=3 \times 10^{-10}$ M) is as active as the natural hormone 1 α ,25-dihydroxyvitamin D₃ ($K_i=1 \times 10^{-10}$ M) in its ability to compete with [³H]-1,25(OH)₂D₃ for binding to the

full-length recombinant rat vitamin D receptor (FIG. 1). There is also little difference between SS-22 ($EC_{50}=1 \times 10^{-9}$ M) in its ability (efficacy or potency) to promote HL-60 cell differentiation as compared to 1 α ,25-dihydroxyvitamin D₃ ($EC_{50}=2 \times 10^{-9}$ M) (See FIG. 2). Also, compound SS-22 ($EC_{50}=1 \times 10^{-10}$ M) has similar transcriptional activity in bone cells as 1 α ,25-dihydroxyvitamin D₃ ($EC_{50}=2 \times 10^{-10}$ M) (See FIG. 3). These results suggest that SS-22 will be very effective in psoriasis because it has direct cellular activity in causing cell differentiation, gene transcription, and in suppressing cell growth. These data also indicate that SS-22 will have significant activity as an anti-cancer agent, especially against leukemia, colon cancer, breast cancer, skin cancer and prostate cancer, as well as against skin conditions such as dry skin (lack of dermal hydration), undue skin slackness (insufficient skin firmness), insufficient sebum secretion and wrinkles. It would also be expected to be very active in suppressing secondary hyperparathyroidism.

Calcium mobilization from bone and intestinal calcium absorption in vitamin D-deficient animals. Using vitamin D-deficient rats on a low calcium diet (0.02%), the activities of SS-22 and 1,25(OH)₂D₃ in intestine and bone were tested. As expected, the native hormone (1,25(OH)₂D₃) increased serum calcium levels at all dosages (FIG. 4). FIG. 4 shows that SS-22 has little, if any, activity in mobilizing calcium from bone. Administration of SS-22 at 260 pmol/day for 4 consecutive days did not result in mobilization of bone calcium, and increasing the amount of SS-22 to 7020 pmol/day was also without any substantial effect.

Intestinal calcium transport was evaluated in the same groups of animals using the everted gut sac method (FIG. 5). These results show that the compound SS-22 promotes intestinal calcium transport only slightly when administered at 260 pmol/day, whereas 1,25(OH)₂D₃ promotes a significant increase at the 260 pmol/day dose. It was only when 7020 pmol/day of SS-22 was administered that significant intestinal calcium transport activity was recorded, an almost 30-fold increase in dosage over the 260 pmol/day dose. Thus, it may be concluded that SS-22 is essentially devoid of intestinal calcium transport activity at the recommended lower doses.

These results illustrate that SS-22 is an excellent candidate for numerous human therapies as described herein, and that it may be particularly useful in a number of circumstances such as suppression of secondary hyperparathyroidism of renal osteodystrophy, autoimmune diseases, cancer, and psoriasis. SS-22 is an excellent candidate for treating psoriasis because: (1) it has significant VDR binding, transcription activity and cellular differentiation activity; (2) it is devoid of hypercalcemic liability at relatively low doses, unlike 1,25(OH)₂D₃; and (3) it is easily synthesized. Since SS-22 has significant binding activity to the vitamin D receptor, but has little ability to raise blood serum calcium, it may also be particularly useful for the treatment of secondary hyperparathyroidism of renal osteodystrophy.

These data also indicate that the compound SS-22 of the invention may be especially suited for treatment and prophylaxis of human disorders which are characterized by an imbalance in the immune system, e.g. in autoimmune diseases, including multiple sclerosis, lupus, diabetes mellitus, host versus graft rejection, and rejection of organ transplants; and additionally for the treatment of inflammatory diseases, such as rheumatoid arthritis, asthma, and inflammatory bowel diseases such as celiac disease, ulcerative colitis and Crohn's disease. Acne, alopecia and hypertension are other conditions which may be treated with the compound SS-22 of the invention.

19

The compounds of the invention of formula I, and particularly formula Ia, are also useful in preventing or treating obesity, inhibiting adipocyte differentiation, inhibiting SCD-1 gene transcription, and/or reducing body fat in animal subjects. Therefore, in some embodiments, a method of preventing or treating obesity, inhibiting adipocyte differentiation, inhibiting SCD-1 gene transcription, and/or reducing body fat in an animal subject includes administering to the animal subject, an effective amount of one or more of the compounds or a pharmaceutical composition that includes one or more of the compounds of formula I. Administration of the compound or the pharmaceutical compositions to the subject inhibits adipocyte differentiation, inhibits gene transcription, and/or reduces body fat in the animal subject. The animal may be a human, a domestic animal such as a dog or a cat, or an agricultural animal, especially those that provide meat for human consumption, such as fowl like chickens, turkeys, pheasant or quail, as well as bovine, ovine, caprine, or porcine animals.

For prevention and/or treatment purposes, the compounds of this invention defined by formula I may be formulated for pharmaceutical applications as a solution in innocuous solvents, or as an emulsion, suspension or dispersion in suitable solvents or carriers, or as pills, tablets or capsules, together with solid carriers, according to conventional methods known in the art. Any such formulations may also contain other pharmaceutically-acceptable and non-toxic excipients such as stabilizers, anti-oxidants, binders, coloring agents or emulsifying or taste-modifying agents.

The compounds of formula I and particularly SS-22, may be administered orally, topically, parenterally, rectally, nasally, sublingually or transdermally. The compound is advantageously administered by injection or by intravenous infusion or suitable sterile solutions, or in the form of liquid or solid doses via the alimentary canal, or in the form of creams, ointments, patches, or similar vehicles suitable for transdermal applications. A dose of from 0.01 μg to 1000 μg per day of the compounds I, particularly SS-22, preferably from about 0.1 μg to about 500 μg per day, is appropriate for prevention and/or treatment purposes, such dose being adjusted according to the disease to be treated, its severity and the response of the subject as is well understood in the art. Since the compound exhibits specificity of action, each may be suitably administered alone, or together with graded doses of another active vitamin D compound—e.g. 1α -hydroxyvitamin D_2 or D_3 , or $1\alpha,25$ -dihydroxyvitamin D_3 —in situations where different degrees of bone mineral mobilization and calcium transport stimulation is found to be advantageous.

Compositions for use in the above-mentioned treatments comprise an effective amount of the compounds I, particularly SS-22, as defined by the above formula I and Ia as the active ingredient, and a suitable carrier. An effective amount of such compound for use in accordance with this invention is from about 0.01 μg to about 1000 μg per gm of composition, preferably from about 0.1 μg to about 500 μg per gram of composition, and may be administered topically, transdermally, orally, rectally, nasally, sublingually, or parenterally in dosages of from about 0.01 $\mu\text{g}/\text{day}$ to about 1000 $\mu\text{g}/\text{day}$, and preferably from about 0.1 $\mu\text{g}/\text{day}$ to about 500 $\mu\text{g}/\text{day}$.

The compounds I, particularly SS-22, may be formulated as creams, lotions, ointments, topical patches, pills, capsules or tablets, suppositories, aerosols, or in liquid form as solutions, emulsions, dispersions, or suspensions in pharmaceutically innocuous and acceptable solvent or oils, and such preparations may contain in addition other pharmaceutically innocuous or beneficial components, such as stabilizers, antioxidants, emulsifiers, coloring agents, binders or taste-modifying agents.

The compounds I, particularly SS-22, may be advantageously administered in amounts sufficient to effect the differentiation of promyelocytes to normal macrophages. Dos-

20

ages as described above are suitable, it being understood that the amounts given are to be adjusted in accordance with the severity of the disease, and the condition and response of the subject as is well understood in the art.

The formulations of the present invention comprise an active ingredient in association with a pharmaceutically acceptable carrier therefore and optionally other therapeutic ingredients. The carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient thereof.

Formulations of the present invention suitable for oral administration may be in the form of discrete units as capsules, sachets, tablets or lozenges, each containing a predetermined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid; or in the form of an oil-in-water emulsion or a water-in-oil emulsion.

Formulations for rectal administration may be in the form of a suppository incorporating the active ingredient and carrier such as cocoa butter, or in the form of an enema.

Formulations suitable for parenteral administration conveniently comprise a sterile oily or aqueous preparation of the active ingredient which is preferably isotonic with the blood of the recipient.


Formulations suitable for topical administration include liquid or semi-liquid preparations such as liniments, lotions, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops; or as sprays.

For nasal administration, inhalation of powder, self-propelling or spray formulations, dispensed with a spray can, a nebulizer or an atomizer can be used. The formulations, when dispensed, preferably have a particle size in the range of 10 to 100 μ .

The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. By the term “dosage unit” is meant a unitary, i.e. a single dose which is capable of being administered to a patient as a physically and chemically stable unit dose comprising either the active ingredient as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers.

We claim:

1. A compound having the formula:



where X_1 , X_2 and X_3 , which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

2. The compound of claim 1 wherein X_3 is hydrogen.

3. The compound of claim 1 wherein X_1 is hydrogen.

4. The compound of claim 1 wherein X_1 , X_2 and X_3 are all t-butyl dimethylsilyl.

24. The method of claim **22** wherein the compound is administered parenterally.

23

25. The method of claim 22 wherein the compound is administered transdermally.


26. The method of claim 22 wherein the compound is administered rectally.

27. The method of claim 22 wherein the compound is administered nasally.


28. The method of claim 22 wherein the compound is administered sublingually.

29. The method of claim 22 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

30. The method of claim 22 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



31. A method of treating an autoimmune disease selected from the group consisting of multiple sclerosis, lupus, diabetes mellitus, host versus graft rejection, and rejection of organ transplants, comprising administering to a subject with said disease an effective amount of a compound having the formula:



where X₁, X₂ and X₃ which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

32. The method of claim 31 wherein the compound is administered orally.

33. The method of claim 31 wherein the compound is administered parenterally.

34. The method of claim 31 wherein the compound is administered transdermally.

24


35. The method of claim 31 wherein the compound is administered rectally.

36. The method of claim 31 wherein the compound is administered nasally.


37. The method of claim 31 wherein the compound is administered sublingually.

38. The method of claim 31 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

39. The method of claim 31 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



40. A method of treating an inflammatory disease selected from the group consisting of rheumatoid arthritis, asthma, and inflammatory bowel diseases, comprising administering to a subject with said disease an effective amount of a compound having the formula:



where X₁, X₂ and X₃, which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

41. The method of claim 40 wherein the compound is administered orally.

42. The method of claim 40 wherein the compound is administered parenterally.

43. The method of claim 40 wherein the compound is administered transdermally.

44. The method of claim 40 wherein the compound is administered rectally.


25

45. The method of claim 40 wherein the compound is administered nasally.


46. The method of claim 40 wherein the compound is administered sublingually.

47. The method of claim 40 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

48. The method of claim 40 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



49. A method of treating a skin condition selected from the group consisting of wrinkles, lack of adequate skin firmness, lack of adequate dermal hydration and insufficient sebum secretion which comprises administering to a subject with said skin condition an effective amount of a compound having the formula:



where X₁, X₂ and X₃ which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

50. The method of claim 49 wherein the compound is administered orally.

51. The method of claim 49 wherein the compound is administered parenterally.

52. The method of claim 49 wherein the compound is administered transdermally.

53. The method of claim 49 wherein the compound is administered topically.

54. The method of claim 49 wherein the compound is administered rectally.


26

55. The method of claim 49 wherein the compound is administered nasally.


56. The method of claim 49 wherein the compound is administered sublingually.

57. The method of claim 49 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

58. The method of claim 49 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



59. A method of treating renal osteodystrophy comprising administering to a subject with renal osteodystrophy an effective amount of a compound having the formula:



where X₁, X₂ and X₃, which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

60. The method of claim 59 wherein the compound is administered orally.

61. The method of claim 59 wherein the compound is administered parenterally.

62. The method of claim 59 wherein the compound is administered transdermally.

63. The method of claim 59 wherein the compound is administered rectally.


64. The method of claim 59 wherein the compound is administered nasally.

65. The method of claim 59 wherein the compound is administered sublingually.


27

66. The method of claim 59 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

67. The method of claim 59 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



68. A method of treating obesity of an animal, inhibiting adipocyte differentiation, inhibiting SCD-1 gene transcription, and/or reducing body fat in an animal comprising administering to an animal in need thereof an effective amount of a compound having the formula:



28

where X₁, X₂ and X₃, which may be the same or different, are each selected from hydrogen or a hydroxy-protecting group.

69. The method of claim 68 wherein the compound is administered orally.

70. The method of claim 68 wherein the compound is administered parenterally.

71. The method of claim 68 wherein the compound is administered transdermally.


72. The method of claim 68 wherein the compound is administered rectally.

73. The method of claim 68 wherein the compound is administered nasally.

74. The method of claim 68 wherein the compound is administered sublingually.

75. The method of claim 68 wherein the compound is administered in a dosage of from about 0.01 $\mu\text{g/day}$ to about 1000 $\mu\text{g/day}$.

76. The method of claim 68 wherein the compound is 2-methylene-(20S,25S)-19,27-dinor-(22E)-1 α ,25-dihydroxyvitamin D₃ having the formula:



77. The method of claim 68 wherein the animal is a human.

78. The method of claim 68 wherein the animal is a domestic animal.

79. The method of claim 68 wherein the animal is an agricultural animal.

* * * * *