



US009499786B2

(12) **United States Patent**
Thomson et al.

(10) **Patent No.:** **US 9,499,786 B2**
(45) **Date of Patent:** ***Nov. 22, 2016**

(54) **ENRICHED POPULATION OF HUMAN
PLURIPOTENT CELLS WITH OCT-4 AND
SOX2 INTEGRATED INTO THEIR GENOME**

(71) Applicant: **Wisconsin Alumni Research
Foundation**, Madison, WI (US)

(72) Inventors: **James A. Thomson**, Madison, WI
(US); **Junying Yu**, Middleton, WI (US)

(73) Assignee: **Wisconsin Alumni Research
Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 447 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **13/766,100**

(22) Filed: **Feb. 13, 2013**

(65) **Prior Publication Data**

US 2013/0210138 A1 Aug. 15, 2013

Related U.S. Application Data

(62) Division of application No. 12/053,440, filed on Mar.
21, 2008, now Pat. No. 8,440,461.

(60) Provisional application No. 60/919,687, filed on Mar.
23, 2007, provisional application No. 60/974,980,
filed on Sep. 25, 2007, provisional application No.
60/989,058, filed on Nov. 19, 2007.

(51) **Int. Cl.**

C12N 15/00 (2006.01)

C12N 5/00 (2006.01)

C12N 5/074 (2010.01)

C12N 5/0735 (2010.01)

(52) **U.S. Cl.**

CPC **C12N 5/0607** (2013.01); **C12N 5/0696**
(2013.01); **C12N 5/0606** (2013.01); **C12N**
2501/60 (2013.01); **C12N 2501/602** (2013.01);
C12N 2501/603 (2013.01); **C12N 2501/604**
(2013.01); **C12N 2501/605** (2013.01); **C12N**
2501/606 (2013.01); **C12N 2501/608**
(2013.01); **C12N 2502/99** (2013.01); **C12N**
2510/00 (2013.01); **C12N 2799/027** (2013.01)

(58) **Field of Classification Search**

CPC **C12N 5/0696**; **C12N 2501/602**; **C12N**
2501/603; **C12N 2510/00**; **C12N 2501/604**;
C12N 2501/606; **C12N 2799/027**; **C12N**
5/0606; **C12N 2501/605**; **C12N 2501/608**;
C12N 15/85; **C12N 2501/60**; **C12N**
2506/1307; **C12N 5/0662**; **C12N 15/86**;
C12N 2840/205; **C12N 2840/206**; **C12N**
2502/99; **A61K 35/545**; **A01N 1/0247**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,925,565	A	7/1999	Berlitz et al.
5,935,819	A	8/1999	Eichner et al.
6,060,273	A	5/2000	Dirks et al.
7,465,580	B2	12/2008	Sugden et al.
7,682,828	B2	3/2010	Jaenisch et al.
8,058,065	B2	11/2011	Yamanaka et al.
8,183,038	B2	5/2012	Thomson et al.
8,278,104	B2	10/2012	Yamanaka et al.
8,440,461	B2	5/2013	Thomson et al.
8,546,140	B2	10/2013	Mack et al.
2002/0090722	A1	7/2002	Dominko et al.
2002/0136709	A1	9/2002	Zahner et al.
2004/0199935	A1	10/2004	Chapman
2005/0079616	A1	4/2005	Reubinoff
2005/0255573	A1	11/2005	Chambers et al.
2006/0110830	A1	5/2006	Dominko et al.
2006/0128018	A1	6/2006	Zwaka et al.
2006/0160218	A1	7/2006	Slack et al.
2006/0263882	A1	11/2006	Fazio et al.
2008/0003560	A1	1/2008	Nakatsuji et al.
2009/0191159	A1*	7/2009	Sakurada et al. 424/93.7
2010/0041054	A1	2/2010	Mack
2010/0279404	A1	11/2010	Yamanaka et al.
2010/0285589	A1	11/2010	Lowry et al.
2011/0217274	A1	9/2011	Reld
2013/0189778	A1	7/2013	Mack
2013/0210138	A1	8/2013	Thomson et al.
2013/0217117	A1	8/2013	Thomson et al.
2014/0038293	A1	2/2014	Mack
2014/0057355	A1	2/2014	Thomson et al.

FOREIGN PATENT DOCUMENTS

CA	2660123	4/2009
EP	1698639	A2 9/2006
EP	2048229	4/2009
EP	2053128	4/2009
EP	2072618	6/2009
JP	2008-502644	3/1996
JP	2007-209240	8/2007

(Continued)

OTHER PUBLICATIONS

Stadtfeld et al. Induced pluripotent stem cells generated without
viral integration. *Science*, vol. 322, pp. 945-949.*
Okita et al. Generation of Mouse Induced Pluripotent Stem Cells
Without Viral Vectors. *Science*, 2008, vol. 322, pp. 949-953.*
Gonzales et al. Generation of mouse-induced pluripotent stem cells
by transient expression of a single nonviral polycistronic vector.
PNAS, 2009, vol. 106, pp. 8918-8922.*
Yamanaka. Induction of pluripotent stem cells from mouse fibro-
blasts by four transcription factors. *Cell Prolif.*, 2008, vol. 41
(Suppl. 1), pp. 51-56.*
Cherry et al. Retroviral Expression in Embryonic Stem Cells and
Hematopoietic Stem Cells. 2000, *Molecular and Cellular Biology*,
vol. 20, pp. 7419-7426.*

(Continued)

Primary Examiner — Deborah Crouch

(74) *Attorney, Agent, or Firm* — Quarles & Brady LLP

(57) **ABSTRACT**

The present invention relates to methods for reprogramming
a somatic cell to pluripotency by administering into the
somatic cell at least one or a plurality of potency-determin-
ing factors. The invention also relates to pluripotent cell
populations obtained using a reprogramming method.

9 Claims, 17 Drawing Sheets

(56)

References Cited

FOREIGN PATENT DOCUMENTS

JP	2007-535923	12/2007
JP	2008-067693	3/2008
JP	2007-525975	12/2013
WO	WO2005/072129	8/2005
WO	WO2005/108585	11/2005
WO	WO2007/047766	4/2007
WO	2007069666 A1	6/2007
WO	WO2007/069666	6/2007
WO	2008013737 A2	1/2008
WO	WO2008/013067	1/2008
WO	2008033469 A1	3/2008
WO	WO2008/118820	10/2008
WO	WO2009/032194	3/2009
WO	WO2009/032456	3/2009
WO	WO2009/061442	5/2009
WO	WO2009/091543	7/2009
WO	WO2009/092042	7/2009
WO	WO2009/133971	11/2009
WO	WO2009/149233	12/2009
WO	WO2010/012077	2/2010
WO	WO2010/028019	3/2010

OTHER PUBLICATIONS

- Daheron et al., LIF/STAT3 Signaling Fails to Maintain Self-Renewal of Human Embryonic Stem Cells. *Stem Cells*, 2004, vol. 22, pp. 770-778.*
- Alberio, et al., Reprogramming Somatic Cells Into Stem Cells, *Reproduction*, 2006, 132:709-720.
- Amit, et al., Clonally Derived Human Embryonic Stem Cell Lines Maintain Pluripotency and Proliferative Potential for Prolonged Periods of Culture, *Developmental Biology*, 2000, 227:271-278.
- Aoi, et al., Generation of Pluripotent Stem Cells From Adult Mouse Liver and Stomach Cells, *Science*, 2008, 321 (5889):699-702.
- Avilion, et al., Multipotent Cell Lineages in Early Mouse Development Depend on SOX2 Function, *Genes & Development*, 2003, 17:126-140.
- Chambers, et al., Functional Expression Cloning of Nanog, a Pluripotency Sustaining Factor in Embryonic Stem Cells, *Cell*, 2003, 113:643-655.
- Cibelli, et al., Cellular Reprogramming for the Creation of Patient-Specific Embryonic Stem Cells, *Stem Cell Reviews*, 2006, 2:289-296.
- Collas, Dedifferentiation of Cells: New Approaches, *Cytherapy*, 2007, 9(3):236-244.
- Cowan, et al., Nuclear Reprogramming of Somatic Cells After Fusion With Human Embryonic Stem Cells, *Science*, 2005, 309:1369-1373.
- Egli, et al., Developmental Reprogramming After Chromosome Transfer Into Mitotic Mouse Zygotes, *Nature*, 2007, 447:679-685.
- Fackler, Scientist at Work / Shinya Yamanaka—Risk Taking is in His Genes, *The New York Times*, Dec. 11, 2007, 4 pages.
- Frolov, et al., Alphavirus-Based Expression Vectors: Strategies and Applications, *Proc. Natl. Acad. Sci. USA*, 1996, 93:11371-11377.
- Gao, et al., Nonviral Gene Delivery: What We Know and What is Next, *AAPS Journal*, 2007, 9(1):E92-E104.
- Hochedlinger, et al., Nuclear Reprogramming and Pluripotency, *Nature*, 2006, 441:1061-1067.
- Ivanova, et al., Dissecting Self-Renewal in Stem Cells With RNA Interference, *Nature*, 2006, 442:533-538.
- Ludwig, et al., Feeder-Independent Culture of Human Embryonic Stem Cells, *Nature Methods*, 2006, 3(8):637-646.
- Ludwig, et al., Derivation of Human Embryonic Stem Cells in Defined Conditions, *Nature Biotechnology*, 2006, 24(2):185-187.
- Maherali, et al., Directly Reprogrammed Fibroblasts Show Global Epigenetic Remodeling and Widespread Tissue Contribution, *Cell Stem Cell*, 2007, 1:55-70.
- Masaki, et al., Heterogeneity of Pluripotent Marker Gene Expression in Colonies Generated in Human iPS Cell Induction Culture, *Stem Cell Research*, 2008, 1(2):105-115.
- Meissner, et al., Direct Reprogramming of Genetically Unmodified Fibroblasts Into Pluripotent Stem Cells, *Nature Biotechnology*, 2007, 25(10):1177-1181.
- Milanesi, et al., BK Virus-Plasmid Expression Vector That Persists Episomally in Human Cells and Shuttles Into *Escherichia coli*, *Molecular and Cellular Biology*, 1984, 4(8):1551-1560.
- Nakagawa, et al., Generation of Induced Pluripotent Stem Cells Without Myc From Mouse and Human Fibroblasts, *Nature Biotechnology*, 2008, 26(1):101-106.
- Nichols, J., et al., Formation of Pluripotent Stem Cells in the Mammalian Embryo Depends on the POU Transcription Factor Oct4, *Cell*, 1998, 95:379-391.
- Nichols, W., et al., Characterization of a New Human Diploid Cell Strain, IMR-90, *Science*, 1977, 196(4285):60-63.
- Park, et al., Reprogramming of Human Somatic Cells to Pluripotency With Defined Factors, *Nature*, 2008, 451:141-146.
- Pralong, et al., Cell Fusion for Reprogramming Pluripotency Toward Elimination of the Pluripotent Genome, *Stem Cell Reviews*, 2006, 2:331-340.
- Ren, et al., Establishment and Applications of Epstein-Barr Virus-Based Episomal Vectors in Human Embryonic Stem Cells, *Stem Cells*, 2006, 24:1338-1347.
- Ren, et al., Establishment of Human Embryonic Stem Cell Line Stably Expressing Epstein-Barr Virus-Encoded Nuclear Antigen 1, *Acta Biochimica et Biophysica Sinica*, 2005, 37(1):68-73.
- Richards, et al., The Transcriptome Profile of Human Embryonic Stem Cells as Defined by SAGE, *Stem Cells*, 2004, 22:51-64.
- Silva, et al., Nanog Promotes Transfer of Pluripotency After Cell Fusion, *Nature*, 2006, 441:997-1001.
- Takahashi, et al., Induction of Pluripotent Stem Cells From Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors, *Cell*, 2006, 126:663-676.
- Takahashi, et al., Induction of Pluripotent Stem Cells From Adult Human Fibroblasts by Defined Factors, *Cell*, 2007, 131:861-872.
- Taranger, et al., Induction of Dedifferentiation, Genomewide Transcriptional Programming, and Epigenetic Reprogramming by Extracts of Carcinoma and Embryonic Stem Cells, *Molecular Biology of the Cell*, 2005, 16:5719-5735.
- Thomson, et al., Embryonic Stem Cell Lines Derived from Human Blastocysts, *Science*, 1998, 282:1145-1147.
- Vodyanik, et al., Human Embryonic Stem Cell-Derived CD34+ Cells: Efficient Production in the Coculture with OP9 Stromal Cells and Analysis of Lymphohematopoietic Potential, *Blood*, 2005, 105(2):617-626.
- Wade-Martins, et al., Long-Term Stability of Large Insert Genomic DNA Episomal Shuttle Vectors in Human Cells, *Nucleic Acids Research*, 1999, 27(7):1674-1682.
- Wernig, et al., In Vitro Reprogramming of Fibroblasts Into a Pluripotent ES-Cell-Like State, *Nature*, 2007, 448:318-324.
- Yamanaka, et al., Strategies and New Developments in the Generation of Patient-Specific Pluripotent Stem Cells, *Cell Stem Cell*, 2007, 1(1):39-49.
- Yu, et al., Human Embryonic Stem Cells Reprogram Myeloid Precursors Following Cell-Cell Fusion, *Stem Cells*, 2006, 24:168-176.
- Yu, et al., Induced Pluripotent Stem Cell Lines Derived from Human Somatic Cells, *Science*, 2007, 318:1917-1920.
- Zwaka, et al., Homologous Recombination in Human Embryonic Stem Cells, *Nature Biotechnology*, 2003, 21:319-321.
- Liao et al., "Enhanced efficiency of generating induced pluripotent stem (iPS) cells from C86. human somatic cells by a combination of six transcription factors," *Cell Research*, 18:600-603, 2008.
- Lindner and Sugden, "The plasmid replicon of Epstein-Barr virus: mechanistic insights into efficient licensed, extrachromosomal replication in human cells," *Plasmid*, 58 (1): 1-12, 2007.
- Lois et al., "Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors," *Science*, 295:868-872, 2002.
- Lowry et al., "Generation of human induced pluripotent stems cells from dermal fibroblasts," *PNAS*, 105(8):2883-2888, 2008.
- Macejak and Sarnow, "Internal initiation of translation mediated by the 5' leader of a cellular mRNA," *Nature*, 353 (6339): 90-94, 1991.

(56)

References Cited**OTHER PUBLICATIONS**

- Okita et al., "Generation of germ line competent induced pluripotent stem cells" *Nature*, 2007, 448:313-317.
- Mackey and Sugden, "The linking regions of EBNA1 are essential for its support of replication and transcription," *Mol. Cell. Biol.*, 19(5):3349-3359, 1999.
- Mali et al. Improved Efficiency and Pace of Generating Induced Pluripotent Stem Cells from Human Adult and Fetal Fibroblasts. *Stem Cells*, 2008, vol. 26, pp. 1998-2005.
- Manzini et al., "Genetically modified pigs produced with a nonviral episomal vector," *PNAS*, 103(47):17672-17677, 2006.
- McBratney et al., "Internal initiation of translation," *Curr. Opin. Cell Biol.*, 5 (6): 961-965, 1993.
- Mikkelsen et al., "Dissecting direct reprogramming through integrative genomic analysis," *Nature*, 454: 49-56, 2008.
- Nanbo et al., "The coupling of synthesis and partitioning of EBV's plasmid replicon is revealed in live cells," *The EMBO Journal*, 26(19):4252-4562, 2007.
- Oster et al., "Myc is an essential negative regulator of platelet-derived growth factor beta receptor expression," *Mol. Cell. Biology*, 20(18):6768-6778, 2000.
- Papapetrou et al., "Stoichiometric and temporal requirements of Oct4, Sox2, Klf4, and c-C99. Myc expression for efficient human iPSC induction and differentiation," *PNAS Early Edition*. Published online before print Jun. 23, 2009, doi: 10.1073/pnas.0904825106.
- Park et al., "Disease-specific induced pluripotent stem cells," *Cell*, 134:877-886, 2008.
- Park et al., "Generation of human-induced pluripotent stem cells," *Nature Protocols*, 3:1180-1186, 2008.
- Pelletier and Sonenberg, "Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA," *Nature*, 334 (6180): 320-325, 1988.
- Piechaczek et al., "A vector based on the SV40 origin of replication and chromosomal S/MARs replicates episomally in CHO cells," *Nucleic Acids Res.*, 27(2):426-428, 1999.
- Polesskaya et al., "Lin-28 binds IGF-2 mRNA and participates in skeletal myogenesis by increasing translation efficiency" *Genes Dev.* 2007; 21: 1125-1138.
- Provost et al., "Viral 2A peptides allow expression of multiple proteins from a single ORF in transgenic zebrafish embryos," *Genesis*, 45 (10): 625-629, 2007.
- Rao, et al. "Assessing iPSC Reprogramming Methods for Their Suitability in Translational Medicine," *J Cell Biochem.* Oct. 2012; 113(10): 3061-3068.
- Pera, Reijo et al., "Gene expression profiles of human inner cell mass cells and embryonic stem cells," *Differentiation*, 78(1):18-23, 2009.
- Reubinoff et al., "Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro," *Nature Biotechnology*, 18:399-404, 2000.
- Schaarschmidt et al., "An episomal mammalian replicon: sequence-independent binding of the origin recognition complex," *EMBO J*, 23(1):191-201, 2004.
- Sears et al., "The amino terminus of Epstein-Barr Virus (EBV) nuclear antigen 1 contains AT hooks that facilitate the replication and partitioning of latent EBV genomes by tethering them to cellular chromosomes," *J Virol.*, 78 (21):11487-11505, 2004.
- Shao et al., "Generation of iPSC cells using defined factors linked via the self-cleaving 2A sequences in a single open reading frame," *Cell Research*, 19:296-306, 2009.
- Shi et al., "A combined chemical and genetic approach for the generation of induced pluripotent stem cells," *Cell Stem Cell*, 2:525-528, 2008.
- Sommer et al., "Induced pluripotent stem cell generation using a single lentiviral stem cell cassette," *Stem Cells*, 27:543-549, 2009.
- Stadtfeld et al., "Defining molecular cornerstones during fibroblast to iPSC cell reprogramming in mouse," *Cell Stem Cell*, 2:230-240, 2008.
- Stojkovic and Phinney, "Programming battle: egg vs. virus," *Stem Cells Express*, available online at <http://stemcells.alphamedpress.org/cgi/content/full/26/1/1>, Nov. 29, 2007.
- Takahashi, et al. Induced pluripotent stem cells in medicine and biology, *Development* 140:2457 (2013).
- Tanabe et al., "Historical overview and future prospects of the iPSC cells," *Regeneration Medicine*, 7(2):83-89, 2008. [Japanese only].
- Warren et al., "Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA," *Cell Stem Cell*, 7:1-13, 2010.
- Wernig et al., "A drug-inducible transgenic system for direct reprogramming of multiple somatic cell types," *Nature Biotechnology*, 26:916-924, 2008.
- Wernig et al., "C-Myc is dispensable for direct reprogramming of mouse fibroblasts," *Cell Stem Cell*, 2(1):10-12, 2008.
- Woltejen et al., "piggyBAC transposition reprograms fibroblasts to induce pluripotent stem cells," *Nature*, 458 (7239):766-770, 2009.
- Yates et al., "A cis-acting element from the Epstein-Barr viral genome that permits stable replication of recombinant plasmids in latently infected cells," *Proc. Natl. Acad. Sci. USA*, 81:3806-3810, 1984.
- Yates et al., "Stable replication of plasmids derived from Epstein-Barr virus in various mammalian cells," *Nature*, 313:812-815, 1985.
- Yeom et al., "Germline regulatory element of Oct-4 specific for the totipotent cycle of embryonic cells," *Development*, 122:881-894, 1996.
- Yoshimizu et al., "Germline-specific expression of the Oct-4/green fluorescent protein (GFP) transgene in mice," *Develop. Growth. Diff.*, 41:675-684, 1999.
- Yu et al., Supporting Online Material for: "Induced pluripotent stem cell lines derived from human somatic cells," *Science*, available at <http://www.sciencemag.org/cgi/data/1151526/DCI/2>, Nov. 20, 2007.
- Yu, et al. "Human induced pluripotent stem cells free of vector and transgene sequences" *Science*, 324, 5928:797-801, 2009.
- Telugu, et al. "Porcine induced pluripotent stem cells analogous to naïve and primed embryonic stem cells of the mouse" *Int. J. Dev. Biol.* 54: 1703-1711 (2010).
- Zhang Efficient Reprogramming of Naïve-Like Induced Pluripotent Stem Cells from Porcine Adipose-Derived Stem Cells with a Feeder-Independent and Serum-Free System.
- Hellen and Wimmer, "Translation of encephalomyocarditis virus RNA by internal ribosomal entry," *Curr Top Microbial Immunol.*, 203:31-63, 1995.
- Goessler et al., "Perspectives of Gene Therapy in Stem Cell Tissue Engineering" *Cells Tissues Organs*, 2006, 183(4): 169-179.
- Jackson et al., "Designing Nonviral Vectors for Efficient Gene Transfer and long-Term Gene Expression" *Molecular Therapy*, 2006; 14(5): 613-626.
- Liu et al., "Maintenance of Pluripotency in Human Embryonic Stem Cells Stably Over-expressing Enhanced Green Fluorescent Protein" *Stem Cells Dev.*, 2004, 13(6): 636-645.
- Nishikawa, "Reprogramming by the numbers" *Nature Biotechnology*, 2007; 25(8): 877-878; published Aug. 2007.
- "pBABE-MN-IRES-E-GFP," vector information accessed online at C64. http://www.lablife.org/p?a=vdb_view&id=g2.CylnpjVjRDXWTULiXTHEx8hmrO-, on Jan. 24, 2011.
- "pBluescript II XR predigested vector," instruction manual accessed at <http://www.genomics.agilent.com/files/Manual/212240.pdf> on Jan. 24, 2011.
- "piRES2-EGFP vector information," accessed at www.clontech.com on Jan. 24, 2011.
- "pLIB vector information," accessed online at www.clontech.com on Jan. 24, 2011.
- "pMXs retroviral vector data sheet," accessed online at www.cellbiolabs.com on Jan. 21, 2011.
- "pMX-STAT5A(1 *6) Retroviral Vector," Cell BioLabs, Inc., available at <http://www.cellbiolabs.com>, 2008.
- Almqvist et al., "Functional interaction of Oct transcription factors with the family of repeats in Epstein-Barr virus oriP," *J of General Virology*, 86 (Part 5): 1261-1267, 2005.
- Altmann et al., "Transcriptional activation by EBV nuclear antigen 1 is essential for the expression of EBV's transforming genes," *PNAS*, 103(38):14188-14193, 2006.

(56)

References Cited**OTHER PUBLICATIONS**

- Baer et al., "Transcriptional properties of genomic transgene integration sites marked by electroporation or retroviral infection," *Biochemistry*, 39:7041-7049, 2000.
- Baker, "A conversation with Shinya Yamanaka, Professor at Kyoto University," *Nature Reports Stem Cells*, pp. 1-2, published online Jun. 7, 2007, accessed online at <http://www.nature.com/stemcells/2007/0706/070607/full/stemcells.2007.9.html> on Oct. 18, 2012.
- Baker, "Reprogramming kinetics," *Nature Reports Stem Cells*, available at <http://www.nature.com/stemcells/2008/0803/080306/full/stemcells2008.44.html>, published online Mar. 6, 2008.
- Bielloch et al., "Generation of induced pluripotent stem cells in the absence of drug selection," *Cell Stem Cell*, 1:245-247, 2007.
- Bingham, "Cosuppression comes to the animals," *Cell*, 90(3):385-387, 1997.
- Bode et al., "The hitchhiking principle: Optimizing episomal vectors for the use in gene therapy and biotechnology," *Gene Ther. Mol. Biol.*, 6:33-46, 2001.
- Bode et al., "The transgeneticist's toolbox: novel methods for the targeted modification of eukaryotic genomes," *Biol. Chem.*, 381:801-813, 2000.
- Boyer et al., "Core transcriptional regulatory circuitry in human embryonic stem cells," *Cell*, 122(6):947-56, 2005.
- Brambrink et al., "Sequential expression of pluripotency markers during direct reprogramming of mouse somatic cells," *Cell Stem Cell*, 7(2):151-159, 2008.
- Calos et al., "Stability without a centromere," *PNAS*, 95:4084-4085, 1998.
- Carey et al., "Reprogramming of murine and human somatic cells using a single polycistronic vector," *PNAS USA*, 106 (1): 157-162, 2009.
- Chang et al., "Polycistronic lentiviral vectors for "Hit and Run" reprogramming of adult skin fibroblasts to induced pluripotent stem cells," *Stem Cells*, 27: 1042-1049, 2009.
- Chou, et al., "Efficient human iPS cell derivation by a non-integrating plasmid from blood cells with unique epigenetic and gene expression signatures" *Cell Research* 21:518 (2011).
- Conese M et al: Gene therapy progress and prospects: Episomally maintained self-replicating systems, *Gene Therapy*, Dec. 2004, pp. 1735-1741, vol. 11, No. 24.
- Cox and Rizzino, "Induced pluripotent stem cells: what lies beyond the paradigm shift," *Experimental Biology and Medicine*, 235:148-158, 2010.
- Egli et al., "Mediators of reprogramming: transcription factors and transitions through mitosis," *Molecular Cell Biology*, 9:505-516, 2008.
- Fellner, T. "Building Bridges from Research to Therapy: Development of Novel Technologies for the Generation of Defined and cGMP-compliant iPSCs" *Bio-Trac / NIH CRM Stem Cell Industry Symposium* May 29, 2013 at Third Party Requestor Lonza Walkersville, Inc. (Walkersville, MD).
- Garrick et al., "Repeat-induced gene silencing in mammals," *Nat. Genet.*, 18:56-59, 1998.
- Graham et al., "SOX2 functions to maintain neural progenitor identity," *Neuron*, 39:749-765, 2003.
- Hanna et al., "Direct reprogramming of terminally differentiated mature B lymphocytes to pluripotency," *Cell*, 133:250-264, 2008.
- Hanna et al., "Reprogramming of somatic cell identity," *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 73: 147-155, 2008.
- Hartenbach et al., "Autoregulated, bidirectional and multicistronic gas-inducible mammalian as well as lentiviral expression vectors," *Journal of Biotechnology*, 120:83-98, 2005.
- He et al., "A simplified system for generating recombinant adenoviruses," *PNAS*, 95:2509-2514, 1998.
- Hermeking et al., "Identification of CDK4 as a target of c-MYC," *PNAS*, 97(5):2229-2234, 2000.
- Huangfu et al., "Induction of pluripotent stem cells by defined factors in greatly improved by small-molecule compounds," *Nature Biotechnology*, 26:795-797, 2008.
- Hung et al., "Maintenance of Epstein-Barr virus (EBV) oriP-based episomes requires EBV C13. encoded nuclear antigen-1 chromosome-binding domains, which can be replaced by highmobility group-I or histone H1," *PNAS*, 98 (4):1865-1870, 2001.
- Jaenisch, "Transgenic animals," *Science*, 240 (4858): 1468-1474, 1988. {00192510}.
- Jenke et al., "Nuclear scaffold/matrix attached region modules linked to a transcription unit are sufficient for replication and maintenance of a mammalian episome," *PNAS*, 101 (31), 11322-11327, 2004.
- Jha et al., "SV 40-mediated immortalization," *Experimental Cell Res.*, 245(1): 1-7, 1998.
- Jia et al., "A nonviral minicircle vector for deriving human iPS cells," *Nature Methods*, 7(3):197-199, 2010.
- Kabouridis, "Biological applications of the protein transduction technology," *Trends in Biotechnology*, 21 (11):498-503, 2003.
- Kaji et al., "Virus-free induction of pluripotency and subsequent excision of reprogramming factors," *Nature*, 458 (7239):771-775, 2009.
- Kameda, T. et al., "A severe de novo methylation of episomal vectors by human ES cells" *Biochem. Biophys. Res. Comm.* 349:1269 (2006).
- Kaufman et al., "Translational efficiency of polycistronic mRNAs and their utilization to express heterologous genes in mammalian cells," *The EMBO Journal*, 6:187-193, 1987.
- Kennedy and Sugden, "EBNA-1, a bifunctional transcriptional activator," *Mol. and Cell. Biol.*, 23(19):6901-6908, 2003.
- Kennedy et al., "Epstein-Barr virus provides a survival factor to Burkitt's lymphomas," *PNAS*, 100:14269-14274, 2003.
- Kim et al., "Construction of a bifunctional mRNA in the mouse by using the internal C84. ribosomal entry site of the encephalomyocarditis virus," *Mol. Cellular Biol.*, 12 (8): 3636-3643, 1992.
- Kim et al., "Pluripotent stem cells induced from adult neural stem cells by reprogramming with two factors," *Nature*, 454:646-650, 2008.
- Leight and Sugden, "Establishment of an oriP replicon is dependent upon an infrequent, epigenetic event," *Mol. and Cell. Biol.*, 21(13):4149-4161, 2001.
- Lewitzky and Yamanaka, "Reprogramming somatic cells towards pluripotency by defined factors," *Current Opinion in Biotechnology*, 18:467-473, 2007.
- Office Action issued in U.S. Appl. No. 12/478,154, mailed Dec. 29, 2010.
- Office Action issued in U.S. Appl. No. 12/478,154, mailed Jul. 20, 2011.
- Office Action issued in U.S. Appl. No. 12/478,154, mailed Mar. 29, 2011.
- Office Action issued in U.S. Appl. No. 12/478,154, mailed May 2, 2013.
- Office Action issued in U.S. Appl. No. 12/478,154, mailed Oct. 23, 2012.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Feb. 29, 2012.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Jan. 28, 2011.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed May 24, 2013.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed May 31, 2011.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Oct. 19, 2010.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Oct. 24, 2012.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Sep. 24, 2014.
- Office Action issued in U.S. Appl. No. 12/539,366, mailed Sep. 21, 2011.
- Office Action issued in U.S. Appl. No. 12/605,220, mailed Feb. 10, 2012.
- Office Action issued in U.S. Appl. No. 12/605,220, mailed May 13, 2011.

(56)

References Cited

OTHER PUBLICATIONS

Office Action issued in U.S. Appl. No. 12/727,061, mailed Jun. 15, 2011.
 Office Action issued in U.S. Appl. No. 13/607,072, mailed Dec. 24, 2013.
 Office Action issued in U.S. Appl. No. 13/607,072, mailed Jul. 17, 2014.
 Office Action issued in U.S. Appl. No. 13/766,100, mailed Nov. 10, 2014.
 Office Action issued in U.S. Appl. No. 13/793,594, mailed Jun. 19, 2014.
 Office Action issued in U.S. Appl. No. 13/794,297, mailed Dec. 2, 2014.
 Office Communication issued in Australian Patent Application No. 2009256202 dated Apr. 29, 2014.
 Office Action issued in Canadian Application No. 2,734,128, mailed May 5, 2014.
 Office Action issued in Chinese Application No. 200980138340.8, mailed Mar. 7, 2013.
 Office Action issued in European Application No. 09759392.5, mailed Jun. 1, 2011.
 Office Action issued in European Application No. 09759392.5, mailed Sep. 5, 2014.
 Office Action issued in European Application No. 09791378.4, mailed Oct. 14, 2014.
 Office Action issued in Japanese Application No. 2011-512634, mailed Nov. 10, 2014.

Office Action issued in Japanese Application No. 2011-512634, mailed Dec. 20, 2013.
 Office Action issued in Japanese Application No. 2011-523094, mailed Jan. 17, 2014.
 Office Action issued in Japanese Application No. 2011-523094, mailed Sep. 19, 2014 {English translation only}.
 Office Action issued in U.S. Appl. No. 12/053,440, mailed Dec. 17, 2010.
 Office Action issued in U.S. Appl. No. 12/053,440, mailed Dec. 18, 2012.
 Office Action issued in U.S. Appl. No. 12/053,440, mailed Jun. 18, 2010.
 Office Action issued in U.S. Appl. No. 12/053,440, mailed Jul. 25, 2012.
 Office Action issued in U.S. Appl. No. 12/053,440, mailed May 20, 2011.
 Third Declaration of Amanda Mack filed in the prosecution of U.S. Appl. No. 12/478,154, filed Jun. 28, 2013.
 U.S. Appl. No. 61/136,615, filed Sep. 19, 2008, by Yamanaka, et al.
 U.S. Appl. No. 61/071,508, filed May 2, 2008, by Yamanaka, et al.
 U.S. Appl. No. 61/136,246, filed Aug. 21, 2008, by Yamanaka, et al.
 U.S. Appl. No. 61/058,858, filed Jun. 4, 2008, by Mack.
 PCT International Preliminary Report on Patentability, issued in International Application No. PCT/US2009/046209, dated Dec. 16, 2010.
 PCT International Search Report and Written Opinion, issued in International Application C102. No. PCT/US2009/053403, mail date Dec. 2, 2009.

* cited by examiner




FIG. 1




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4A





FIG. 4B

MOI: ~7.1



87.1%

FIG. 4C




FIG. 5




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B





FIG. 7C

SK46



iPS(SK46)




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B

1

ENRICHED POPULATION OF HUMAN PLURIPOTENT CELLS WITH OCT-4 AND SOX2 INTEGRATED INTO THEIR GENOME

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional of U.S. patent application Ser. No. 12/053,440, filed Mar. 21, 2008 and claims the benefit of U.S. Provisional Patent Application No. 60/919,687, filed Mar. 23, 2007; U.S. Provisional Patent Application No. 60/974,980, filed Sep. 25, 2007; and U.S. Provisional Patent Application No. 60/989,058, filed Nov. 19, 2007, each of which is incorporated herein by reference as if set forth in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was made with government support under RR000167 and GM069981 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND

Embryonic stem (ES) cells can grow indefinitely while maintaining pluripotency and can differentiate into cells of all three germ layers (Evans & Kaufman, *Nature* 292:154-156 (1981)). Human ES cells will be useful in treating a host of diseases, such as Parkinson's disease, spinal cord injury and diabetes (Thomson et al., *Science* 282:1145-1147 (1998)). Scientists have sought technical solutions to avoid the current method of generating ES cells from blastocyst cells and to avoid anticipated tissue rejection problems following transplantation into patients. One desirable way to accomplish these solutions would be to generate pluripotent cells directly from somatic cells of a post-natal individual.

Somatic cells can be reprogrammed by transferring their nuclear contents into oocytes (Wilmut et al., *Nature* 385: 810-813 (1997)) or by fusion with ES cells (Cowan et al., *Science* 309:1369-1373 (2005)), indicating that unfertilized eggs and ES cells contain factors that confer totipotency or pluripotency in somatic cells.

Likewise, Yu et al. showed that cells derived by in vitro differentiation from an H1 Oct4 knock-in ES cells did not express EGFP, but that EGFP expression was restored upon cell-cell fusion with human ES cells. Yu et al., *Stem Cells* 24:168-176 (2006), incorporated herein by reference as if set forth in its entirety). Therefore, Yu et al. demonstrated that differentiated cells can become pluripotent via cell-cell fusion with human ES cells. Regardless of the differentiated cell type, upon fusion with undifferentiated human ES cells, ES cell specific antigens and marker genes were expressed and differentiation-specific antigens were no longer detectable in the fused hybrid cells. Advantageously, EGFP expression was re-established in the hybrid cells, providing a convenient marker for re-establishment of pluripotent stem cell status. When the hybrid cells formed embryoid bodies (EBs), genes characteristic of all three germ layers and extra-embryonic tissues were up-regulated, indicating that the hybrid cells had a potential to differentiate into multiple lineages.

Although the transcriptional determination of pluripotency is not fully understood, several transcription factors, including Oct 3/4 (Nichols et al., *Cell* 95:379-391 (1998)), Sox2 (Avilion et al., *Genes Dev.* 17:126-140 (2003)) and

2

Nanog (Chambers et al., *Cell* 113:643-655 (2003)) are involved in maintaining ES cell pluripotency; however, none is sufficient alone to specify ES cell identity.

Chambers & Smith (EP 1 698 639 A2, (2002)) maintained pluripotent murine cells without a feeder layer or feeder cell extract and without a gp130 cytokine by introducing vectors that encode or activate differentiation-suppressing factors, but did not convert differentiated cells into a pluripotent state.

More recently, Takahashi & Yamanaka introduced four factors (i.e., Oct3/4, Sox2, c-Myc and Klf4) into mouse ES cells and mouse adult fibroblasts cultured under conditions suitable for mouse ES cell culture to obtain induced pluripotent stem (iPS) cells that exhibited mouse ES cell morphology and growth properties and expressed mouse ES cell marker genes (Takahashi & Yamanaka, *Cell* 126:663-676 (2006)). Notably, exogenous Oct-4 introduced into the mouse fibroblasts resulted in only marginal Oct-4 expression. Subcutaneous transplantation of iPS cells into nude mice resulted in tumors containing a variety of tissues from all three germ layers. Following injection into blastocysts, iPS cells contributed to mouse embryonic development. However, c-Myc, which was necessary for pluripotent induction, is an oncogene. Likewise, Klf4 is an oncogene. These data demonstrate that pluripotent cells can be directly generated from mouse fibroblast cultures by adding only a few defined factors using a retroviral transduction. However, as described infra, the set of factors used to produce iPS cells from differentiated mouse cells was insufficient to reprogram human somatic cells to pluripotency using lentiviral vectors without introducing additional changes to the cells.

One could hypothesize that factors that can reprogram human somatic cells differ from those factors that can reprogram somatic cells from model organisms (including mice) because ES cells from mice and humans require distinct sets of factors to remain undifferentiated, illustrating the significance of species-specific differences, even among mammals. For example, the leukemia inhibitory factor (LIF)/Stat3 pathway, a key to mouse ES cell proliferation, does not support human ES cell proliferation and appears inactive in conditions that support human ES cells (Daheron L, et al., *Stem Cells* 22:770-778 (2004); Humphrey R, et al., *Stem Cells* 22:522-530 (2004); and Matsuda T, et al., *EMBO J.* 18:4261-4269 (1999)).

Similarly, while bone morphogenetic proteins (BMPs) together with LIF support mouse ES cell self-renewal at clonal densities in serum-free medium (Ying Q, et al., *Cell* 115:281-292 (2003)), they cause rapid human ES cell differentiation in conditions that would otherwise support self-renewal, such as culture on fibroblasts or in fibroblast-conditioned medium (Xu R, et al., *Nat. Biotechnol.* 20:1261-1264 (2002)). Indeed, inhibition of BMP signaling in human ES cells is beneficial (Xu R, et al., *Nat. Methods* 2:185-190 (2005)).

Still further, fibroblast growth factor (FGF) signaling is important to self-renewal of human ES cells, but apparently not for mice (Xu et al., (2005), supra; and Xu C, et al., *Stem Cells* 23:315-323 (2005)).

Accordingly, the art still seeks a set of potency-determining factors suited at least for use in methods for reprogramming primate (including human and non-human) somatic cells to yield pluripotent cells. Such cells, obtained without relying upon embryonic tissues, would be suited for use in applications already contemplated for existing, pluripotent, primate ES cells.

BRIEF SUMMARY

The present invention is broadly summarized as relating to methods for reprogramming differentiated, somatic, pri-

mate cells into pluripotent cells, and more specifically into iPS cells. As used herein, "iPS cells" refer to cells that are substantially genetically identical to their respective differentiated somatic cell of origin and display characteristics similar to higher potency cells, such as ES cells, as described herein. The cells can be obtained from various differentiated (i.e., non-pluripotent and multipotent) somatic cells.

iPS cells exhibit morphological (i.e., round shape, large nucleoli and scant cytoplasm) and growth properties (i.e., doubling time; ES cells have a doubling time of about seventeen to eighteen hours) akin to ES cells. In addition, iPS cells express pluripotent cell-specific markers (e.g., Oct-4, SSEA-3, SSEA-4, Tra-1-60, Tra-1-81, but not SSEA-1). iPS cells, however, are not immediately derived from embryos and can transiently or stably express one or more copies of selected potency-determining factors at least until they become pluripotent. As used herein, "not immediately derived from embryos" means that the starting cell type for producing iPS cells is a non-pluripotent cell, such as a multipotent cell or terminally differentiated cell, such as somatic cells obtained from a post-natal individual.

In the methods described herein, at least two potency-determining factors can be introduced into, and expressed in, differentiated somatic cells, whereupon the somatic cells convert in culture to cells having properties characteristic of pluripotent cells, such as human ES cells (i.e., express at least Oct-4, SSEA-3, SSEA-4, TRA-1-60 or TRA-1-81, but not SSEA-1, and appear as compact colonies having a high nucleus to cytoplasm ratio and prominent nucleolus), that can differentiate into cells characteristic of all three germ layers, and that contain the genetic complement of the somatic cells of a post-natal individual. Apart from genetic material introduced to encode the potency-determining factors, the reprogrammed (i.e., converted) cells are substantially genetically identical to the somatic cells from which they were derived.

As used herein, a "potency-determining factor" refers to a factor, such as a gene or other nucleic acid, or a functional fragment thereof, as well as an encoded factor or functional fragment thereof, used to increase the potency of a somatic cell, so that it becomes pluripotent. The potency-determining factors optionally can be present only transiently in the reprogrammed cells or can be maintained in a transcriptionally active or inactive state in the genome of the reprogrammed cells. Likewise, the potency-determining factors can be present in more than one copy in the reprogrammed cells, where the potency-determining factor can be integrated in the cell's genome, can be extra-chromosomal or both. The potency-determining factors can include, but are not limited to, Stella (SEQ ID NO:1); POU5F1 (Oct-4; SEQ ID NO:2); Sox2 (SEQ ID NO:3); FoxD3, UTF1, Rex1, ZNF206, Sox15, Myb12, Lin28 (SEQ ID NO:4); Nanog (SEQ ID NO:5); DPPA2, ESG1, Otx2 and subsets thereof. In some embodiments, as few as two potency-determining factors, e.g., Oct-4 and Sox2, can be sufficient. Efficiency in obtaining reprogrammed cells, however, can be improved by including additional potency-determining factor, such as Lin28, Nanog or both.

In a first aspect, the invention relates to a replenishable, enriched population of pluripotent cells obtained from a post-natal individual, especially from a living individual, but optionally from a deceased individual. Cells in the enriched cell population express at least one cell-type-specific marker, including, but not limited to, Oct-4, SSEA3, SSEA4, Tra-1-60, Tra-1-81 or combinations thereof and have other hallmarks of pluripotent cells, such as ES cells. In addition, the pluripotent cells may express alkaline phosphatase

(ALP). Furthermore, the pluripotent cells may have a genome substantially genetically identical to that of a pre-existing, differentiated cell from the individual. Likewise, the pluripotent cells may have a genome that encodes at least one of the potency-determining factors, which may be transcriptionally active or inactive after reprogramming. Additionally, the potency-determining factors may be in a form of a reprogramming sequence in which a polynucleotide encoding the potency-determining factor is operably linked to a heterologous promoter. As used herein, "heterologous promoter" means a promoter that is operably linked to a polynucleotide for which the promoter does not normally initiate transcription.

In a second aspect, the invention relates to methods and compositions for identifying potency-determining factors required to reprogram somatic cells into pluripotent cells.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although suitable methods and materials for the practice or testing of the present invention are described below, other methods and materials similar or equivalent to those described herein, which are well known in the art, can also be used.

Other objects, advantages and features of the present invention will become apparent from the following specification taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

FIG. 1 illustrates a site downstream from a human Oct4 promoter into which a knock-in construct was introduced. In cells containing the knock-in construct, enhanced green fluorescent protein (EGFP) and neomycin phosphotransferase (NEO) are expressed when the Oct4 promoter is active. These cells can be used to evaluate which factors can reprogram somatic cells into pluripotent cells.

FIGS. 2A-B illustrate human H1 ES cell differentiation. FIG. 2A shows schematics of myeloid precursor derivation and purification from human ES cells. FIG. 2B shows phenotypic analysis of differentiated cells obtained after Percoll® separation. Gray line: isotype control; black line: antibody staining. Abbreviations: hESC, human embryonic stem cell; MPO, myeloperoxidase; PHEMA, poly(2-hydroxyethyl methacrylate).

FIG. 3 illustrates the Oct-4 region containing the knock-in construct of FIG. 1.

FIGS. 4A-C illustrate lentiviral transduction of somatic cells. FIG. 4A shows a schematic diagram of lentiviral construct. FIG. 4B shows Percoll®-purified cells were transduced with EGFP-expressing lentiviral vectors at various MOI. EGFP expression was analyzed by flow cytometry three days after transduction without drug selection. FIG. 4C shows lentiviral transduction of Percoll®-purified cells after several additional days of culture on Matrigel®. EGFP expression was analyzed two days after lentiviral transduction.

FIG. 5 illustrates transgene overexpression in cells differentiated for seven days on Matrigel®. No significant change in morphology was observed in cells overexpressing Nanog or EGFP (control). Morphology of Oct-4-expressing cells changes dramatically, and many of these cells survived neomycin selection, but none of these cells showed typical human ES cell morphology, indicating that a drug-selectable

5

population of Oct-4-expressing ES cells does not persist through the culture period necessary for myeloid differentiation.

FIGS. 6A-B illustrate reprogramming of Oct4KICD45+ A cells through introduction of fourteen potency-determining factors. FIG. 6A shows the established clones display undifferentiated human ES cell morphology and express EGFP under direction of the endogenous Oct4 promoter. FIG. 6B shows flow cytometry analysis of human ES cell-specific cell surface antigen expression in established clones. Gray line: isotype control; black line: antibody staining.

FIGS. 7A-C illustrate reprogramming efficiency, as evidenced by colony formation, after introduction of various sets of potency-determining factors. FIG. 7A shows the identified set of fourteen potency-determining factors was introduced into cells in combinations, wherein each combination excluded one of the fourteen factors. By evaluating the ability of the potency-determining factors to reprogram the tested cells to an ES-like state, the inventors determined whether the excluded potency-determining factor was essential to the reprogramming. For example, a set of potency-determining factors termed M1 that lacked Oct-4 (depicted as M1-Oct-4) was unable to form a significant number of ES-like colonies. As such, it was concluded that Oct-4 was important for somatic cell reprogramming. FIG. 7B shows a set of potency-determining factors (narrowed from FIG. 7A) evaluated for further testing was narrowed from fourteen to four (M4, being Oct-4, Sox2, Lin28 and Nanog). These four potency-determining factors were tested by serially excluding one of the four from the combination. Where a combination of three potency-determining factors (e.g., M4-Oct-4) was unable to reprogram the tested cells to form a significant number of stable ES-like colonies, the inventors concluded that the omitted gene is important for somatic cell reprogramming. In FIG. 7B, the light gray bars indicate the total number of reprogrammed colonies formed having typical human ES cell morphology; dark gray bars indicate the number of large colonies with minimal differentiation. FIG. 7C shows a set of potency-determining factors (narrowed from FIG. 7B) evaluated for further testing was narrowed from four to two (i.e., Oct-4 and Sox2). Oct-4, Sox2, Lin28 and Nanog were tested by serially excluding two of the four from the combination.

FIGS. 8A-B illustrate reprogramming in human adult skin fibroblasts. FIG. 8A shows bright-field images of human adult skin cell (p5) (left) and reprogrammed cells (right). FIG. 8B shows flow cytometry analysis of human ES cell-specific markers in human adult skin cells (p5) (bottom) and reprogrammed cells (top). Gray line: isotype control; black line: antibody staining.

FIGS. 9A-B illustrate the effect on reprogramming of relative expression of Oct-4 and Sox2. FIG. 9A shows Western blot analysis of Oct-4 and Sox2 in 293FT cells; lane 1, pSin4-EF2-Oct4-IRES1-Sox2 (OS-IRES1); lane 2, pSin4-EF2-Oct4-IRES2-Sox2 (OS-IRES2); lane 3, pSin4-EF2-Oct4-F2A-Sox2 (OS-F2A); lane 4, pSin4-EF2-Oct4-IRES1-puro (O); lane 5, pSin4-EF2-Sox2-IRES1-puro (S); lane 6, no plasmid (control). FIG. 9B shows reprogramming in mesenchymal cells derived from OCT4 knock-in human ES cells using linked potency-determining factors; gene combinations are the same as in FIG. 9A, with the addition of pSin4-EF2-Nanog-IRES1-puro (N) and pSin4-EF2-Lin28-IRES1-puro (L).

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The inventors hypothesized that potency-determining factors present in primate ES cells play an important role in

6

maintaining pluripotency and that differentiated somatic cells could be reprogrammed to a state of pluripotency through expression of potency-determining factors.

Cell types pass through various levels of potency during differentiation, such as totipotency, pluripotency and multipotency. Of particular interest herein are pluripotent cells. As used herein, "pluripotent cells" refer to a population of cells that can differentiate into all three germ layers (e.g., endoderm, mesoderm and ectoderm). Pluripotent cells express a variety of pluripotent cell-specific markers, have a cell morphology characteristic of undifferentiated cells (i.e., compact colony, high nucleus to cytoplasm ratio and prominent nucleolus) and form teratomas when introduced into an immunocompromised animal, such as a SCID mouse. The teratomas typically contain cells or tissues characteristic of all three germ layers. One of ordinary skill in the art can assess these characteristics by using techniques commonly used in the art. See, e.g., Thomson et al., supra. Pluripotent cells are capable of both proliferation in cell culture and differentiation towards a variety of lineage-restricted cell populations that exhibit multipotent properties. Multipotent somatic cells are more differentiated relative to pluripotent cells, but are not terminally differentiated. Pluripotent cells therefore have a higher potency than multipotent cells. As used herein, "reprogrammed pluripotent primate stem cells" (and similar references) refer to the pluripotent products of somatic cell reprogramming methods. Such cells are suitable for use in research and therapeutic applications currently envisioned for human ES cells.

The present invention broadly relates to novel methods for reprogramming differentiated somatic cells into higher-potency cells, such as pluripotent cells, by administering at least two potency-determining factors into somatic cells to achieve a higher level of potency in the reprogrammed cells than in the somatic cells. Advantageously, the present invention allows the generation of pluripotent cells, such as iPS cells, from somatic cells without requiring an addition of cell surface receptors for introducing the potency-determining factors to the somatic cells. As used herein, "reprogramming" refers to a genetic process whereby differentiated somatic cells are converted into de-differentiated, pluripotent cells, and thus have a greater pluripotency potential than the cells from which they were derived. That is, the reprogrammed cells express at least one of the following pluripotent cell-specific markers: SSEA-3, SSEA-4, TRA-1-60 or TRA 1-81. Preferably, the reprogrammed cells express all these markers.

Potency-determining factors that can reprogram somatic cells include, but are not limited to, factors such as Oct-4, Sox2, FoxD3, UTF1, Stella, Rex1, ZNF206, Sox15, Myb12, Lin28, Nanog, DPPA2, ESG1, Otx2 or combinations thereof. In the examples, a set with as few as two of the fourteen factors was sufficient to reprogram the tested cells; this set included Oct-4 and Sox2. Addition of other potency-determining factors to Oct-4 and Sox2, however, increased the efficiency with which reprogrammed cells were obtained. c-Myc and Klf4, however, are not essential as potency-determining factors. Preferably, the potency-determining factor may be a transcription factor.

Suitable somatic cells can be any somatic cell, although higher reprogramming frequencies are observed when the starting somatic cells have a doubling time about twenty-four hours. Somatic cells useful in the invention are non-embryonic cells obtained from a fetal, newborn, juvenile or adult primate, including a human. Examples of somatic cells that can be used with the methods described herein include, but are not limited to, bone marrow cells, epithelial cells,

fibroblast cells, hematopoietic cells, hepatic cells, intestinal cells, mesenchymal cells, myeloid precursor cells and spleen cells. Another type of somatic cell is a CD29⁺ CD44⁺ CD166⁺ CD105⁺ CD73⁺ and CD31⁻ mesenchymal cell that attaches to a substrate. Alternatively, the somatic cells can be cells that can themselves proliferate and differentiate into other types of cells, including blood stem cells, muscle/bone stem cells, brain stem cells and liver stem cells. Multipotent hematopoietic cells, suitably myeloid precursor or mesenchymal cells, are specifically contemplated as suited for use in the methods of the invention.

Likewise, suitable somatic cells are receptive, or can be made receptive using methods generally known in the scientific literature, to uptake of potency-determining factors including genetic material encoding the factors. Uptake-enhancing methods can vary depending on the cell type and expression system. Exemplary conditions used to prepare receptive somatic cells having suitable transduction efficiency are known in the art and are described in the examples below. One method for making cells receptive to potency-determining factors is described below in connection with the electroporation methods.

A potency-determining factor may be introduced as a reprogramming sequence in which a polynucleotide sequence encoding the potency-determining factor is operably linked to a heterologous promoter, which may become inactive after somatic cells are reprogrammed. The heterologous promoter is any promoter sequence that can drive expression of a polynucleotide sequence encoding the potency-determining factor in the somatic cell, such as, e.g., an Oct4 promoter.

The relative ratio of potency-determining factors may be adjusted to increase reprogramming efficiency. For example, linking Oct-4 and Sox2 in a 1:1 ratio on a single vector increased reprogramming efficiency in cells by a factor of four (FIG. 9A-B) when compared to reprogramming efficiency wherein the potency-determining factors were provided to cells in separate constructs and vectors, where the uptake ratio of the respective potency-determining factors into single cells was uncontrolled. Although the ratio of potency-determining factors may differ depending upon the set of potency-determining factors used, one of ordinary skill in possession of this disclosure can readily determine an optimal ratio of potency-determining factors.

Pluripotent cells can be cultured in any medium used to support growth of pluripotent cells. Typical culture medium includes, but is not limited to, a defined medium, such as TeSR™ (StemCell Technologies, Inc.; Vancouver, Canada), mTeSR™ (StemCell Technologies, Inc.) and StemLine® serum-free medium (Sigma; St. Louis, Mo.), as well as conditioned medium, such as mouse embryonic fibroblast (MEF)-conditioned medium. As used herein, a “defined medium” refers to a biochemically defined formulation comprised solely of biochemically-defined constituents. A defined medium may also include solely constituents having known chemical compositions. A defined medium may further include constituents derived from known sources. As used herein, “conditioned medium” refers to a growth medium that is further supplemented with soluble factors from cells cultured in the medium. Alternatively, cells can be maintained on MEFs in culture medium.

The inventors used a serial analysis of gene expression (SAGE) library to obtain transcriptome profiles of genes abundant in ES cells. Specifically, a SAGE library was used to identify potency-determining factors that regulate pluripotency and self-renewal in ES cells. SAGE libraries are well-known to one of ordinary skill in the art and are

publicly available or can be specifically constructed by companies, such as Agencourt Bioscience Corp. (Beverly, Mass.).

In another aspect, the invention provides an enriched population of pluripotent cells substantially genetically identical to cells of a post-natal individual. The cells can be obtained by reprogramming somatic cells isolated from the post-natal individual. In some embodiments, the cell population is a purified population, representing at least 60%, 70%, 80% and advantageously greater than 95% of the cells in the population, and any and all whole or partial integers therebetween. The reprogrammed cells are euploid, exhibit cell morphology characteristic of pluripotent cells and express pluripotent cell-specific markers, such as, e.g., Oct-4, SSEA-3, SSEA-4, Tra-1-60, Tra-1-81 or combinations thereof, and form teratomas when introduced into an immunocompromised animal.

Yet another aspect provides methods and compositions for identifying and using potency-determining factors sufficient to reprogram somatic cells into pluripotent cells. As noted herein, the reprogrammed pluripotent cells contain the genetic complement of, and are substantially genetically identical to somatic cells obtained from a post-natal individual. Generally, methods for identifying potency-determining factors include the steps of introducing genetic material encoding one or a plurality of putative potency-determining factors into somatic cells receptive to uptake of the genetic material under conditions effective to express the factors encoded on the introduced genetic material at levels sufficient to reprogram the cells to a less differentiated, higher-potency state; and observing a population of pluripotent cells after introduction of the genetic material. The pluripotent cells can be characterized by cell morphology, pluripotent cell-specific markers or both. Advantageously, the pluripotent cells can be identified by expression in the treated cells of a marker provided in the cells so as to be expressed only upon reprogramming of the cells to a pluripotent state. Through this approach, potency-determining factors capable of reprogramming somatic cells into pluripotent cells can be identified, as is described in the examples below.

Genetic material encoding a potency-determining factor can be introduced by transfection or transduction into the somatic cells using a vector, such as an integrating- or non-integrating vector. Of particular interest herein are retroviral vectors. Retroviral vectors, particularly lentiviral vectors, are transduced by packaging the vectors into virions prior to contact with a cell. After introduction, the DNA segment(s) encoding the potency-determining factor(s) can be located extra-chromosomally (e.g., on an episomal plasmid) or stably integrated into cellular chromosome(s).

A viral-based gene transfer and expression vector is a genetic construct that enables efficient and robust delivery of genetic material to most cell types, including non-dividing and hard-to-transfect cells (primary, blood, stem cells) in vitro or in vivo. Viral-based constructs integrated into genomic DNA result in high expression levels. In addition to a DNA segment that encodes a potency-determining factor of interest, the vectors include a transcription promoter and a polyadenylation signal operatively linked, upstream and downstream, respectively, to the DNA segment. The vector can include a single DNA segment encoding a single potency-determining factor or a plurality of potency-determining factor-encoding DNA segments. A plurality of vectors can be introduced into a single somatic cell. The vector can optionally encode a selectable marker to identify cells that have taken up and express the vector. As an example,

when the vector confers antibiotic resistance on the cells, antibiotic can be added to the culture medium to identify successful introduction of the vector into the cells. Integrating vectors can be employed, as in the examples, to demonstrate proof of concept. Retroviral (e.g., lentiviral) vectors are integrating vectors; however, non-integrating vectors can also be used. Such vectors can be lost from cells by dilution after reprogramming, as desired. A suitable non-integrating vector is an Epstein-Barr virus (EBV) vector. Ren C, et al., *Acta. Biochim. Sin.* 37:68-73 (2005); and Ren C, et al., *Stem Cells* 24:1338-1347 (2006), each of which is incorporated herein by reference as if set forth in its entirety.

The vectors described herein can be constructed and engineered using art-recognized techniques to increase their safety for use in therapy and to include suitable expression elements and therapeutic genes. Standard techniques for the construction of expression vectors suitable for use in the present invention are well-known to one of ordinary skill in the art and can be found in such publications such as Sambrook J, et al., "Molecular cloning: a laboratory manual," (3rd ed. Cold Spring Harbor Press, Cold Spring Harbor, N.Y. 2001), incorporated herein by reference as if set forth in its entirety.

The ability to identify and enrich for pluripotent cells can be facilitated by providing a non-lethal marker in the somatic cells, such as Green Fluorescent Protein (GFP), Enhanced Green Fluorescent Protein (EGFP) or luciferase, under the control of a promoter active only after the somatic cell has converted to a pluripotent state. A selectable marker gene is used to identify the reprogrammed cells expressing the marker through visible cell selection techniques, such as fluorescent cell sorting techniques. Alternatively, the reprogrammed cells can be produced without a selectable marker. In the examples below, a marker was provided in the genome of the somatic cells downstream of the promoter that regulates Oct-4 expression. The endogenous Oct4 promoter is active in undifferentiated, pluripotent ES cells. A drug-selectable population of Oct-4-expressing ES cells did not persist through the culture period necessary for myeloid differentiation. However, because some Oct-4 expression can persist into early stages of differentiation, it is appropriate to enrich the population for pluripotent cells by selecting colonies having characteristic ES cell morphology and by maintaining the cells under ES cell maintenance culture conditions. It is not intended that all cells in the reprogrammed cell culture have the desired level of potency. Given the inefficiencies of cell sorting technology, the variations in levels of gene expression and other biological effects, some cells in the enriched population may not be pluripotent. However, at a practical level, the reprogrammed cell population derived from somatic cells is enriched for pluripotent cells.

The non-lethal marker can be constructed to enable its subsequent removal using any of a variety of art-recognized techniques, such as removal via Cre-mediated, site-specific gene excision. For example, it may become desirable to delete the marker gene after the pluripotent cell population is obtained, to avoid interference by the marker gene product in the experiment or process to be performed with the cells. Targeted deletions can be accomplished by providing structure(s) near the marker gene that permits its ready excision. That is, a Cre/Lox genetic element can be used. The Lox sites can be built into the cells. If it is desired to remove the marker from the pluripotent cells, the Cre agent can be added to the cells. Other similar systems also can be used. Because Cre/Lox excision can introduce undesirable chro-

mosomal rearrangements and can leave residual genetic material after excision, the inventors recognize the desirability of introducing the potency-determining factors into the somatic cells using non-integrating, episomal vectors and obtaining cells from which the episomal vectors are lost (e.g., at a rate of about 5% per generation) by subsequently withdrawing the drug selection used to maintain the vectors during the reprogramming step.

The following examples are provided as further non-limiting illustrations of methods for identifying potency-determining genes or factors for converting somatic cells into pluripotent cells. In some examples, human H1 Oct4 knock-in ES cells were differentiated in stromal cell co-culture to yield cells suited for use as reprogrammable somatic cells. These cells are a model for cells isolated from a post-natal individual for use in a somatic cell reprogramming method.

The methods were repeated with other differentiated cell types. One cell type was human fetal lung fibroblast cells, IMR-90. See, Nichols W, et al., *Science* 196:60-63 (1977), incorporated herein by reference as if set forth in its entirety. IMR-90 cells are being extensively characterized by the ENCODE Consortium, are readily available from American Type Culture Collection (ATCC; Manassas, Va.; Catalog No. CCL-186), and have published DNA fingerprints that allow independent confirmation of the origin of reprogrammed clones. In addition, these cells proliferate robustly in Eagle's Minimal Essential Medium-10% FBS for more than twenty passages before undergoing senescence, but grow slowly in human ES cell culture conditions, a difference that provides a proliferative advantage to reprogrammed clones and aids in their selection by morphological criteria alone. Other differentiated cell types used in the methods were human post-natal foreskin fibroblast cells (ATCC; Catalog No. CRL-2097) and human adult skin cells (ATCC; Catalog No. CRL-2106).

The cells were made receptive for transduction with a viral expression system as described below. The somatic cells were transduced with polynucleotides encoding potency-determining factors thought to be associated with pluripotency, such that the somatic cells were reprogrammed to pluripotent cells. It is not yet determined whether all fourteen potency-determining factors provided in transduction vectors were taken up and expressed in the somatic cells. Having identified a set of fourteen potency-determining factors, and a subset of at least two of the fourteen factors, sufficient to reprogram somatic cells, the inventors provide one of ordinary skill in art the with the ability to identify one or more specific subsets of the potency-determining factors that are also capable of somatic reprogramming, thereby facilitating identification of other subsets of such potency-determining factors. Accordingly, the methods described below facilitate the identification of the potency-determining factors involved in reprogramming somatic cells into pluripotent cells.

It is specifically envisioned that the set of potency-determining factors sufficient to reprogram somatic cells can vary with the cell type of the somatic cells. It is noted that exposure to a set of fourteen potency-determining factors resulted in conversion to a pluripotent status in cultures of the indicated somatic cells. As shown below, one can identify a set of potency-determining factors sufficient to reprogram other cell types by repeating the methods described below using different combinations of potency-determining factors, which may include some or all of the fourteen factors as well as other potency-determining factors. Con-

sequently, one can produce pluripotent cell lines/populations that are substantially genetically identical to a pre-existing, differentiated, somatic cell.

EXAMPLES

In the following examples, differentiated cells received vectors that encoded various potency-determining factors. Some of the cells contained in their genome a marker gene that encodes EGFP positioned downstream from the regulated Oct4 promoter, which is active only in pluripotent cells. The production of this useful tool is described in Yu et al., *supra*, which demonstrated that differentiated cells can become pluripotent via cell-cell fusion with human ES cells.

Example 1

Lentiviral Vector Packaging and Production

Transgene-expressing lentivirus vector was produced in 293FT cell lines (Invitrogen). 293T is a fast-growing, highly transfectable clonal variant derived from transformed 293 embryonal kidney cells, which contains the large T antigen for high-level expression of the packaging proteins that contribute to higher viral titers. For routine maintenance and expansion, these cells were cultured in 293FT medium (DMEM/10% FBS, 2 mM L-glutamine and 0.1 mM MEM Non-Essential Amino Acids) in the presence of 500 µg/ml geneticin. For packaging, 293FT cells were collected by trypsinization. Following removal of trypsin by centrifugation, these cells were aliquoted into T75 flasks (15×10⁶ cells/flask, and 6 flasks per construct) in 293FT medium without geneticin.

Co-transfection of lentiviral vector and two helper plasmids was carried out with Superfect® transfection reagent (Qiagen) immediately following cell aliquoting (lentiviral vector: MD.G:pCMVdeltaR8.9:Superfect®=5 µg:5 µg:10 µg:40 µl in 400 µl of Iscove's Modified Dulbecco's Medium (IMDM) (1×)/flask incubated at room temperature for 10 minutes). The next day, the culture medium containing the transfection mixture was replaced with fresh 293FT medium supplemented with 1 mM sodium pyruvate (8 ml/flask). Lentivirus-containing supernatant was collected around 48 to 72 hours after transduction (~48 ml per construct). The 293FT cell debris was removed from the supernatant by centrifugation at 3000 rpm (1750 g) for 15 minutes at 4° C. To concentrate the lentivirus, the supernatant was filtered through 0.4 µm cellulose acetate (CA) membrane (Cornington, 115 ml low-protein binding), and ultracentrifuged in 70 ml sterilized bottles (Beckman, Cat#355622, polycarbonate for 45Ti rotor only) at 33,000 rpm (50,000 g) for 2.5 hours at 4° C. Lentivirus along with any remaining cell debris formed a visible pellet at the bottom of the centrifuge tube. Following supernatant removal, PBS (~300 µl for each construct) was added to resuspend the pellet by rocking the centrifuge tubes at 4° C. for 8 to 14 hours, or at room temperature for 2 hours. The remaining cell debris was removed by centrifugation at 5000 rpm (2700 g) for 5 minutes, and the resuspended lentivirus was aliquoted and stored at -80° C. The titer obtained generally ranged between 10⁷ to 10⁸ viral particles (vp)/ml after concentration. The sequence for a lentivirus (pSIN4-EF2-Stella-puro; SEQ ID NO:6, with the sequence for Stella from 3604 to 4083) harboring Stella (SEQ ID NO:1) is provided in the Sequence Listing. The same sequence was used for all other potency-determining factors (e.g., SEQ ID NOS: 2-5),

except that the sequence for Stella (SEQ ID NO:1) was replaced with the sequence of another potency-determining factor.

To efficiently introduce potency-determining factors into myeloid cells, inventors modified the lentiviral expression system (FIG. 4A). Inventors reduced the size of the original lentiviral construct (>11 kb) by removing sequences neighboring 5' and 3' LTRs through serial deletion analysis. These modifications minimized the negative effect on the packaging efficiency. The titer obtained routinely ranged between 10⁵ to 10⁶ vp/ml of supernatant, and 10⁷ to 10⁸ vp/ml after concentration (through ultracentrifugation). Restriction sites were introduced into the backbone for convenient exchanges of the coding regions for specific transgenes.

Example 2

Reprogramming of Myeloid Precursor Cells after Lentiviral Transduction and Expression of Potency-Determining Factors

To identify genes capable of reprogramming differentiated cells back to a state of pluripotency, efficient transduction of the cells is required. Inventors first tested the lentiviral transduction efficiency immediately after Percoll® purification of a human H1 Oct4 knock-in ES cells (FIG. 2).

H1.1 human ES cells (WiCell Research Institute; Madison, Wis.) were maintained on irradiated mouse embryonic fibroblasts (MEFs) in DMEM/F12 culture medium consisting of 80% Dulbecco's modified Eagle's medium (no pyruvate, high glucose formulation; Invitrogen; Carlsbad, Calif.) supplemented with 20% KnockOut serum replacer, 1% non-essential amino acids (Gibco), 1 mM L-glutamine, 0.1 mM β-mercaptoethanol (Sigma) and 4 ng/ml basic fibroblast growth factor (bFGF) (all from Invitrogen unless otherwise noted), as previously described (see Amit et al., *Dev Biol.* 227:271-278 (2000); and Thomson et al., *Science* 282:1145-1147 (1998), each of which is incorporated herein by reference as if set forth in its entirety). Feeder-free culture on Matrigel® (BD Biosciences; Bedford, Mass.) with chemically defined TeSR™ medium (StemCell Technologies, Inc.) was carried out as described in Ludwig et al. Ludwig T, et al, *Nat. Methods.* 3:637-646 (2006); and Ludwig T, et al., *Nat. Biotechnol.* 24:185-187 (2006), each of which is incorporated herein by reference as if set forth in its entirety.

The H1 Oct4 knock-in ES cell line was generated from the H1.1 human ES cells according to a method described by Zwaka & Thomson. U.S. Patent Publication No. 2006/0128018 and Zwaka T & Thomson J, *Nat. Biotechnol.* 21:319-321 (2003), each of which is incorporated herein by reference as if set forth in its entirety. Briefly, a gene targeting vector was constructed by inserting a cassette, an IRES-EGFP, an IRES-NEO and a simian virus polyadenylation sequence (approximately 3.2 kilobases (kb)) into the 3' untranslated region of the fifth exon of the human Oct-4 (octamer-binding transcription factor 4) gene, also known as POU domain, class 5, transcription factor 1 (POU5F1). This cassette was flanked in the 5' direction by a 6.3 kb homologous arm and by a 1.6 kb (6.5 kb in the alternative targeting vector) homologous arm in the 3' region (FIG. 1). The cassette was inserted at position 31392 of the Oct-4 gene (SEQ ID NO:2). The long arm contained a sequence from 25054-31392. The short arm contained a sequence from 31392-32970. In an alternative targeting vector, the short arm was substituted by a longer homologous region (31392-32970 in AC006047 plus 2387-7337 in gene accession number AC004195). Isogenic homologous

DNA was obtained by long-distance, genomic PCR and subcloned. All genomic fragments and the cassette were cloned into the multiple cloning site (MCS) of a cloning vector, pBluescript® SK II (GenBank Accession Number X52328; Stratagene; La Jolla, Calif.).

For electroporation, cells were harvested with collagenase IV (1 mg/ml, Invitrogen) for 7 minutes at 37° C., washed with medium and re-suspended in 0.5 ml culture medium (1.5–3.0×10⁷ cells). To prepare the cells for electroporation, cells were added to 0.3 ml phosphate-buffered saline (PBS; Invitrogen) containing 40 mg linearized targeting vector DNA. Cells were then exposed to a single 320 V, 200 µF pulse at room temperature using a BioRad Gene Pulser® II (0.4 cm gap cuvette). Cells were incubated for ten minutes at room temperature and were plated at high-density on Matrigel®. G418 selection (50 mg/ml; Invitrogen) was started 48 hours after electroporation. After one week, G418 concentration was doubled. After three weeks, surviving colonies were analyzed individually by PCR using primers specific for the NEO cassette and for the POU5F1 gene just downstream of 3' homologous region, respectively. PCR-positive clones were re-screened by Southern blot analysis using BamHI digested DNA and a probe outside the targeting construct.

The H1 Oct4 knock-in ES cell line expressed both EGFP and neomycin phosphotransferase (neo) from an endogenous Oct4 promoter/regulatory region using dual internal ribosome-entry sites (IRES) (FIG. 3). Expression of EGFP and neo in the H1 Oct4 knock-in ES cells indicated an active, endogenous Oct4 promoter/regulatory region.

H1 Oct4 knock-in ES cells were maintained through co-culture with mouse OP9 bone marrow stromal cells (FIG. 2A) maintained on gelatin-coated 10 cm plastic dishes (BD Biosciences) consisting of: DMEM medium (Invitrogen) supplemented with 20% non-heat-inactivated defined fetal bovine serum (FBS; HyClone Laboratories; Logan, Utah) (10 ml/dish). The OP9 cultures were split every 4 days at a ratio of 1:7. For use in human ES cell differentiation, after OP9 cells reached confluence on the fourth day, half of the medium was changed, and the cells were cultured for an additional four days.

For reprogramming, H1 Oct4 knock-in ES cells were differentiated into attached cells (i.e., CD29+CD44+CD166+CD105+CD73+CD31-). Briefly, human H1 Oct4 knock-in ES cells (p76 to 110) were added to the OP9 monolayer (1.5×10⁶/10-cm dish) in 20 ml of DMEM medium supplemented with 10% FBS (HyClone Laboratories) and 100 µM monothioglycerol (MTG; Sigma; St. Louis, Mo.). The human ES/OP9 cell co-culture was incubated for nine days with changes of half of the medium on days 4, 6 and 8. After incubation, the co-culture was dispersed into individual cells by collagenase IV treatment (1 mg/ml in DMEM medium, Invitrogen) for 20 minutes at 37° C., followed by trypsin treatment (0.05% Trypsin/0.5 mM EDTA, Invitrogen) for 15 minutes at 37° C. Cells were washed twice with medium and re-suspended at 2×10⁶/ml in DMEM medium supplemented with 10% FBS, 100 µM MTG and 100 ng/ml GM-CSF (Leukine; Berlex Laboratories Inc.; Richmond, Calif.). Cells were further cultured in flasks coated with poly (2-hydroxyethyl methacrylate) (pHEMA; Sigma) for 10 days with changes of half of the medium every 3 days. During adhesion-preventing pHEMA culture, cells that would otherwise be adherent formed floating aggregates, while the cells of interest grew as individual cells in suspension. Large cell aggregates were removed by filtration through 100 µm cell strainers (BD Biosciences), while small aggregates and dead cells were

removed by centrifugation through 25% Percoll® (Sigma). The differentiated cells recovered from the cell pellet expressed CD33, MPO, CD11b and CD11c molecules, which are characteristic for bone marrow myeloid cells (FIG. 2B). Inventors routinely produce 6-10×10⁶ differentiated cells from 1×10⁶ H1 ES cells (human H1 Oct4 knock-in ES cells). See also, Yu J, et al., Science 318:1917-1920 (2007), including the supplemental materials available at the Science website on the World Wide Web, incorporated herein by reference as if set forth in its entirety.

Lentivirus encoding a potency-determining factor (MOI: 3 to 10) was added to the cell culture after addition of polybrene carrier at a final concentration of 6 µg/ml (Sigma). The lentivirus-containing medium was replaced with fresh medium the next day, and cells were cultured further in appropriate medium. Drug selection, if needed, started the third day after transduction. As shown in FIG. 5B, the transduction efficiency was very low (~18.4% at MOI of 10). Moreover, the expression of EGFP was barely above background. Similar results have been obtained with routine plasmid or Epstein-Barr virus nuclear antigen (EBNA)-based plasmid transfections (data not shown).

On the other hand, cells having high transduction efficiency were prepared as follows. The Percoll®-purified H1 Oct4 knock-in ES cells were allowed to differentiate further to mesenchymal-like cells for an additional seven days in the presence of GM-CSF on Matrigel®, as described above. Many cells attached to the plate during this culture period. The attached cells (referred to herein below as Oct4KICD45+ A cells, or simply as CD45+ A cells) showed significantly higher transduction efficiency (FIG. 4C) and were used for this reprogramming experiment. While the cells were not CD45+ at the time of the experiments, the cells were obtained from CD45+ cells. As noted elsewhere herein, cell surface markers on the attached cells were characterized as CD29+, CD44+, CD166+, CD105+, CD73+, and CD31-.

Inventors tested the hypothesis that differentiated cells could be reprogrammed to a state of pluripotency by expressing potency-determining factors in Oct4KICD45+ A cells (FIG. 3), and obtained promising results. Because Nanog and Oct-4 are the best characterized potency-determining factors, inventors examined the effect of their over-expression in the cells.

The Oct4KICD45+ A cells were first dissociated to individual cells with trypsin and replated onto Matrigel® at ~10⁵ cells/well of 6-well plates in TeSR™ medium. Transgene-expressing lentiviral transduction was carried out the next day. Nanog-expressing Oct4KICD45+ A cells showed similar morphology to that of EGFP transfected cells (FIG. 5). Nanog over-expression, however, significantly enhanced Oct4KICD45+ A cell proliferation, similar to that observed in human ES cells. Following neomycin selection for an active endogenous Oct4 promoter/regulatory region, no Nanog- or EGFP-transfected cells survived. Importantly, these results indicate that a drug-selectable population of Oct-4-expressing ES cells does not persist through the culture period necessary for differentiation. Oct-4 expression resulted in dramatic morphological changes (FIG. 5), and many of these cells survived neomycin selection. None of these cells, however, exhibited morphology typical of human ES cells. The Oct4KICD45+ A cells co-expressing Nanog and Oct-4 showed morphological changes similar to those observed in cells expressing Oct-4 alone. Thus, it appears that the two key potency-determining factors, Nanog and Oct-4, alone were not sufficient to convert differentiated cells to pluripotency.

Cells were analyzed using cell-sorting methods before and after exposing the somatic cells to the factors. Adherent cells were individualized by trypsin treatment (0.05% Trypsin/0.5 mM EDTA, Invitrogen), and fixed in 2% para-formaldehyde for 20 minutes at room temperature. The cells were filtered through a 40- μ m mesh, and resuspended in FACS buffer (PBS containing 2% FBS and 0.1% sodium azide). Cells grown in suspension were stained in the FACS buffer supplemented with 1 mM EDTA and 1% normal mouse serum (Sigma). Intracellular myeloperoxidase (MPO) staining was performed using Fix & Perm® reagents (Caltag Laboratories; Burlingame, Calif.). About 100 μ l of cell suspension containing 5×10^5 cells was used in each labeling. Both primary and secondary antibody incubation (where applied) were carried out at room temperature for 30 minutes. Control samples were stained with isotype-matched control antibodies. After washing, the cells were resuspended in 300-500 μ l of the FACS buffer, and analyzed on a FACSCalibur flow cytometer (BDIS; San Jose, Calif.) using CellQuest™ acquisition and analysis software (BDIS). A total of 20,000 events were acquired. All of the antibodies used in the flow cytometry analysis are listed in Table 1. The final data and graphs were analyzed and prepared using FlowJo software (Tree Star, Inc.; Ashland, Oreg.).

TABLE 1

Antibodies for flow cytometry.				
ANTIGEN	LABEL	Clone/ Product#	ISOTYPE	VENDOR
SSEA-3	None	MAB4303	ratIgM	Chemicon
SSEA-3	None	14-8833-80	ratIgM	eBioscience
SSEA-4	None	MAB4304	mIgG3	Chemicon
SSEA-4	APC	FAB1435A	mIgG3	R&D systems
Tra-1-60	None	MAB4360	mIgM	Chemicon
Tra-1-81	None	MAB4381	mIgM	Chemicon
CD29	PE	MCA2298PE	IgG	AbD Serotec
Tra-1-85	APC	FAB3195A	mIgG1	R&D Systems
CD140a	PE	556002	mIgG2a	BD Pharmingen
CD56	PE	340724	mIgG2b	BDIS
CD73	PE	550257	mIgG1	BD Pharmingen
CD105	PE	MHCD10504	mIgG1	Caltag
CD31	FITC	557508	mIgG1	BD Pharmingen
CD34	FITC	555821	mIgG1	BD Pharmingen

BD Pharmingen (San Diego, CA)

BD Immunocytometry Systems (BDIS) (San Jose, CA)

Caltag Laboratories (Burlingame, CA)

Chemicon International (Temecula, CA)

AbD Serotec (Raleigh, NC)

NA—not applicable.

To further evaluate the potency-determining factors involved in reprogramming these cells, inventors explored the transduction of pools of ES cells enriched with various combinations of potency-determining factors. An exemplary pool of potency-determining factors for reprogramming myeloid precursors included the fourteen potency-determining factors described in Table 2 below.

TABLE 2

Human ES cell-enriched genes.			
GENE SYMBOL	UNIGENE ID	ENTREZ ID	ACCESSION
POU5F1 (Human Oct-4)	Hs.249184	5460	NM_002701
Sox2	Hs.518438	6657	NM_003106
Nanog	Hs.329296	79923	NM_024865

TABLE 2-continued

Human ES cell-enriched genes.			
GENE SYMBOL	UNIGENE ID	ENTREZ ID	ACCESSION
5 FoxD3	Hs.546573	27022	NM_012183
UTF1	Hs.458406	8433	NM_003577
Stella	Hs.131358	359787	NM_199286
Rex1	Hs.335787	132625	NM_174900
ZNF206	Hs.334515	84891	NM_032805
10 Sox15	Hs.95582	6665	NM_006942
Mybl2	Hs.179718	4605	NM_002466
Lin28	Hs.86154	79727	NM_024674
DPPA2	Hs.351113	151871	NM_138815
ESG1	Hs.125331	340168	NM_001025290
15 Otx2	Hs.288655	5015	NM_172337

The expression of at least some of these fourteen factors in the Oct4KICD45+ A cells resulted in colonies with typical morphology of pluripotent cells, such as human ES cells (FIG. 6A—left-hand photos). After neomycin selection from $\sim 10^5$ starting Oct4KICD45+ A cells, over ten colonies having the distinct ES cell morphology initially appeared. More than half of these colonies were subsequently lost to differentiation, suggesting either that over-expression of one or more introduced genes had a negative effect on the cells or that the cells continued to depend upon the foreign transgenes and gene silencing. Nevertheless, surviving colonies expressed the endogenous Oct4 promoter-driven EGFP (FIG. 7A-right-hand photos), indicating that the endogenous Oct4 promoter/regulatory region was reactivated.

In this embodiment, EGFP expression occurs when the native Oct4 promoter/regulatory region is active. In other words, undifferentiated cells are identified by a green color that disappears when the cells differentiate. Thus, the expression of endogenous Oct-4 in the primate ES cells was selectable. These colonies also expressed Oct-4, SSEA3, SSEA4, Tra-1-60 and Tra-1-81 pluripotent cell-specific markers (FIG. 6B). Similar results were obtained in reprogrammed colonies obtained using chemically defined TeSR™ medium.

Inventors randomly picked six colonies from two separate transfections with the same pool of fourteen ES cell-enriched potency-determining factors, and propagated five stable colonies for at least eight weeks. Thus, inventors identified a novel approach for reprogramming primate somatic cells to become higher potency cells by administering fourteen potency-determining factors into the somatic cells.

When these cells were exposed to other combinations of potency-determining factors (i.e., Sox2, c-Myc, Oct 3/4 and Klf4) using the lentiviral delivery system described herein, reprogramming and conversion of the cells were not observed.

Inventors used the techniques described herein to screen for subsets of the fourteen tested factors that are sufficient to reprogram the tested cells. Inventors' set of fourteen sufficient factors was subsequently narrowed to a set of six, and then four genes sufficient to reprogram these cells (FIGS. 7A-B; described further below). The four genes shown to be sufficient in combination to yield stable pluripotent cell were Oct-4, Nanog, Sox2 and Lin28, as shown in FIG. 7B.

Example 3

Reprogramming of Mesenchymal-Like Cells with a Limited Set of Four Potency-Determining Factors after Lentiviral Transduction

To identify a more limited set of potency-determining factors capable of reprogramming differentiated cells back

17

to pluripotency, the above-identified methods were repeated with a combination of Pou5F1 (Oct-4), Nanog, Sox2 and Lin28. Inventors used the techniques described above to screen these potency-determining factors for their ability to reprogram cells.

A different cell type was used in this example to further demonstrate the utility of the methods. The cell type was a mesenchymal-like clonal cell directly differentiated from human H1 Oct4 knock-in ES cells, as described above. As used herein, "clonal" refers to a characteristic of a population of cells derived from a common ancestor (i.e., derived from a single cell, not derived from a cell aggregate). That is, in a "clonal population," the cells display a uniform pattern of cell surface markers and morphological characteristics, as well as being substantially genetically identical.

Briefly, human H1 Oct4 knock-in ES cells (p76 to p110) were induced to differentiate in co-culture with mouse OP9 bone marrow stromal cells. See, Vodyanyk M, et al., Blood 105:617-626 (2005), incorporated herein by reference as if set forth in its entirety. Small aggregates of human H1 Oct4 knock-in ES cells were added to OP9 cells in alpha MEM supplemented with 10% FCS and 100 μ M MTG (Sigma). On the next day (day 1) of culture, the medium was changed, and the cultures were harvested on the days indicated below.

On day 2 of co-culture, mesodermal commitment was detected by a peak expression of transcription factors for mesendoderm (GSC, MIXL1 and T (BRACHYURY)) and early mesoderm (EVX1, LHX1 and TBX6) with Nimble-Gen® (Madison, Wis.) microarrays. During days 3-5, specification of endoderm and mesodermal lineages was observed. This stage was accompanied with sustained expression of genes involved in epithelial-mesenchymal transition (EMT; SNAIL and SLUG) and cell expansion (HOXB2-3). It also coincided with a maximal cell proliferation rate in human H1 Oct4 knock-in ES cells/OP9 co-culture.

Differentiation of specific mesendodermal lineages was observed on days 5-7 of co-culture, when markers of developing endoderm (AFP and SERPINA1), mesenchymal (SOX9, RUNX2 and PPARG2) and hemat endothelial (CDH5 and GATA1) cells were detected. However, muscle-inductive factors (MYOD1, MYF5 and MYF6) were not expressed throughout seven days of co-culture. Moreover, neuroectoderm (SOX1 and NEFL) or trophectoderm (CGB and PLAC) markers were not detected, indicating that OP9 cells provided an efficient inductive environment for directed hESC differentiation toward the mesendodermal pathway.

Also on day 2, a single-cell suspension of the human ES cell-derived cells was harvested by successive enzymatic treatment with collagenase IV (Gibco-Invitrogen) at 1 mg/ml in DMEM/F12 medium for 15 minutes at 37° C. and 0.05% Trypsin-0.5 mM EDTA (Gibco-Invitrogen) for 10 minutes at 37° C. Cells were washed 3 times with PBS-5% FBS, filtered through 70 μ m and 30 μ m cell strainers (BD Labware; Bedford, Mass.) and labeled with anti-mouse CD29-PE (AbD Serotec; Raleigh, N.C.) and anti-PE paramagnetic monoclonal antibodies (Miltenyi Biotech; Auburn, Calif.). The cell suspension was purified with magnet-activated cell sorting (MACs) by passing it through a LD magnetic column attached to a Midi-MACS separation unit (Miltenyi Biotech) to obtain a negative fraction of OP9-depleted, human H1 Oct4 knock-in ES cell-derived cells. Purity of human H1 Oct4 knock-in ES cell-derived cells was verified using pan anti-human TRA-1-85 monoclonal antibodies (R&D Systems; Minneapolis, Minn.).

18

Purified human H1 Oct4 knock-in ES cell-derived cells were plated at density of 2×10^4 cells/ml on semisolid, serum-free medium composed of StemLine™ serum-free medium (Sigma) supplemented with 5-100 ng/ml basic fibroblast growth factor (bFGF; PeproTech; Rocky Hill, N.J.) and 1% methylcellulose (StemCell Technologies, Inc.) with or without 10-20 ng/ml PDGF-BB (PeproTech). PDGF-BB improved growth of mesenchymal cells, but was not essential for colony formation. After 14-21 days of culture, large, compact mesenchymal colonies formed, resembling embryoid bodies (EBs). Mesenchymal colonies were detected on day 7; however, 14-21 days were required to reveal actively growing colonies.

Individual mesenchymal colonies were transferred to wells of a collagen- or fibronectin-coated, 96-well plate pre-filled with 0.2 ml/well StemLine® serum-free medium supplemented with 5-100 ng/ml bFGF. After 3-4 days of culture, adherent cells from individual wells were harvested by trypsin treatment and expanded on collagen- or fibronectin-coated dishes in StemLine® serum-free medium with 5-100 ng/ml bFGF.

Transgene-expressing lentiviral transduction was then carried out as described above. Inventors tested the hypothesis that differentiated mesenchymal-like cells could be reprogrammed to a state of pluripotency by expressing a limited set of potency-determining factors (e.g., Oct-4, Nanog, Sox2 and Lin28). The expression of at least these four potency-determining factors resulted in colonies having cells with typical morphology of pluripotent cells, such as human ES cells (FIG. 7B; dark gray bars). As shown in FIG. 7B, the greatest number of colonies having cells with typical morphology of pluripotent cells was obtained using the full complement of Oct-4, Nanog, Sox2 and Lin28. However, when one of Oct-4, Nanog, Sox2 or Lin28 was absent, the number of ES-like colonies was significantly attenuated (e.g., Nanog or Lin28) or absent (e.g., Oct-4 or Sox2).

In this embodiment, EGFP expression occurred when the native Oct4 promoter/regulatory region was active. In other words, undifferentiated cells were identified by a green color that was absent from differentiated cells. Thus, the expression of endogenous Oct-4 in the cells was selectable. Reprogrammed colonies also expressed Oct-4, SSEA3, SSEA4, Tra-1-60 and Tra-1-81 pluripotent cell-specific markers (data not shown).

Inventors randomly picked six colonies from two separate transfections with the same pool of fourteen ES cell-enriched potency-determining factors, and propagated five stable colonies for at least eight weeks. Thus, inventors identified a novel approach for reprogramming primate somatic cells to become higher potency cells by administering four potency-determining factors into the somatic cells.

When these cells were exposed to other combinations of potency-determining factors (i.e., Sox2, c-Myc, Oct 3/4 and Klf4) using the lentiviral delivery system described herein, reprogramming and conversion of the cells were not observed.

Example 4

Reprogramming of Mesenchymal-Like Cells with a Limited Set of Two Potency-Determining Factors after Lentiviral Transduction

To identify an even more limited set of potency-determining factors capable of reprogramming differentiated cells back to pluripotency, the above-identified methods

were repeated in the mesenchymal-like cells of Example 3 with a combination of two of the following four potency-determining factors: Oct-4, Nanog, Sox2 and Lin28. Inventors used the techniques described above to screen these potency-determining factors for their ability to reprogram cells.

Transgene-expressing lentiviral transduction was then carried out as described above. Inventors tested the hypothesis that differentiated mesenchymal-like cells could be reprogrammed to a state of pluripotency by expressing fewer than four potency-determining factors. The expression of at least Oct-4 and Sox2 (FIG. 7C) resulted in colonies having cells with typical morphology of pluripotent cells, such as human ES cells. Nanog and Lin28, singly and in combination, had a beneficial effect in clone recovery by improving reprogramming efficiency in human ES cell-derived mesenchymal cells to a state of pluripotency, but were essential neither for the initial appearance of reprogrammed cells nor for the expansion of reprogrammed cells.

Example 5

Reprogramming of a Differentiated Cells after Lentiviral Transduction and Expression of Four Potency-Determining Factors

To further demonstrate the utility of the limited set of potency-determining factors in reprogramming differentiated cells back to pluripotency, the above-identified methods were repeated with ATCC Catalog No. CCL-186 (IMR-90; ATCC), which are human fetal lung fibroblast cells (see also, Birney E, et al., Nature 447:799-816 (2007)).

Transgene-expressing lentiviral transduction was carried out as described above. That is, IMR-90 cells (0.9×10^6 well), were transduced with a combination of Oct-4, Sox2, Nanog and Lin28. Inventors tested the hypothesis that differentiated fibroblast cells could be reprogrammed to a state of pluripotency by expressing a limited set of potency-determining factors (e.g., Oct-4, Sox2, Nanog and Lin28). Following transduction, cells were transferred to three 10-cm dishes seeded with irradiated mouse embryonic fibroblasts (MEFs). By day 12 post-transduction, small colonies with human ES cell morphology became visible. On day 20 post-transduction, a total of 198 colonies were visible on 3 plates. Forty-one of the colonies were picked, thirty-five of which were successfully expanded for an additional three weeks. Six of these colonies were then selected for continued expansion and analysis, and the other twenty-nine were frozen.

The introduction of at least Oct-4, Sox2, Nanog and Lin28 resulted in colonies with typical morphology of pluripotent cells like human ES cells that had a normal karyotype. Cells from each colony likewise expressed telomerase activity and expressed human ES cell-specific surface antigens (i.e., SSEA-3, SSEA-4, Tra-1-60 and Tra-1-81). For each of the colonies, the expression of endogenous OCT4 and NANOG was at levels similar to that of pluripotent cells, although the exogenous expression of these genes did vary. Moreover, EB and teratoma formation demonstrated that the reprogrammed cells had a developmental potential to give rise to differentiated derivatives of all three primary germ layers.

DNA fingerprint analysis confirmed that these colonies were derived from IMR-90 cells and that they were not derived from human ES cells lines (e.g., H1, H7, H9, H13 and H14).

Similar to the data obtained with differentiated mesenchymal cells, the greatest number of colonies having cells with typical morphology of pluripotent cells, such as human ES cells was obtained using the full complement of Oct-4, Nanog, Sox2 or Lin28. However, when Oct-4, Nanog, Sox2 or Lin28 were absent, the number of ES-like colonies was significantly attenuated (e.g., Nanog or Lin28) or absent (e.g., Oct-4 or Sox2).

The colonies selected for expansion and detailed characterization proliferated for at least twelve weeks and retained typical characteristics of normal pluripotent cells, even though no selection for the activation of a pluripotency-specific gene was applied during reprogramming.

Reprogrammed cells were identified based on morphology alone (i.e., having a compact colony with high nucleus to cytoplasm ratio and prominent nucleolus). Reprogrammed cells also expressed Oct-4, SSEA3, SSEA4, Tra-1-60 and Tra-1-81 pluripotent cell-specific markers.

Example 6

Reprogramming of Differentiated Cells after Lentiviral Transduction and Expression of Three Potency-Determining Factors

To further demonstrate the utility of the limited set of potency-determining factors in reprogramming differentiated cells back to pluripotency, the above-identified methods were repeated with the IMR-90 cells, described above. In this set of experiments, fewer potency-determining factors were used than in Example 5.

Transgene-expressing lentiviral transduction was carried out as described above. IMR-90 cells were transduced with a combination of three of the following: Oct-4, Sox2, Nanog and Lin28. Inventors tested the hypothesis that differentiated fibroblast cells could be reprogrammed to a state of pluripotency by expressing the even more limited set of potency-determining factors. The expression of at least three factors resulted in colonies with typical morphology of pluripotent cells like human ES cells. Reprogrammed colonies having cells with typical morphology of pluripotent cells were obtained using the full complement of Oct-4, Sox2 and Nanog with or without Lin28. Therefore, the presence or absence of Lin28 did not affect reprogramming. However, when any of Oct-4, Nanog or Sox2 was absent, the number of reprogrammed colonies was significantly attenuated or absent.

To examine for the presence of Oct-4, Sox2, Nanog and Lin28 provirus in the reprogrammed cells, PCR with transgene-specific primer pairs (see, Table 3; one gene-specific primer and one lentiviral vector-specific primer) was carried out using genomic DNA from IMR-90 clones as template. The reactions employed the pfx DNA polymerase (Invitrogen, amplification buffer was used at $2\times$, and enhancer solution was used at $3\times$), and the following conditions: initial denaturation for 1 minute at 95°C ; 35 cycles of 94°C . for 30 seconds, 55°C . for 30 seconds, 68°C . for 2 minutes; and followed by 68°C . for 7 minutes. PCR analysis for the transgenes showed that either all four transgenes or three transgenes (i.e., Oct-4, Sox2 and Nanog) integrated into the pluripotent cells following exposure to transgene-expressing lentivirus vectors.

TABLE 3

Primer sets for assessing provirus integration.			
Genes	Size(bp)	Sequences (5' to 3')	
OCT4	656	OCT4-F1	CAGTGCCCGAAACCCACAC (SEQ ID NO: 7)
		SP3	AGAGGAAGTGTTCCTTCACGACA (SEQ ID NO: 8)
NANOG	732	NANOG-F2	CAGAAGGCCTCAGCACCTAC (SEQ ID NO: 9)
		SP3	AGAGGAAGTGTTCCTTCACGACA (SEQ ID NO: 8)
SOX2	467	SOX2-F1	TACCTCTTCTCCCACTCCA (SEQ ID NO: 10)
		SP3	AGAGGAAGTGTTCCTTCACGACA (SEQ ID NO: 8)
LIN28	518	LIN28-F1	AAGCGCAGATCAAAGGAGA (SEQ ID NO: 11)
		SP3	AGAGGAAGTGTTCCTTCACGACA (SEQ ID NO: 8)
OCT4endo	113	OCT4-F2	AGTTTGTGCCAGGGTTTTTG (SEQ ID NO: 12)
		OCT4-R2	ACTTCACCTTCCCTCCAACC (SEQ ID NO: 13)

Reprogrammed cells were identified based on morphology alone (i.e., having a compact colony with high nucleus to cytoplasm ratio and prominent nucleolus). Reprogrammed cells also expressed Oct-4, SSEA3, SSEA4, Tra-1-60 and Tra-1-81 pluripotent cell-specific markers.

Example 7

Reprogramming of Differentiated Cells after Lentiviral Transduction and Expression of Three Potency-Determining Factors

To further demonstrate the utility of the limited set of potency-determining factors in reprogramming differentiated cells to pluripotency, the above-identified methods were repeated with ATCC Catalog No. CRL-2097 (ATCC), which are human post-natal foreskin fibroblast cells.

Transgene-expressing lentiviral transduction was carried out as described above. Post-natal fibroblast cells (0.6×10^6 /well) were transduced with a combination of Oct-4, Sox2, Nanog and Lin28. Inventors tested the hypothesis that differentiated, post-natal, fibroblast cells could be reprogrammed to a state of pluripotency by expressing a limited set of potency-determining factors and obtained promising results. Following transduction, cells were transferred to three 10-cm dishes seeded with irradiated MEFs. By day 15 post-transduction, small colonies with pluripotent cell morphology became visible. On day 20 post-transduction, a total of 57 colonies were visible on the plates. Twenty-nine of the colonies were picked, twenty-seven of which were successfully expanded for an additional three weeks. Four of these colonies were then selected for continued expansion and analysis, and the other twenty-three were frozen.

The expression of Oct-4, Sox2, Nanog and Lin28 resulted in colonies having cells with typical morphology of pluripotent cells, such as human ES cells, and a normal karyotype. Reprogrammed colonies likewise expressed telomerase activity and expressed pluripotent cell-specific markers (i.e., SSEA-3, SSEA-4, Tra-1-60 and Tra1-81). For each, endogenous OCT4 and NANOG was expressed at levels similar to that observed in human pluripotent cells,

although the exogenous expression of these genes varied. Moreover, EB and teratoma formation demonstrated that the reprogrammed cells had a developmental potential to give rise to differentiated derivatives of all three primary germ layers. However, in contrast to the iPS cells obtained from IMR-90 cells, iPS cells derived from CRL-2097 cells showed a variation in the lineages apparent in teratomas examined at five weeks. Two of the iPS cell colonies showed neural differentiation; whereas the other two colonies showed multiple foci of columnar epithelial cells, reminiscent of primitive ectoderm.

DNA fingerprint analysis confirmed that these colonies were derived from the original cell line and confirmed that they were not derived from human ES cells lines (e.g., H1, H7, H9, H13 and H14).

Similar to the data obtained after transduction of differentiated mesenchymal cells, the greatest number of colonies having cells with typical morphology of human pluripotent cells were obtained using the full complement of Oct-4, Sox2, Nanog and Lin28. Interestingly, one cell line lacked Lin28, confirming that Lin28 was not essential for reprogramming somatic cells.

The colonies selected for expansion and detailed characterization proliferated for at least twelve weeks and retained typical characteristics of normal human pluripotent cells, even though no selection for the activation of a pluripotency-specific gene was applied during reprogramming.

Reprogrammed cells were identified based on morphology alone (i.e., having a compact colony with high nucleus to cytoplasm ratio and prominent nucleolus). Reprogrammed cells also expressed Oct-4, SSEA3, SSEA4, Tra-1-60 and Tra-1-81 pluripotent cell-specific markers.

When these cells were exposed to other combinations of factors (i.e., Sox2, c-Myc, Oct 3/4 and Klf4) using the lentiviral delivery system described herein, reprogramming and conversion of the cells were not observed.

Example 8

Reprogramming of Differentiated Cells after Lentiviral Transduction and Expression of Four Potency-Determining Factors

To further demonstrate the utility of the limited set of potency-determining factors in reprogramming differentiated cells to pluripotency, the above-identified methods were repeated with ATCC Catalog No. CRL-2106 (SK46; ATCC), which are human adult skin cells.

Transgene-expressing lentiviral transduction was carried out as described above. That is, skin cells (2.0×10^5 /well) were transduced with a combination of Oct-4, Sox2, Nanog and Lin28. Inventors tested the hypothesis that adult skin cells could be reprogrammed to a state of pluripotency by expressing a limited set of potency-determining factors and obtained promising results. Following transduction, cells were transferred to three 10-cm dishes seeded with irradiated mouse embryonic fibroblasts (MEFs). After 10 days in human ES cell culture medium human ES cell culture medium conditioned with irradiated MEFs was used to support cell growth. By day 18 post-transduction, small colonies with pluripotent cell morphology became visible.

The expression of Oct-4, Sox2, Nanog and Lin28 resulted in colonies having cells with typical morphology of pluripotent cells (see, FIG. 8A), such as human ES cells (i.e., having a compact colony with high nucleus to cytoplasm ratio and prominent nucleolus). As shown in FIG. 8B, the reprogrammed cells also expressed cell surface markers

23

typical of pluripotent cells; SK46 cells (control), however, did not. However, the reprogrammed colonies from adult skin cells appeared later than the cells in Example 7 and had a lower reprogramming efficiency than the cells in Example 7.

Example 9

Increasing Reprogramming Efficiency by Linking Potency-Determining Factors on a Single Construct

To increase the reprogramming efficiency, the above-identified methods were repeated using the construct shown in FIG. 4A; however, either Oct-4 or Sox2 were inserted in the transgene section, and Sox2 optionally replaced the puromycin resistance gene. The constructs were then expressed either in 293FT cells or in OCT4 knock-in human H1 ES cells (p6).

Transgene-expressing lentiviral transduction was carried out as described above. That is, 293FT cells or mesenchymal cells (~2x10⁵ cells/well of 6-well plate, seeded overnight) were transduced with various transgene combinations. Cells were transferred to 10 cm MEF dish (1 well of 6-well plate to 1x10 cm MEF dish) following the overnight incubation with lentivirus. Geneticin selection (50 µg/ml) for an active, endogenous, OCT4 promoter was carried out between day 11 to 15 post transduction. iPS colonies were counted on day 16.

FIG. 9A demonstrates that Oct-4 and Sox2 expression occurred in 293FT cells following transfection (see, e.g., lanes 1-3). In FIGS. 9A-B, pSin4-EF2-Oct4-IRES1-Sox2 is abbreviated as OS-IRES1; pSin4-EF2-Oct4-IRES2-Sox2 is

24

abbreviated as OS-IRES2; pSin4-EF2-Oct4-F2A-Sox2 is abbreviated as OS-F2A; pSin4-EF2-Oct4-IRES1-puro is abbreviated as 0; and pSin4-EF2-Sox2-IRES1-puro is abbreviated as S.

FIG. 9B shows that reprogramming efficiency increased in mesenchymal cells derived from OCT4 knock-in human H1 ES cells (p6) when Oct-4 and Sox2 were provided on the same construct (IRES1 is a very low-efficiency internal ribosome entry site; whereas IRES2 is a high-efficiency internal ribosome entry site). OS-IRES2+N+L (the high-efficiency IRES) showed an approximate four fold increase in reprogramming efficiency when compared to O+S, O+S+N+L or OS-IRES1 (the low-efficiency IRES)+N+L. Therefore, providing the potency-determining factors in one construct that provides for approximately equal expression levels of each can improve reprogramming efficiency.

It is understood that certain adaptations of the invention described in this disclosure are a matter of routine optimization for those skilled in the art, and can be implemented without departing from the spirit of the invention, or the scope of the appended claims.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. It is understood, however, that examples and embodiments of the present invention set forth above are illustrative and not intended to confine the invention. The invention embraces all modified forms of the examples and embodiments as come within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 480

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(480)

<223> OTHER INFORMATION: Stella

<400> SEQUENCE: 1

```
atggacccat cacagtttaa tccaacctac atcccagggt ctccacaaat gtcaccgaa    60
gaaaattccc gggacgattc aggggcctct caaatctcct ccgagacggt gataaagaac    120
cttagtaact tgactatcaa cgctagtagc gaatctgttt cccctctatc ggaagcttta    180
ctccgtcgag agtctgtagg agcagcagtc ctccaggaaa tcgaagatga gtggctttac    240
agcaggagag gagtaagaac attgctgtct gtgcagagag aaaagatggc aagattgaga    300
tacaatgttac tcggcggagt tcgtacgcac gaaagaagac caacaacaaa ggagcctaag    360
ggagttaaga aggaatcaag accattcaaa tgtccctgca gtttctgcgt gtctaatgga    420
tgggatcctt ctgagaatgc tagaataggg aatcaagaca ccaagccact tcagccataa    480
```

<210> SEQ ID NO 2

<211> LENGTH: 1083

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(1083)

-continued

<223> OTHER INFORMATION: Oct-4

<400> SEQUENCE: 2

```

atggcgggac acctggcttc ggatttcgcc ttctcgcccc ctccaggtgg tggaggtgat    60
gggccagggg ggccggagcc gggctggggt gatcctcgga cctggctaag cttccaaggc    120
cctcctggag ggccaggaat cgggccgggg gttgggccag gctctgaggt gtgggggatt    180
cccccatgcc ccccgccgta tgagtctgtg ggggggatgg cgtactgtgg gccccaggtt    240
ggagtggggc tagtgcccca aggcggcttg gagacctctc agcctgaggg cgaagcagga    300
gtcgggggtg agagcaactc cgatggggcc tccccggagc cctgcacctg caccctgggt    360
gccgtgaagc tggagaagga gaagctggag caaaaccggg aggagtccca ggacatcaaa    420
gctctgcaga aagaactcga gcaatttgcc aagctcctga agcagaagag gatcacctg    480
ggatatacac aggccgatgt ggggctcacc ctgggggttc tatttgggaa ggtattcagc    540
caaacgacca tctgccgctt tgaggctctg cagcttagct tcaagaacat gtgtaagctg    600
cggcccttgc tgcagaagtg ggtggaggaa gctgacaaca atgaaaatct tcaggagata    660
tgcaaagcag aaacctctgt gcaggcccca aagagaaagc gaaccagtat cgagaaccga    720
gtgagaggca acctggagaa tttgttctg cagtgccga aaccacact gcagcagatc    780
agccacatcg ccagcagct tgggctcgag aaggatgtgg tccagtggtg gttctgtaac    840
cggcgccaga agggcaagcg atcaagcagc gactatgcac aacgagagga ttttgaggct    900
gtggggtctc ctttctcagg gggaccagtg tcctttctc tggccccagg gccccathtt    960
ggtacccag gctatgggag ccctcacttc actgcactgt actcctcggc ccctttccct   1020
gagggggaag cctttccccc tgtctccgtc accactctgg gctctcccat gcattcaaac   1080
tga                                                    1083

```

<210> SEQ ID NO 3

<211> LENGTH: 954

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(954)

<223> OTHER INFORMATION: Sox2

<400> SEQUENCE: 3

```

atgtacaaca tgatggagac ggagctgaag ccgccgggccc cgcagcaaac ttcggggggc    60
ggcggcgcca actccaccgc ggcggcgggc ggcggcaacc agaaaaacag cccggaccgc    120
gtcaagcgcc ccatgaatgc cttcatggtg tgggtccgcg ggcagcggcg caagatggcc    180
caggagaacc ccaagatgca caactcggag atcagcaagc gcctgggcgc cgagtggaaa    240
cttttgtcgg agacggagaa gcggccggtc atcgacgagg ctaagcggct gcgagcgctg    300
cacatgaagg agcaccggga ttataaatac cggccccggc ggaaaaccaa gacgctcatg    360
aagaaggata agtacacgct gcccggcggg ctgctggccc ccggcggcaa tagcatggcg    420
agcgggggtc ggggtgggcgc cggcctgggc gggggcgtga accagcgcat ggacagttac    480
gcgcacatga acggctggag caacggcagc tacagcatga tgcaggacca gctgggctac    540
ccgcagcacc cgggcctcaa tgcgcacggc gcagcgcaga tgcagcccat gcaccgctac    600
gacgtgagcg cctcgagta caactccatg accagctcgc agacctacat gaacggetcg    660
cccacctaca gcatgtccta ctgcagcag ggcaccctg gcatggctct tggctccatg    720
ggttcggtag tcaagtccga ggccagctcc agccccctg tggttacctc ttctcccac    780

```

-continued

tccagggcgc cctgccaggc cggggacctc cgggacatga tcagcatgta tctcccgggc	840
gccgaggtgc cggaaccgcg cgcgccagc agacttcaca tgtcccagca ctaccagagc	900
ggcccggtgc cgggcacggc cattaacggc acactgcccc tctcacacat gtga	954

<210> SEQ ID NO 4
 <211> LENGTH: 630
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1) .. (630)
 <223> OTHER INFORMATION: Lin28

<400> SEQUENCE: 4

atgggctccg tgtccaaacca gcagtttgca ggtggctgcg ccaaggcggc agaagaggcg	60
cccgaggagg cgccggagga cgcggcccg gcgccggacg agcctcagct gctgcacggt	120
gcgggcatct gtaagtgggt caacgtgcgc atggggttcg gcttctctgc catgaccgcc	180
cgcgcggggg tcgcgctega ccccccagtg gatgtctttg tgcaccagag taagctgcac	240
atggaagggt tccggagctt gaaggagggt gaggcagtgg agttcacctt taagaagtca	300
gccaagggtc tggaatccat ccgtgtcacc ggacctggtg gagtattctg tattgggagt	360
gagaggcggc caaaaggaaa gagcatgcag aagcgcagat caaaaggaga caggtgctac	420
aactgtggag gtctagatca tcatgccaa gaaatgcaag tgccacccca gcccaagaag	480
tgccacttct gccagagcat cagccatatt gtagcctcat gtccgctgaa ggcccagcag	540
ggccctagtg cacagggaaa gccaacctac ttctgagagg aagaagaaga aatccacagc	600
cctacctctg tcccggaggc acagaattga	630

<210> SEQ ID NO 5
 <211> LENGTH: 918
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1) .. (918)
 <223> OTHER INFORMATION: Nanog

<400> SEQUENCE: 5

atgagtgtgg atccagcttg tccccaaagc ttgccttgct ttgaagcatc cgactgtaaa	60
gaatcttcac ctatgcctgt gatttgtggg cctgaagaaa actatccatc cttgcaaattg	120
tcttctgctg agatgcctca cacggagact gtctctctc ttcctctctc catggatctg	180
cttattcagg acagccctga ttcttccacc agtcccaaag gcaaacaacc cacttctgca	240
gagaatagtg tcgcaaaaaa ggaagacaag gtcccagtc aaaaacagaa gaccagaact	300
gtgttctctt ccaccagct gtgtgtactc aatgatagat ttcagagaca gaaatacctc	360
agcctccagc agatgcaaga actctccaac atcctgaacc tcagctacaa acagggtgaag	420
acctggttcc agaaccagag aatgaaatct aagagggtgc agaaaaacaa ctggccgaag	480
aatagcaatg gtgtgacgca gaaggcctca gcacctacct accccagcct ctactcttcc	540
taccaccagg gatgcctggt gaacccgact gggaaacctc caatgtggag caaccagacc	600
tggaacaatt caacctggag caaccagacc cagaacatcc agtcctggag caaccactcc	660
tggaacacto agacctggtg cacccaatcc tggaacaato aggcctggaa cagtccttcc	720
tataactgtg gagaggaatc tctgcagtcc tgcattgcagt tccagccaaa ttctcctgcc	780

-continued

agtgacttgg aggctgcttt ggaagctgct ggggaaggcc ttaatgtaat acagcagacc	840
actaggtatt ttagtactcc acaaacctatg gatttattcc taaactactc catgaacatg	900
caacctgaag acgtgtga	918

<210> SEQ ID NO 6
 <211> LENGTH: 8060
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Lentivirus harboring Stella
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (3604)..(4083)
 <223> OTHER INFORMATION: Sequence for Stella

<400> SEQUENCE: 6

cggaccgccca ctgccaatta cctgtgggttt catttactct aaacctgtga ttccctctgaa	60
ttatttttcat tttaaagaaa ttgtatttgt taaatatgta ctacaaactt agtagttgga	120
agggctaatt cactcccaaa gaagacaaga tacccttgat ctgtggatct accacacaca	180
aggctacttc cctgattagc agaactacac accagggccca ggggtcagat atccactgac	240
ctttggatgg tgctacaagc tagtaccagt tgagccagat aaggtagaag aggccaataa	300
aggagagaac accagcttgt tacacctgt gagcctgcat gggatggatg acccgagag	360
agaagtgtta gagtggaggt ttgacagccg cctagcattt catcacgtgg cccgagagct	420
gcatccggag tacttcaaga actgctgata tcgagcttgc tacaaggagc ttcccgctgg	480
ggactttcca gggaggcgtg gcctgggagg gactggggag tggcgagccc tcagatcctg	540
catataagca gctgcttttt gcctgtactg ggtctctctg gttagaccag atctgagcct	600
gggagctctc tggctaacta gggaaaccac tgcttaagcc tcaataaagc ttgccttgag	660
tgcttcaagt agtgtgtgcc cgtctgttgt gtgactctgg taactagaga tccctcagac	720
ccttttagtc agtgtggaat atctctagca gtggcgcccg aacagggact tgaaagcgaa	780
agggaaacca gaggagctct ctgcagcag gactcggtt gctgaagcgc gcacggcaag	840
aggcgagggg cggcgactgg tgagtacgcc aaaaattttg actagcgag gctagaagga	900
gagagatggg tgcgagagcg tcagtattaa gcgggggaga attagatcgc gatgggaaaa	960
aattcggtta aggccagggg gaaagaaaa atataaatta aaacatatag tatgggcaag	1020
caggagagcta gaacgattcg cagttaatcc tggcctgtta gaaacatcag aaggctgtag	1080
acaaatactg ggacagctac aaccatccct tcagacagga tcagaagaac ttagatcatt	1140
atataataca gtacgaaccc tctatttgtt gcatcaaagg atagagataa aagacaccaa	1200
ggaagcttta gacaagatag aggaagagca aaacaaaagt aagaccaccg cacagcaagc	1260
ggccgctgat cttcagacct ggaggaggag atatgaggga caattggaga agtgaattat	1320
ataaatataa agtagtaaaa attgaacct taggagtagc accaccaag gcaaagagaa	1380
gagtgggtgca gagagaaaaa agagcagtgg gaataggagc tttgttcctt gggttcttgg	1440
gagcagcagg aagcactatg ggcgcagcgt caatgacgct gacggtacag gccagacaat	1500
tattgtctgg tatagtgcag cagcagaaca atttgctgag ggctattgag gcgcaacagc	1560
atctgttgca actcacagtc tggggcatca agcagctcca ggcaagaatc ctggctgtgg	1620
aaagatacct aaaggatcaa cagctcctgg ggatttgggg ttgctctgga aaactcattt	1680
gcaccactgc tgtgccttgg aatgctagtt ggagtaataa atctctggaa cagatttgg	1740
atcacacgac ctggatggag tgggacagag aaattaacaa ttacacaagc ttaatacact	1800

-continued

ccttaattga	agaatcgcaa	aaccagcaag	aaaagaatga	acaagaatta	ttggaattag	1860
ataaatgggc	aagtttgttg	aattggttta	acataacaaa	ttggctgttg	tatataaaat	1920
tattcataat	gatagtagga	ggcttgtag	gtttaagaat	agtttttgct	gtactttcta	1980
tagtgaatag	agttaggcag	ggatattcac	cattatcggt	tcagaccac	ctcccaaccc	2040
cgaggggacc	cgacaggccc	gaaggaatag	aagaagaagg	tggagagaga	gacagagaca	2100
gatccattcg	attagtgaac	ggatctcgac	ggatctcgca	caaatggcag	tattcatcca	2160
caattttaaa	agaaaggggg	ggattggggg	gtacagtgcg	ggggaagaa	tagtagacat	2220
aatagcaaca	gacatacaaa	ctaaagaatt	acaaaaacaa	attacaaaaa	ttcaaaattt	2280
tcgggtttat	tacagggaca	gcagagatcc	actttggctg	ataagctttg	caaagatgga	2340
taaagtttta	aacagagagg	aatctttgca	gctaattggac	cttctaggtc	ttgaaaggag	2400
tgggaatttg	ctccgggtgc	cgtcagtggg	cagagcgcac	atcgcccaca	gtccccgaga	2460
agttgggggg	aggggtcggc	aattgaaccg	gtgcctagag	aaggtggcgc	ggggtaaac	2520
gggaaagtga	tgctgtgtac	tggtctcgcc	tttttcccg	gggtggggga	gaaccgtata	2580
taagtgcagt	agtcgccgtg	aacgttcttt	ttcgcaacgg	gtttgccgcc	agaacacagg	2640
taagtgccgt	gtgtggttcc	cgcggggcctg	gcctctttac	gggttatggc	ccttgctgtc	2700
cttgaattac	ttccacctgg	ctgcagtacg	tgattcttga	ttccgagctt	cgggttgga	2760
gtgggtggga	gagttcgagg	ccttgcgctt	aaggagcccc	ttcgccctcg	gcttgagttg	2820
aggcctggcc	tgggcgctgg	ggcgcccgcg	tgcaaatctg	tgggcacctt	cgcgcctgtc	2880
tcgctgcttt	cgataagtct	ctagccattt	aaaatttttg	atgacctgct	gcgacgcttt	2940
ttttctggca	agatagtctt	gtaaatgcgg	gccaaatctt	gcacactggg	atttcggttt	3000
ttggggccgc	ggcgggcgac	ggggcccggtg	cgtcccagcg	cacatgttcg	gcgaggcggg	3060
gcctgcgagc	gcggccaccg	agaatcggac	gggggtagtc	tcaagctggc	cggcctgctc	3120
tgggtgcctg	cctcgcgccg	cgtgtatcgt	ccccgccctg	ggcggaagg	ctggcccggg	3180
cggcaccagt	tgctgtgagc	gaaagatggc	cgttcccgg	ccctgctgca	gggagctcaa	3240
aatggaggac	gcggcgctcg	ggagagcggg	cgggtgagtc	acccacacaa	aggaaaagg	3300
cctttccgtc	ctcagccgtc	gcttcatgtg	actccacgga	gtaccgggcg	ccgtccaggc	3360
acctcgatta	gttctcgagc	ttttggagta	cgtcgtcttt	aggttggggg	gaggggtttt	3420
atgcgatgga	gtttccccc	actgagtggt	tggagactga	agttaggcca	gcttggcact	3480
tgatgtaatt	ctccttgga	tttgcccttt	ttgagtttgg	atcttggttc	attctcaagc	3540
ctcagacagt	ggttcaaa	gtttttcttc	catttcaggt	gtcgtgagga	attcgcgac	3600
gccatggacc	catcacagtt	taatccaacc	tacatcccag	ggtctccaca	aatgctcacc	3660
gaagaaaatt	cccgggacga	ttcagggggc	tctcaaatct	cctccgagac	gttgataaag	3720
aaccttagta	acttgactat	caacgctagt	agcgaatctg	tttcccctct	atcggaagct	3780
ttactccgtc	gagagtctgt	aggagcagca	gtcctcaggg	aaatcgaaga	tgagtggctt	3840
tacagcagga	gaggagtaag	aacattgctg	tctgtgcaga	gagaaaagat	ggcaagattg	3900
agatacatgt	tactcggcgg	agttcgtacg	catgaaagaa	gaccaacaaa	caaggagcct	3960
aaggaggtta	agaaggaatc	aagaccattc	aaatgtccct	gcagtttctg	cgtgtcta	4020
ggatgggato	cttctgagaa	tgctagaata	gggaatcaag	acaccaagcc	acttcagcca	4080
taataggttt	aaacatcgat	ggcctcgaga	aagcggccgc	ccccacgctg	aaactagttc	4140

-continued

gaaggatccg	catgcatcta	gggcggccaa	ttccgcccct	ctccctcccc	ccccctaac	4200
gttactggcc	gaagccgctt	ggaataaggc	cgggtgtcgt	ttgtctatat	gtgattttcc	4260
accatattgc	cgtcttttgg	caatgtgagg	gcccggaaac	ctggccctgt	cttcttgacg	4320
agcattccta	ggggctcttc	ccctctcgcc	aaaggaatgc	aaggctctgt	gaatgtcgtg	4380
aaggaagcag	ttcctctgga	agcttcttga	agacaaacaa	cgtctgtagc	gaccttttgc	4440
aggcagcgga	acccccacc	tggcgacagg	tgcctctcgc	gccaaaagcc	acgtgtataa	4500
gatacacctg	caaaggcggc	acaaccccag	tgccacgttg	tgagttggat	agttgtggaa	4560
agagtcaaat	ggctctcttc	aagcgtattc	aacaaggggc	tgaaggatgc	ccagaaggta	4620
ccccattgta	tgggatctga	tctggggcct	cgggtgcacat	gctttacatg	tgtttagtcg	4680
aggttaaaaa	aacgtctagg	ccccccgaac	cacggggacg	tggttttcct	ttgaaaaaca	4740
cgatgataag	cttgccacaa	cccacaagga	gacgaccttc	catgaccgag	tacaagccca	4800
cgggtgcgct	cgccaccgcg	gacgacgtcc	cccgggccgt	acgcaccctc	gccgcccgt	4860
tcgcccacta	ccccgccacg	cgccacaccg	tcgacccgga	ccgccacatc	gagcgggtca	4920
ccgagctgca	agaactcttc	ctcacgcgcg	tcgggctcga	catcgccaag	gtgtgggtcg	4980
cggacgacgg	cgccgcggtg	gcgggtctga	ccacgccgga	gagcgtcgaa	gcgggggcgg	5040
tgttcgccga	gatcgcccg	cgcacggccg	agttgagcgg	ttcccggctg	gccgcgcagc	5100
aacagatgga	aggcctctcg	gcgcgcgacc	ggcccaagga	gcccgcgtgg	ttcctggcca	5160
ccgtcggcgt	ctcgcccgac	caccagggca	agggtctggg	cagcgccgtc	gtgtcccccg	5220
gagtgagggc	ggccgagcgc	gccgggggtg	ccgccttcct	ggagacctcc	gcgccccgca	5280
acctccccct	ctacgagcgg	ctcggtctca	ccgtcaccgc	cgacgtcgag	tgcccgaagg	5340
accgcgcgac	ctgggtcatg	accgcgaagc	ccgggtccct	ataataggcg	gccgctcgag	5400
acctagaaaa	acatggagca	atcacaaagta	gcaatacagc	agctaccaat	gctgattgtg	5460
cctggctaga	agcacaagag	gaggaggagg	tgggttttcc	agtcacacct	caggtacctt	5520
taagaccaat	gacttacaag	gcagctgtag	atcttagcca	cttttttaaa	gaaaaggggg	5580
gactggaagg	gctaattcac	tcccaacgaa	gacaagatct	gctttttgct	tgtactgggt	5640
ctctctggtt	agaccagatc	tgagcctggg	agctctctcg	ctaactaggg	aacctactgc	5700
ttaagcctca	ataaagcttg	ccttgagtgc	ttcaagtagt	gtgtgccctg	ctgttgtgtg	5760
actctggtaa	ctagagatcc	ctcagaccct	tttagtcagt	gtggaaaatc	tctagcagta	5820
gtagttcatg	tcactcttatt	attcagttatt	tataacttgc	aaagaaatga	atatcagaga	5880
gtgagaggcc	ttgacattat	aatagattta	gcagggaattg	aactaggagt	ggagcacaca	5940
ggcaaagtcc	tagagctcgc	tgatcagcct	cgactgtgcc	ttctagttgc	cagccatctg	6000
ttgtttgccc	ctccccctg	ccttccttga	ccctggaagg	tgccactccc	actgtccctt	6060
cctaataaaa	tgaggaaatt	gcacgcgatt	gtctgagtag	gtgtcattct	attctggggg	6120
gtgggggtgg	gcaggacagc	aagggggagg	attgggaaga	caatagcagg	catgctgggg	6180
atgcggtggg	ctctatggct	tctgaggcgg	aaagaaccag	ctgggggcgc	gccctcgag	6240
gccgccatgg	tcatagctgt	ttgacgtcag	gtggcacttt	tcggggaaat	gtgcgcggaa	6300
ccccattttg	tttatttttc	taaatacatt	caaatatgta	tccgctcatg	agacaataac	6360
cctgataaat	gcttcaataa	tattgaaaaa	ggaagagtat	gagtattcaa	catttccgtg	6420
tcgcccttat	tccctttttt	gcggcatttt	gccttcctgt	ttttgctcac	ccagaaacgc	6480
tggtgaaagt	aaaagatgct	gaagatcagt	tgggtgcacg	agtggttac	atcgaactgg	6540

-continued

```

atctcaacag cggttaagatc cttgagagtt ttgcgccga agaactgttt ccaatgatga 6600
gcacttttaa agttctgcta tgtggcgccg tattatcccg tattgacgcc gggcaagagc 6660
aactcggctc cgcatacac tattctcaga atgacttggg tgagtactca ccagtcacag 6720
aaaagcatct tacggatggc atgacagtaa gagaattatg cagtgtgcc ataaccatga 6780
gtgataaac tgccggccaac ttacttctga caacgatcgg aggaccgaag gagctaaccg 6840
cttttttgca caacatgggg gatcatgtaa ctgcgcttga tcgttgggaa ccggagctga 6900
atgaagccat accaaacgac gagcgtgaca ccacgatgcc tgtagcaatg gcaacaacgt 6960
tgcgcaaact attaactggc gaactactta ctctagcttc ccggcaacaa ttaatagact 7020
ggatggaggc ggataaagtt gcaggaccac ttctgcgctc ggcccttcgg gctggctggt 7080
ttattgctga taaatctgga gccggtgagc gtgggtctcg cgggtatcatt gcagcactgg 7140
ggccagatgg taagccctcc cgtatcgtag ttatctacac gacggggagt caggcaacta 7200
tggatgaacg aaatagacag atcgttgaga taggtgcctc actgattaag cattggtaac 7260
tgtcagacca agtttactca tatatacttt agattgattt aaaacttcat ttttaattta 7320
aaaggatcta ggtgaagatc ctttttgata atctcatgac caaaatccct taacgtgagt 7380
tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa aggatcttct tgagatcctt 7440
ttttctcgcg cgtaactctc tgcttgcaaa caaaaaaacc accgctacca gcggtggttt 7500
gtttgccgga tcaagagcta ccaactcttt ttccgaaggt aactggcttc agcagagcgc 7560
agataccaaa tactgtcctt ctagtgtagc cgtagttagg ccaccacttc aagaactctg 7620
tagcacccgc tacatacctc gctctgctaa tcctgttacc agtggctgct gccagtggcg 7680
ataagtcgtg tcttaccggg ttggactcaa gacgatagtt accggataag gcgcagcggg 7740
cgggctgaac ggggggttcg tgcacacagc ccagcttggg gcgaacgacc tacaccgaac 7800
tgagatacct acagcgtgag cattgagaaa gcgccacgct tcccgaaggg agaaaggcgg 7860
acaggtatcc ggtaagcggc agggctcgaa caggagagcg cacgaggag cttccagggg 7920
gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca cctctgactt gagcgtcgat 7980
ttttgtgatg ctgcgcaggg gggcggagcc tatggaaaaa cgccagcaac gcggcctttt 8040
tacggttcct ggccttttgc                                     8060

```

```

<210> SEQ ID NO 7
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 7

```

```

cagtgcgccg aacccacac                                     19

```

```

<210> SEQ ID NO 8
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

```

```

<400> SEQUENCE: 8

```

```

agaggaaactg cttccttcac gaca                                     24

```

```

<210> SEQ ID NO 9

```

-continued

```

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

cagaaggcct cagcacctac                                20

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

tacctcttcc tcccactcca                                20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 11

aagcgcat caaaaggaga                                  20

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 12

agtttgtgcc aggggtttttg                              20

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 13

acttcacctt ccctccaacc                                20

```

We claim:

1. An enriched population of human pluripotent cells having a normal karyotype, wherein the human pluripotent cells comprise the genome of a single somatic cell of a postnatal individual human and further comprise an integrated non-native polynucleotide sequence encoding potency-determining factors comprising Oct-4 and Sox2.

2. The population of claim 1, wherein the potency-determining factors additionally comprise at least one of Nanog and Lin28.

3. The population of claim 1, wherein the population is obtained by exposing the somatic cell to a plurality of potency-determining factors under conditions sufficient to reprogram the somatic cell to a pluripotent cell.

4. The population of claim 3, wherein the potency-determining factors additionally comprise at least one of Nanog and Lin28.

5. The population of claim 3, wherein the exposing step comprises introducing a vector encoding one or more potency-determining factors into the somatic cell.

6. The population of claim 5, wherein the vector is a viral-based vector.

7. The population of claim 2, wherein the potency-determining factors are introduced to the somatic cell as a polynucleotide sequence comprising a nucleic acid sequence encoding a potency-determining factor operably linked to a heterologous promoter.

8. An enriched population of human pluripotent cells having a normal karyotype, comprising the genome of a somatic cell of a postnatal individual human, and further comprising integrated non-native polynucleotide sequences encoding potency-determining factors Oct-4, Sox2, and at least one of Nanog and Lin28.

50

55

60

65

9. A human pluripotent cell line derived from a somatic cell of a postnatal individual human, the somatic cell having a human somatic cell genome, by a process comprising

- (a) contacting the human somatic cell to a plurality of potency-determining factors comprising Oct4 and Sox2 under conditions sufficient to reprogram the human somatic cell to a pluripotent cell comprising the human somatic cell genome and further comprising an integrated non-native polynucleotide sequence encoding the plurality of potency-determining factors; and
- (b) culturing the pluripotent cell to obtain the human pluripotent cell line.

* * * * *