



US010144941B2

(12) **United States Patent**
Fox et al.(10) **Patent No.:** US 10,144,941 B2
(45) **Date of Patent:** Dec. 4, 2018(54) **METHOD AND COMPOSITIONS FOR IMPROVED LIGNOCELLULOSIC MATERIAL HYDROLYSIS**2008/0182249 A1 7/2008 Fox
2008/0286749 A1 11/2008 Fox
2010/0304405 A1 12/2010 Fox(71) Applicant: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

FOREIGN PATENT DOCUMENTS

(72) Inventors: **Brian Grant Fox**, Madison, WI (US); **Taichi Takasuka**, Madison, WI (US); **Adam Joel Book**, Madison, WI (US); **Cameron Robert Currie**, Madison, WI (US)WO 2008/028147 3/2008
WO 2008/127997 10/2008
WO 2009/064954 5/2009
WO 2010/141604 12/2010
WO 2010/141604 A2 12/2010(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

OTHER PUBLICATIONS

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 149 days.

(21) Appl. No.: **14/851,812**(22) Filed: **Sep. 11, 2015**(65) **Prior Publication Data**

US 2016/0032340 A1 Feb. 4, 2016

Related U.S. Application Data

(63) Continuation of application No. 13/709,971, filed on Dec. 10, 2012.

Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN08184, provided with alignment with SEQ ID No. 1.*

(60) Provisional application No. 61/579,897, filed on Dec. 23, 2011, provisional application No. 61/579,301, filed on Dec. 22, 2011.

Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN08183, provided with alignment with SEQ ID No. 2.*(51) **Int. Cl.**

C12N 1/20 (2006.01)
C12P 19/14 (2006.01)
C12N 9/42 (2006.01)
C12N 9/24 (2006.01)
C12P 19/02 (2006.01)
A23K 10/12 (2016.01)

(52) **U.S. Cl.**

CPC **C12P 19/14** (2013.01); **A23K 10/12** (2016.05); **C12N 1/20** (2013.01); **C12N 9/2434** (2013.01); **C12N 9/2437** (2013.01); **C12N 9/2491** (2013.01); **C12P 19/02** (2013.01); **C12Y 302/01004** (2013.01); **C12Y 302/01025** (2013.01); **C12Y 302/01091** (2013.01)

(58) **Field of Classification Search**

CPC C12P 19/14; C12P 19/02; A23K 10/12; C12N 1/20; C12N 9/2434; C12N 9/2491; C12N 9/2437; C12Y 302/01091; C12Y 302/01025; C12Y 302/01004

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,601,358 B2 10/2009 Fox
8,088,601 B2 1/2012 Fox
8,268,581 B2 9/2012 Fox

A method of digesting a lignocellulosic material is disclosed. In one embodiment, the method comprises the step of exposing the material to an effective amount of *Streptomyces* sp. ActE secretome such that at least partial lignocellulosic digestion occurs.

45 Claims, 70 Drawing Sheets
(25 of 70 Drawing Sheet(s) Filed in Color)
Specification includes a Sequence Listing.

(Continued)

Primary Examiner — Karen Cochrane Carlson*(74) Attorney, Agent, or Firm* — Quarles & Brady LLP(57) **ABSTRACT**

(56)

References Cited**OTHER PUBLICATIONS**

- Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN12581, provided with alignment with SEQ ID No. 13.*
- Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN13266, provided with alignment with SEQ ID No. 14.*
- Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN13437, provided with alignment with SEQ ID No. 15.*
- Lucas et al. Aug. 2011; Complete sequence of *Streptomyces* sp. SirexAA-E; Gen Bank Accession No. CP002993, EMBL AEN13760, provided with alignment with SEQ ID No. 16.*
- Adams, et al., Cellulose-degrading Bacteria Associated with the Invasive Woodwasp Sirex Noctilio, *The ISME Journal*, 2011, 5:1323-1331.
- Adav, et al., Quantitative iTRAQ Secretome Analysis of Celulolytic Thermobifida Fusca, *Journal of Proteome Research*, 2010, 9:3016-3024.
- Altschul, et al., Basic Local Alignment Search Tool, *J. Mol. Biol.*, 1990, 215:403-410.
- Anne, et al., *Streptomyces lividans* as Host for Heterologous Protein Production, *FEMS Microbiology Letters*, 1993, 114(2):121-128.
- Balan, et al., Lignocellulosic Biomass Pretreatment Using AFEX, *Methods in Molecular Biology*, 2009, 581:61-77.
- Baldrian, et al., Degradation of Cellulose by *Basidiomycetous* Fungi, *FEMS Microbiol. Rev.*, 2008, 32:501-521.
- Bayer, et al., From Cellulosomes to Cellulosomics, *The Chemical Record*, 2008, 8(6):364-377.
- Bergeron, et al., Putative Origin of Clonal Lineages of Amylostereum Areolatum, the Fungal Symbiont Associated with Sirex Noctilio, Retrieved from Pinus Sylvesteris, in Eastern Canada, *Fungal Biology*, 2011, 115(8):750-758.
- Bignell, et al., Isolation of Facultatively Aerobic Actinomycetes from the Gut, Parent Soil and Mound Materials of the Termites Proculitermes Aburiensis and Cubitermes Severus, *FEMS Microbiology Ecology*, 1991, 85:151-160.
- Blommel, et al., Flexi Vector Cloning, *Methods in Molecular Biology*, 2009, 498:55-73.
- Bradford, A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding, *Analytical Biochemistry*, 1976, 72(1-2):248-254.
- Cantarel, et al., The Carbohydrate-Active EnZymes Database (CAZy): An Expert Resource for Glycogenomics, *Nucleic Acids Research*, 2009, 37:D233-D238.
- Cereghino, et al., Production of Recombinant Proteins in Fermenter Cultures of the Yeast *Pichia Pastoris*, *Current Opinion in Biotechnology*, 2002, 13(4):329-332.
- Chater, et al., The Complex Extracellular Biology of *Streptomyces*, *FEMS Microbiol. Rev.*, 2010, 34:171-198.
- Chen, et al., Proteomic and Transcriptomic Analysis of Extracellular Proteins and mRNA Levels in Thermobifida Fusca Grown on Cellobiose and Glucose, *Journal of Bacteriology*, 2007, 189(17):6260-6265.
- Cheng, et al. Transition of Cellulose Crystalline Structure and Surface Morphology of Biomass as a Function of Ionic Liquid Pretreatment and Its Relation to Enzymatic Hydrolysis, *Biomacromolecules*, 2011, 12(4):933-941.
- Chundawat, et al., Restructuring the Crystalline Cellulose Hydrogen Bond Network Enhances Its Depolymerization Rate, *J. Am. Chem. Soc.*, 2011, 133(29):11163-11174.
- Crawford, Lignocellulose Decomposition by Selected *Streptomyces* Strains, *Applied and Environmental Microbiology*, 1978, 35(6):1041-1045.
- Deboy, et al., Insights into Plant Cell Wall Degradation from the Genome Sequence of the Soil Bacterium *Cellvibrio japonicus*, *Journal of Bacteriology*, 2008, 190(15):5455-5463.
- Deng, et al., Development and Application of a PCR-Targeted Gene Disruption Method for Studying CeIR Function in Thermobifida Fusca, *Applied and Environmental Microbiology*, 2010, 76(7):2098-2106.
- Eastwood, et al., The Plant Cell Wall-Decomposing Machinery Underlies the Functional Diversity of Forest Fungi, *Science*, 2011, 333:762-765.
- Forsberg, et al., Cleavage of Cellulose by a CBM33 Protein, *Protein Science*, 2011, 20:1479-1483.
- Galm, et al., Comparative Analysis of the Biosynthetic Gene Clusters and Pathways for Three Structurally Related Antitumor Antibiotics Bleomycin, Tallysomycin and Zorbamycin, *J. Nat Prod.*, 2011, 74(3):526-536.
- Goodfellow, et al., Ecology of Actinomycetes, *Annual Review of Microbiology*, 1983, 37:189-216.
- Herpoel-Gimbert, et al., Comparative Secretome Analyses of Two Trichoderma Reesei RUT-C30 and CL847 Hypersecretory Strains, *Biotechnology for Biofuels*, 2008, 1:18, 12 pages.
- Hess, et al., Metagenomic Discovery of Biomass-Degrading Genes and Genomes from Cow Rumen, *Science*, 2011, 331:463-467.
- Hulcr, et al., Presence and Diversity of *Streptomyces* in Dendroctonus and Sympatric Bark Beetle Galleries Across North America, *Microb. Ecol.*, 2011, 61:759-768.
- Hyatt, et al., Prodigal: Prokaryotic Gene Recognition and Translation Initiation Site Identification, *BMC Bioinformatics*, 2010, 11:119.
- Ishaque, et al., Cellulase Complex of a Mesophilic *Streptomyces* Strain, *Canadian Journal of Microbiology*, 1980, 26(2):183-189.
- Jarboe, et al., Metabolic Engineering for Production of Biorenewable Fuels and Chemicals: Contributions of Synthetic Biology, *Journal of Biomedicine and Biotechnology*, vol. 2010, Article ID 761042, 18 pages.
- Kestler, et al., VennMaster: Area-Proportional Euler Diagrams for Functional GO Analysis of Microarrays, *BMC Bioinformatics*, 2008, 9:67, 12 pages.
- Klepzig, et al., Symbioses: A Key Driver of Insect Physiological Processes, Ecological Interactions, Evolutionary Diversification, and Impacts on Humans, *Environ. Entomol.*, 2009, 38(1):67-77.
- Kukor, et al., Acquisition of Digestive Enzymes by Siricid Woodwasps from Their Fungal Symbiont, *Science*, 1983, 220:1161-1163.
- Langston, et al. Oxidoreductive Cellulose Depolymerization by the Enzymes Cellobiose Dehydrogenase and Glycoside Hydrolase 61, *Applied and Environmental Microbiology*, 2011, 77(19):7007-7015.
- Li et al. Influence of Physico-Chemical Changes on Enzymatic Digestibility of Ionic Liquid and AFEX Pretreated Corn Stover, *Bioresource Technology*, 2011, 102(13):6928-6936.
- Luyten, et al., Extensive Variation in Intracellular Symbiont Community Composition Among Members of a Single Population of the Wood-Boring Bivalve *Lyrodus Pedicellatus* (Bivalvia: Teredinidae), *Applied and Environmental Microbiology*, 2006, 72(1):412-417.
- Lykidis, et al., Genome Sequence and Analysis of the Soil Celulolytic Actinomycete Thermobifida Fusca YX, *Journal of Bacteriology*, 2007, 189(6):2477-2486.
- Lynd, et al., Microbial Cellulose Utilization: Fundamentals and Biotechnology, *Microbiology and Molecular Biology Reviews*, 2002, 66(3):506-577.
- Marushima, et al., CebR as a Master Regulator for Cellulose/Celooligosaccharide Catabolism Affects Morphological Development in *Streptomyces griseus*, *Journal of Bacteriology*, 2009, 191(19):5930-5940.
- McCarthy, et al., Actinomycetes as Agents of Biodegradation in the Environment—A Review, *Gene*, 1992, 115(1-2):189-192.
- Medema, et al., antiSMASH: Rapid Identification, Annotation and Analysis of Secondary Metabolite Biosynthesis Gene Clusters in Bacterial and Fungal Genome Sequences, *Nucleic Acids Research*, 2011, 39:W339-W346.
- Merino, et al., Progress and Challenges in Enzyme Development for Biomass Utilization, *Adv. Biochem. Engin. Biotechnol.*, 2007, 108:95-120.
- Miller, Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar, *Analytical Chemistry*, 1959, 31(3):426-428.
- Natale, et al., Sec- and Tat-mediated Protein Secretion Across the Bacterial Cytoplasmic Membrane—Distinct Translocases and Mechanisms, *Biochimica et Biophysica Acta*, 2008, 1778:1735-1756.

(56)

References Cited**OTHER PUBLICATIONS**

- Nijland, et al., Optimization of Protein Secretion by *Bacillus Subtilis*, Recent Patents on Biotechnology, 2008, 2(2):79-87.
- Pasti, et al., Cellulolytic Activity of Actinomycetes Isolated from Termites (*Termitidae*) Gut, FEMS Microbiology Letters, 1985, 26(1):107-112.
- Pasti, et al., Lignin-Solubilizing Ability of Actinomycetes Isolated from Termite (*Termitidae*) Gut, Applied and Environmental Microbiology, 1990, 56(7):2213-2218.
- Peralta-Yahya, et al., Advanced Biofuel Production in Microbes, Biotechnol., J., 2010, 5:147-162.
- Quinlan, et al., Insights into the Oxidative Degradation of Cellulose by a Copper Metalloenzyme that Exploits Biomass Components, PNAS, 2011, 108(37):15079-15084.
- Raman, et al., Impact of Pretreated Switchgrass and Biomass Carbohydrates on Clostridium Thermocellum ATCC 27405 Cellulosome Composition: A Quantitative Proteomic Analysis, PLoS One, 2009, 4(4):e5271, 13 pages.
- Adams, A. S. et al. ISME Journal 5, 1323-1331 (2011).
- Takasuka, T.E., et al. Scientific Report, 3, 1-10 (2013).
- Adam, A., et al. Cellulolytic associations of *Sirex noctilio* within the context of a multipartite symbiosis. Entomological Society of America Meeting (Nov. 16, 2008).
- Adam, A., et al. Cellulose-degrading microbial symbionts of the woodwasp, *Sirex noctilio*. The GLBRC 2008 Retreat Poster Session Abstracts (Oct. 15-18, 2008).
- Riederer, et al., Global Gene Expression Patterns in Clostridium Thermocellum as Determined by Microarray Analysis of Chemostat Cultures on Cellulose or Cellobiose, Applied and Environmental Microbiology, 2011, 77(4):1243-1253.
- Rutherford, et al., Artemis: Sequence Visualization and Annotation, Bioinformatics, 2000, 16(10):944-945.
- Saloheimo, et al., The Cargo and the Transport System: Secreted Proteins and Protein Secretion in *Trichoderma Reesei* (*Hypocrea Jeckerina*), Microbiology, 2012, 158:46-57.
- Schafer, et al., Hemicellulose-Degrading Bacteria and Yeasts from the Termite Gut, Journal of Applied Bacteriology, 1996, 80(5):471-478.
- Scharf, et al., Multiple Levels of Synergistic Collaboration in Termite Lignocellulose Digestion, PLoS ONE, 2011, 6(7):e21709, 7 pages.
- Schlatter, et al., Resource Amendments Influence Density and Competitive Phenotypes of *Streptomyces* in Soil, Microb. Ecol., 2009, 57:413-420.
- Schlochtermeier, et al., Biochemical and Electron Microscopic Studies of the *Streptomyces reticuli* Cellulase (Avicelase) in Its Mycelium-Associated and Extracellular Forms, Applied and Environmental Microbiology, 1992, 58(10):3240-3248.
- Schlochtermeier, et al., The Gene Encoding the Cellulase (Avicelase) Cell From *Streptomyces reticuli* and Analysis of Protein Domains, Molecular Microbiology, 1992, 6(23):3611-3621.
- Schuster, et al., Biology and Biotechnology of *Trichoderma*, Appl. Microbiol. Biotechnol., 2010, 87:787-799.
- Semedo, et al., *Streptomyces drozdowiczii* sp. nov., a Novel Cellulolytic *Streptomyces* from Soil in Brazil, International Journal of Systematic and Evolutionary Microbiology, 2004, 54:1323-1328.
- Shannon, et al., Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks, Genome Research, 2003, 13:2498-2504.
- Suen, et al., An Insect Herbivore Microbiome with High Plant Biomass-Degrading Capacity, PLoS Genetics, 2010, 6(9):e1001129, 14 pages.
- Susi, et al., Biological Control of Wood Decay Against Fungal Infection, Journal of Environmental Management, 2011, 92(7):1681-1689.
- Tabata, et al., Molecular Phylogeny of Species in the Genera *Amylostereum* and *Echinodontium*, Mycoscience, 2000, 41:585-593.
- Teather, et al., Use of Congo Red-Polysaccharide Interactions in Enumeration and Characterization of Cellulolytic Bacteria from the Bovine Rumen, Applied and Environmental Microbiology, 1982, 43(4):777-780.
- Tian, et al. Comparisons of SPORL and Dilute Acid Pretreatments for Sugar and Ethanol Productions from Aspen, Biotechnol. Prog., 2011, 27(2):419-427.
- Tolonen, et al., Proteome-Wide Systems Analysis of a Cellulosic Biofuel-Producing Microbe, Molecular Systems Biology, 2011, 7:461, 12 pages.
- Trop, et al., The Specificity of Proteinases from *Streptomyces griseus* (Pronase), Biochem. J., 1970, 116:19-25.
- Turini, et al., The Action of Phenylmethylsulfonyl Fluoride on Human Acetylcholinesterase, Chymotrypsin and Trypsin, Journal of Pharmacology and Experimental Therapeutics, 1969, 167(1):98-104.
- Vaaje-Kolstad, An Oxidative Enzyme Boosting the Enzymatic Conversion of Recalcitrant Polysaccharides, Science, 2010, 330:219-222.
- Vuong, et al., Glycoside Hydrolases: Catalytic Base/Nucleophile Diversity, Biotechnology and Bioengineering, 2010, 107(2):195-205.
- Wachinger, et al., Identification of Mycelium-Associated Cellulase from *Streptomyces reticuli*, Applied and Environmental Microbiology, 1989, 55(10):2653-2657.
- Walter, et al., Physiological Studies of Cellulase (Avicelase) Synthesis in *Streptomyces reticuli*, Applied and Environmental Microbiology, 1996, 62(3):1065-1069.
- Walter, et al., The Synthesis of the *Streptomyces reticuli* Cellulase (Avicelase) is Regulated by Both Activation and Repression Mechanisms, Mol. Gen. Genet., 1996, 251:186-195.
- Wang, et al., Sulfite Pretreatment to Overcome Recalcitrance of Lignocellulose (SPORL) for Robust Enzymatic Saccharification of Hardwoods, Biotechnol. Prog., 2009, 25(4):1086-1092.
- Wilson, Microbial Diversity of Cellulose Hydrolysis, Current Opinion in Microbiology, 2011, 14:1-5.
- Wilson, Biochemistry and Genetics of Actinomycete Cellulases, Critical Reviews in Biotechnology, 1992, 12(1-2):45-63.
- Wood, et al., Production of Recombinant Bacterial Endoglucanase as a Co-Product with Ethanol During Fermentation Using Derivatives of *Escherichia coli* KO11, Biotechnology and Bioengineering, 1997, 55(3):547-555.
- Zhang, et al., Extracellular Accumulation of Recombinant Proteins Fused to the Carrier Protein YebF in *Escherichia coli*, Nature Biotechnology, 2006, 24(1):100-104.
- Zylstra, et al., Sirex Noctilio in North America: The Effect of Stem-Injection Timing on the Attractiveness and Suitability of Trap Trees, Agricultural and Forest Entomology, 2010, 12:243-250.
- Adams, et al., Cellulolytic Associations of *Sirex Noctilio* Within the Context of a Multipartite Symbiosis, Department of Entomology, University of Wisconsin, Poster, 2008, 23 pages.
- Book et al. "Cellulolytic *Streptomyces* strains associated with herbivorous insects share a phylogenetically linked capacity to degrade lignocellulose," Appl. Environ. Microbiol. 2014, 80(15):4692.

* cited by examiner




Figure 1A-B




Figure 2A




Figure 2B




Figure 2C

Figure 3A-B





Figure 4






Figure 5A-B




Figure 6

Figure 7A





Figure 7B






Figure 8A-B

Figure 9




Figure 10




Figure 11




Figure 12




Figure 13





Figure 14






Figure 15A-B

Figure 17

Protein band ^a	Locus	Catalytic domain	CBM	Functional class ^b
a	SACTE_3159	CBM33	CBM2	cellulase
b	SACTE_0265	GH10	CBM2	xylanase
c	SACTE_4755	GH64		beta-1,3-glucanase
d	SACTE_0482	GH5	CBM2	cellulase
e	SACTE_0237	GH6	CBM2	cellulase
f	SACTE_0236	GH48	CBM2	cellulase
g	SACTE_3717	GH9	CBM2	cellulase
h	SACTE_2347	GH5	CBM2	mannanase

^a Protein bands labeled in Figure 3A were identified by MALDI-TOF mass spectrometry. ^b Function identified by assays of individual fractions from ion exchange chromatography.

Figure 18

Figure 18: Spectra count of proteins identified on each substrate, where top 95 % spectra covered were highlighted green, light purple, purple, blue, orange, pink, light blue and yellow on glucose, cellobiose, cellulose, xylan switchgrass, AFEX-SG, IL-SG and chitin, respectively.

Locus	CAZy	Identified Proteins (414)	glucose	cellobiose	cellulose	xylan	switchgrass	AFEX-SG	IL-SG	chitin
SACTE_0237	GH16	ACTE_1, 4-beta cellobiohydrolase	0	263	3895	8	3390	3300	3520	0
SACTE_0236	GH48	ACTE_Glycoside hydrolase, 48F	0	24	3296	0	577	928	1043	0
SACTE_2347	GH5,CE3	ACTE_Cellulose-binding family II/chitobiase, carbohydrate-binding domain	0	28	224	0	225	249	205	0
SACTE_3159	CBM33,2	ACTE_Cellulose-binding domain, family II, bacterial type	0	4	564	0	493	646	847	8
SACTE_0482	GH5	ACTE_Cellulose-binding family II/chitobiase, carbohydrate-binding domain	0	13	156	0	32	113	153	0
SACTE_0265	GH10	ACTE_Glycoside hydrolase, family 10	0	23	20	0	118	320	204	8
SACTE_0357	CE4	ACTE_Glycoside hydrolase/deacetylase, beta/alpha-barrel	0	13	87	0	34	195	190	3
SACTE_4439		ACTE_Catalase, N-terminal	0	55	82	0	20	40	40	271
SACTE_0562	GH74	ACTE_Cellulose-binding family II/chitobiase, carbohydrate-binding domain	0	0	22	0	29	48	30	0
SACTE_0358	GH11	ACTE_Glycoside hydrolase, family 11, active site	0	0	37	0	165	324	119	7
SACTE_4343		ACTE_Bacterial extracellular solute-binding protein, family 5	0	204	93	0	52	43	30	19
SACTE_1546		ACTE_Bacterioferritin	0	317	32	0	32	22	24	35
SACTE_1310	PL3	ACTE_Pectate lyase, catalytic	0	57	26	0	24	98	48	25
SACTE_4638		ACTE_Chondroitin AC/alginate lyase	0	31	23	0	34	59	59	14
SACTE_5668		ACTE_Alpha/beta hydrolase fold-1	0	29	20	0	0	0	9	39
SACTE_3717	GH9	ACTE_Carbohydrate-binding, CenC-like	0	49	21	0	382	437	143	0
SACTE_3590		ACTE_Phospholipase C, phosphatidylinositol-specific , X domain	0	5	23	0	0	4	7	0
SACTE_5978	PL1	ACTE_Galactose-binding domain-like	0	5	20	0	187	295	10	0
SACTE_2172		ACTE_Citrate synthase-like, core	0	26	26	0	47	12	33	46
SACTE_4571	GH18	ACTE_EF-Hand 1, calcium-binding site	0	8	25	5	28	14	10	1758
SACTE_6428	CBM33	ACTE_Chitin-binding, domain 3	0	3	20	0	6	5	8	0
SACTE_2313	CBM33	ACTE_Chitin-binding, domain 3	0	11	19	0	12	0	11	308
SACTE_0366	GH78	ACTE_Six-hairpin glycosidase-like	0	4	27	0	0	0	5	20
SACTE_1604		ACTE_Pyridine nucleotide-disulphide oxidoreductase, class I, active site	0	472	16	0	22	18	26	57
SACTE_4702		ACTE_Protein of unknown function DUF756	0	26	16	0	2	26	26	26
SACTE_2059		ACTE_Manganese/iron superoxide dismutase, C-terminal	0	63	13	6	33	16	0	62
SACTE_2556		ACTE_Enolase, N-terminal	0	22	84	10	10	4	19	0
SACTE_4730	GH30	ACTE_Ricin B lectin	0	88	12	1	32	25	53	7
SACTE_4755	GH64	ACTE_Twin-arginine translocation pathway, signal sequence	2	24	12	0	28	40	40	24
SACTE_4945		ACTE_Extracellular solute-binding protein, family 3	0	28	28	0	4	22	9	9
SACTE_1041		ACTE_ABC-type glycine betaine transport system, substrate-binding domain	0	21	11	8	6	6	10	17
SACTE_0880		ACTE_S-adenosylmethionine synthetase, central domain	0	15	10	6	3	0	33	33
SACTE_4246	GH18	ACTE_Galactose-binding domain-like	0	39	10	7	9	11	11	1115
SACTE_0464	GH16	ACTE_Ricin B lectin	0	7	10	6	4	8	9	9
SACTE_5330		ACTE_Peptidase S1/S6, chymotrypsin/Hap	0	43	10	6	9	11	11	14
SACTE_1130		ACTE_Galactose-binding domain-like	0	43	9	0	11	6	0	0
SACTE_2232	GH3	ACTE_Glycoside hydrolase, family 3, C-terminal	0	22	9	0	0	0	0	9
SACTE_5335		ACTE_ABC transporter, substrate-binding protein, aliphatic sulphonates	0	28	9	4	11	9	9	35
SACTE_0132		ACTE_hypothetical protein	0	0	9	0	6	0	6	0
SACTE_2289		ACTE_Bacterial extracellular solute-binding, family 1	0	13	9	0	0	7	0	0
SACTE_5230		ACTE_Xylose isomerase-like, TIM barrel domain	0	45	8	0	150	183	348	6
SACTE_5457	GH46	ACTE_Glycosidase hydrolase, family 46	0	461	8	0	21	29	12	57
SACTE_5847	GH87	ACTE_Galactose-binding domain-like	0	26	8	0	75	63	26	0
SACTE_1422	GH23	ACTE_Twin-arginine translocation pathway, signal sequence	0	19	8	0	0	2	19	0
SACTE_3306		ACTE_6-phosphogluconate dehydrogenase related protein	0	46	8	0	11	9	9	18
SACTE_3803		ACTE_Pyridoxal phosphate-dependent transferase, major domain	0	265	8	0	11	0	23	0
SACTE_4676		ACTE_Bacterial extracellular solute-binding protein, family 5	0	25	8	0	11	9	0	6
SACTE_0383	GH92	ACTE_Alpha-1,2-mannosidase, putative	0	0	8	8	0	0	0	0
SACTE_0642		ACTE_Peptidase S33, tripeptidyl-peptidase C-terminal	2	24	8	0	35	0	0	0
SACTE_3779	GH31	ACTE_Glycoside hydrolase, family 31	0	5	8	0	0	0	0	0

Figure 18 (continued)

SACTE_5166	GH43	ACTE_Twin-arginine translocation pathway, signal sequence	0	82	7
SACTE_4363	GH55	ACTE_hypothetical protein	0	72	7
SACTE_1369		ACTE_Glyceraldehyde 3-phosphate dehydrogenase, NAD(P) binding domain	33	326	7
SACTE_0746		ACTE_Twin-arginine translocation pathway, signal sequence	0	0	7
SACTE_4468		ACTE_Bacterial extracellular solute-binding, family 1	0	4	7
SACTE_1364		ACTE_Phosphoglucose isomerase, conserved site	2	347	6
SACTE_5519		ACTE_YD repeat	9	0	6
SACTE_4612		ACTE_hypothetical protein	0	8	5
SACTE_0244		ACTE_N-acetylumamoy-L-alanine amidase, family 2	6	27	5
SACTE_4198		ACTE_NAD(P)-binding domain	4	66	4
SACTE_3458		ACTE_Surface protein from Gram-positive cocci	8	21	4
SACTE_1367		ACTE_Triosephosphate isomerase	0	351	4
SACTE_0379	GH2	ACTE_Galactose-binding domain-like	0	72	4
SACTE_2033		ACTE_Nucleoside diphosphate kinase, core	0	7	4
SACTE_5371		ACTE_Neuraminidase	0	4	4
SACTE_5629	GH93	ACTE_Ricin B lectin	0	88	4
SACTE_5589		ACTE_Gamma-glutamyltranspeptidase	9	33	3
SACTE_4958	GH18	ACTE_Galactose-binding domain-like	0	37	3
SACTE_4231		ACTE_Serine/cysteine peptidase, trypsin-like	0	4	3
SACTE_5240		ACTE_hypothetical protein	0	36	3
SACTE_0844	GH18	ACTE_Glycoside hydrolase, subgroup, catalytic core	0	19	3
SACTE_0604	GH109	ACTE_NAD(P)-binding domain	3	2	3
SACTE_5267	GH43	ACTE_Twin-arginine translocation pathway, signal sequence	2	2	3
SACTE_6439	GH43	ACTE_Glycoside hydrolase, family 43	0	66	2
SACTE_4727		ACTE_Isocitrate/isopropylmalate dehydrogenase	0	47	2
SACTE_1302	PL1	ACTE_Twin-arginine translocation pathway, signal sequence	0	8	2
SACTE_3711		ACTE_Glycosyl transferase, family 20	0	23	2
SACTE_6549		ACTE_Peptidase S11, D-alanyl-D-alanine carboxypeptidase A	0	0	2
SACTE_1859		ACTE_Bacterial stress protein	9	395	0
SACTE_3700		ACTE_Bacterial stress protein	383	207	0
SACTE_3078		ACTE_Ketose-bisphosphate aldolase, class-II	0	314	0
SACTE_3038		ACTE_Bacterial stress protein	174	253	0
SACTE_2065		ACTE_Peptidase S8/553, subtilisin/kevin/sedolisin	33	4230	0
SACTE_1949		ACTE_Peptidase M4, thermolysin	0	364	0
SACTE_1701		ACTE_Htaa	4	78	0
SACTE_1250		ACTE_Peptidase S8/553, subtilisin, active site	0	8	0
SACTE_0528	GH115	ACTE_Twin-arginine translocation pathway, signal sequence	0	0	0
		ACTE_Aconitase/3-isopropylmalate dehydratase large subunit, alpha/beta/alpha, subdomain 1/3	45	28	0
SACTE_5220		ACTE_Heat shock protein DnaJ-like protein djlA	441	0	0
SACTE_3197		ACTE_Exoribonuclease, phosphorolytic domain 2	9	13	0
SACTE_4908		ACTE_Glycoside hydrolase, family 31	0	2	0
SACTE_5231	GH31	ACTE_Glycoside hydrolase, family 19, catalytic	0	23	0
SACTE_3064	GH19	ACTE_Peptidoglycan binding-like	0	25	0
SACTE_4515	GH25	ACTE_Twin-arginine translocation pathway, signal sequence	0	35	0
SACTE_5418	GH23	ACTE_Peptidoglycan binding-like	0	3	0
SACTE_4493		ACTE_Superoxide dismutase, Nickel-type	37	30	0
SACTE_0081	GH19	ACTE_Glycoside hydrolase, family 19	0	30	0
SACTE_0634		ACTE_Periplasmic binding protein	0	3	0
SACTE_4669		ACTE_Pyridoxal phosphate-dependent transferase, major domain	32	24	0
SACTE_4483		ACTE_Cyclic nucleotide-binding-like	0	0	0
SACTE_6308		ACTE_Periplasmic binding protein	9	25	0
SACTE_4078		ACTE_MGS-like	32	7	0
SACTE_0926		ACTE_Vitamin B6 biosynthesis protein	35	33	0
SACTE_4607		ACTE_Protein of unknown function DUF1557	0	19	0




Figure 18 (continued)

SACTE_4178		ACTE_FAD-dependent pyridine nucleotide-disulphide oxidoreductase	0	0	0	0	0	0	0	0	0	0
SACTE_3666		ACTE_Histidine phosphatase superfamily, clade-1	0	0	0	0	0	0	0	0	0	0
SACTE_1858		ACTE_Bacterial stress protein	0	25	0	0	0	0	0	0	19	0
SACTE_5630		ACTE_Aldo/keto reductase	0	0	0	0	0	0	0	0	0	0
SACTE_5751		ACTE_Twin-arginine translocation pathway, signal sequence	0	0	0	0	0	0	0	0	0	0
SACTE_1603		ACTE_Peptidase M17, leucyl aminopeptidase, C-terminal	0	20	0	0	0	0	0	0	0	0
SACTE_3452		ACTE_Cobaltochelatase, CobN subunit	32	0	0	0	0	0	0	0	0	0
SACTE_4459		ACTE_Peptidoglycan binding-like	0	0	0	0	0	0	0	0	0	0
SACTE_5682		ACTE_Galactose-binding domain-like	0	24	0	0	0	0	0	0	0	0
SACTE_1239		ACTE_Enoyl-[acyl-carrier-protein] reductase (NADH)	0	0	0	0	0	0	0	0	0	0
SACTE_1356		ACTE_Transaldolase, active site	0	6	0	0	0	0	0	0	0	0
SACTE_2768		ACTE_Surface protein from Gram-positive cocci	0	13	0	0	0	0	0	0	54	0
SACTE_3335		ACTE_Single-strand DNA-binding	0	25	0	0	0	0	0	0	0	0
SACTE_0169		ACTE_Glyceraldehyde 3-phosphate dehydrogenase, active site	0	31	0	0	0	0	0	0	0	0
SACTE_6558		ACTE_Quinoprotein amino dehydrogenase, beta chain-like	8	16	0	0	0	0	0	0	0	0
SACTE_1136		ACTE_Peptidoglycan recognition protein	3	1	0	0	0	0	0	0	0	0
SACTE_1434		ACTE_S'-Nucleotidase/apyrase	32	24	0	0	0	0	0	0	0	0
SACTE_0365		ACTE_Peptidase S8/S53, subtilisin/kexin/sedolisin	0	35	0	0	0	0	0	0	0	0
SACTE_2585		ACTE_Phosphoribosyltransferase	8	28	0	0	0	0	0	0	7	4
SACTE_4738	GH16	ACTE_Galactose-binding domain-like	0	338	0	0	0	0	0	0	3	0
SACTE_4102		ACTE_Fumarate reductase/succinate dehydrogenase, FAD-binding site	20	12	0	0	0	0	0	0	0	0
SACTE_1680		ACTE_Cupredoxin	0	3	0	0	0	0	0	0	0	0
SACTE_1003		ACTE_NAD(P)-binding domain	0	0	0	0	0	0	0	0	0	0
SACTE_5263		ACTE_Dimeric alpha-beta barrel	3	35	0	0	9	2	0	0	7	0
SACTE_0549	GH16	ACTE_Concanavalin A-like lectin/glucanase	0	14	0	0	9	0	0	0	0	0
SACTE_1325		ACTE_Dihydrodipicolinate synthetase	0	0	0	0	9	0	0	0	0	0
SACTE_1324		ACTE_hypothetical protein	0	0	0	0	9	0	0	0	0	0
SACTE_2068		ACTE_Peptidase M1, aminopeptidase N actinomycete-type	8	18	0	0	8	0	0	0	0	0
SACTE_4038		ACTE_Cystathione beta-synthase, core	8	0	0	0	8	0	0	0	0	0
SACTE_0222		ACTE_Peptidase M4, propeptide, PepSY	0	23	0	0	8	0	0	0	0	0
SACTE_1281		ACTE_Surface protein from Gram-positive cocci	0	3	0	0	8	0	0	0	0	0
SACTE_3389		ACTE_Peptidase M24B, X-Pro dipeptidase/aminopeptidase P, conserved site	8	26	0	0	7	6	2	3	11	0
SACTE_3164		ACTE_hypothetical protein	0	0	0	0	7	2	0	0	7	0
SACTE_5880		ACTE_Twin-arginine translocation pathway, signal sequence	0	18	0	0	7	22	0	0	0	0
SACTE_4081		ACTE_NAD(P)-binding domain	33	7	0	0	7	0	0	0	0	0
SACTE_2819		ACTE_Chaperonin ClpA/B, conserved site	8	0	0	0	7	0	0	0	0	0
SACTE_3589		ACTE_S'-Nucleotidase/apyrase	0	0	0	0	7	0	0	0	0	0
SACTE_2062		ACTE_DSBA oxidoreductase	0	22	0	0	6	5	0	0	51	0
SACTE_5859		ACTE_hypothetical protein	0	13	0	0	6	16	27	31	0	0
SACTE_5144		ACTE_Twin-arginine translocation pathway, signal sequence	0	2	0	0	6	0	0	0	0	0
SACTE_4739	GH3	ACTE_Na-Ca exchanger/integrin-beta4	0	1	0	0	6	0	0	0	0	0
SACTE_4768		ACTE_Glutamyl/glutaminyl-tRNA synthetase, class Ic, catalytic domain	0	0	0	0	6	0	0	0	0	0
SACTE_2049		ACTE_Ribose 5-phosphate isomerase, actinobacteria	32	18	0	0	5	0	0	0	7	0
SACTE_0639		ACTE_Formyl transferase, N-terminal	0	53	0	0	5	0	0	0	4	0
SACTE_0324	GH55	ACTE_Galactose-binding domain-like	0	3	0	0	5	4	0	8	2	0
SACTE_5685	GH13	ACTE_Glycoside hydrolase, subgroup, catalytic core	0	16	0	0	5	24	5	0	0	0
SACTE_0364	GH87	ACTE_Galactose-binding domain-like	0	0	0	0	5	9	4	0	0	0
SACTE_1312	Pl1	ACTE_Pectate lyase/Amb allergen	0	0	0	0	5	4	3	0	0	0
SACTE_3227		ACTE_Peptidase M18, aminopeptidase I	0	24	0	0	5	2	0	0	0	0
SACTE_4960		ACTE_Aminotransferase class-III	32	5	0	0	5	0	0	0	0	0
SACTE_0687		ACTE_Protein of unknown function DUF885, bacterial	7	0	0	0	5	0	0	0	0	0
SACTE_3765		ACTE_hypothetical protein	0	4	0	0	5	0	0	0	0	0
SACTE_4039		ACTE_IMP dehydrogenase/GMP reductase	6	3	0	0	4	0	0	2	12	0

Figure 18 (continued)

SACTE_5260		ACTE_FAD-dependent pyridine nucleotide-disulphide oxidoreductase	0	31	0	4	0	0	0	0	9
SACTE_4145		ACTE_Basic membrane lipoprotein	0	43	0	4	2	0	0	4	8
SACTE_1137		ACTE_Peptidase S45, penicillin amidase	0	0	0	4	9	4	0	0	2
SACTE_1715		ACTE_Intermediate filament, C-terminal	0	35	0	4	0	0	0	0	0
SACTE_0325		ACTE_Bacterial stress protein	0	7	0	4	0	0	0	0	0
SACTE_5146		ACTE_FMN-binding split barrel, related	0	0	0	4	0	0	0	0	0
SACTE_4910		ACTE_Dihydripicolinate reductase, N-terminal	0	0	0	4	0	0	0	0	0
SACTE_0213		ACTE_Luciferase-like	0	0	0	4	0	0	0	0	0
SACTE_1311	CE8	ACTE_Pectinesterase, catalytic	0	0	0	4	0	0	0	0	0
SACTE_1313	CE12	ACTE_Lipase, GDSL	0	0	0	4	0	0	0	0	0
SACTE_5657		ACTE_MaoC-like dehydratase	0	34	0	3	4	0	0	0	3
SACTE_4436		ACTE_Uncharacterised protein family UPF0182	0	9	0	3	0	0	0	0	3
SACTE_4718		ACTE_Delta-1-pyrroline-5-carboxylate dehydrogenase 1	3	0	0	3	28	4	5	0	0
SACTE_5482		ACTE_Aldo/keto reductase, conserved site	0	4	0	3	0	0	4	0	0
SACTE_5606		ACTE_D-hydantoinase	0	1	0	3	3	2	0	0	0
SACTE_1995		ACTE_Gamma-glutamyl phosphate reductase GPR	0	0	0	3	2	0	0	0	0
SACTE_4618		ACTE_Phosphofructokinase, pyrophosphate dependent	0	0	0	3	0	0	0	0	0
SACTE_1240		ACTE_NAD(P)-binding domain	3	0	0	3	0	0	0	0	0
SACTE_6512	CBM13	ACTE_Cadherin-like	0	35	0	3	0	0	0	0	0
SACTE_5114		ACTE_Ornithine carbamoyltransferase	0	1	0	3	0	0	0	0	0
SACTE_2698		ACTE_hypothetical protein	0	1	0	3	0	0	0	0	0
SACTE_6206		ACTE_hypothetical protein	0	0	0	3	0	0	0	0	0
SACTE_4914		ACTE_Ribosomal protein L19	0	0	0	3	0	0	0	0	0
SACTE_2318		ACTE_HAD-superfamily hydrolase, subfamily II A	0	0	0	3	0	0	0	0	0
SACTE_1862		ACTE_Transketolase, C-terminal/Pyruvate-ferredoxin oxidoreductase, domain II	0	0	0	3	0	0	0	0	0
SACTE_4083		ACTE_Malate dehydrogenase, active site	36	18	0	2	0	0	0	0	19
SACTE_2645		ACTE_Penicillin/cephalosporin acylase	0	33	0	2	0	0	0	0	2
SACTE_1895		ACTE_Mandelate racemase/muconate lactonizing enzyme, N-terminal	0	0	0	2	9	6	3	0	0
SACTE_5740		ACTE_Twin-arginine translocation pathway, signal sequence	0	0	0	2	3	0	0	0	0
SACTE_0133	CBM35.6	ACTE_Galactose-binding domain-like	0	0	0	2	3	0	0	0	0
SACTE_5493		ACTE_Periplasmic binding protein/LacI transcriptional regulator	0	12	0	2	0	0	0	0	0
SACTE_1002		ACTE_Twin-arginine translocation pathway, signal sequence	0	4	0	2	0	0	0	0	0
SACTE_2381		ACTE_Exoribonuclease, phosphorolytic domain 2	0	1	0	2	0	0	0	0	0
SACTE_5764	GH18	ACTE_Carbohydrate-binding domain, family 5/12	0	3	0	0	0	0	0	0	365
SACTE_6494	GH18	ACTE_Glycoside hydrolase, chitinase active site	0	0	0	0	0	0	0	0	69
SACTE_0860	GH18	ACTE_Carbohydrate-binding domain, family 5/12	0	0	0	0	0	0	0	0	44
SACTE_3097		ACTE_Heat shock protein Hsp70	39	4	0	0	0	0	0	0	25
SACTE_2384		ACTE_Phosphotransferase system, Etc component, type 1	0	0	0	0	0	0	0	0	19
SACTE_3685		ACTE_Molybdate/tungstate binding	0	7	0	0	4	0	0	0	14
SACTE_4472		ACTE_Glucosamine-6-phosphate isomerase, subgroup	0	0	0	0	0	0	0	0	12
SACTE_1702		ACTE_Htar	0	0	0	0	0	0	0	0	12
SACTE_0450		ACTE_Dak kinase	0	36	0	0	7	8	8	8	11
SACTE_5342		ACTE_Nitrite/sulphite reductase, hemoprotein beta-component, ferredoxin-like	0	6	0	0	3	0	4	9	0
SACTE_0264		ACTE_Protein of unknown function DUF541	0	7	0	0	0	0	0	0	9
SACTE_3319		ACTE_Pyridine nucleotide-disulphide oxidoreductase, class-II, active site	0	0	0	0	0	0	4	6	0
SACTE_3012		ACTE_AMP-binding, conserved site	38	0	0	0	2	2	3	6	0
SACTE_2323		ACTE_Fumarylacetacetate, C-terminal-related	0	0	0	0	0	0	0	0	6
SACTE_2490		ACTE_S-adenosyl-L-homocysteine hydrolase, conserved site	0	0	0	0	5	0	0	0	5
SACTE_0985		ACTE_N-acetyl-gamma-glutamyl-phosphate reductase	0	0	0	0	0	0	0	0	5
SACTE_3327		ACTE_NAD(P)-binding domain	32	1	0	0	23	15	43	4	0
SACTE_4843		ACTE_Aminotransferase class-III	0	8	0	0	2	0	0	0	4
SACTE_4243		ACTE_Endoribonuclease L-PSP/chorismate mutase-like	0	13	0	0	0	0	0	0	4
SACTE_2260		ACTE_Acyl-CoA dehydrogenase/oxidase, N-terminal	0	0	0	0	5	0	0	0	3
SACTE_4946		ACTE_ABC transporter, conserved site	0	3	0	0	0	0	0	0	3

Figure 18 (continued)

SACTE_6131	ACTE_Purple acid phosphatase-like, N-terminal	0	0	0	0	0	0
SACTE_5455	ACTE_Amidohydrolase 1	0	38	0	0	0	0
SACTE_1073	ACTE_Proteasome, alpha subunit	0	0	0	2	2	11
SACTE_1901	GH51 ACTE_Alpha-L-Arabinofuranosidase, C-terminal	0	0	0	0	8	9
SACTE_1074		0	0	0	0	7	0
SACTE_0841	ACTE_Winged helix-turn-helix transcription repressor DNA-binding	0	3	0	0	8	0
SACTE_1897	ACTE_hypothetical protein	0	0	0	0	4	0
SACTE_4728	ACTE_Aminotransferase, class IV	0	0	0	0	0	0
SACTE_1619	ACTE_Glutamine synthetase, beta-Grasp	0	0	0	0	0	0
SACTE_6303	ACTE_Serine/cysteine peptidase, trypsin-like	0	0	0	0	4	0
SACTE_1650	ACTE_Bacterial extracellular solute-binding, family 1	6	2	0	0	0	0
SACTE_0782	ACTE_Transcription regulator PadR N-terminal-like	0	0	0	0	0	0
SACTE_3219	ACTE_Methionyl-tRNA synthetase, class Ia, N-terminal	14	7	0	0	2	2
SACTE_6051	ACTE_Catalase	0	8	0	0	0	0
SACTE_1473	ACTE_Dienelactone hydrolase	0	0	0	0	0	0
SACTE_0534	ACTE_Bacterial extracellular solute-binding, family 1	0	0	0	0	0	0
SACTE_1344	ACTE_Pyridoxal phosphate-dependent transferase, major domain	13	0	0	0	2	0
SACTE_4624	GH16 ACTE_Concanavalin A-like lectin/glucanase	0	26	0	0	37	11
SACTE_3777		0	4	0	0	35	0
SACTE_5741	ACTE_Aldo/keto reductase	0	0	0	0	8	0
SACTE_2762	ACTE_NAD(P)-binding domain	0	0	0	0	8	0
SACTE_1162	ACTE_hypothetical protein	0	0	0	0	7	0
SACTE_1640	GH13 ACTE_Glycoside hydrolase, carbohydrate-binding	0	0	0	0	7	0
SACTE_2544		0	5	0	0	6	0
SACTE_2213	ACTE_NAD(P)-binding domain	0	0	0	0	5	0
SACTE_4566	ACTE_ATPase, F1 complex, alpha subunit, C-terminal	72	0	0	0	3	0
SACTE_4568	ACTE_ATPase, F1 complex, beta subunit	32	0	0	0	3	0
SACTE_6063	ACTE_Protein of unknown function DUF336	0	0	0	0	3	0
SACTE_3962	ACTE_Peptidase S1A, chymotrypsin	0	0	0	0	3	0
SACTE_2518	ACTE_Acyl-CoA dehydrogenase/oxidase, N-terminal	0	0	0	0	3	0
SACTE_1738	ACTE_N-acetyl-gamma-glutamyl-phosphate reductase	0	0	0	0	3	0
SACTE_5109	ACTE_Luciferase-like	0	29	0	0	2	0
SACTE_5881	ACTE_Multicopper oxidase, type 2	0	0	0	0	2	0
SACTE_3741	ACTE_Twin-arginine translocation pathway, signal sequence	0	0	0	0	2	0
SACTE_1638	ACTE_Tautomerase	0	0	0	0	2	0
SACTE_1419	ACTE_Ribosomal protein S1, RNA-binding domain	126	0	0	0	0	0
SACTE_1888	ACTE_Acyl carrier protein-like	60	0	0	0	0	0
SACTE_2468	ACTE_Bacterial NAD-glutamate dehydrogenase	36	0	0	0	0	0
SACTE_3896	STRACTE_03857	35	0	0	0	0	0
SACTE_3716	STRACTE_03676	28	0	0	0	0	0
SACTE_4281	STRACTE_04237	27	0	0	0	0	0
SACTE_3955	STRACTE_03916	25	0	0	0	0	0
SACTE_4565	STRACTE_04519	23	0	0	0	0	0
SACTE_4031	ACTE_Chaperonin Cpn60	22	0	0	0	0	0
SACTE_3086	ACTE_hypothetical protein	22	0	0	0	0	0
SACTE_4591	ACTE_Thiolase-like	20	4	0	0	0	0
SACTE_4376	STRACTE_04332	19	0	0	0	0	0
SACTE_4194	ACTE_Twin-arginine translocation pathway, signal sequence	18	0	0	0	0	0
SACTE_1934	ACTE_ATP-grasp fold, subdomain 2	18	0	0	0	0	0
SACTE_4501	STRACTE_04456	17	0	0	0	0	0
SACTE_4959	STRACTE_04913	16	0	0	0	0	0
SACTE_2103	ACTE_Protein of unknown function DUF756	16	0	0	0	0	0
SACTE_4795	ACTE SCP-like extracellular	15	0	0	0	0	0
SACTE_4073	ACTE_NAD(P)-binding domain	15	0	0	0	0	0

Figure 18 (continued)

SACTE_2558	STRACTE_02525
SACTE_4399	ACTE_Protein of unknown function DUF3107
SACTE_1368	ACTE_Phosphoglycerate kinase
SACTE_4224	STRACTE_04182
SACTE_3067	STRACTE_03033
SACTE_4462	STRACTE_04417
SACTE_3361	STRACTE_09325
SACTE_3995	ACTE_RNA polymerase, alpha subunit, C-terminal
SACTE_0514	STRACTE_00503
SACTE_6342	ACTE_Biotin/lipoyl attachment
SACTE_4610	STRACTE_04564
SACTE_1068	STRACTE_01051
SACTE_4926	ACTE_Ferritin-related
SACTE_4830	STRACTE_04784
SACTE_3392	STRACTE_03356
SACTE_3037	ACTE_Tellurium resistance
SACTE_2403	STRACTE_02370
SACTE_5983	STRACTE_05928
SACTE_4283	STRACTE_04239
SACTE_3385	STRACTE_03349
SACTE_1328	STRACTE_01307
SACTE_0800	ACTE_Glycine cleavage system P-protein, N-terminal
SACTE_4205	ACTE_Cys/Met metabolism, pyridoxal phosphate-dependent enzyme
SACTE_4415	STRACTE_04372
SACTE_2431	STRACTE_02398
SACTE_1586	STRACTE_01564
SACTE_1201	STRACTE_01182
SACTE_1426	ACTE_Extracellular ligand-binding receptor
SACTE_5081	ACTE_S-methyltetrahydropteroyltriglutamate--homocysteine S-methyltransferase
SACTE_4873	STRACTE_04827
SACTE_3960	ACTE_Translation elongation factor EFG/EF2
SACTE_2756	STRACTE_02721
SACTE_2238	ACTE_Putative agmatinase
SACTE_4616	ACTE_Acetate/butyrate kinase
SACTE_3956	ACTE_DNA-directed RNA polymerase, beta subunit, bacterial-type
SACTE_3948	STRACTE_03909
SACTE_2801	STRACTE_02766
SACTE_0810	ACTE_Alpha-D-phosphohexomutase, alpha/beta/alpha domain II
SACTE_0669	STRACTE_00659
SACTE_5028	STRACTE_04982
SACTE_4567	ACTE_ATPase, F1 complex, gamma subunit
SACTE_4397	ACTE_Ferritin/ribonucleotide reductase-like
SACTE_4191	ACTE_L-Aspartase-like
SACTE_4030	ACTE_Chaperonin Cpn10, subgroup
SACTE_3961	STRACTE_03922
SACTE_3438	ACTE_Uracil phosphoribosyl transferase
SACTE_3371	STRACTE_03335
SACTE_3088	ACTE_Chaperonin ClpB
SACTE_2755	STRACTE_02720
SACTE_2729	STRACTE_02694
SACTE_2036	ACTE_Aminoacyl-tRNA synthetase, class 1a, anticodon-binding
SACTE_1285	STRACTE_01266
SACTE_1006	STRACTE_00988
SACTE_0548	ACTE_Aldo/keto reductase

Figure 18 (continued)

Figure 19

>SACTE_0237|1, 4-beta cellobiohydrolase|GH6 (SEQ ID NO:17)
atgagccgcacagccgcaccacccigegccatcccgaacagcaactcaigccggggccctcgtcgeccgcagccggggctcc
gcccgcggcggccaccctcggtgccaccggccggccggccgtgcaccgtcactacaaggatccagaaccagtgaaacggc
gggcgcaccgcgtccggccggccggccgtcaccacaaacacggggacgcacatctccgggtggcagtcactcgccgtgg
gtcagccagggtggaaacgcaccccgatctcagagccgtccggteaccgccaaggacgcggctacaacgcggccctggccacc
gggcgcacggcccttcgggttcaacgcgacggcaacgcggccgtcgccggccggccgtcaagctgaacgcgtcaccgtcaa
cggcggccaccacggccggccgtccggccgtacggccgtacccacggacccgacggacccgaccggccggccggcaaccgtgtgg
aacccctaccaggagccaaaggcttatgtgaaccggagttgtcgccgaacgcggccggccggccggccggccggccggcc
ccagccacccggcgtcggtggaccgcacgcggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
ctgacgcagaaggctccggcgaactcgctcgccaggctcgatcatacaaccgtccccggccggccggccggccatcgccgg
tgagctggaccgcaccggatcgccgtaccaagccggatcatcgacccgcgtcgccggccggccggccggccggccatcgccgg
ctgccgtatcgccatcgccgtaccccgatcgactcgccgtccggccggccggccggccggccatcgccggccatcgccgg
cgteatgaaggccaaacggcaactacgtcaaggccgtcgatcgccgtcaacaacgtcgccggccatcgccggccatcgccgg
acggggccaccacgggtggatcgccgtggacgacaactcgccgtccggccggccatcgccggccatcgccggccatcgccgg
cgaccgtcaacgcgtccggccgtaccccgatcgccgtccggccggccatcgccggccatcgccggccatcgccggccatcgccgg
aacggcacgtcggtccggccgtccggccatcgccgtccggccatcgccggccatcgccggccatcgccggccatcgccgg
ggccgtccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
ggccgtccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
ggccgtccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
ttccagctcgccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
aacatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccggccatcgccgg
gctctag

>SACTE_0236|glycoside hydrolase family 48|GH48 (SEQ ID NO:18)
gtggccgcctcgccctcccttggaatgaccgcagccggccacggaggccaggccgcggccgtcgctgcagcgtcgactac
acgaccagtgtactgggtatcgccgttcaaccacccgcacactccctcgaccacccgggtcccgccgcgtcgacggctggaccctgacgt
acgactacgcggaccaggccatcgccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ctggaaacccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
cgccgtcaacggcgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ggggaccgggtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggcc
ctcgccaccgcaccctcccgccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggcc
gcctcgccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
agcaggccaggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
caccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
gaagggtacgcacccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ctccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
atgtacgcgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
gcccacgaactccctcgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ggacgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
ttccagcgccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
cgaccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc
tactggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtccggccgtcc

Figure 19 (continued)

cctgcgtactccatgtcgacaagtacitcaagaagatcgccgactcgctggccgaccacccgtccggccgttcggcaaggaca
gcccgcactacctgtatgtctggfactacgcgtggggggccggccaccgcacactcgccgggttgcgtccggatcggccac
gcccacggggataccagaacccgtggccgtacgcgtcgactcgccgaccctaagccaaatcgccacccggagcgcag
gactggccaaaggactggccaaactggacttacactcgactggccgtccacgcgacgggtgcatecgccccgggtgcgaccaaca
getggaaaggccagtcacgcgcacccccggccggcacgcgcacccgtactacgcgtactacgacgagaagcccggtgacccacgacc
gcccgtccaaaccagggtggccgttccaggcgtggccatggagcgcgtccgagactaccacgcgtccgggtgacccaggcgaagg
cgctgtcgtcacaatgggtgcactggccctgtccgagacgcacccgtcaacccggacggccacccatctgtgcctccaccccccactgtgg
cgccggccggacccgttgaaacgcgtccgtccacccgggtgcacacccgcacgcgtcc
gctgcgtccggccgtacggccggactcgactacgcgtccgtccacgcgtccgggtgcacccggccggccggccggcc
ctcgacggcatgtggcgtccaccaccaggcggacgcacccggccgtggccgggtgcacccggccgactacaaacccgggtgcacgcaccc
gtcgtccggccgtggcgtccacgcgtccggccgtggccggccgttcacccgggttcacccgggttcacccgggttcacccgggttcac
aggacgcacccgaaactggcccccagggtgcagggtactcggtacccggccgtccacccgggttcacccgggttcacccgggttcac
cgacatcgcaactggccctggggcgtacccggccgtccatcggtacccgggttcacccgggttcacccgggttcacccgggttcac

>SACTE_3159|chitin-binding domain 3 protein|CBM33.2 (SEQ ID NO:19)

atggctagacgcagcagactatccctcgccagggtggccaccctgcgtgggcgetggccteaccgcactctggccggcaaa
ggggaggcgcacggtgtgegatgaccccgatgcgtaccatctgtgcceagtcgcacgcctgtccggcacecgggcgcgtgaac
ccccacgaacccggctgecgggacgegctgagccagagcgggegagaacgegctgtacaactggttgcgcgtgcactccaacgcg
ggggccggggatatgtccggacggcagcctgtgcagtgtccggtgaccgtccccgtacgacttccgcctacaacgcgg
ccccgcggactggcccccggacacatctgaccccggtgcgcgtcaagggtcagtagaccaactggggcgeccacccggtgactc
cggtctacctgaccaagccggctggcacccacgtccgaactctgttggacgaccitcagttgtacagaccgtaaagaacccgccc
gcagcaggcgggcgggaccaacggggcactactgtggacactgtccgtcgccgtccggccgttcgggtacggcgtatgttc
atccagtgggtgcgttcggacagtcaggagaacttcttcgtctggacatcgcttcggacggccaaacggcggatgtacgggaatcg
ggggcacggcacccccacccactccgaccccgactcgaccccgaccccgacggacccggagactccgggttcgtcatggccgt
ctacaacgtcgtcagtcgtggccgggttcggcaggctccgtcgagggtatgaaccacggtaacggcaacggcgtacccggc
tgtcagtgacccgggttcgggacgeagatcaacagegtgtggacccggcgttcgtccgtacccgggttcggcaggccacccggc
cgacgtggaccacaaccgtgtcatcgcccccggacggcagttgcacccgttcgtccgtacccggcgtacccggc
ggacgtccgtgtgtacccgt

>SACTE_0482|glycoside hydrolase family 5|GH5 (SEQ ID NO:20)

Figure 19 (continued)

gccccggagagcgcggcggtccggccggcacctgcggcgccgactacagcggcagggcggtgctgaccgagtccgggg
cgtgtcaagaaccggatcageaccccccatttcctccaccggctga

>SACTE_0265|glycoside hydrolase family 10|GH10 (SEQ ID NO:21)
 atggecaagaaaatccccgcggcgacttcgttgcacggcacttcgttgcacgggggggtgtcgcggccggcggtgttcgtcgc
 cggcaeggccggaggcaggcaggcaccctgggtgacgcggcgccgcaaggccggtaacttcggcaccggcggtcgccgcaaccac
 ctccggcgaggcaccgtacgcgtccacgtggacgcaccgttgcacttcggcaccggagaacgagatgaagtgggacgcggcgag
 ggcaggcccaacttcgttgcacggcggccgaccagatcgtcagtcacgcaccggagaacgggaatgaagggtgcgcggcacaccc
 tgggtggcactcgacgtccgggggtgggtcgccggcgttggggccaccgttgcgcggcgatgacaaccatcaccaggat
 gatgacgcaactacaaggcaagatccatcttgcggacgttgtgaacgaggccatcaggacggcaacagcgggtgcceggcgacgttc
 cttccaggacaagttgggtgacggcgttgcacggaggcgatcgcaccggccgtacggcgatccgcaccggcgaaacgttacaac
 actacaacaccgcggccggaaacgcgaagggcggcggatcgacgggttgcacggacttcggcaccggccgtacggcgatccgc
 actgggttcccgacgttccacacgttgcacggacttcggcggccgttgcacggacttcggcaccgttgcacggacttcggc
 cgttgcacgttcccgacgttgcacggatcgggttccggcggccgttgcacggacttcggcaccgttgcacggacttcggc
 cccgttgcacggccgttcccgacgttgcacggacttcggcggccgttgcacggacttcggcaccgttgcacggacttcggc
 aacagaaggccggcgtacgcgggtgtccgcgcgttgcggccggccggccggccggccggccggccggccggccggccgg
 gggccggaaacccggcaggcgttgcacggcgttgcacggacttcggcggccgttgcacggacttcggcggccggccgg
 ggccgttcccgacgttgcacggacttcggcggccgttgcacggacttcggcggccgttgcacggacttcggcggccgg
 gtggaaacggcggaaacccggtgttgcacgggttgcacggacttcggcggccgttgcacggacttcggcggccgg
 gtcatgacgcaacacccggccgttgcacggacttcggcggccgttgcacggacttcggcggccgttgcacggacttcggc
 g

>SACTE_2347|cellulose-binding family II|GH5,CE3 (SEQ ID NO:22)

gtgagaacacgcatacgcacagcagcgaacgcggccacagccccctggcccttcgttgcgggggttgcgccttcgc
 cggggccctcgccggccacccggccggggccggggccggggccggggccggccctccacatcagcgcac
 gggccgttgcgtcaaggcaacgcgttgcacttcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cgcgcacatcaaggcgacccggcgacacacgggttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cgcgcacatcgccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gccggaaaccctcgaccacccggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gcaacggccctggggcaacccggcgatccggggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gccacacgtatgggtggacgcggccaaactggggccggacttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cgacccggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gacccggacttccttcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gccggaggatgggttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gagttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 gcaacccctcggttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 agtggccggccgggttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 ccaacggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc
 cggccgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgcgttgc

>SACTE_0357|polysaccharide deacetylase|CE4 (SEQ ID NO:23)

Figure 19 (continued)

gaccgtcaccggcaacgtactggcgctgtactgtcccgccctgcgcggaggactcgacgggaccgactcgacacgacggctcgacccc
ggctccccggactggccgaggcccgagaaggaccgggttcacgcaggcagaaacgaegagcgggacgggtacttgcac
ccggcggtgcgcaggccaaaggacacgggtggggacgcgtcgccactcgacgcgcacatgcacgaggcgcc
ggggacagcagcgtcggttccatccggcagcgcgcgtcgccggaggcgaaccgcctcgcgggggtfjacggaggcgcta
ctcgacgcgttcggacgcgggtgtggcgtggcagggaggcccactcgacaccagcgggtcgacacccggcage
gggtctctcgccacggaaatcgaaacctggatccgcgtggactggcagggtacggcagacatccacatcgctga

>SACTE_5647|coagulation factor 5/8 type domain protein|GH87 (SEQ ID NO:31)
atgacccccacegcacagcacccctgtcaggcgctgggtgcgcgcgttcgcacccgcgtggcaccgcgcgg
gtcgctcgccgtgcggccggccaggccgggggggcggcaccctcccggtggcatatcgccgggggcgcgcgg
ccgticacggcaggcaggaggccagggtacgcgcggccaccgtcactcgccggggggggccggc
ggcgccggccggcaggccgtacgcgtcgacgggggggtgagtatcggtggacttccatcaccggggccaca
cggcggccgggacttccatccacgtcgacgggggggactgggtttccatcaccgtcgatgcacatc
gtatttgcgtggccacaacgcggcgggggacgggggggggggggactggcttcgc
gggtggacttcgcgggtactacgggggtttccatcacaacaacccggggggggggggggggg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gg
gggggggggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggg
gggggggggggggggggggggggggg
gggggggggggggggggggggggg
gggggggggggggggggggggg
gggggggggggggggggggg
gggggggggggggggggg
gggggggggggggg
gggggggggg
gggggggg
gggggg
gggg
gg
gg

>SACTE_5978|Pectate lyase/Amb allergen|PL1 (SEQ ID NO:32)
atgaggagaccagtgcctcgactcagcgccggggggccaccctggcgtgggtgcgcggaccggccactgtgg
gggggggtgcggcaggcaccgggggtcaccggatacgcgaccgcaggaaacggccgacccaccggccgc
gtcgggccaccaacccggaccgcgtacccgcggccatcgccatcgccgtgggtccaccatccac
accatcaaccacggcaacaccggacaagggtcggcagcgtcgcaaccgcggccggaggcattcg
gtcgacatcgccgtgggtttccatcaccgtcgccggcggcacaaccggcgcacgggttccctcc
ccctccgg
gg
gg
gggggggggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggggg
gggggggggggggggggggggggggggggg
gggggggggggggggggggggggggg
gggggggggggggggggggggggg
gggggggggggggggggggggg
gggggggggggggggggg
gggggggggggg
gggggg
gggg
gg
gg

Figure 19 (continued)

ggtggaggaccccgctggccatgtttcgacggcatcgacatcgactggagfaccacaacgcctgcggcctgacctgtgacacca
ggccccgcgcgtgaagaacctgttccctcgcgccaaagtgcggcgeagaagaacctggtcacccgcgtacccgcga
cggtcgacggcgcaagatcgacgcgcgcactacggggcgccgeactccctcgactggfacaacgtgtatgcgtacgactct
tcggcgtctgggaggcgaagggcccgactcccgctgaacgcgtacgcggcatccgcaggacggctcaactccg
ccgcgcgcategccaagctgaaggccaaggggteccggctcgaaagctgtctcggeateggcttatacgccgggtggacgg
cgtgaccaggcccaccggccacccggcacatcgccgcacccgcgtacgcgcactggcaccactgtggagctacgacacccggcga
agaccatcgccggccacccgcacatcgccgcacccgcgtacgcgcactggcaccactgtggagctacgacacccggcga
ccatcacctcaagaatggcgtctgggaaacagecaggcctcgccgggtcgcttcgtggagttcagggcagacaccgc
gctgtgagcgcacatggacagcggcactaactag

>SACTE_2313|chitin-binding domain 3 protein|CBM33 (SEQ ID NO:50)
atgcggaaaaggcgaagcgcggccgtataggcctggcgatgtcgccaccaggcagtgcggcagccacgg
acaccgattccccatcagcagacagaagctgttgtgcacacggcaccgtcaccggcgtcgccaaacatcgatgtggagccgeagagcg
cgaggggcccgaaagggttcccgccgeaggteccggcgacggcaagatcgccggcggaaacagcttgcgcgegotcgac
cccgcgccggggcaactggccgcaccaggcgtaccggcgccagggctacaacttcgtggcgttcaccgcggccacgc
gaccgacttcggtaactacatccaaggacggcgtggactccacaaggcgtcaccaggccggctggagtcgc
cgccgtatggaaaccaggcagccggccaccggcaccatccccacccagaatcgccggcaagcacatcate
ggccgttggaaacgtggcgtacccgccaacgcgttacgcgtcggtcgttgtga

>SACTE_4246|Carbohydrate-binding CenC domain protein|GH18 (SEQ ID NO:51)
gtggccgcctcgccggcggccctgaccgtgaccggctgtcgccaccgcacaggccgcacatcaacgtgcgc
gggtcgagagcggcctcagcggctggaccgttcccgccgaccccgatccatcgccgttgaaaggccaaactcc
tcaaggecaccggagcggccaggacaacgcgttcaaccggccatcaccatcggcgttgc
ggcgcaggccgggiacgcctactcgccggagggcaccggcaccaccgcgttccatcgccgttgaaaggcc
cccgatcgccacggatccatcaccaccggccgttgcaccatccatcgccgttgc
gggacgacgtcgccgtaccggaccggacggccggccggatcccgaggagccggccggatcccg
tggccgtccggcaccaccatcgccgttccatcgccgttgcaccatcgccgttgc
accaggcaccaccaccggcaccatcgccgttgcaccaccatcgccgttgc
aacggccgggttgcgttgcaccatcgccgttgcaccaccatcgccgttgc
aacggccgggttgcgttgcaccatcgccgttgcaccaccatcgccgttgc
aaccggccgggttgcgttgcaccatcgccgttgcaccaccatcgccgttgc
aaccggccgggttgcgttgcaccatcgccgttgcaccaccatcgccgttgc
cggtcaacaggccgttgcgttgcaccatcgccgttgcaccatcgccgttgc
ctggagaacggccgttgcgttgcaccatcgccgttgcaccatcgccgttgc
cccgacatcgccgttgcgttgcaccatcgccgttgcaccatcgccgttgc
tacatcaccatcgccgttgcgttgcaccatcgccgttgcaccatcgccgttgc
tggaggccggccctggccctccagggttgcgttgcaccatcgccgttgc
cgttgcgttgcaccatcgccgttgcaccatcgccgttgcaccatcgccgttgc
tgaccatcgccgttgcaccatcgccgttgcaccatcgccgttgcaccatcgccgttgc

>SACTE_3064|Chitinase|GH19 (SEQ ID NO:52)
gtgatcagacgcgtatggcctgttcaccggcgttgcgtccggcgacgcgttcc
caccgtggccctggaaacctggcgttgcgttgcgttgcaccatcgccgttgc
tggacgagaacggcgatcgccgttgcaccatcgccgttgcaccatcgccgttgc
ccctcgccgttgcgttgcaccatcgccgttgcaccatcgccgttgcaccatcgcc
gctgagcggcgttgcaccatcgccgttgcaccatcgccgttgcaccatcgccgttgc

Figure 19 (continued)

>SACTE_0562|cellulose-binding family II|GH74 (SEQ ID NO:55)
gtgtatccatgcctccacggccctggcggttcaggcgttgcactcatggAACCCCTGCGGGTGGGGACGGTGTGGAGCAGAAGC
atcgatgttccatgcggcacaccataccttggAAAGAACGCCGGATCGACGGGGGGCTCGTGGGGACCAGTCGGCAAGCAGTGGAAAGGCCCTGCTG
gactgggtggactgggaccegtggggctggacggcggtggagcgttgcacttcggacacgggtcgacccgacaacgtgtacggcc
gtggggacgtaccaacagctgggacccgacccgacggcggtctgcgttccggaccggggcccttggaaaggcgccacc
ctcccggtcaagctggccgaacatggccggacggcatggggggcggtcgccgttccggacacaagaactcgtgttac
ggggcgcccgccggcaacggcccttgggttccaccggacggggggatcgttccggaggttgcggccctttcccaacccgggaa
ctacggcggaccggcgtggacaccagccgttccggacacggacaaccaggcgttccggatcgacggcgttccggc
ggggcagegecaccaggacatctacgttccgggttccggacacaggagaacaccgttaccgttccacggacggcggegecaccctgg
tcggcgatccggcccgccacccgggttaccctcggttccggacacaaggcggttactcgacttcggcgacccgttaccgttac
caccggcgcccttacggacggggcaaggccggatctgggttccggacacgggttccggcccttggcaggacgttccgggttgg
cgaggccgttccggactacgggttccggatccgggttccggcagaacccgggttaccgttaccgttaccgttaccgttacc
gtggcccgacaccaggatctccgttccacggacagcggttccggaccctggacttccgggttaccgggttaccgttacc
accgttacacgttccgttccgttccgggttaccctcggttccggacaccgggttaccgttaccgggttaccgttacc
atggacggggcggtggagatcgaccgggttccgttccgggttaccgttaccgggttaccgttaccgggttaccgttacc
cttccggggaccgttccgttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgttacc
ccccgaacctcgactcgaccaccaggccgttccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ccacatcggttccaccgacaacggggccaacttccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ggggcggacgggttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
accggcgttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
gaccggacggggggcgaccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
aggggacatcggttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
cgccggcgttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
cgccgggttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ccggccgttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
cgacagcgccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
cgccgttccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
gttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ggagcaacggccaaacggccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ggagcaacggccaaacggccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc

>SACTE_4343|extracellular solute-binding protein family 5| (SEQ ID NO:56)
atgcgggttccatgcggccaaagagcgccaaatgggtcgggggagcggttaccgttaccgggttaccgttaccgggttaccgttacc
aggacaaacgggttccatggggcgccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
aacacgttaccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
aacgttaccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ccggacttaccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
ctacgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc
accatcggttaccgttaccgggttaccgttaccgttaccgggttaccgttaccgggttaccgttaccgggttaccgttacc

Figure 19 (continued)

ggcgeccaaccagteggtgctggagatgcttgcaggccaaacggcgccgacgtcactccttcatacagaaggtaagaaca
aggaggacggcgccgtatggcgtccggccacgggtgtacaagtcctcgccgcaagtcataaggccggccca
cgacgtccttcctcgctcgcaagtccgacgagctgtggacatcgccgtcaagctggaggacacgcgcgtctccgacgactacttcgc
tcgcgtcaaccttacccaaacgtggacttctacacgggectgtatctacgggcatggcgtcccgaccgagatgttcacccgtcttgc
gtccggcccccctcccggtgtcgatcgatcgtggcaagagatgtcaaggagccgggtccgcateggcccccgcgcagatctaca
ccggcgagggtctcgccgtacttgtcccggtcgagagccgcgtga

>SACTE_5668|Serine Protease| (SEQ ID NO:60)

atgacgaaacgtcaggcatttcgtcgactcgccacgggtcgccgtgtttccggccacccgttcgtccgcgtccaccgcgc
cgccggccctggggccgcgcgcgtcaatgtggaccgtcgccgtggacgaaaggctgtccgcgtccgcgtccaccgttcgc
ccactggccatcgacaggccgtcaggacggcaggteacgtcgcctcgccgtccgcgtccgcgtccgcgtccgcgtcc
gtcgccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
cgtacgacgtatcgccgttcgcacccgcgggggtcgccaggacgacccggcgtggactgtgtccgcgtccgcgtcc
ccgcgcacccgtgccccgtcccgccggacgacgggaccaacccggaaacgcggccgtccgcgtccgcgtccgcgtcc
caggggacgtcgccgttcatggacacggcgtcaggacccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
actactcggtactctacggcaccttccggcgttacgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
cgtcgaccggacgggtctgttacgaggacaacccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
gcaagaacgcgacccactacccggcttcggcaccgacccggcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
gcacccggccggccggccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
ggccgaggcggtccgcgttacgtgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
gcggggacaacggacttcggcttacccggcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
ctggcaggccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
ggccgggtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
ttagcatgcaccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
gcctcgacgcgcacccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
cccccggccggaggccggccggccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
gggggtga

>SACTE_6428|chitin-binding domain 3 protein|CBM33 (SEQ ID NO:61)

atgaaitgtcatgtatcgcatcaacttacgcggctggacacaggctgagccgtctgttcgtcgccgcgtctgtctgtcccggtggac
ggcgcacggccggccacggccatcgccgtcgccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
cccgccatggccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
aacaggatccgcggcaactccggcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
gtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
actactccgggtgtacgttacccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
ggacgttacggtcccgatcgacgttacccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc
tgcacatggacacgtacttccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgc

>SACTE_0366|alpha-L-rhamnosidase|GH78 (SEQ ID NO:62)

gtgtatcgacgaaagacgtactgtcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
ccggccgcacccggccggccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc
cccgccgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtccgcgtcc

Figure 19 (continued)

ctggcccgccccgacgtctggacacggcaaggcgtgtcccgacgtcggtctgtccctacgcggccggcgtggctccc
gtacgegetaccactggtcgtggcgtgtgggaccaggacggacgcgtcteggcgtggagcggccgtctgggagacccggc
cttggcggaggccgactggtcgggggtggatcgccgcggccggcgtacccctccctggaggccggcgtctggatct
ggttcccgagggggatcggccgtggggcgtcccgccggccacccgggtggatcccgccggggatcccgaggggcgtaccc
gcgeccgcgtgtcatgaccgcggacgcgggttcacccgcgtggatcgggtgtcccgccgtaccggccgcacggcc
ggagaactggcgtgtcccggtggatcggcgtacggcgtacccgcgtcccggtgtcccgccgtacggccaccaacgc
gtggacggcccgccgggtctcgccggggggatcggcgtacccgcgtacggcgtacccgcgtcccgccgtacggc
gcacccgcggggggatcggcgtcccgccggatcggcgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
gggtcccgcccccgtggggcggggatcggcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
cgccggggccggcgtactcgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
ggctggaccgactaccgcaagcggcgtcccgccgtacccgcgtacccgcgtcccgccgtacccgcgtcccgccgtac
cctcgccgggggtggatcggcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
gtcacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
tggcaggcggggatcggcgtcccgccgtggatcggcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
gggtggaaagggtgtacccgcgtcccgccgtggatcggcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
cgaacccggccgggggttaactggtgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
acccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
tacacgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
egccccggccggacccgggtgtggccgggtcatccacccctcgccgtcccgccgtacccgcgtcccgccgtacccgc
ccggccgtccacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
tggaccggccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
cgacgaccagctggccgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
ccgggggtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
gagtacccctcgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
gacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
cgacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
cgatcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
ggacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
ctgcgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
ggacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgcc
ggcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
cgactccctgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgcgtcccgccgtacccgc
ggcgggggtgtgggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggg
cttgcggccgggtgtgggggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggggggggtgtgggg

Figure 20

>SACTE_0237|1, 4-beta cellobiohydrolase|GH6|GI:344313496 (SEQ ID NO:1)
MSRTSRTLRRSRTALMAAGALVAAAAGSAAAAAPFGATAAAAAGCTVDYKIQNQW
NGGLTASVSVTNNGDAJSGWQLQWSFAGGEQVSQGWNTVSQSGSAVTAKDAGYNA
ALATGASASFGFNATGNNSVVPATFKLNGVTCNGGTTGPTDPTDPTDPTDPAGN
RVDNPYQGAKVYVNPEWSANAAEPPGDRIADQPTGVWLDRIAIEGANGSMGLRDH
LDEALTQKGSGELVVQVVIYNLPGRDCAAALASNGLGPTEIGRYKTEYIDPIAEILGDPK
YAGRIVTTVEIDSLPNLVTNAGGRPTATPACDVMKANGNYVKVGVGYALNKLGDAPN
VYNYIDAGHHGWIGWDDNFAGASAEIFHEAATAEGATVNDVHGIFTNTANYSLKEENF
SIDDAVNGTSVRQSKWVDWNRYTDELSFAQAFRNELVSGFNSGIMLIIDTSRNGWGG
ANRPSGPGANTSVDTYVDGGRYDRRIHLGNWCNCQAGAGLGERPQAPEPGIDAYVWM
KPPGESDGSSSEIPNDEKGKGFDRMCDPTYTGNARNNNNNMSGALGGAPVSGKWFSAQFQ
ELMKNAYPAL*

>SACTE_0236|glycoside hydrolase family 48|GH48|GI:344313495 (SEQ ID NO:2)
VAALALPLGMTAAGTEAQAAAACSVDYTTSDWGSFTTELTLNRGSAIDGWTLT
YDYAGNQLTSGWSGTWSQSGKTVSVKNAAWNGAIAAGAAVTGAQFTYSGANTAP
TTFAVNGETVCAGAHQPPIAVLTSPAAGAVFSAGDPVPLAATAAAADGATISKVEFYDDT
TLLGTDTTSPSYEAGQLAAGSHSVYARAYDSLGSADSPAGITVVTGPAAVVSPLAQ
GVQQGRSGTFDVSLSTAPAADVTVTAARSAGNTGLSVTGGSTLTFTPANWSTPQKVTV
TADGSGTGAATFTVTAPGHGKAEVTVTQLAAKEYDARFLDLYGKITDPANGYFSPEGI
PYHSVETLIVEAPDHGHETTSEAYSYLIWLQAMYGKITGDWTKFNGAWDTMETYMIPT
HADQPTNSFYDASKPATYAPEHDTPNEYPAVLDGSASSGSDPIAAELKSAYGTDDIYGM
HWIQRDVNVGYGNAPGTCAAGPTQAGPSYINTFQRGSQESVWETVTHPTCDNFTYGG
ANGYLDLFTGDSSYAKQWKFTNAPDADARAVQAAYWADVWAKEQGKAGEVADTVG
KAAKMGDYLRYSMFDKYFKKIGDCVGPTTCPAGSGKDASHYLMSWYYAWGGATDTS
AGWSWRIGSSHAGGGYQNPMAYALSSVADLKPKSATGAQDWAKSLDRQLDFYQWL
QSDEGAIAGGATNSWKGSYAQPPAGTPTFYGMYYDEKPVYHDPPSNQWFGFQAWSME
RVAEYYHESGDAQAKAVLDKWVDWALSETTVNPDTYLMPSTLQWSGAPDTWNASN
PGANAQLHVTADYTDDVGVAGAYARTLTYAAKSGDTEAEATAEALLDGMWQHHQ
DDAGVAVPETRADYNRFDDPVYVPGGWTGAMPNGDTVDEDSTFLSIRSFYKDDPNWP
QVQAYLDGGAAPVFTYHRFWAQADIALALGAYADLLE*

>SACTE_3159|chitin-binding domain 3 protein|CBM33,2|GI:344316337 (SEQ ID NO:3)
MARRSRLISLAALVATLLGALGLTALWPGKAEAHGVAMTPGSRTYLCQLDALSGTGAL
NPTNPACRDLALSQGANALYNWFAVLDSNAGGRGAGYVPGSLSAGDRSPYDFSAY
NAARADWPRTHLTSGATLKVQYSNWAHPGDFRVYLTKPGWAPTSELAWDDLQLVQ
TVSNPPQGGAGTNGGHYYWDLALPSGRSGDALMFQWVRSDSQENFFSCSDIVFDGG
NGEVTVGIGGTGTPPTPTPTDPEHSGSCMAVYNVSSWAGGFQASVEVMNHGT
EPRNGWAQWKGPGSGTQINSVWNGSLSTGSDGTVRDVDHNRVIAPDGSVTGFTAT
STGNDYPAGTIGCVTS*

>SACTE_0482|glycoside hydrolase family 5|GH5|GI:344313735 (SEQ ID NO:4)

Figure 20 (continued)

VKRFLALLATCATVLGLTALTGPQAVAAAGCTADYTTSQWQGGFQAAVKVTNLGTPV
TGWLKTFTLPDAGQKVQGWNAAWSQSGSAVTAAAGADWNGLATGASAEAGFVGSE
TGANPPPTAFALNGVACTGSTGEPPAGSDGGTPVDVNGQLHVCVNLCNQYDRPVQLR
GMSTHGIQWFDAKYDAASLDALANDWKS DLLRIAMYVQEDGYETDPAGFTRRVNDLV
DMAEARGMYALIDFHTLPGDPNVNLDRAKTFFASVAARNAGKKNVIYEIANEPNGVT
WTAVKSYAEQVIPVIRAADPDAVVIVTRGWSLGVSDGSDESEVVNSPVNATNIMYAF
HFYAASHKDARYRSTSRAAARLPLFVTEFGTVSATGGGAMDRASTTAWLDDQLKIS
YANWTYSDAPESSAAF RPGTCGGGDYSGSGLTESGALLKNRISTPDSFPTG*

>SACTE_0265|glycoside hydrolase family 10|GH10|GI:344313522 (SEQ ID NO:5)
MAKKIPARARRALSVLTAGVLAAGVVSAGTAAAGTLGDAAAAKGRYFGTAVAAN
HLGEAPYASTLDAQFDSTPENEMKWDAVEGSRNSFTTAADQIVSHAQSKGMKVRG
HTLVWHSQLPGVVGLGATDLRAAMNNHITQVMTHYKGKIHSWDVVNEAFQDGNSG
ARRSSPFQDKLGDGFIEEAFRARTVDPTAKLCYNDYNTDGRNAKSDAVYAMAKDFKQ
RGVPIDCVGFQSFSNSPVPDSYRANLQRFDLGLDVQITELDIEGSGSAQAANYTSVV
NACLA VTRCTGLTVWGVTDKYSWRSSGTPLLFDGDYNKKPAYDAVLAALGGTPDGGG
DDGGGDNGGGNTGSCTATYTQTATWNGGYNGETVKA GSSGITTWSVPVTVPSSQQV
SALWNGAPTWNAGNTVMTVKPTYNGT LAAGASTSF GFTVMTNGNTSAPA AVGACTAS*

>SACTE_2347|cellulose-binding family II|GH5,CE3|GI:344315549 (SEQ ID NO:6)
VRTAIRTARRPQPLALLRGLAFLGLALAGALGPATARAADLPQRAEARAAGLHISDG
RLVEGNNDVFMRGINHAHTWYPGETQSLADIKATGANTVRVLSDG YRWSENS PED
VASHARCKAERLICVLEVHD TTG YGEDAAAGTL DHAAD YWIGLKDVLDGEEDYVVNI
GNEPWGNADPAGWTAPTTAAIQKLRAAGFAHTIMVDAPN WQDWEGVMRADARS VY
DADPTGNLIFS IHMYSVYDTAAK VTDYLNAF DAGLPLLIGEFGGPADQYGD PDEDTM
MATAEELGLGYLAWSWSGNTDPVLDLVLDFDPTRLSSWGERVLHGP DGTET SREATV
FGGGQGGDTEAPTAGPTPTASGV TATSVT LGS WAATDDVGV TAYDV VRVTGGSETK
VASSAATSVT V TGL SAGT AYSAV YAR DAAG NRS ARSGT VSVT DEGG SVPGGAC SVG
YRVIGEWPGGFQGEITLRNTGAAAVDGWT LGFA ADG QTV TNMWGGT ATQSGG AVSV
TPASYTSTIAAGGSVTVGFTGTLGANAAPAAFTLNGATCTAA*

>SACTE_0357|polysaccharide deacetylase|CE4|GI:344313612 (SEQ ID NO:7)
MSITPRPSL RAMVTGLAVAASALAGGA VTAA PPARAA CNGYVGL FDDGPSAAQTPAL
LSALKQNGLRATMFNQGNYAASNP AQVKAQV DAGM WVG NH SYSHPLTQQSQAQM
DSEISRTQQIAAGGGGTPKLF RPPYGETNATL RSVEAKYGLTEVIWDVDSQDWNGAST
DAIVQAVSRL TAGQVILMHEWPANTLAI PRIA QTLSAKGLCSGMISPQTGRAVAPDGG
GNGGGGGGGCTATLSAGEK WGD RYLN NVAVGSSNWT VMN VPS GERV MTTWN
VSASYPSAQVLVAKPNGSGNNWGA TIQANGNW TWPTVSC TTS*

>SACTE_0358|Endo-1,4-beta-xylanase|GH11|GI:344313613 (SEQ ID NO:8)
MNPLVYTERRRRGRLTSLAGSVCALVAAAAA MLLPGTASADTVVTTNQTGNNNNGYY
YSFWTDGGGQVSMNLASGGSYSTS WTNTGNFVAGKGWSTG GRKS VTYSGTFNPSGNA
YLTLYGWSTNPLVEYYIVDNWGT YRPTGTFKGTVSSDG GTYDIETRTNAPSIEGT KTF

Figure 20 (continued)

KQFWSVRQSKRTGGTITTGNHFDAWARNGMNLGTMNYMILATEGYQSSGSSNITVSEG
GSGGGGDNGGGGGGGCTATLSAGEKWGDRYNLNAVGSSNWTVMNVPSAEKV
LSTWNISASYPSSQVLVAKPNGSGNNWGATIQANGNWPTVSCITTS*

>SACTE_1310|Pectate lyase|PL3|GI:344314542 (SEQ ID NO:9)
MSERAASPRTHRERRPGRRRIATALTAALGLTGAALATGVMLQPAGAATTAAIPAWPSAT
GSQSVSKTIEVSGTYDGGLKRFTGSGDLGDDQDEGQDPIFKLKDGTATKVNVLGTPAAD
GIHCSGSCTIQNVWWEDVGEDAASFKGTSVYGGAKKASDKVFQFNGAGKL
VVTKFQVADFGKLVRCNCQSKYKREIVNDVDVTAPGKSLVGINTNYGDTAALRSV
RVHGDSKKIKPCVRYTGNSTGAEPKETGSGPDGTYCKYTASDLSYD*

>SACTE_3717|glycoside hydrolase family 9|GH9|GI:344316877 (SEQ ID NO:10)
MWCHPYLRLRTSGRKVSSVNALPPPAPARPVPRSRVYGRRLGMSAAALLCAGALAVP
GTAMADDAEPPGPGEQITNGDFATGTSAPWWTPNASAAVSEGRCLCVEVPAGTANAW
DVIVGQNDVPIVAGESYELSYTARSTVPLTVQTRVQEAVEPYTTVLAATADPVGAEDETRV
ARTFTASVQDQPAASVQLQIGGGERATTFCCLDDVSLRGGAEPVVYVPTGSPVRVNQVG
YLRGPKSGTVVTDAEAPLTWTKAEDGSTAATGTTVPRGEDPSSRRVHTDFGDLTT
AGDGYTEVDGEVSEPFSSIRGDLYDSLRSDALAYFYHNRSGIEIDADLVGEQYARPAGHI
GVAPNKGDTDVPCCRPGVCDYRLDVSQGGWYDAGDHGKYVNVNGGISVAQLMATYERTL
TAPDAESAELGDGALRPERDNGVPDILDEARWEMDFLIKMQVPAGEQLAGMVHHKM
HDAEWWTGLPMKPHLDQQRELHPPSTAATLNAAATAAQCARLYAPFDAADFADRCRA
AETAWDAAKRHPDVLADPNQDGIGGGAYNDDDSDEFWAAAELFTTGKDIYRQAVL
SSAWHGDAGAVFPAGGGISWGSTAGLVLTATVPNALTSDQLAQVRTVVTEGADRY
AAQSREQAYGLPYAPRGEDYVWGSNSQVLLNMVVLATAHDLTGDAAVYQDAVLRGAD
YLLGRNPLNQSYVTGYGERDHSNHQHRFWAHQNDPSLPNPAPGSIAGGPNLTAIASGDP
VAAEKLSGCAPAMCYVDDIGSWATNEITINWNAPLAFIASYLDAGEGGQTAAARTCQ
VTYSSHWNNSGSTVTVVENTGSDPVSPWALTWLLPGEQRSLHTWSAEFDQHGRTVSA
RPLSWNRTLAPGAAVDFGFNTSAAGSSPEPGAFKLNRCAGSAG*

>SACTE_4638|conserved hypothetical protein|GI:344317777 (SEQ ID NO:11)
MRTGSIARVGLAAALAALLTAFMAPAMAGKHDATDSPSAAAAPASFTHPGVLVSRP
QLDFVRGKVQAGAQWPWKGAYDQMLASPYASLSRTAKPRAVVECGYSNPNNGCTDER
EDALAAYTLSLAWYISQDGARYAQKAIQIMDAWSGVIKDHTNSNAPLQTCWAGSSWPR
AAEIIKYTYGNWPASGRFGTMLRDVYLPKVANGNSNSNGNWELSMTEAAIGIAVFLEDR
GAYDRAVAKFRGRVPAYIYVTADGSLPKAAPGSGLDTREKIINYWQQQSTFVDGLSQET
CRDLHTGYGLSAISHAETSRIQGQDLYPEVADRLRHALGLHAKYQLGEKVPSSLCGGS
LKDSLGPVTEVGFNALHNRMGYAMTNTQLTERQRPAASNNLFVAWETLTHADNPN*

>SACTE_4738|glycoside hydrolase family 16|GH16|GI:344317876 (SEQ ID NO:12)
MPSRTTLIATTAALVALAAPMAFAAPAPAPDPAVEAAAAAWDTDRAASAYAANPAAV
TASGSENPASGPGAAATGDATTRWSSDFADNAWIRVDLGSTIRINQVKLEWEAAYGKK
YVLEVSKDGTNWTPFYTEDAGTGGTVTAHTYPQEVTRGLYVRMRGVERATAWGYSLFS

Figure 20 (continued)

FQVYGGEPAPASTTRSNLALNHPAYGDLYQHAGNSPAFVTDGGWPADLKADRSRWSS
DWNAWRWGVVDLGATSTINSVDLYWEAAYAVDYEIQVSDDNRTWRTVHRPSAAEVA
ARRADVKAPEAVGRHDTINLPTPATGRYVRMLGKERRSFYNPAPSTAQFGYSLYEFQ
VWGTGGSADAAYPALPKNPGGAYRTTFDDFTGSGLDRSKWRVVRTGTEMGPVNGES
QAYVDSPDNIRTENGALVLESKYCKGCTPTPNGTFDFTSRVDTNTKFDFTYGKVSARM
KLPVGDFWPAFWLLGSDVDDPAVSWPGSGETDIMENIGYGDWTSSGLHGPGYSADG
NIGASQTYPNGGRADEWHTYGEWEWTPEGMTFTVDDRVVQQTSRQKLESTRGKWFH
NQYVILNLALGGAYPGGYNQVTQPYWGLPQSSVDRIAQGGIKAEDWVRVEQK*

>SACTE_4755|conserved hypothetical protein|GH64|GI:344317893 (SEQ ID NO:13)
VISRRMFLTGAAASATALTYPLWTGTLSPRTSAAAATCELALENRSLPGTVHAYVTGHE
QGTD SWVLLRADGSVYRPESPGAPQTPLPVDCAIPLNGAGAGPVLTPQMYGARVYF
VRDDKLDFYLNPGPSLVEPAFATPTDPNYGRTWSFCEFTFPNPQQLYANISYVDLVTALPI
GLTLEGDSTHTVAPLPDGAVQRIADDLTAQAAADGQPWDKLVTRGSDGQVLRVVSPQ
NLMAPYFDRPDEMPFRDLFAAQIDEVWEKYRSTDRLRIDLOGGRGTLAGRSGDTLTFE
GGHTFSKPTSKDIFTCNHGPFTNNPSDSDKKALLARIAAGFNRSIMLSHPSQPNCTSVA
DYYQDAVTNHWSRVVHANSPIGYAFPYDDVRPDGEVDVSGAANDGNPRRFTSVG*

>SACTE_5457|Chitosanase|GH46|GI:344318578 (SEQ ID NO:14)
VLPHNRTARRTTRLRTGGLAAAALGLALMALPVTAHAGAPTQPAAHHEAAATGL
DDPAKKDIAMQLVSSAENSTLDWKAQYGYIEDIGDGRGYTAGHGFCSGTGDMALVER
YTD RSPGNVLASYLPALREV DGTDSHDGLDPGFPRDWAEAAKDPVFQQAQNDERDRV
YFDPAVRQAKDDGLGTLGQFAYYDAIVMHGGGGDSTSFGSIRQRALAEAEPPSRGGDE
VAYLDAFLDARVWAMRQEEAHSDTSRVDTAQRVFLRDGNLNLDPLDWQVYGDHF
G*

>SACTE_5647|coagulation factor 5/8 type domain protein|GH87|GI:344318749 (SEQ ID NO:15)
MTPPHRHRLFRRSVSASLSLALTAVGTAAAVLAGAPAAQAAAVPAPSPVGISRGAA
VPFTEQEAEYAATNGTLIGPDRRYGSLPSEASGRQAVTLDAAGEYVEFTLTAPANAMTF
RYSLPDNAAGTGRDASLDLRVNGSVLKVSPVTSKYGWYYGGYPFNNNPGDTNPHFY
DETRTMFGSTLPAGTKVRLQVASTAGSPSFTVDLADFEQVAAPVGKPSGALDVVSDFG
ADPTGAADSTAKIQA AVDAGRTQGKVYYIPQGTFQVRDHIVVDQVTLRGAGPWYSVLT
GRHPTDRSKAVGVYKGYSAQCGGSRNVLKDFAIQDIQERVNDQVNAIGGAMSDSV
DNVWMQHTKCGAWMDGPMDFNFTIKNSRILDQTADGVNFHYGVTNSTVTNTRNNTG
DDGLAMWAENVPNVKNKFTFNTVILPILANNIVTYGGKDITISDNVMADTTNGGGLHI
ANRYPGVNSGQGTAVAGTHTAARNTLIRTGNSDFNWNGVGAIWFSGLNEPISNATINI
TDSEVLDSSYAAIHIEGASNGLHFKNVKIDGAGTYALQIQAPGTATFENVVATHIAQSN
PIHNCVGSGFQITRGSGNSGWYADPPACTGVWPDPVWTNGGVPGGGPTNPTDPTDPT
DPTDPTDPPEETGNLARGRTVTETSHTDVGAANTVDGNADTYWESRNNAFPQSUTVD
LGAAKAVKRVVVLKLPAAWATRTQTLSVSGSTDNGTYNSLKASAGYTFNPSSGNTAT
VSLPGTPVRYLRLTFTQNTGWPAALSELEAYTS*

>SACTE_5978|Pectate lyase/Amb allergen|PL1|GI:344319072 (SEQ ID NO:16)

Figure 20 (continued)

MRRPVALRSAAGATLALAAATGALMAMPEAASAATGGVTGYATQNGGTTGGAGGQ
TVRATTGTAIHAALCGRASSSTPLTIQVEGTINHGNTDKVSGSSCTAAGVIELKQISNVT
IVGVGGGAVFVDQVGIHVRESSNIIQNVTVKNVKKSGSPTSNGGDAIGMEKDVRNVWVD
HTTLEASGGESEGFDGLFDMKAGTQYVTLSYSILRNNSRGGLVGSSESDSLNGFITYHHN
LYENIDSRAPLLRRGGVAHIYNHYVGLSKSGINSRAGARAKVDNNYFEDSKDVLGTFYT
DAAGYWQVSGNVDNVTWSGRSSDNNPAGPDPQSNTSVSIPYAYTLDGANCVPSVVSR
TAGANTGLKVSDGSCSPQTDPDPTDPTPPGTNLSLGAGSDGSSKASGTS
YGDVRDGDMSTYWSPSGSTGSVIKWSATTVKINVREAAGSTGSITSWKVGNADTG
AVLASGSGAGVITFPQTSRKITEITGSTGTPKVAEFETYAG*

>SACTE_5230|xylose isomerase||GI:344318358 (SEQ ID NO:33)
MPERFTPTPEDKFTFGLWTVGWRGNPDFGEPTRPVLDPVESVERLAELGAHGVTFHDD
DLIPFGSDDRERARLVGRFREALERTGLKVPMAATTNLFTPVFKDGGFTSNDRDVRRFA
LRKVIRNIIDLAVELGAQTYVAWGREGAESGAAKDVRSAALDRMKEAFDLLGDYVTEQ
GYDLRFAIEPKPNEPRGDILLPTIGHALAFIERLERPELVGVNPETGHEQMAGLNFPHGIA
QALWAGKLHFIDLNGQSGIKYDQDFRGAGDLRQAFWLVDLLETAGWDGSRHFDFKP
VRTDGIDGVWESAKNCMRNYLILKERAFAFRADPAVQEALTASRLDELARP TADDGLK
ALLADRTAYEDFDATAAAERSMAFEALDQLAMDHLLNVR*

>SACTE_4571|glycoside hydrolase family 18|GH18|GI:344317711 (SEQ ID NO:34)
MTSALRATQGLQSTNHPRLSDLTRGAPLSTESPRESSRLRWRLGPGRATRAKAVAGFTA
LLLPLAACMVGLASPAQAATSATATYLKKSDWGSFEGQWTVKNTGTTALSSWTIEWDF
PSGTAVGSAWDASVTSSGTHWTAKNLGWNGTVAPGASISFGFNGTGSGSPTGCKLNGA
SCDGGGTVPGDSAPSKPGPTASGITDTSVKLSWSAATDDKGKINYDVLRDGAKVATV
TTTYTDTGLTKGTDYSYSVQARDTADQTGPVSGAVAVRTGGNDNPGPGTGSKVNLG
YFTNWGVYGRNYHVKNLVTSGSAEKITHINYAFGNVQGGKCTIGDSYADYDKAYTAD
QSVDGVAUTWDQPLRGNFNQLRKLAKYPHIKVIWSFGGWTWSGGFGAAAQNPAAF
QSCYDLVEDPRWADVFDGIDWEYPNACGLTCDSGPAAALKNSSLARAKFGAKNLV
TAAITADGSDGGKIDAADYAGAAQSFDWNVMTYDFFGAWEAKGPTAPHSPNAYAG
IPQDGFSAAAIAKLKAKGVPASKLLLIGFYGRGWTGVTQAAPGGTATGAAPGTYEA
GIEDYKVLKTSCPATGTIAGTAYAHCGTNWWSYDTPATITSKMAWANSQGLGGAFFWE
FSGDTANGELVSAMDSGLN*

>SACTE_2313|chitin-binding domain 3 protein|CBM33|GI:344315516 (SEQ ID NO:35)
MRKRASAAVIGLAIAGVSMFATSSASSHGYTDSPISRQKLCANGTVTCGNIQWEPQSV
EGPKGFPAAGPADGKICAGGNSSFAALDDPRGGNWPATQVTGGQGYNFRWQFTARHA
TTDFRYYITKDGWDSTKPLTRAALESQPFMTVPYGNQQPATLTHQGTIPTQKSGKHIIL
AVWNVADTANAFYACSDVKF*

>SACTE_4246|Carbohydrate-binding CenC domain protein|GH18|GI:344317395 (SEQ ID NO:36)
VAALAAGALTGTGLVGTAAQADINVAKNAGFESGLSGWTCTGGSGATVSSPVHGGSA
ALKATPSGQDNAKCTQTVAVKPNSTYALSSWVQGGYAYLGASGTGTTDVSTWTPGST
GWTQLRTSFTGPSTTSVQVYTHGWWYQQAAYYADDVAVTGPDPGGGGTEEP GPAIPGAP
AGLAVGTTSSSVALSWNAVGATGYTVYRDGTKATTITGTSATVSGLAADTAYQFSV
SATNAAGESVRSATVSGRTAKKDETGPGPSTSVPKHAVTGYWQNFNNGAAVQKLSDV

Figure 20 (continued)

PANYDIIAVSFADAAGTPGAVTFNLDAGLNGYTVAQFKADIKAKQAAGKNVIISVGGE
KGTVSVNSDASANAFADSPLYLIQEYGFNGVDIDLEGLNSTYMTKALRSLSSKVGSGL
VITMAPQTIDMQSTSGEYFKTALNIKDIITVVMQYYNSGSGMLGCDGKVYSQGSVDFLT
ALACIQLLEGLAPSQVGLGVPASTRGAGSGYVAPSVVNAALDCLAKGTGCGSFKPSRT
YPDIRGAMTWSTNWDATAGNAWSNAVPHVHGLP*

>SACTE_3064|Chitinase|GH19|GI:344316244 (SEQ ID NO:37)
VIRRVMGLLTALAAVVATLVFLPAATASAATCAPAWNASSVTGGGSASYNHNWSA
KWWTQNERTPGTSDDWADQQGACGSGGGGTDPNPSGFVSEAFQNQMFPSRNSFYTYSG
LTAALSAYPAFANTGSDTVKKQEAAGFLANVSHETGGLVHIVEQNTANYPHYCDSQS
YGCPAGQAAYYGRGPIQLSWNFNYKAAGDALGIDLGNPWFQEQNASVAKTGLWY
WNTQSGPGTMTPHNAIVNGSGFGETIRSINGSIECNGGNPGQVQSRVNTYQSFVQILGTT
PGSNLSC*

>SACTE_5764|Chitinase|GH18|GI:344318865 (SEQ ID NO:38)
MRRRSRSVRALVTAAVITVAAAGMAVLGSGTAQAATPLPDHVFAPYFESWTGESPAAM
AAESGAKHLTMAFLQTTAKGSCPYWNGDTGLPIAQASFADIDTIQAGGGDVIPSGGG
YTADTTGTEIADSCTDWDQIAAYQKVTTYDVSRLDMIEVDSDLDDTAGIDRRNKAIC
KLQDWADANGRDLEISYTLPTTRGLASSGLAVLRNAVNGARVDVVNLMTFDYYDN
ASHDMAADTETAAQGLYDQLAKLYPGRATQLWSMVGVTEMGPVDDFGPAETFTLAN
AARVYDWAVAKGINTLSFWALQRDNGGCPGGPAADDCSGJQQNTWDFTRVFAPFTSG
TTAPDDDFSVTATPASGTVTAGGSATTVKAVTKGAAQQVGLTVSGVPAGVTASLSPS
SVTAGGRSTLTATTQAAVSGTYRISVTGTPSGHATAYTLLTVGGTGSQCTAGPWAG
GTVYTGGQQVSYKGHTWKAKWWTTGEEPGTTGEWGVWQDLGAC*

>SACTE_4439|Catalase||GI:344317584 (SEQ ID NO:39)
VTQGPLTTEAGAPVADNQNSETAGPGGPVLVQDQALLEKLAHFNRERIPERVVHARGA
GAYGTFTLTRDVSQWTRAKFLSEVGKETETFLRFSTVAGNLGSADAARDPRGWALKFY
TEEGNYDLVGNNTPVFFIKDAIKFPDFIHTQKRDPTYTGSQEADNVWDFWGLSPESTHQV
TWLFGDRGIPASFRHMNGYGSHTFQWNNEAGEVFVWKYHFKTDQGINKLTTEAVRLS
GVDPDSHQRLRESIERGDFPTWTVQVQIMPAAEAATYRFNPFDLTKVWPHEDYPPIEIG
KLELNRPENIFAEVEQSIFSPAHHFVPGIGPSPDKMLQGRLFAYGDAHRYRVGINADHLP
VNRPHATEARTNSRDGYLYDGRHKGTKNYEPNSFGGPVQDRPLWQPVSVTGGTGNH
EAAVHAEDNDVFQAGNLYRIMSEDEKGRLIDNLAGFIAKVSRDDIADRAINNFRQADA
DFGKRLEVAVQALRG*

>SACTE_0562|cellulose-binding family II|GH74|GI:344313814 (SEQ ID NO:40)
VYAMPSTAPAAVQSGEDAPVRSSPRFAALLAALALTGSLIGTPAVARSDEAPAATE
ASDVSIAADTYTWKNARIDGGGFVPGIVFNRSEKNLAYARTDIGGAYRWDQSGKQWKP
LLDWVDWDRWGWTGVVSLASDTVDPPDNVYAAVGTYTNSWDPTDGAVLSSDRGAS
WKAATLPFKLGGNMPGRGMGERLAVDPNKNSVLYLGAPSGNGLWRSTDAGVSWSEV
TAFPNPGNYAQDPSDTSGYGNNDNQGIVWTFDERSGSAGSATQDIYVGVADKENTVYR
STDGGATWSRIPGQPTGYLAHKGVLDSATGHLYLTS DTGGPYDGGKGRIWRYDTASG
AWQDVSPVAEADAYYGFSGLSVDRQKPGTLMA TAYSSWPDTQIFRSTDSGATWTQA
WDYTGYPNRSNRYTLDVSSVPWLWGASPAPPETAPKLGWMTEALEIDPDFSDRM MY

Figure 20 (continued)

GTGATVYGTEDLTSWDSGGTFRITPMVKIEETAVNDLASPPSGAPLLSALGDIGGFRHT
DLDAPDLMYTSPNLDSTTSLDFAESSPGTVVRVGNSDAAPHIGFSTDNGANWFQGSEP
SGVTGGGTVAADAGSGFVWSPEGAGVHHTGGTSWTASTGIPAGATVESDRKNPEK
FYGFEAGTFYVSTDGGATFTAETGLPAEGNVRFQALPGTEGDIWLAGGSDTGAYGLW
RSTDGATFTKSAGVEQADSVFGKAAPGASYRTFVSAKIGCVRGIFRSTDAGASWTR
INDDAHQWGBTGAATGDPRVYGRVVNSTNGRGIQVGETSDSGGGTDPGTDPTDPG
TDPGPEQPADAACAVTYAVTNQWPFGQADVTVTNTGDAAYNGWKLGSFPQQQIS
QIWNASHRQDGVKVTVDAGWNGTVAPGSSAGFGFTGSWAGSNAEPAAFTLDQACT
VG*

>SACTE_4343|extracellular solute-binding protein family 5||GI:344317489 (SEQ ID NO:41)
MRGAKSAKWVAGAAIIALAATACGGGDSDSDNGAKGAVDADGIFSVEVGEQPQNPLQP
ANTMESNGSIVTDAIFSQLVDYDPDGKLEMINAESVETTDKLWTVKLKKDWKFHDGT
PVTADSYVKAWNWAANIENAQTNASWFADIKYADVHPDGEAKPKSDAMSGLKKV
DDYTFTIELNSAVPYFSYKLGTVFSPLPESFYADPKAAGEKPVGNAYKFVSWDHKKQ
IKVVRNDDYKGPDKAKNGGVIFKNYTTLETAYEDLKSGNVDLRQIGPKDLPVYRADL
EDRAVDKAYSQVTLGVAMYTDQWKNTDPKVLQGLSMAIDRDTITKTVLQGTREPAT
GWVAKGVLYQENVAGDVTKYDPAKAKALIKEGGGVPGNEIFIQFNADGGHKEWIEA
VCNSITQATGVKCTGDSKADFQADLNARDAKQVKSFYRSGWVLDYPVNANFISDLFRT
GAAGNNNGFFSNKDLDAKIKAADSAASLDDSVKAYQEIEKELVNMPSIPLWYYKVNAG
YSENVKNDYAQDGDPLTEVQVIK*

>SACTE_1546|bacterioferritin||GI:344314774 (SEQ ID NO:42)
MQGDPEVLEFLNEQLTAELTAIJNQYFLHAKMQDHRGWTKLAKHTRAESFDEMKAEL
TDRILLLDGLPNYQRLFHVVRVGQTTEMFQADRQVEVEAIDRLRRGVDLMRAKSDITS
NIFERILEDEEHHDYLDTQLELIEKLGEPLYLAQVIEQVEL*

>SACTE_3590|phosphatidylinositol-specific phospholipase C X region||GI:344316754 (SEQ ID NO:43)
MSPYTATRRTFLT GALAAATGVVLGGTPALAAPARVLGTQDW MGALADSTPLRRLTIP
GTHNAGARYGGPWTECQNTTVAEQLGSGIRFLDVR CRITGDAFAIHGASYQNL MFGD
VLIACRDFLAHPSETVLMRVKQEYSEESDAAFRQIFDLYLDGKGWRPLFRLDPTLPDL
GGARGKVVLADNGGLPGVRYADPAVFDIQDDYMAEPFGKYPKIEAQFRKAAQQPGK
LFMNYVSTAALLPPRSNADRLNPQVHTFLDGSEAAGWTGLGIVPLDYPATRPGLVESLI
RHNPVA*

>SACTE_2172|citrate synthase I||GI:344315379 (SEQ ID NO:44)
VSEHTNNAVVLRYGDDEYTPVIDSTVGDKGFDIGKL RANTGLVTLDSGYGNTAA YKS
AITYLDGEQGILRYRGYPIEQLAESSTFLEVAYTLINGDLPKVDELSAFKNEITQHLLHE
DVKRFFDGFPRDAHPMAMLSSVVSALSTFYQDSHNPFDEEQRHLSTIRLLAKLPTIAAYA
YKKSIGHPFVYPRNDLGYVENFLRMTFSVPAQEYVDPIVVSALEKLLILHADHEQNCST
STVRLVGSSQANMFASISAGISALWGPLHGGANQSVLEMLEGIQANGGDVDSFIQKVKN
KEDGVRLMGFGHRYKSFDPRAKIKA AAHDV LSSLGKSDELL DIALKLEE HALSDDYF

Figure 20 (continued)


VSRNLYPNVDFYTGLIYRAMGFPTEMFTVLFALGRPGWIAQWHEMIKEPGSICRPRQI
YTGEVLRDFVPVESR*

>SACTE_5668|TAP domain protein|GI:344318769 (SEQ ID NO:45)
MTKragilvavgatvaglvtaVPSAASAPGAPGAAAPLKWtACGTkAYPTQQCATV
RAPLDHDPGRQVTLaLARIPTAKTSQGPPLLNPGGPGGSGLSMAGFVASSLPakla
AQyDViGFDPRGVGRSSPAldcVpkhfdPVRDtvPGSPRDERTNRERAASFADACGeK
HGDLLPFMDTVStAKLDVIRRALGARQInYFGSYGTYLGAVYAKLFPERVRLVLD
SIVDPDGWVYEDNLGQDYAFDARHKAFAAWVAKNDATYRLGTDPAKVEAAWYRMR
AAVKKHPAAGKVGPSELEDTFLPGGYYNGYWPQLAEAFAAVNDKDEDALATAyDDF
AAVDASGDNgySVYTAVQCRDTGWPKSWTWRNNTWQAHRKAPFMSWNNTWYNAP
CATWPVAPLRPVVRTNREIPPALLFQATDDAATPYEGGLSMHRKLKGSLVVEEGGN
HGISLSGNDCLDAHLIAyLTDGTLPRSGSGADAVCDALPEPEAAATAKAATGQKGS
TLHSLLGFRG*

>SACTE_6428|chitin-binding domain 3 protein|CBM33|GI:344319509 (SEQ ID NO:46)
MNCHDRINLRGWTRLSGLFVAVLCLLPWTGTAEAHGSVVDPASRNYGCWLWGSD
FQNPAMAQEDPMCWQAQADPNAMWNWNGLYRNESAGNFPAVIPDGQLCSGRTE
GGRYNALDTVGAWQATDITDDFTVRLEDQASHGADYFRVYVTEQGFDPТАQPLTWGA
LDLVAETGRYGPSTSYEIPVSTSGYTGRHVYTIWQASHMDQTYFLCSDVNFG*

>SACTE_0366|alpha-L-rhamnosidase|GH78|GI:344313621 (SEQ ID NO:47)
VISRRRLSTTAATAALAAVSSPAARAAPADTAAGRLRVTGPTVEYVRRPLGLDVSRP
RLSWPLASDHDPDHGQSAYQVRVATSPDRLARPDPWDGKVVSPSTSVLVPYAGPALVSR
TRYHWSVRVWDQDGRVSAWSEPSWWETGLDEADWSAGWIGAPAALTSSPSLEAAS
WIWFPEGDPAVGAPAATRWFRGRVIEPEGVTRARLVMtADDGFTALVDGVQVARTEP
DGPAENWRRPVVVDVT AHLSPGSRVVAVTATNAVDGPAGLLGALELTADGAVTLAT
GTGWRATDREP DGDWASGGYDDTGWPAAA VLPWGSGPWGEVRAALSPATQLTEF
RLGRKRVARARLYSTALGLYEVFLNGARVGEDRLAPGWTDYRKRVQYQTYDVTALLR
SGGNALGVTLAGPWYAGNIAWFGPHQYGERPAVLAQLEVFTDGSIERVLSGTGWAAA
TGPVTATDLMAGEEYDARLETDGWSRAGFDASGWLAEEAVEGVTAVPVAADVACR
VERELTAREVTEPEPGVYVFDLGQNMQVTVRLLVSGPAGTTVRLRHAEVLPDGTLYT
ANLRTARATDTYTLRGGGPETYEPRFTFHGFRYVEVTGFPGRPGPDAVVGRVIHTSAPF
TMAFSTDVPMULDRLHSNITWGQRGNFLSVPTDTPARDERLGWTGDINVFAPTAAYTME
SARFLGKWLQLDRDDQLADGAFPNVAPDLPVGSGAAGWGDAGVTVPWALYQAYGD
VRVLEQSWSSMVAWLEYLQAHSDGLLRPADGYGDWLNIEDETPKDVIGTAYFAHSAD
LTARTAEVLGKDGPYRTLSGRVRDAFRAAYVGDDGRVKGDTQTAYVLALSMDLLEP
GDRAPAADRLVALIEAKDWHLSTGFLGTPRLLPVLTDTGHTDVAYRLTRRTFPSWGY
QIDRGATTMWERWDSVRPDGGFQDAGMNSFNHYAYGSVGEWMYANIAGIAPAAPGF
REIRVRPRPGGGVHRAEARFDsLYGPVTRWTSDDGGFALRVVLPANTTAEVWVPGGD
GRSSVRGTAVFLREDGCAVFAAGSGIHRFTAPA*

Figure 21 A-B



CellLoc_CRM3a

Page 1 of 4

5' ATGGGACATCACCACATCATCACCATCACCATGCATCCGAAACCTGTACTTCCAGGCATC
60
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 M G H H H H H H H H A S E N L Y F Q A I
o
5' GCCATGGatccgaacaatgacgactggctgcattttttaaggtaaaaatagtggacatg
120
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 A M D P N N D D W L H V E G N K I V D M
o
5' tacggtaatcaggcttcggctgaccggctgcactggttttggattcaataccggtaaccat
180
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 Y G N Q V W L T G C N W F G F N T G T N
o
5' gtgttttgcggatcatggaggctgcaatatggaggaaaggcccataagggttatggcgacaga
240
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 V F D G V W S C N M R E A L K G M A D R
o
5' ggaataaaattttttggagatataccatattttcaacagaatttgctgttatcaatggctcaagga
300
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 G I N F L R I P I S T E L L Y Q W S Q G
o
5' atatatccccaaagcaaatgttaatgattttgtaaatccggagctgaaaaggaaacacgc
360
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 I Y P K A N V N D F V N P E L K G K N S
o
5' ctttagcttttgactttgccgttcatgtgtgcacaaaggatccggaaataagataatggtg
420
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 L E L F D F A V Q C C K E F G I K I M V
o
5' gatatacacagtcggcaacagatgcctatggggcatatgtatctttatggtatgacgg
480
o ++++++||||| ++++++| ++++++| ++++++| ++++++|
o **|||||** **|||||**
1 D I H S P A T D A M G H M Y P L W Y D G

Figure 22 (SEQ ID NOs: 63 & 64)

CelLcc_CBM3a

Page 2 of 4




5'	caatttacaacagagatatggattcaacttggagtgggtgacggaaagatataaaaat	
o	540
o		
1	Q F T T E I W I S T L E W L T E R Y K N	
o	
5'	gatgacacaattttgcactggacctaaaaatgaggctcacggcacccggcagcgaa	
o	600
o		
1	D D T I L A L D L K N E P H G T P G S E	
o	
5'	ttaatggccaaatggatgggtccacggatttgaacaactggaaagcatgctgtgaaaca	
o	660
o		
1	L M A K W D G S T D L N N N W K H A A E T	
o	
5'	tgcgcaaagagaatccttgcaataaatccgaatattttatttgtggtagaaggagtgaa	
o	720
o		
1	C A K R I L A I N P N I L I V V E G V E	
o	
5'	gttatccaaagcctggctatgattataccgcagtggacgaaatggggaaaagagagtaaa	
o	780
o		
1	V Y P K P G Y D Y T A V D E W G K E S K	
o	
5'	tatttctataactggggggaggaaatttaagaggagtcaaggattatccattgacatt	
o	840
o		
1	Y F Y N W W G G N L R G V R D Y P I D L	
o	
5'	ggcaaggcatcagaaggcagctgtataactcacctcacgattacggccccctgtacataaa	
o	900
o		
1	G K H Q K Q L V Y S P H D Y G P L V H K	
o	
5'	caaccttggtttatgaaggcttaacaaagaaactttgtataatgattgtggagagat	
o	960
o		
1	Q P W F Y E G F N K E T L Y N D C W R D	

Figure 22 continued (SEQ ID NOs: 63 & 64)

CelLcc_CBM3a

Page 3 of 4

1140

1200

1260

1320

1380

1440

1500

1560

1620

aataaaatatcccacacttttgtgtataatgc当地
N K I S H T F W C Y N A N S G D T G G L
gtataactatgatttattaccctggcgaagaaaaatatgc当地
V Y Y D F I T W D E E K Y A L L K P A L
tggcagacagaggacggaaagttaataggccattgacataccttggccaat
W Q T E D G K F I G L D H Q I P L G S N
ggagGTTAAACGGACTCCCCTAAAGTGCACCCATACCAATAACGGCAGTCCGACT
G G L N A T P T K G A T P T N T A T P T
AAGTCGGCAACGGCAACGCCACTCGCCCCAGCGTACCGACCAAATACTCCGACTAAC
K S A T A T P T R P S V P T N T P T N T
CCGGCGAACACCCCACTAACCGCTAACCTGAAGGTTCAATTAACTCCAACCCAAC
P A N T P V S G N L K V E F Y N S N P S
GACACAACGAATAAGCATCAATCCGAGTCAAAGTCACGAACACTGGCAGITCAGCTATC
D T T N S I N P Q F K V T N T G S S A I
GATCTGTCGAAACTGACCCCTCGTTACTACTATAACGGTTGATGGCAAAAAGATCAGACC
D L S K L T L R Y Y Y T V D G Q K D Q T
TTTGCTGGGACCATGCAACATCGGTAGCAATGGTTCTTATAACGGCATTACTTCT
F W C D H A A I I G S N G S Y N G J T S

CelLcc_CBM3a from CjaA
CelLcc_CBM3a from CjaA
CelLcc_CBM3a from CjaA

Figure 22 continued (SEQ ID NOs: 63 & 64)

CellLoc_CBM3a

Page 4 of 4









5'	AATGTAAAAGGCACCTTGTAAGATGTCAAGTAGCACCAACAATGCTGATACTACCTG	
o		1680
o	 1681-1740: 40aa Unlabelled CBM3 from CjaA	
1	N V K G T F V K M S S S T N N A D T Y L	
o		1740
o	 1801-1860: 60aa Unlabelled CBM3 from CjaA	
1	E I S F T G G T L E P G A H V Q I Q G R	
o		1800
o	 1861-1873: 13aa Unlabelled CBM3 from CjaA	
1	F A K N D W S N Y T Q S N D Y S F K S A	
o		1860
o	 1874-1878: 5aa Unlabelled CBM3 from CjaA	
1	S Q F V E W D Q V T A Y L N G V L V W G	
o		1872
o	 1879-1882: 4aa Unlabelled CBM3 from CjaA	
1	K E P G	

Figure 22 continued (SEQ ID NOs: 63 & 64)




Figure 23




Figure 24




Figure 25 A-B

Fraction Number from Figure 7




Figure 26 A-B

Figure 26 C-D





Figure 27A








Figure 27B

Figure 28



MS #	ID	Function	CAZY	CBM	AA	Best BLAST
3	SACTE_4738	glycoside hydrolase family 16	GH16	CBM33	627	beta-1,3-glucanase
4	SACTE_3199	chitin-binding domain 3 protein	CBM33.2	CBM33.2	363	cellulose oxygenase
5	SACTE_3159	chitin-binding domain 3 protein	CBM33.3	CBM33.2	363	
6	SACTE_3199	chitin-binding domain 3 protein	CBM33.2	CBM33.2	363	
7	SACTE_3159	chitin-binding domain 3 protein	CBM33.2	CBM33.2	363	
8	SACTE_0265	glycoside hydrolase family 10	GH10	CBM2	459	xylanase
9	SACTE_5083	putative RNA polymerase	N/A	N/A	438	
10	SACTE_0237	1, 4-beta cellobiohydrolase	GH6	CBM2	587	1,4-beta cellobiohydrolase
11	SACTE_0482	glycoside hydrolase family 9	GH6	CBM2	457	endo-1,4-beta-glucanase
12	SACTE_4755	beta-1,3-glucanase	GH64		409	beta-1,3-glucanase
13	SACTE_0482	glycoside hydrolase family 9	GH6	CBM2	457	
14	SACTE_0237	1, 4-beta cellobiohydrolase	GH6	CBM2	587	1,4-beta cellobiohydrolase
15	SACTE_0549	glucan endo-1,3-beta-D-glucosidase	GH16	CBM54	307	beta-1,3-glucanase
16	SACTE_0236	glycoside hydrolase family 48	GH48	CBM2,37	955	cellulose 1,4-beta-cellubiosidase
17	SACTE_0236	glycoside hydrolase family 48	GH48	CBM2,37	955	
18	SACTE_3457	chitatanase	GH46		290	chitatanase
19	SACTE_0236	glycoside hydrolase family 48	GH48	CBM2,37	955	cellulose 1,4-beta-cellubiosidase
20	SACTE_3717	carbohydrate-binding, CenC-like	GH9	CBM2,4	809	endo-1,4-beta-glucanase
21	SACTE_2347	cellulose-binding family II	GH5,CE3	CBM2,37	563	secreted beta-mannosidase
22	SACTE_2347	cellulose-binding family II	GH5,CE3	CBM2,37	563	secreted beta-mannosidase
23	SACTE_2347	cellulose-binding family II	GH5,CE3	CBM2,37	563	secreted beta-mannosidase
24	SACTE_3629	Ricin B lectin	GH93	CBM42,13	593	exo-alpha-L-1,3-arabinanase
25	SACTE_4363	putative secreted protein	GH55	CBM56,54,57	606	endo-beta-1,3-glucanase

Figure 28 (continued)

A**B**

Gene Locus	CAZy	MW (kDa)	Microarray rank ^a			Present in secretomes	Diagnostic substrate			
			cellulose	xylan	chitin		MUG	MUC	MUM	MUX2
SACTE_0265	GH10	49.8	20	519	3530	yes	-	-	-	+
SACTE_0358	GH11	37.2	13	160	593	yes	-	-	-	+
SACTE_2548	GH1	90.8	4197	4135	5330	no	-	-	-	-
SACTE_2286	GH2	55.3	28	2533	3012	no	+	-	-	-
SACTE_4737	GH1	52	702	791	1718	no	+	-	-	-

^a Out of 6152 genes total, ranking by transcript intensity, with highest rank equal 1.

Figure 29A-B

Figure 30





Figure 31






Figure 32

Cellulose/Glucose
(percent of total carbon)






Figure 33



glucose \cap sigmacell = 4
glucose \cap xylan = 45
sigmacell \cap xylan = 16
glucose \cap sigmacell \cap xylan = 4

glucose / (sigmacell \cup xylan) = 117
sigmacell / (xylan \cup glucose) = 9
xylan / (glucose \cup sigmacell) = 46



switchgrass \cap afexSG = 36
switchgrass \cap ilSG = 35
afexSG \cap ilSG = 27
switchgrass \cap afexSG \cap ilSG = 27

switchgrass / (afexSG \cup ilSG) = 19
afexSG / (ilSG \cup switchgrass) = 0
ilSG / (switchgrass \cup afexSG) = 8

Figure 34A-B

1

**METHOD AND COMPOSITIONS FOR
IMPROVED LIGNOCELLULOSIC
MATERIAL HYDROLYSIS**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a continuation application of U.S. patent application Ser. No. 13/709,971, filed Dec. 10, 2012, which claims benefit from U.S. Provisional Application 61/579,301 filed Dec. 22, 2011 and U.S. Provisional Application 61/579,897 filed Dec. 23, 2011, all of which are incorporated herein by reference for all purposes.

**STATEMENT REGARDING FEDERALLY
SPONSORED RESEARCH OR DEVELOPMENT**

This invention was made with government support under DE-FC02-07ER64494 awarded by the US Department of Energy and GM094584 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND

Cellulose is the most abundant organic polymer on Earth and represents a vast source of renewable energy. Most of this energy is stored in the recalcitrant polysaccharide cellulose, which is difficult to hydrolyze because of the highly crystalline structure, and in hemicellulose, which presents challenges because of its structural diversity and complexity. Plant cell walls are approximately composed in pinewood of lignin (30% by weight), hemicellulose (glucomannan, 20%, arabinoxylan, 10%), and crystalline cellulose (40%), which presents a major barrier to efficient use. In terrestrial ecosystems, cellulolytic microbes help drive carbon cycling through the deconstruction of biomass into simple sugars. The deconstruction is largely accomplished through the action of combinations of secreted glycoside hydrolases (GHs), carbohydrate esterases (CEs), polysaccharide lyases (PLs), and carbohydrate binding modules (CBMs) (Baldrian and Valaskova, 2008; Cantarel, et al., 2009; Lynd, Weimer, et al., 2002; Schuster and Schmoll, 2010). Consequently, organisms from many lignocellulose-rich environments and their enzymes are being studied for new insights into overcoming this barrier.

In order to obtain the hydrolysis of crystalline cellulose, enzymes must cleave three types of glycosidic bonds. These enzymes are endocellulases, which cleave beta-1,4 glycosidic bonds that reside within intact cellulose strands in the crystalline face, non-reducing-end exocellulases, which remove cellobiose units from the non-reducing end of cellulose strands, and reducing-end exocellulases, which remove glycosyl units from the reducing-end of a cellulose strand. The endocellulolytic reaction is essential because it creates the non-reducing and reducing ends that serve as the starting point for exocellulolytic reactions. The exocellulolytic reactions are essential because they remove glycosyl groups in a processive manner from the breakages in the cellulose strand introduced by the endocellulases, thus amplifying the single initiating reaction of the endocellulases.

Trichoderma reesei and *Clostridium thermocellum* are well-characterized cellulose-utilizing organisms (Merino and Cherry, 2007; Bayer et al., 2008; Wilson, 2011). *T. reesei* is a slow-growing eukaryote fungus that secretes enzymes containing glycoside hydrolase (GH) domains fused to car-

2

bohydrate binding domains, while *C. thermocellum* is a strictly anaerobic prokaryote that predominantly assembles GHs and carbohydrate-binding molecules (CBMs) into a large complex called the cellulosome. Enzymes from these free-living organisms cleave polysaccharides using general acid-base catalyzed hydrolytic reactions (Vuong and Wilson, 2010). Moreover, fungal and microbial communities associated with termites (Scharf et al., 2011) shipworms (Luyten et al., 2006), and rumen (Hess et al., 2011) contribute these types of hydrolytic enzymes to their respective anaerobic niches.

Some free-living aerobes such as *Cellvibrio japonicus* (Ueda 107) (DeBoy et al., 2008), *Streptomyces* (Schlochtermeier et al., 1992; Wilson, 1992; Forsberg et al., 2011), 15 *Thermoascus aurantiacus* (Langston et al., 2011; Quinlan et al., 2011) and *Serratia marcescens* (Vaaje-Kolstad et al., 2010) also grow on biomass polysaccharides. Recent work with some of these organisms has identified that the structurally related fungal GH61 (Langston et al., 2011; Quinlan 20 et al., 2011) and bacterial CBM33 (Forsberg et al., 2011) families of proteins catalyze a previously unrecognized oxidative breakage of glycosidic bonds. This reaction is thought to be an endo-cleavage, with the oxidation reaction yielding gluconate and keto-sugars instead of the typically 25 observed reducing and non-reducing sugars obtained from hydrolytic cellulases.

Actinobacteria in the genus *Streptomyces* are an ecologically important group, especially in soil environments, where they are considered to be vital players in the decomposition 30 of cellulose and other biomass polymers (Cantarel et al., 2009; Crawford et al., 1978; Goodfellow and Williams, 1983; McCarthy and Williams, 1992). *Streptomyces* are able to utilize a wide range of carbon sources, form spores when resources are depleted, and produce antimicrobial 35 secondary metabolites to reduce competition (Goodfellow and Williams, 1983; Schlatter et al., 2009).

Although a large number of *Streptomyces* species can grow on biomass, only a small percentage (14%) have been shown to efficiently degrade crystalline cellulose (Wachinger, Bronnenmeier, et al., 1989). Furthermore, the secreted 40 cellulolytic activities of only a few species have been biochemically characterized, and still fewer species have been examined to identify key biomass degrading enzymes (Ishaque and Kluepfel, 1980; Semedo et al., 2004). *Streptomyces reticuli* is one of the best-studied cellulose- and chitin-degrading soil-dwelling *Streptomyces*; functional 45 analyses of several important cellulases and other hydrolytic enzymes have been reported (Wachinger, Bronnenmeier, et al., 1989; Schlochtermeier, Walter, et al., 1992; Walter and Schrempp, 1996).

Furthermore, polysaccharide monooxygenase (PMO) 50 activity with cellulose was identified using the CBM33 protein from *Streptomyces coelicolor* (Forsberg, et al., 2011), which suggests *Streptomyces* may use both hydrolytic and oxidative enzymes to deconstruct biomass. With the tremendous amount of sequence data collected in the past few years, and despite the view that *Streptomyces* make 55 important contributions to cellulose degradation in the soil, genome-wide analyses of cellulolytic *Streptomyces* have not been reported.

In addition to their putative roles in carbon cycling in the soil, *Streptomyces* may also potentiate biomass deconstruction in insects through symbiotic associations (Bignell, Anderson, et al., 1991; Pasti and Belli, 1985; Pasti, Pometto, 60 et al., 1990; Schafer, et al., 1996). Recent work has identified cellulose degrading *Streptomyces* associated with the pine-boring woodwasp *Sirex noctilio*, including *Streptomyces* sp.

SirexAA-E (ActE) (Adams, et al., 2011). *S. noctilio* is a highly destructive wood-feeding insect that is found throughout forests in Eurasia and North Africa and is spreading invasively in North America and elsewhere (Bergeron, et al., 2011). While the wasp itself does not produce cellulolytic enzymes, evidence supports the role of a symbiotic microbial community that secretes biomass-degrading enzymes to facilitate nutrient acquisition for developing larvae in the pine tree (Kukor and Martin, 1983).

The white rot fungus, *Amylostereum areolatum*, is the best-described member of this community, and the success of *Sirex* infestations is thought to arise from the insect's association with this cellulolytic fungal mutualist. However, work with pure cultures has suggested that ActE and other *Sirex*-associated *Streptomyces* are more cellulolytic than *A. areolatum* (Adams, et al., 2011).

Optimal activity in the CBM33 enzymes apparently requires the addition of a transition metal ion such as Cu(II), Fe(III), or Mn(II) and an external reducing agent. In the laboratory, the reducing agent can be provided by ascorbate. In natural systems, the reducing function is most likely provided by another redox active protein such as cellobiose dehydrogenase (Langston et al., 2011; Quinlan et al., 2011) or some other presently unknown protein.

Needed in the art are improved compositions and organisms for digestion of lignocellulosic materials.

BRIEF SUMMARY

The invention relates generally to methods and compositions for digesting lignocellulosic material and more particularly to methods that involve exposing the material to secretome derived from *Streptomyces* sp. ActE.

In a first aspect, the present invention is summarized as a method of digesting a lignocellulosic material comprising the step of exposing the material to an effective amount of *Streptomyces* sp. ActE secretome preparation such that at least partial lignocellulosic digestion occurs.

In some embodiments of the first aspect, the preparation is a supernatant preparation obtained from a *Streptomyces* sp. ActE culture. In some embodiments of the first aspect, the preparation is obtained from *Streptomyces* sp. ActE grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass and chitin. In some embodiments of the first aspect, the lignocellulosic material is selected from the group consisting of materials that comprise at least 75% cellulose, cellulose/hemicelluloses, xylose, biomass and chitin.

In a second aspect, the present invention is summarized as a purified preparation comprising the *Streptomyces* sp. ActE secretome.

In some embodiments of the second aspect, the preparation is a supernatant preparation obtained from a *Streptomyces* sp. ActE culture. In some embodiments of the second aspect, *Streptomyces* sp. ActE is grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass and chitin.

In a third aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_0237 (GH6) (SEQ ID NOs:1 and 17) gene or expression product thereof.

In a fourth aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_0236 (GH48) (SEQ ID NOs:2 and 18) gene or expression product thereof.

In a fifth aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_3159 (CBM33) (SEQ ID NOs:3 and 19) gene or expression product thereof.

In a sixth aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_0482 (GH5) (SEQ ID NOs:4 and 20) gene or expression product thereof.

In a seventh aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_0265 (GH10) (SEQ ID NOs:5 and 21) gene or expression product thereof.

In a eighth aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_2347 (GH5) (SEQ ID NOs:6 and 22) gene or expression product thereof.

In a ninth aspect, the present invention is summarized as a composition useful for digesting lignocellulosic material comprising SActE_0237 (GH6) (SEQ ID NOs: 1 and 17), SActE_0236 (GH48) (SEQ ID NOs: 2 and 18), SActE_3159 (CBM33) (SEQ ID NOs: 3 and 19), SActE_0482 (GH5) (SEQ ID NOs: 4 and 20) and gene or expression product thereof.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for cellulose utilization. In these embodiments the composition can additionally comprise at least one member selected from SActE_0265 (GH10) (SEQ ID NOs: 5 and 21) and SActE_2347 (GH5) (SEQ ID NOs: 6 and 22) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for xylan release. By "release," we mean degradation, such as hydrolysis, and release of an important or desired product. In these embodiments the composition can additionally comprise at least one member selected from SActE_0265 (GH10) (SEQ ID NOs: 5 and 21), SActE_0358 (GH11) (SEQ ID NOs: 8 and 24), SActE_0357 (CE4) (SEQ ID NOs: 7 and 23), SActE_5978 (PL1) (SEQ ID NOs: 16 and 32) and SActE_5230 (xylose isomerase) (SEQ ID NOs:33 and 48) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for chitin release. In these embodiments the composition can additionally comprise at least one member selected from SActE_4571 (GH18) (SEQ ID NOs:34 and 49), SActE_2313 (CBM33) (SEQ ID NOs:35 and 50), SActE_4246 (GH18), (SEQ ID NOs:36 and 51) SActE_3064 (GH19) (SEQ ID NOs:37 and 52), and SActE_5764 (GH18) (SEQ ID NOs:38 and 53) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for biomass degradation. In these embodiments the composition can additionally comprise SActE_5457 (GH46) (SEQ ID NOs: 14 and 30) gene or expression products thereof.

5

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for mannan release. In these embodiments the composition can additionally comprise SactE_2347 (GH5) (SEQ ID NOS: 6 and 22) gene or expression products thereof.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for beta-1,3-glucan release. In these embodiments the composition can additionally comprise at least one member selected from SActE_4755 (GH64) (SEQ ID NOS:13 and 29) and SActE_4738 (GH16) (SEQ ID NOS:12 and 28) genes or expression products thereof. In a preferred embodiment, the composition comprises both of the genes or expression products.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for pectin cleavage. In these embodiments the composition can additionally comprise SActE_1310 (PL3) (SEQ ID NOS:9 and 25) gene or expression products derived thereof.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for alginate release. In these embodiments the composition can additionally comprise SActE_4638 (SEQ ID NOS:11 and 27) gene or expression products derived thereof.

In some embodiments of the third, fourth, fifth, sixth, seventh, eighth, and ninth aspects, the composition is optimized for galactose release. In these embodiments the composition can additionally comprise SactE_5647 (GH87) (SEQ ID NOS:15 and 31) gene or expression products derived thereof.

In a tenth aspect, the present invention is summarized as a composition useful for xylan degradation comprising SActE_0265 (GH10) (SEQ ID NOS:5 and 21) and SActE_0358 (GH11) (SEQ ID NO:8 and 24) gene or expression products thereof.

In some embodiments of the tenth aspect, the composition additionally comprises SActE_0265 (GH10) (SEQ ID NOS:5 and 21), SActE_0358 (GH11) (SEQ ID NOS:8 and 24), SActE_0357 (CE4) (SEQ ID NOS:7 and 23), SActE_5978 (PL1) (SEQ ID NOS:16 and 32), and SActE_5230 (xylose isomerase) (SEQ ID NOS:33 and 48) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In an eleventh aspect, the present invention is summarized as a composition useful for biomass degradation comprising SActE_0237 (GH6) (SEQ ID NOS:1 and 17), SActE_0482 (GH5) (SEQ ID NOS:4 and 20), SActE_3159 (CBM33) (SEQ ID NOS:3 and 19), SActE_0236 (GH48) (SEQ ID NOS:2 and 18), SActE_3717 (GH9) (SEQ ID NOS:10 and 26), SActE_0265 (GH10) (SEQ ID NOS:5 and 21), SActE_0358 (GH11) (SEQ ID NOS:8 and 24), SActE_2347 (GH5) (SEQ ID NOS:6 and 22) and SActE_1310 (PL3) (SEQ ID NOS:9 and 25) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In a twelfth aspect, the present invention is summarized as a composition useful for cellulose degradation comprising SActE_0237 (GH6) (SEQ ID NOS:1 and 17), SActE_0482 (GH5) (SEQ ID NOS:4 and 20), SActE_3159 (CBM33) (SEQ ID NOS:3 and 19), SActE_0236 (GH48) (SEQ ID NOS:2 and 18), SActE_2347 (GH5) (SEQ ID NOS:6 and 22), and SActE_0265 (GH10) (SEQ ID NOS:5 and 21) genes

6

or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In a thirteenth aspect, the present invention is summarized as a method for digesting a lignocellulosic material, comprising exposing the material to a sufficient amount of a composition of any one of the third to eighth aspects of the invention, wherein the exposed material is at least partially digested.

10 In a fourteenth aspect, the present invention is summarized as a purified preparation of *Streptomyces* sp. ActE, wherein the *Streptomyces* sp. ActE has been grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass, and chitin.

15 In a fifteenth aspect, the present invention is summarized as a purified preparation of *Streptomyces* sp. ActE, wherein the *Streptomyces* sp. ActE has been grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon in the substrate is derived from pretreated lignocellulosic material.

20 In some embodiments of the fifteenth aspect, the pre-treated material has been exposed to pretreatment selected from the group consisting of acid hydrolysis, steam explosion, ammonia fiber expansion (AFEX), organosolve, sulfite pretreatment to overcome recalcitrance of lignocellulose (SPORL), ionic liquids, metal-catalyzed hydrogen peroxide, 25 alkaline wet oxidation and ozone pretreatment. In some embodiments of the fifteenth aspect, the pretreated material is wood.

25 These and other features, objects, and advantages of the present invention will become better understood from the description that follows. In the description, reference is made to the accompanying drawings, which form a part hereof and in which there is shown by way of illustration, not limitation, embodiments of the invention. The description of preferred embodiments is not intended to limit the invention to cover all modifications, equivalents and alternatives. Reference should therefore be made to the claims recited herein for interpreting the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

30 The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

35 The present invention will be better understood and features, aspects and advantages other than those set forth above will become apparent when consideration is given to the following detailed description thereof. Such detailed description makes reference to the following drawings, 40 wherein:

45 FIGS. 1A-B are sets of pictures showing growth of ActE in minimal medium containing filter paper as the sole carbon source. (A) Growth of ActE, *Streptomyces coelicolor*, and *Streptomyces griseus* in minimal medium for 7 days at 30° C. and pH 6.9. The expanded image shows small colonies of *S. coelicolor* and *S. griseus* forming on the surface of the paper. (B) Growth of ActE and *Trichoderma reesei* Rut-C30 for 7 days at 30° C. and pH 6.0.

50 FIGS. 2A-C are sets of graphs demonstrating reactions of ActE secretomes and SPEZYME-CP. (A) HPLC of sugars released from cellulose (1, cellobiose; 2, cellobiose; 3, glucose) and quantification of glucose equivalent (insert).

(B) Reducing sugars released from xylan and mannan by the secretomes of ActE grown on cellulose and xylan. (C) Total reducing sugar released from ionic liquid-switchgrass (IL-SG) or AFEX-switchgrass (AFEX-SG) in reactions of the ActE cellulose, AFEX-SG, and IL-SG secretomes and Spezyme CP. Data represent the mean \pm s.d. from three experiments; * indicates P<0.01 compared with SPEZYME CP.

FIGS. 3A-B are tables illustrating composition of ActE secretomes identified by LC-MS/MS. (A) Carbohydrate Active Enzyme (CAZy) genes account for 2.6% of the 6357 predicted protein-coding sequences in the ActE genome. (B) Identity of most abundant proteins in the cellulose secretome proteins is sorted according to decreasing spectral counts (accounting for 95% of total spectral counts); corresponding spectral counts from other secretomes are also shown.

FIG. 4 is a systematic diagram showing genome-wide changes in expression during growth of ActE on AFEX-treated switchgrass (AFEX-SG) versus glucose. Nodes are genes (circles) or KEGG/CAZy functional categories (yellow triangles); edges indicate that the gene belongs to the indicated functional group as defined by either KEGG or CAZy analysis. Gene node sizes reflect expression intensity determined by microarray from growth on AFEX-SG as a log₂ ratio, where the genome-wide average transcriptional intensity was ~10.5 for both substrates. Node colors represent expression changes as the log₂ ratio of AFEX-SG/glucose transcript intensities.

FIGS. 5A-B are diagrams with a table showing expression of ActE CAZy genes on various carbon sources. (A) Hierarchical clustering of expression for 167 CAZy genes from the ActE genome during growth on the indicated substrates. (B) Identity of CAZy genes with distinct changes in expression observed in group 1 CAZy genes during growth in different carbon sources.

FIG. 6 is a set of scanning electron microscopy (SEM) images showing ActE grown on different carbon sources including glucose, cellulose, xylan, switchgrass, ammonia fiber expansion-treated switchgrass (AFEX-SG) and ionic liquid-treated switchgrass (IL-SG). ActE cells were grown in minimum medium with the indicated substrate as a sole carbon source for 7 days at 30° C. The scale bar indicates 5 μ m.

FIGS. 7A-B are sets of graphs demonstrating fractionation of the ActE cellulose secretome and assays of reactions with different polysaccharides. (A) Anion exchange chromatography was performed using the ActE cellulose secretome, and fractions were collected and analyzed by SDS-PAGE. Lowercase letters indicate protein identified by MALDI-TOF MS shown in FIG. 17. (B) Results from hydrolysis assays for reaction with filter paper (FP), xylan, mannan and beta-1,3 glucan as detected by DNS assay of each fraction. The percentage reactivity relative to the maximum activity observed for each substrate is shown. Error bars indicate the standard deviation, with n=3 for technical replicates.

FIGS. 8A-B are sets of diagrams showing temperature and pH profiles of the ActE secretome obtained from growth on AFEX-treated corn stover. (A) The effect of temperature on the deconstruction of AFEX-treated switchgrass (AFEX-SG) and ionic liquid-treated switchgrass (IL-SG). The relative activity of the ActE secretome was compared to the maximal rates determined for reaction with AFEX-SG (blue star), and IL-SG (red star) at pH 6.0. (B) The effect of pH on the AFEX-SG and IL-SG deconstruction activities in the indicated ActE secretomes. The maximal rates observed for AFEX-SG and IL-SG were at pH 7.0 (blue star) and pH 8

(red star), respectively. Reactions were carried out at 40° C. and the 0.1 M buffers used were citrate (pH 4.5), phosphate (pH 6-8), CHES (pH 9-10), and CAPS (pH 11). The reaction was performed for 20 h and the reducing sugar content was measured by DNS assay.

FIG. 9 is a systematic diagram showing genome-wide changes in expression during growth of ActE on substrate cellobiose versus glucose visualized as a Cytoscape interaction network. Nodes are genes (circles) or KEGG/CAZy functional categories (yellow triangles); edges indicate that the gene belongs to the indicated functional group as defined by either KEGG or CAZy analysis. Gene node sizes reflect expression intensity determined by microarray from growth on substrate as a log 2 ratio. Node colors represent expression changes as the log 2 ratio of substrate/glucose transcript intensities, where the genome-wide average transcriptional intensity was ~10.5 for both substrate and glucose. Transcripts with less than two-fold changes in expression intensity are colored white; transcripts with greater than two-fold increase in expression intensity during growth on substrate are shown as a red gradient; transcripts with greater than two-fold increase in expression intensity during growth on glucose are shown as a blue gradient.

FIG. 10 is a systematic diagram showing genome-wide expression changes for growth on the substrate cellulose versus glucose visualized as a Cytoscape interaction network. Other information is the same as that described in FIG. 9.

FIG. 11 is a systematic diagram showing genome-wide expression changes for growth on the substrate xylan versus glucose visualized as a Cytoscape interaction network. Other information is the same as that described in FIG. 9.

FIG. 12 is a systematic diagram showing genome-wide expression changes for growth on the substrate switchgrass versus glucose visualized as a Cytoscape interaction network. Other information is the same as that described in FIG. 9.

FIG. 13 is a systematic diagram showing genome-wide expression changes for growth on the substrate IL-treated switchgrass versus glucose visualized as a Cytoscape interaction network. Other information is the same as that described in FIG. 9.

FIG. 14 is a systematic diagram showing genome-wide expression changes for growth on the substrate chitin versus glucose visualized as a Cytoscape interaction network. Other information is the same as that described in FIG. 9.

FIGS. 15A-B are diagrams with a table showing expression of 167 predicted CAZy genes in ActE, highlighting group 2 genes. These genes showed no signal above the average genomic expression intensity (log 2=10.5). (A) Clustering of genes with similar expression profiles. (B) Additional information on group 2 genes including expression profile, SACTE_locus ID, CAZy family, and annotated function.

FIGS. 16A-B are diagrams with a table showing expression of 167 predicted CAZy genes in ActE, highlighting group 3 genes. (A) Clustering of genes with similar expression profiles. (B) Additional information on group 3 genes including expression profile, SACTE_locus ID, CAZy family, and annotated function.

FIG. 17 is a table illustrating proteins separated by ion exchange chromatography and identified by mass spectrometry.

FIG. 18 is a table showing spectra count of proteins identified on each substrate, where top 95% spectra covered were highlighted green, light purple, purple, blue, orange,

pink, light blue and yellow on glucose, cellobiose, cellulose, xylan, switchgrass, AFEX-SG, IL-SG and chitin, respectively.

FIG. 19 shows the nucleic acid sequences of the ActE genes.

FIG. 20 shows the amino acid sequences of the ActE genes.

FIG. 21A-B are graphs illustrating a comparison of specific activities of *Streptomyces* sp. ActE secretomes with SPEZYME CP. (A) depicts relative specific activity of ActE secretomes prepared from growth on cellulose or xylan and SPEZYME CP (100%) for reducing sugar release from xylan or mannan. (B) depicts relative activity (pH 6.0, 40° C.) of ActE cellulose secretome and CelLcc_CBM3a, an engineered *C. thermocellum* endo/exoglucanase, compared to SPEZYME CP. Total amounts of protein included in all reactions were equivalent.

FIG. 22 illustrates nucleotide and amino acid sequence of CelLcc_CBM3a. Construct described in US Patent Application Publication No.: US2010/037094 (Fox and Elsen).

FIG. 23 is a graph illustrating SDS-PAGE of *Streptomyces* sp. ActE secretomes obtained from growth on minimal medium containing different substrates (SG, switchgrass; CS, corn stover; UBLPKP, unbleached lodgepole pine kraft pulp; BSKP, bleached spruce kraft pulp; LP-SPORL, lodgepole pine pretreated by sulfite pretreatment to overcome recalcitrance of lignocellulose (SPORL)). Culture secretomes were separated after 7 days of growth at 30° C. by centrifugation and concentrated by ultrafiltration. Sample loading was normalized to total protein. The identities of proteins were determined from samples extracted from the SDS-PAGE gel. Among the 162 proteins accounting for 95% of spectral counts from the glucose secretome, most were intracellular proteins originating from cell lysis during growth, and were not detected in the polysaccharide secretomes.

FIG. 24 is a graph illustrating SDS-PAGE of time-dependent changes in the *Streptomyces* sp. ActE secretome obtained from growth on minimal medium containing cellulose. Culture secretomes were collected after 7 days by centrifugation and concentrated by ultrafiltration. The concentrated secretomes were incubated at 25° C. for the indicated times and analyzed. Protein bands with time-dependent decrease in intensity were excised from the gel and identified by LC-MS/MS.

FIGS. 25A-B illustrate synergy of recombined fractions from ion exchange chromatography. All reactions were prepared to contain the same total amount of protein.

FIGS. 26A-D are sets of graphs illustrating mannanase activity demonstrated in fractions containing various naturally truncated versions of SACTE_2347 (GH5). (A-B) depict proteins found in previous assayed fractions. (C) depicts Coomassie Blue staining of 12% polyacrylamide gel (PAGE) separation of different mannanase isoforms. Three polypeptide bands corresponding to SACTE_2347 (GH5) with molecular masses of ~57, ~45, and ~37 kDa. (D) depicts a zymogram performed in the presence of 0.5% mannan. The strong clearing zone in fraction F1 associated with the ~37 kDa isoform demonstrates how size reduction can increase the specific activity of a protein.

FIGS. 27A-B are sets of graphs illustrating ion exchange fractionation of *Streptomyces* sp. ActE secretome. (A) depicts an SDS-PAGE analysis of the fractionation of an ActE secretome by ion exchange chromatography. (B) depicts catalytic assays of the separate fractions at 40° C. for 20 h in 0.1 M phosphate buffer, pH 6.0, showing different

enzymes are capable of reacting with xylan, mannan, and cellulose. The reactivity of fractions marked with stars is also described in FIG. 25A.

FIG. 28 is a SDS-PAGE graph and a list illustrating mass spectral assignment of polypeptides from the *Streptomyces* sp. ActE secretome separated by ion exchange chromatography. FIG. 28A depicts an SDS PAGE of separated fractions annotated with identities of polypeptides determined by LC-MS analysis. FIG. 28B depicts information on the 10 identified proteins including gene locus, function, CAZY GH and CBM assignments, number of amino acid (AA) residues, and best BLAST result for relationship to another known enzyme. The reactivity of fractions marked with stars is also described in FIG. 25A.

FIGS. 29A-B are SDS-PAGE graphs and a table that 15 demonstrates the existence of xylanases from *Streptomyces* sp. ActE. Five ActE proteins were produced using cell-free translation as described in US Patent Application Publication No.: US2010/037094 (Fox and Elsen). (A) depicts a stain-free gel image of proteins produced by wheat germ cell-free translation (indicated by asterisks). (B) depicts a summary of protein information, expression and secretion data, and diagnostic assay results. Small molecule assays (MUG, methylumbelliferyl glucoside; MUC, methylumbelliferyl 20 celllobioside; MUM, methylumbelliferyl mannoside and MUX2, methylumbelliferyl xylobioside) were performed in 0.1 M phosphate buffer, pH 6.0, at 30° C. SACTE_0265 and SACTE_0358, highly expressed and secreted proteins during growth on xylan, are confirmed by these assays to be 25 xylanases. Results from three other non-secreted ActE enzymes are provided as controls.

FIG. 30 is a graph illustrating quantification of total 30 secreted protein obtained from *Streptomyces* sp. ActE grown on different substrates (AFEX-CS, AFEX corn stover; UBLPKP, unbleached lodgepole pine kraft pulp; BSKP, bleached spruce kraft pulp; LP-SPORL, lodgepole pine pretreated by SPORL).

FIG. 31 is a graph illustrating the temperature versus 35 activity profile of the *Streptomyces* sp. ActE secretome obtained from growth on cellulose. Hydrolysis activities were measured by DNS assay. Greater than 80% of maximal rates for cellulase and mannanase activity were observed at the range of 31-43° C., while greater than 80% of maximal rate for xylanase activity was observed in the range of 35-59° C.

FIG. 32 is a graph illustrating the pH versus activity profile of the *Streptomyces* sp. ActE secretome obtained from growth on cellulose. The maximal rate was observed at approximately pH 6. Buffers used in this study were 0.1 M citrate (pH 4.5), phosphate (pH 6-8), CHES (pH 9-10) and CAPS (pH 11).

FIG. 33 is a SDS-PAGE graph illustrating ActE induction 40 in medium containing as little as 20% cellulose.

FIGS. 34A-B are a set of Venn diagrams representing 45 95% of total proteins identified in LC-MS/MS analyses generated using VennMaster-0.37.5 (Kestler et al., 2008). (A) depicts secretomes obtained from growth on glucose, SigmaCell™, and xylan. (B) depicts secretomes obtained from growth on switchgrass, ammonia fiber expansion (AFEX)-SG, and IL-SG. For clarification, glucose \cap SigmaCell=4 represents the intersection of the two sets, while glucose/(SigmaCell \cap xylan)=117 represents the proteins 50 uniquely associated with growth on glucose as compared to SigmaCell. Other results are interpreted in a similar manner.

While the present invention is susceptible to various modifications and alternative forms, exemplary embodiments thereof are shown by way of example in the drawings and are herein described in detail. It should be understood,

11

however, that the description of exemplary embodiments is not intended to limit the invention to the particular forms disclosed, but on the contrary, the intention is to cover all modifications, equivalents and alternatives falling within the spirit and scope of the invention as defined by the appended claims.

DESCRIPTION OF EXEMPLARY EMBODIMENTS

In General

The present invention comprises many embodiments. In one embodiment, the invention is a method of digesting a lignocellulosic material, comprising the step of exposing the material to an effective amount of *Streptomyces* sp. ActE secretome preparation such that at least partial lignocellulosic digestion occurs. In one embodiment of that method, the preparation is a supernatant preparation obtained from a *Streptomyces* sp. ActE culture. In another embodiment of that method, the preparation is obtained from *Streptomyces* sp. ActE grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass, and chitin. In another embodiment of that method, the lignocellulosic material is selected from the group consisting of materials that comprise at least 75% cellulose, cellulose/hemicelluloses, xylose, biomass and chitin.

In one embodiment, the invention is a purified preparation comprising the *Streptomyces* sp. ActE secretome. In one embodiment, the preparation is a supernatant preparation obtained from a *Streptomyces* sp. ActE culture. In another embodiment of the preparation, *Streptomyces* sp. ActE is grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass, and chitin.

In one embodiment, the invention is a composition useful for digesting lignocellulosic material comprising one gene or expression product thereof selected from the group consisting of SActE_0237 (GH6) (SEQ ID NOS:1 and 17), SActE_0236 (GH48) (SEQ ID NOS:2 and 18), SActE_3159 (CBM33) (SEQ ID NOS:3 and 19), SActE_0482 (GH5) (SEQ ID NOS:4 and 20), SActE_0265 (GH10) (SEQ ID NOS:5 and 21), and SActE_2347 (GH5) (SEQ ID NOS:6 and 22) genes or expression products thereof. In one embodiment, the composition additionally comprises at least one member selected from the group consisting of SActE_0357 (CE4) (SEQ ID NOS:7 and 23), SActE_0358 (GH11) (SEQ ID NOS:8 and 24), SActE_1310 (PL3) (SEQ ID NOS:9 and 25), SActE_3717 (GH9) (SEQ ID NOS:10 and 26), SActE_4638 (SEQ ID NOS:11 and 27), SActE_4738 (GH16) (SEQ ID NOS:12 and 28), SActE_4755 (GH64) (SEQ ID NOS:13 and 29), SActE_5457 (GH46) (SEQ ID NOS:14 and 30), SActE_5647 (GH87) (SEQ ID NOS:15 and 31), and SActE_5978 (PL1) (SEQ ID NOS:16 and 32) genes or expression products derived thereof.

In one embodiment, the invention is a composition useful for cellulose degradation comprising SActE_0236 (GH48) (SEQ ID NOS:2 and 18), SActE_3159 (CBM33) (SEQ ID NOS:3 and 19), SActE_0482 (GH5) (SEQ ID NOS:4 and 20) and SActE_0237 (GH6) (SEQ ID NOS:1 and 17) genes or expression product thereof. In one embodiment, the composition additionally comprises at least one member selected from the group consisting of SActE_0357 (CE4) (SEQ ID

12

NOS:7 and 23), SActE_0358 (GH11) (SEQ ID NOS:8 and 24), SActE_1310 (PL3) (SEQ ID NOS:9 and 25), SActE_3717 (GH9) (SEQ ID NOS:10 and 26), SActE_4638 (SEQ ID NOS:11 and 27), SActE_4738 (GH16) (SEQ ID NOS:12 and 28), SActE_4755 (GH64) (SEQ ID NOS:13 and 29), SActE_5457 (GH46) (SEQ ID NOS:14 and 30), SActE_5647 (GH87) (SEQ ID NOS:15 and 31), and SActE_5978 (PL1) (SEQ ID NOS:16 and 32) genes or expression products derived thereof.

10 In one embodiment, the invention is a method for digesting a lignocellulosic material, comprising exposing the material to a sufficient amount of a composition of any combinations of genes or expression products derived thereof as disclosed above, wherein the exposed material is at least partially digested.

15 In one embodiment, the invention is a purified preparation of *Streptomyces* sp. ActE, wherein the *Streptomyces* sp. ActE has been grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass and chitin.

20 In one embodiment, the invention is a purified preparation of *Streptomyces* sp. ActE, wherein the *Streptomyces* sp. ActE has been grown on a substrate wherein at least 40%, preferably 85%, of *Streptomyces* sp. ActE's carbon in the substrate is derived from pretreated lignocellulosic material. In one embodiment of the preparation, the pretreated material has been exposed to pretreatment selected from the group consisting of acid hydrolysis, steam explosion, ammonia fiber expansion (AFEX), organosolve, sulfite pretreatment to overcome recalcitrance of lignocellulose (SPORL), ionic liquids (IL), metal-catalyzed hydrogen peroxide treatment, alkaline wet oxidation and ozone pretreatment. In another embodiment of the preparation, the pretreated material is wood.

Specific Embodiments

25 Applicants have been interested in insects that utilize plant biomass and their associated microbial and fungal communities. *Sirex noctilio*, a wood boring wasp, is found in pine forests throughout Eurasia and North Africa and is spreading throughout North America and elsewhere (Bergeron et al., 2011). Although the destructive nature of the *Sirex* infestation is generally considered to arise from a symbiotic relationship between *S. noctilio* and *Amylostereum areolatum*, a white rot basidiomycete (Kukor and Martin, 1983; Klepzig et al., 2009; Bergeron et al., 2011), the role of cellulolytic microbes has not been previously considered in the context of the infestation or symbiosis. *Streptomyces* sp. SirexAA-E [*Streptomyces* sp. ActE, also referred to herein as "ActE" (Adams et al., ISME J. 5:1321-1231, 2011)], was isolated from the ovipositor mycangium of *S. noctilio* (Adams et al., 2011). Applicants hypothesized that ActE is inoculated into insect feeding tunnels upon infestation along with the symbiotic fungus. Thus, Applicants were interested to learn how ActE might contribute to the *Sirex* community.

30 The present invention will be more fully understood upon consideration of the following non-limiting Examples. All papers and patents disclosed herein are hereby incorporated by reference as if set forth in their entirety.

35 As used herein, the term "ActE" refers to *Streptomyces* sp. SirexAA-E, as described in Adams et al., ISME J. 5:1321-1231, 2011. A representative sample of *Streptomyces* sp. ActE has been deposited according to the Budapest Treaty for the purpose of enabling the present invention. The

13

repository selected for receiving the deposit is the American Type Culture Collection (ATCC) having an address at 10801 University Boulevard, Manassas, Va. USA, Zip Code 20110. The ATCC repository has assigned the patent deposit designation PTA-12245 to the *Streptomyces* sp. ActE strain.

As used herein, the term "secretome" refers to the plurality of secreted enzymes. For example, ActE secretome refers to the secreted enzymes from *Streptomyces* sp. SirexAA-E.

As used herein, the term "lignocellulosic material" refers to any material that is composed of cellulose, hemicellulose, and lignin, wherein the carbohydrate polymers (cellulose and hemicelluloses) are tightly bound to the lignin.

As used herein, the term "biomass" refers to a renewable energy source, and comprises biological material from living or recently living organisms. As an energy source, biomass can either be used directly, or converted into other energy products such as biofuel. Biomass includes plant or animal matter that can be converted into fibers or other industrial chemicals, including biofuels. Industrial biomass can be grown from numerous types of plants, including miscanthus, switchgrass, hemp, corn, poplar, willow, sorghum, sugarcane, bamboo, and a variety of tree species, ranging from *eucalyptus* to oil palm (palm oil). Thus, biomass can include wood biomass and non-wood biomass.

The present invention has multiple embodiments. All embodiments are related to Applicants' discovery of improved lignocellulosic digestion and utilization using proteins and genes obtained from the *Streptomyces* sp. ActE secretome.

ActE Isolates and Secretomes

Streptomyces sp. SirexAA-E may be isolated from ovipositor mycangia of *S. noctilio*. In Adams, et al, *S. noctilio* were collected from a population in Pennsylvania, USA. Infested trees were cut and transported to USDA Pest Survey, Detection, and Exclusion Lab in Syracuse, N.Y., USA (Zylstra et al. (2010) Agric. Forest. Entomol. in press). Four adult females and six larvae from the Pennsylvania population were sampled, and cultures of bacteria derived from these insect samples were screened for cellulose degradation.

Prior to sampling for bacteria, all insects were typically surface sterilized in 95% ethanol for 1 minute and then rinsed twice in sterile phosphate-buffered saline (1×PBS). Larval guts and adult ovipositors and mycangia were removed surgically. These segments and the body were ground separately in 1 ml 1×PBS using a sterilized mortar and pestle. 50 µl of three 100-fold dilutions of each insect part were plated onto yeast and malt extract agar (Becton, Dickinson and Company, Sparks, Md., USA), acidified yeast malt extract agar (for gut dissections only), 10% tryptic soy agar (Becton, Dickinson and Company, Sparks, Md., USA), and agar supplemented with chitin (MP Biomedicals, Solon, Ohio). Petri dishes were stored at room temperature in darkness for at least three days until visible colonies formed, except for Petri dishes with chitin agar that were stored for at least one month.

All isolates were typically screened for production of cellulolytic enzymes on carboxymethyl cellulose (CMC) (Teather R M, Wood P J (1982); incorporated herein by reference as if set forth in its entirety). Isolates that tested positive on CMC were then studied further. Assays on CMC, AFEX-treated corn stover at three pH levels, and microcrystalline cellulose were typically performed to assess growth and degradation ability of each insect-derived bacterial isolate. Isolates capable of degrading CMC were further analyzed genetically to identify isolates with high

14

Carbohydrate Active Enzyme (CAZy) content relative to one another and relative to known organisms. *Streptomyces* sp. ActE was selected based on its CMC degradation and CAZy gene profile.

In one embodiment, secretomes from ActE would be used alone in a first reaction to convert biomass into a hydrolyzed solution of sugars that would be used in a second reaction with a fermentation organism to convert the sugars into usable biofuels. The first and second reaction could occur simultaneously.

In a second embodiment, secretomes from ActE would be combined with secretomes from other organisms, or with enzymes or enzyme compositions, such as Spezyme CP, to increase the activity of both preparations by synergy of the enzymes contained in each preparation.

Preferably, the ActE secretomes would be prepared as supernatants from ActE cultures.

In one embodiment, the supernatant is prepared by centrifugation of the ActE culture for 10 min at 3,000×g, which will pellet the remaining insoluble polysaccharide and adhered ActE cells. The supernatant fraction is filter-sterilized, preferably using a 0.22 µm filter, in order to remove any remaining cells. The supernatant is concentrated, preferably using a 3 kDa cut-off ultrafiltration membrane. The concentration of total protein is determined by Bradford assay (Bradford, 1976). In one preferred embodiment, the proteomic composition of the ActE secretome is that described in FIG. 3 or FIG. 18.

The secretomes obtained from growth on specific lignocellulosic materials, such as cellulose, xylan, cellulosic hemi-cellulosic biomass, and chitin, will have distinct compositions of individual enzymes and also distinct reactivity with different polysaccharides. The cellulosic hemi-cellulosic biomass may be non-wood biomass or wood biomass. For example, the secretome prepared from ActE grown on cellulose has unique enzymes and enhanced reactivity with cellulose and mannan. Also, the secretome prepared from ActE grown on xylan possesses high xylan degradation activity, whereas the secretome from ActE grown on chitin possesses uniquely high chitin degradation activity. Example A discloses the specific secretomes.

When ActE is grown on switchgrass, AFEX-pretreated switchgrass or ionic liquid pretreated switchgrass, the secretome has a protein composition that partially matches that obtained from growth on either cellulose or xylan. However, switchgrass, AFEX-pretreated switchgrass or ionic liquid pretreated switchgrass elicit the appearance of new proteins in the secretome that enhance the degradative ability of the secretome for the plant biomass materials. Applicants envision that the present invention would also apply to other pretreatment methods comprising acid hydrolysis, steam explosion, organosolve, sulfite pretreatment to overcome recalcitrance of lignocellulose (SPORL), metal-catalyzed hydrogen peroxide treatment, alkaline wet oxidation and ozone pretreatment.

The inventors' preliminary data shows synergistic filter paper degrading activity between the ActE secretome and other cellulases from a different organism. Also, addition of a beta-glucosidase to the secretome helps to break down the oligosaccharides (e.g., cellobiose, cellooligosaccharides) released from filter paper into simpler sugars.

Preferably, the secretome would be prepared as a concentrated solution by ultrafiltration. The concentrated material would be mixed with the substrate at weight percentages varying from 0.1% to 20% w/w, with the remainder of the solution containing a buffer substance that controls pH. Trace metals would be added to the reaction. The material

15

would be incubated at the appropriate temperature to allow the reaction to occur, with mixing of the reaction materials. The sample might be equilibrated with air or O₂ gas throughout the reaction time period.

The secretome obtained from growth of ActE on cellulose provides all necessary enzymes for most efficient breakdown of cellulose to cellobiose and mannan to mannose. Weak reaction is observed for breakdown of xylan to xylose and a mixture of mannobiase and mannose.

The secretome obtained from growth of ActE on xylan provides all necessary enzymes for most efficient breakdown of xylan to xylobiose and xylose. Weak reaction is observed for breakdown of cellulose to cellobiose and for breakdown of mannan to mannose.

The secretome obtained from growth of ActE on chitin provides all necessary enzymes for most efficient breakdown of chitin to N-acetylglucosamine. Weak reaction is observed for breakdown of xylan to xylose. Weak reaction is observed for breakdown of cellulose to cellobiose and for breakdown of mannan to mannose.

The secretome obtained from growth of ActE on switchgrass biomass provides all of the necessary enzymes for breakdown of cellulose, xylan, and mannan contained in switchgrass to the constituent monosaccharides and disaccharides. Growth of ActE on switchgrass exposed to different chemical pretreatments changes the composition of enzymes present, which alters the rate of production and yield of the constituent monosaccharides and disaccharides.

The secretome obtained from growth of ActE on cellulose provides the necessary enzymes for breakdown of cellulose to cellobiose. ActE uses cellobiose as the growth substrate, so no enzymes are present to convert cellobiose to glucose.

In order to obtain glucose, a cellobiase or beta-glucosidase would be added. This is a standard practice in biofuels enzymology.

In order to convert cellobiose to glucose, a cellobiase or beta-glucosidase would be added. Addition of cellulases from other organisms can improve the rate of hydrolysis of cellulose, e.g., addition of CelLcc_CBM3a, an engineered enzyme from *C. thermocellum* covered in Fox and Elsen Patent Application No.: PCT/US2010/037094.

The secretome obtained from growth of ActE on cellulose provides all of the necessary enzymes for breakdown of cellulose to cellobiose in a soluble form. One skilled in the art might purify these proteins directly from the secretome without use of tags or recombinant approaches.

As previously noted, the dominance of cellobiose as a product of cellulose deconstruction by ActE might help to channel cellulolytic activity to only a subset of the diverse microbes found in the *Sirex* community. Exploiting this community interaction, along with establishing control of the highly regulated patterns of gene expression observed in ActE provides the basis for a new biotechnological route for lignocellulosic digestion. For example, use of ActE secretomes to produce cellobiose will restrict the use of cellulose as a fermentation substrate to only those organisms capable of cellobiose uptake followed by intracellular conversion to glucose and subsequent glycolytic pathway intermediates. This might be achieved by coupling ActE enzymes with a yeast fermentation strain engineered to contain a specific cellobiose transporter and an intracellular cellobiose phosphorylase, leading to the intracellular production of glucose and glucose-1-phosphate.

ActE secretomes can be mixed with cellulosic biomass to convert it to cellobiose and xylose, as in the biofuels industry. For example, one might (1) mix the secretome with paper waste to convert it to a mixture of readily fermentable

16

oligo-, di-, and monosaccharides; (2) mix with animal feeds to increase the digestibility of the biomass to promote animal growth; (3) mix with cotton-based textiles for smoothing or other refinements; (4) mix with waste from the shrimp industry to process solid chitin to soluble constituents; (5) mix with mannan-enriched materials to convert them to mannose and mannobiase. One would also find the secretome useful for commercial food processing or treatment of cellulosic bezoar found in the human stomach.

One embodiment of the present invention is an isolation or purified preparation of *Streptomyces* sp. ActE.

An isolation of ActE was originally reported by Adams et al., (2011) ISME j doi:10.1038/ismej.2011.14, where it was stated that “*Sirex noctilio* were collected from infested scots pine, *Pinus sylvestris* L, in Onondaga County, NY, USA in 2008”, and “Microbial isolates were obtained from four adult females and six larvae collected in 2008, and were screened for cellulase activity.” These isolates were screened for cellulolytic ability by growing them on CMC, AFEX-treated corn stover, and microcrystalline cellulose.

Applicants envision that one would wish to prepare ActE isolates on specific nutrient sources for optimization for particular digestion profiles. Therefore, one may wish to prepare ActE on substrates wherein at least 40%, preferably 85% of *Streptomyces* sp. ActE's carbon source in the substrate is derived from a material selected from the group consisting of cellulose, cellulose/hemicelluloses mixture, hemicelluloses, xylan, non-wood biomass, wood biomass, and chitin.

In a preferred embodiment, ActE would be grown aerobically to maximize the secretion of enzymes that include both oxidative and hydrolytic enzymes capable of the rapid deconstruction of biomass. Since ActE cannot utilize mannan for growth, but efficiently liberates mannose from biomass, mannose would become available for growth of the inoculum of a fermentation organism in co-culture. The likely fact that ActE produces at least one antibiotic that would help maintain culture sterility is another possible advantage to establishment of an effective co-culture.

The high capacity for mannan hydrolysis coupled with the inability of ActE to use mannan as a growth substrate offers unique potential opportunity for expansion of deconstruction enzymology to the use of woody substrates. The deconstruction of woody substrates is considered to be more challenging for biofuels production despite the fact that woody substrates are also considerably more highly enriched in mannan than grass substrates. This unique potential opportunity will be enhanced by ongoing plant engineering research efforts to redefine the proportion of xylan and mannan in plant hemicellulose. The availability of plant material enriched in mannan will be coupled to vigorous conversion to mannose by ActE secretomes, providing a targeted, simply fermented C6 sugar for exclusive use by the fermentation organism.

When sufficient titer of enzymes and fermentation organism have been achieved, facilitated by the vigorous, obligate aerobic growth of ActE and corresponding deconstruction of biomass, the fermentation could be initiated by removal of the air source from the culture vessel. In the anoxic conditions, ActE would cease to grow, and perhaps even lyse to become a protein source for the fermentation organism, which will continue to grow on biomass that is simultaneously being deconstructed by the loading of highly active hydrolytic enzymes originally produced by ActE during the aerobic growth phase.

Applicants envision adding an ActE isolate directly to biomass slurry. More preferably an ActE isolate would be

added to the pretreated biomasses in the enzyme hydrolysis step, because ActE is able to grow at wide range of pH. ActE can be genetically modified so that the proteolysis proof secretome will be achieved. Growth on switchgrass elicits the appearance of new proteins in the secretome that enhance the degradative ability of the secretome for the plant biomass materials. Applicants envision that the present invention would apply to the biomasses pretreated by many pretreatment methods comprising AFEX, ionic liquid pretreated, acid hydrolysis, steam explosion, organosolve, sulfite pretreatment to overcome recalcitrance of lignocellulose (SPORL), metal-catalyzed hydrogen peroxide, alkaline wet oxidation and ozone pretreatment.

In one preferred embodiment of the present invention, at least one key enzyme in the secretome can be overexpressed by genetic modification of the ActE strain. Table 1 provides various combinations of genes that can be overexpressed. For example, one may wish to overexpress core cellulose deconstructing enzymes, SACTE_0237 (SEQ ID NOs:1 and 17), SACTE_0482 (SEQ ID NOs:4 and 20), SACTE_0236 (SEQ ID NOs:2 and 18), or SACTE_3159 (SEQ ID NOs:3 and 19) together with one or more of SACTE_2347 (SEQ ID NOs:6 and 22), and SACTE_0265 (SEQ ID NOs:5 and 21). One may wish to overexpress core xylan deconstructing enzymes, SACTE_0265 (SEQ ID NOs:5 and 21), SACTE_0358 (SEQ ID NOs:8 and 24), SACTE_0357 (SEQ ID NOs:7 and 23), SACTE_5978 (SEQ ID NOs:16 and 32), and SACTE_5230 (SEQ ID NOs:33 and 48). One may wish to overexpress core mannan deconstruction enzymes, such as SACTE_2347 (SEQ ID NOs:6 and 22). Additionally, SACTE_4755 (SEQ ID NOs:13 and 29) and SACTE_4738 (SEQ ID NOs:12 and 28) may be overexpressed for beta-1,3-glucan deconstruction. One may also overexpress all or some of the aforementioned genes for efficient biomass deconstruction.

In another embodiment of the present invention, at least one key enzyme in the secretome can be overexpressed and secreted by genetic modification of a different microbial host such as *Streptomyces lividans*, which is used for industrial secretion of proteins (Anne and Van Mellaert. (1993)), or *T. reesei*, which is used for secretion of enzymes in the biofuels industry (Saloheimo and Pakula, Microbiology, Epub date 2011 Nov. 5).

In another embodiment of the present invention, at least one key enzyme in the secretome can be overexpressed by genetic modification of a different microbial host such as *S. cerevisiae* or *E. coli* such that the expressed protein will be retained inside of the host cell. The host cells would then be harvested and used as a delivery agent without need for purification of the entrained enzyme, as described in Wood et al., 1997. This version of the invention may be useful in the enzymatic pretreatment of agricultural crop materials for consumption by ruminant animals.

Combinations of ActE Genes and Expression Products

Selected minimal genes in each subset were chosen based on the combination of genomic, transcriptomic and secretomic results (See Examples and Table 1). For example, in the cellulose minimal gene set, expression of these genes was relatively enriched in cellulose grown cells, compared to glucose grown cells, also corresponding proteins were highly secreted in response to the cellulose in culture medium. Selected minimal genes were annotated to have cellulose utilization function. A larger set of genes for cellulose utilization were selected based on the enrichment of gene expression in cellulose-grown cells relative to glucose-grown cells, and a functional annotation supports cellulose utilization of these genes. Additionally, neighbor-

hood genes to these selected genes on genome were included as genes regulated under same promoter. Similarly, both minimal and a large set of genes for xylan, chitin, and biomasses were elected.

In one embodiment, the present invention is a composition useful for digesting lignocellulosic material comprising genes or expression products thereof selected from the group consisting of: (a) SActE_0237 (SEQ ID NOs:1 and 17), SActE_0236 (SEQ ID NOs:2 and 18), SActE_3159 (SEQ ID NOs:3 and 19), SActE_0482 (SEQ ID NOs:4 and 20), SActE_0265 (SEQ ID NOs:5 and 21), and SActE_2347 (SEQ ID NOs:6 and 22), and (b) SActE_0357 (CE4) (SEQ ID NOs:7 and 23), SActE_0358 (GH11) (SEQ ID NOs:8 and 24), SActE_1310 (PL3) (SEQ ID NOs:9 and 25), SActE_3717 (GH9) (SEQ ID NOs:10 and 26), SActE_4638 (SEQ ID NOs:11 and 27), SActE_4738 (GH16) (SEQ ID NOs:12 and 28), SActE_4755 (GH64) (SEQ ID NOs:13 and 29), SActE_5457 (GH46) (SEQ ID NOs:14 and 30), SActE_5647 (GH87) (SEQ ID NOs:15 and 31), and SActE_5978 (PL1) (SEQ ID NOs:16 and 32). In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In one embodiment, one would use at least one member of (a) to digest a preferred lignocellulosic material.

In another embodiment, one would use at least the first four members [SActE_0237 (SEQ ID NOs:1 and 17), SActE_0236 (SEQ ID NOs:2 and 18), SActE_3159 (SEQ ID NOs:3 and 19), and SActE_0482 (SEQ ID NOs:4 and 20)] of (a) to digest a preferred lignocellulosic material.

In another embodiment, one would use at least one member of (a) and at least one member from (b), to digest a preferred lignocellulosic material.

In a preferred embodiment, one would use all the members of (a) and (b), to digest a preferred lignocellulosic material.

In other embodiments, the combination of genes or expression products thereof in the present invention is dependent on the specific lignocellulosic material to be digested. In one embodiment, a composition optimized for cellulose utilization may include any combinations of ActE genes and expression products disclosed above with at least one member selected from SActE_0265 (GH10) (SEQ ID NOs:5 and 21) and SActE_2347 (GH5) (SEQ ID NOs:6 and 22) genes or expression products thereof.

In another embodiment, a composition optimized for xylan utilization may include any combinations of ActE genes and expression products disclosed above with at least one member selected from SActE_0265 (GH10) (SEQ ID NOs:5 and 21), SActE_0358 (GH11) (SEQ ID NOs:8 and 24), SActE_0357 (CE4) (SEQ ID NOs:7 and 23), SActE_5978 (PL1) (SEQ ID NOs:16 and 32) and SActE_5230 (xylose isomerase) (SEQ ID NOs:33 and 48) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In another embodiment, a composition optimized for chitin utilization may include any combinations of ActE genes and expression products disclosed above with at least one member selected from SActE_4571 (GH18) (SEQ ID NOs:34 and 49), SActE_2313 (CBM33) (SEQ ID NOs:35 and 50), SActE_4246 (GH18) (SEQ ID NOs:36 and 51), SActE_3064 (GH19) (SEQ ID NOs:37 and 52), and SActE_5764 (GH18) (SEQ ID NOs:38 and 53) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In another embodiment, a composition optimized for biomass utilization may include any combinations of ActE genes and expression products disclosed above with SActE_5457 (GH46) (SEQ ID NOs:14 and 30) genes or expression products thereof.

In another embodiment, a composition optimized for mannan utilization may include any combinations of ActE genes and expression products disclosed above with SActE_2347 (GH5) (SEQ ID NO:6 and 22) genes or expression products thereof.

In another embodiment, a composition optimized for beta-1,3-glucan utilization may include any combinations of ActE genes and expression products disclosed above with at least one member selected from SActE_4755 (GH64) (SEQ ID NOs:13 and 29) and SActE_4738 (GH16) (SEQ ID NOs:12 and 28) genes or expression products thereof.

In another embodiment, a composition optimized for pectin release utilization may include any combinations of ActE genes and expression products disclosed above with SActE_1310 (PL3) (SEQ ID NOs:9 and 25) gene or expression products derived thereof.

In another embodiment, a composition optimized for alginate release utilization may include any combinations of ActE genes and expression products disclosed above with SActE_4638 (SEQ ID NOs:11 and 27) gene or expression products derived thereof.

In another embodiment, a composition optimized for galactose release utilization may include any combinations of ActE genes and expression products disclosed above with SActE_5647 (GH87) (SEQ ID NOs:15 and 31) gene or expression products derived thereof.

In another embodiment, the present invention is summarized as a composition useful for xylan degradation comprising SActE_0265 (GH10) (SEQ ID NOs:5 and 21) and SActE_0358 (GH11) (SEQ ID NOs:8 and 24) genes or expression products thereof.

In another embodiment, the present invention is summarized as a composition useful for xylan degradation comprising SActE_0265 (GH10) (SEQ ID NOs:5 and 21), SActE_0358 (GH11) (SEQ ID NOs:8 and 24), SActE_0265 (GH10) (SEQ ID NOs:5 and 21), SActE_0358 (GH11) (SEQ ID NOs:8 and 24), SActE_0357 (CE4) (SEQ ID NOs:7 and 23), SActE_5978 (PL1) (SEQ ID NOs:16 and 32), and SActE_5230 (xylose isomerase) (SEQ ID NOs:33 and 48) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In another embodiment, the present invention is summarized as a composition useful for biomass degradation comprising SActE_0237 (GH6) (SEQ ID NOs:1 and 17), SActE_0482 (GH5) (SEQ ID NOs:4 and 20), SActE_3159 (CBM33) (SEQ ID NOs:3 and 19), SActE_0236 (GH48) (SEQ ID NOs:2 and 18), SActE_3717 (GH9) (SEQ ID NOs:10 and 26), SActE_0265 (GH10) (SEQ ID NOs:5 and 21), SActE_0358 (GH11) (SEQ ID NOs:8 and 24), SActE_2347 (GH5) (SEQ ID NOs:6 and 22) and SActE_1310 (PL3) (SEQ ID NOs:9 and 25) genes or expression products thereof. In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In one embodiment, the present invention is a composition useful for digesting lignocellulosic material comprising genes or expression products thereof selected from the group consisting of: (a) SActE_0237 (SEQ ID NOs:1 and 17), SActE_0236 (SEQ ID NOs:2 and 18), SActE_3159 (SEQ ID NOs:3 and 19), SActE_0482 (SEQ ID NOs:4 and 20), SActE_0265 (SEQ ID NOs:5 and 21), and SActE_2347

(SEQ ID NOs:6 and 22) (for cellulose); (b) SActE_0265 (SEQ ID NOs:5 and 21), SActE_0357 (SEQ ID NOs:7 and 23), SActE_0358 (SEQ ID NOs:8 and 24), SActE_5230 (SEQ ID NOs:33 and 48) and SActE_5978 (SEQ ID NOs:16 and 32) (for xylan); (c) SActE_2313 (SEQ ID NOs:35 and 50), SActE_3064 (SEQ ID NOs:37 and 52), SActE_4246 (SEQ ID NOs:36 and 51), SActE_4571 (SEQ ID NOs:34 and 49) and SActE_5764 (SEQ ID NOs:38 and 53) (for chitin); (d) SActE_2347 (SEQ ID NOs:6 and 22) (for mannan); and (e) SActE_0236 (SEQ ID NOs:2 and 18), SActE_0237 (SEQ ID NOs:1 and 17), SActE_0265 (SEQ ID NOs:5 and 21), SActE_0358 (SEQ ID NOs:8 and 24), SActE_1310 (SEQ ID NOs:9 and 25), SActE_2347 (SEQ ID NOs:6 and 22) and SActE_3159 (SEQ ID NOs:3 and 19) (for biomass). In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In one embodiment, one would use at least two members of (a), (b), (c), (d) or (e) to digest a preferred lignocellulosic material.

In another embodiment, one would use at least three members.

In a preferred embodiment, one would use all members of (a), (b), (c), (d) or (e).

In another embodiment, one would add gene expression products from the list in Table 1 to a substrate to be digested. For example, for preferred cellulose digestion, one would select at least two members of (a), as described above, and at least one member of the “additional useful genes” in Table 1.

In the case of cellulose degradation, the inventors believe SACTE_3159 (SEQ ID NOs:3 and 19), SACTE_0237 (SEQ ID NOs:1 and 17), SACTE_0482 (SEQ ID NOs:4 and 20), and SACTE_0236 (SEQ ID NOs:2 and 18) act cooperatively to create nicks and hydrolyze cellobiose units from crystalline cellulose.

ActE key genes can be transferred into known cellulolytic organisms in order to enhance the cellulolytic ability of these organisms. A cellulolytic fungus, *T. reesei*, has been studied for industrial applications, and can be genetically modified. Applicants' data support synergism of cellulolytic ability of enzymes from different species. A chromosomal gene transfer can be performed into *T. reesei* by protoplast transformation with a high copy plasmid carrying one or more of the ActE cellulolytic key genes.

A chromosomal or a non-chromosomal gene transfer can be made into a yeast species such as *Saccharomyces cerevisiae*. For non-chromosomal gene transfer, a high copy plasmid carrying a cassette of five minimal genes (SACTE_0236 (SEQ ID NOs:2 and 18), SACTE_0237 (SEQ ID NOs:1 and 17), SACTE_0482 (SEQ ID NOs:4 and 20), SACTE_3717 (SEQ ID NOs:10 and 26) and SACTE_3159 (SEQ ID NOs:3 and 19)) would be used to confer cellulolytic and mannanolytic capability to the yeast strain. Similar approaches could be used to confer xylanolytic and chitinolytic capability using combinations of the genes described herein.

One might wish to recombinantly express the disclosed enzymes in *E. coli* in order to achieve high yield of each enzyme. As is shown in the synergistic result in Example 18, cellulose degradation can be improved by combination of ActE enzymes to enzymes from other organisms.

FIG. 18 shows Spectra count of proteins identified on each substrate, where top 95% most abundant proteins were highlighted green, light purple, purple, blue, orange, pink,

light blue and yellow on glucose, cellobiose, cellulose, xylan, switchgrass, AFEX-SG, IL-SG and chitin, respectively.

Applicants envision that one would use a composition comprising at least one member of the abundant proteins, e.g., those highlighted proteins in FIG. 18, for digesting the corresponding lignocellulosic materials. For example, to digest a cellulose material, one would choose at least one gene or expression products thereof selected from the group consisting of SACTE_0237 (SEQ ID NOS:1 and 17), SACTE_0236 (SEQ ID NOS:2 and 18), SACTE_2347 (SEQ ID NOS:6 and 22), SACTE_3159 (SEQ ID NOS:3 and 19), SACTE_0482 (SEQ ID NOS:4 and 20), SACTE_0265 (SEQ ID NOS:5 and 21), SACTE_0357 (SEQ ID NOS:7 and 23), SACTE_4439 (SEQ ID NOS:39 and 54), SACTE_0562 (SEQ ID NOS:40 and 55), SACTE_0358 (SEQ ID NOS:8 and 24), SACTE_4343 (SEQ ID NOS:41 and 56), SACTE_1546 (SEQ ID NOS:42 and 57), SACTE_1310 (SEQ ID NOS:9 and 25), SACTE_4638 (SEQ ID NOS:11 and 27), SACTE_5668 (SEQ ID NOS:45 and 60), SACTE_3717 (SEQ ID NOS:10 and 26), SACTE_3590 (SEQ ID NOS:43 and 58), SACTE_2172 (SEQ ID NOS:44 and 59), SACTE_4571 (SEQ ID NOS:34 and 49), SACTE_5978 (SEQ ID NOS:16 and 32), SACTE_6428 (SEQ ID NOS:46 and 61), SACTE_2313 (SEQ ID NOS:35 and 50), and SACTE_0366 (SEQ ID NOS:47 and 62). In a preferred embodiment, the composition comprises at least three or four of the genes or expression products.

In one preferred embodiment, one would use all the highlighted proteins for digesting the corresponding lignocellulosic materials.

In another embodiment, one would add gene expression products from the list in Table 1 to a substrate to be digested. For example, for preferred cellulose digestion, one would select at least one member of the abundant proteins, as described above, and at least one member of the “additional useful genes” in Table 1.

TABLE 1

ActE genes or expression products useful for lignocellulosic degradation.		
Gene or Expression Product Combinations	Preferred subsets	Additional Useful Genes
SACTE_0236, SACTE_0237, SACTE_3159, SACTE_0482 and SACTE_3717	Cellulose degradation	SACTE_0229, SACTE_0230, SACTE_0231, SACTE_0232, SACTE_0233, SACTE_0234, SACTE_0235, SACTE_0480, SACTE_0481, SACTE_0483, SACTE_0562, SACTE_0563, SACTE_0733, SACTE_0734, SACTE_2286, SACTE_2287, SACTE_2288, SACTE_2289, SACTE_3158, SACTE_4737, and SACTE_6428
SACTE_0265, SACTE_0357, SACTE_0358, SACTE_5230 and SACTE_5978	Xylan degradation	SACTE_0364, SACTE_0365, SACTE_0366, SACTE_0368, SACTE_0369, SACTE_0370, SACTE_0527, SACTE_0528, SACTE_5227, SACTE_5228, SACTE_5229, SACTE_5858, and SACTE_5859
SACTE_2313, SACTE_3064, SACTE_4246, SACTE_4571 and SACTE_5764 SACTE_2347	Chitin degradation	SACTE_0080, SACTE_0081, SACTE_0844, SACTE_0846, SACTE_0860, SACTE_3063, SACTE_4858, SACTE_6493 and SACTE_6494
	Mannan degradation	

TABLE 1-continued

ActE genes or expression products useful for lignocellulosic degradation.		
Gene or Expression Product Combinations	Preferred subsets	Additional Useful Genes
SACTE_1310	Pectin degradation	
SACTE_4638	Alginate release	
10 SACTE_5647	Galactose release	SACTE_5648
SACTE_4738 and SACTE_4755	Beta-1,3-glucan degradation	SACTE_4737, SACTE_4739 and SACTE_4756
15 SACTE_0236, SACTE_0237, SACTE_0265, SACTE_0358, SACTE_0482, SACTE_1310, SACTE_2347, SACTE_3159 and SACTE_3717	Cellulose and hemicelluloses degradation	SACTE_3065, SACTE_4730, SACTE_4755, and SACTE_5166
20		

In one embodiment, the present invention is a method for digesting a lignocellulosic material, comprising exposing the material to a sufficient amount of a composition of enzymes, wherein the exposed material is at least partially digested. The enzymes may be ActE secretomes, and ActE secretomes may be prepared and isolated using the methods described above.

30 In another embodiment, the composition of enzymes for a method for digesting a lignocellulosic material may include ActE secretomes in a combination with secretomes from other organisms, or with enzymes or enzyme compositions, such as Spezyme CP, to increase the activity of both preparations by synergy of the enzymes contained in each preparation.

35 In another embodiment, the composition of enzymes for a method for digesting a lignocellulosic material may be any combinations of ActE genes and expression products as described above.

EXAMPLES

Materials and Methods

45 Genome Analysis.
The complete genome sequence of *Streptomyces* sp. Sir-exAA-E (ActE, taxonomy ID 862751) was determined by the Joint Genome Institute, project ID 4086644. Gene annotation models were predicted using Prodigal (Hyatt, et al., 2010), examined using Artemis (Rutherford, et al., 2000), and are available at NCBI with the following accession numbers, GenBank: CP002993.1; RefSeq: NC_015953.1. Carbohydrate-active enzymes were annotated by comparison of all translated open-reading frames to the CAZy database (Cantarel, et al., 2009). We collected CAZy annotated genes from the CAZy database. We then used BLASTP to compare all ActE protein-coding sequences to the CAZy database and to the pfam database. These two annotations were then crosschecked, and proteins annotated by both databases were identified as our final CAZy annotation. Secreted proteins were identified by SignalP, TatP, and SecretomeP analyses. BLAST was used to identify sequence orthologs in other organisms. Secondary metabolite gene clusters were identified by AntiSmash analysis (Medema, et al., 2011). CebR boxes were identified by using BLAST comparison of the *S. griseus* CebR box sequence to the ActE

genome (Marushima, Ohnishi, et al., 2009). Networks of expression and functional categories were visualized using Cytoscape (Shannon, et al., 2003)

Biomass Substrates.

Switchgrass and AFEX-treated switchgrass were obtained from Great Lakes Bioenergy Research Center. Extensively washed ionic liquid-treated switchgrass was the generous gift of Dr. Masood Hadi (Joint BioEnergy Institute). Wood kraft pulp preparations were the generous gift of Dr. Xuejun Pan (University of Wisconsin Department of Biosystems Engineering).

Growth of Organisms.

ActE, *S. coelicolor*, *S. griseus* and *T. reesei* RUT-C30 were grown at pH 6.0 and ActE was also grown at pH 6.9 in M63 minimal medium, where 1 L contains: 10.72 g K₂HPO₄; 5.24 g KH₂PO₄; 2 g (NH₄)₂SO₄; 0.5 mL iron sulfate (1 mg/mL in 0.01 M HCl); 1 mL 1 M MgSO₄; 1 mL thiamine solution (1 mg/mL) supplemented with glucose, cellulose (either Whatman #1 filter paper or Sigmacell-20, Sigma/Aldrich, St. Louis, Mo. as indicated), xylan, chitin, switchgrass, AFEX-treated switchgrass (Balan et al., 2009), or ionic liquid-treated switchgrass as the sole carbon source (0.5% w/v). Cultures were incubated for 7 days at 30° C. with shaking. In this medium at pH 6.9, ActE has doubling times of 2.5 h for growth on xylan and switchgrass, 8 h for glucose and 13 h for cellulose as determined by time-dependent increases in total protein present in the culture medium.

RNA microarray. ActE was grown in minimal medium plus the indicated substrate for 7 days. The cell pellet was separated from the culture medium by centrifugation for 10 min at 3000×g. Microarray experiments were carried out as reported previously (Riederer, et al., 2011). The total RNA was extracted from the cell pellet and purified. The University of Wisconsin Gene Expression Center carried out the syntheses of cDNA and array hybridizations. Four-plex arrays were constructed by Nimblegen and hybridized with 10 µg of labeled cDNA. ArrayStar (v4.02, DNASTAR, Madison, Wis.) was used to quantify and visualize data. All analyses were based on three or more biological replicates per carbon source. Quantile normalization and robust multi-array averaging (RMA) were applied to the entire data set. Unless otherwise specified, expression levels are based on log 2 values and statistical analysis of the datasets were performed using the moderated t-test.

Preparation of Secretomes.

Supernatants obtained from different culture media were prepared by centrifugation of the culture medium for 10 min at 3000×g, which removed the remaining insoluble polysaccharide and adhered cells. The supernatant fraction was then passed through a 0.22-µm filter in order to remove any remaining cells. For enzymatic assays, the secretomes were concentrated using a 3-kDa cut off ultrafiltration membrane. The concentration of secretome protein was determined by Bradford assay, and the typical yield was ~150-300 mg of total secreted protein per liter of culture medium.

Extracellular Protein Profiles.

Extracellular proteins from culture secretomes were precipitated with trichloroacetic acid (TCA), resuspended in denaturing sample buffer (SDS and 2-mercaptoethanol), and separated by SDS-PAGE in 4-20% gels. Protein bands of interest were excised from the gel, digested with trypsin, desalting with C18 pipette tips (Millipore, Billerica, Mass.) and identified by MALDI-TOF (MDS SCIEX 4800 MALDI TOF/TOF, Applied Biosystems, Foster City, Calif.). Addi-

tional samples from the same culture secretomes were analyzed by LC-MS/MS to identify highly abundant proteins in the sample.

Ion Exchange Separation of the ActE Secretome.

The ActE cellulose secretome was diluted with cold deionized water until the ionic strength was less than 50 mM. The diluted sample was loaded onto an AKTApürifier™ chromatography station equipped with a 16/10 MonoQ FF ion exchange column. The column was washed with 100 mL of 10 mM phosphate, pH 6.0, to remove unbound proteins. The bound proteins were eluted in a linear, 200 mL gradient of NaCl from 0 to 0.8 M in the same buffer. Fractions from the gradient elution were collected and separated by SDS PAGE. The proportional contribution of individual proteins in each fraction was estimated from SDS PAGE. Individual protein bands from each fraction were cut from the gel and submitted for LC-MS/MS analysis to confirm their identities.

LC-MS/MS Analyses.

These experiments were performed at the University of Wisconsin Biotechnology Center. Samples were prepared by TCA precipitation of 100 ng of total secreted protein from 7-day old culture supernatants. Protein samples were digested with trypsin (sequencing grade trypsin, Promega, Madison, Wis.) and were desalting using C18 pipette tips (Millipore, Billerica, Mass.). High-energy collision dissociation (HCD) MS analyses employing a capillary LC-MS/MS were performed on an electrospray ionization FT/ion-trap mass spectrometer (LTQ Orbitrap XL, Thermo Fisher Scientific, San Jose, Calif.). The MS and MS/MS spectra were searched against the spectra obtained from the ActE proteome by using Scaffold (Scaffold_3_00_06, Proteome Software, Portland, Oreg.).

Enzyme Activity Measurements.

Reducing sugar assays were carried out by mixing secretome preparations with polysaccharide-containing substrates including cellulose (either Whatman #1 filter paper or Sigmacell-20 as indicated), xylan, chitin, mannan, switchgrass, AFEX pretreated switchgrass, or ionic-liquid pretreated switchgrass²⁴. After incubation in 0.1 M sodium phosphate, pH 6 at 40° C. for 20 h, the reducing sugar content was detected by dinitrosalicylic acid assay (Miller, 1959) and calibrated by using glucose, xylose, or mannose as standards. Purified polysaccharide preparations had negligible background response in the absence of added enzymes. Cellooligosaccharides were assayed by a coupled enzyme assay (K-GATE system, Megazyme, Bray Ireland). SPEZYME CP was obtained from Genencor with batch number #4901522860. The distributions of soluble sugar oligomers obtained from secretome reactions were determined using a Shimadzu Liquid Chromatograph HPLC system (Shimadzu Scientific Instruments, Columbia, Md.) equipped with a refractive index detector (RID-10A) and a Phenomenex Rezex RPM-monosaccharide column. The temperature was maintained at 85° C. and Milli-Q water was used as the mobile phase at 0.6 mL min⁻¹ flow rate. Glucose, cellobiose, cellotriose, cellotetraose, cellopentaose, and cellobhexaose (Sigma) were used as standards. The integrated areas of peaks were analyzed by EZ start 7.2 SP1 software (Shimadzu).

Fractions obtained from the ion exchange separation of the ActE cellulose secretome were combined as unary, binary, ternary, and quaternary assemblies where the total protein concentration was fixed and the individual fractions contributed all, halves, thirds, or quarters of the total protein. The most active fraction was assembled from a ternary combination of fractions containing the following enzymes:

fraction 1, SACTE_3159 (CBM33/CBM2 oxidative endo-cellulase, 95%) and SACTE_4738 (GH16 β -1,3 endoglucanase, 5%); fraction 2, SACTE_0237 (GH6 exocellulase, 60%), SACTE_0482 (GH5 endocellulase, 25%), SACTE_0237 (β -1,3 glucanase, 10%) and SACTE_3159 (oxidative endocellulase, <5%); and fraction 3, SACTE_0236 (GH48 exocellulase, 75%), SACTE_3717 (GH9 endocellulase, 20%) and SACTE_5457 (GH46 chitinase, 5%).

Cellobionic and gluconic acids were assayed by a coupled enzyme assay (K-GATE system, Megazyme, Bray Ireland), either with or without the addition of a large excess of β -glucosidase (Cat. No. 31571, Lucigen, Middleton, Wis.).

Two lots of Spezyme CP were obtained from Genencor (#4900901244, Jan. 27, 2010 and #4901522860, Sep. 2, 2011). The specific activity of these two preparations was indistinguishable.

HPLC Analysis.

The distributions of soluble sugar oligomers obtained from secretome reactions without and with the addition of excess β -glucosidase (Lucigen) were determined using a Shimadzu Liquid Chromatograph HPLC system (Shimadzu Scientific Instruments, Columbia, Md.) equipped with a refractive index detector (RID-10A) and a Phenomenex Rezex RPM-monosaccharide column. The temperature was maintained at 85° C. and milli-Q water was used as the mobile phase at 0.6 mL min⁻¹ flow rate. Glucose, cellobiose, cellotriose, cellotetraose, and cellopentaose (Sigma) were used as standards. The integrated areas of peaks were analyzed by EZ start 7.2 SP1 software (Shimadzu).

For the experiments shown in FIG. 21, the ActE secretome (1 μ g total protein); CellLcc_CBM3a (1 μ g); ActE secretome (0.5 μ g) and CellLcc_CBM3a (0.5 μ g); or Spezyme CP (1 μ g total protein) were used. The products of the enzyme reactions detected by HPLC were: ActE secretome, 95% cellobiose, 5% glucose; CellLcc_CBM3a reaction, 90% cellobiose, 10% glucose; ActE & CellLcc_CBM3a, 5% cellotetraose, 80% cellotriose, 15% cellobiose; Spezyme CP, 33% cellobiose, 67% glucose. All products could be converted to glucose in the presence of excess β -glucosidase.

CellLcc_CBM3a.

The nucleotide and amino acid sequence of CellLcc_CBM3a is shown in FIG. 22. CellLcc_CBM3a is an engineered exoglucanase composed of the catalytic core of *C. thermocellum* CelL (Cthe_0405, residues 32 to 429) fused to a *C. thermocellum*-derived linker sequence and the CBM3a domain from Cthe_3077, the CipA scaffoldin. This construct was created to better understand the performance of enzymes that are normally targeted to the clostridial cellulosome. The replacement of the dockerin domain in Cthe_0405 with the CBM3a domain abrogates the need for a cellulosomal attachment to obtain maximal catalytic activity from CellLcc_CBM3a on solid substrates. The indicated nucleotide sequence was sub-cloned into wheat germ cell-free translation (Makino et al., 2010) and *E. coli* expression vectors (Blommel et al., 2009) for protein production. CellLcc_CBM3a was purified by standard immobilized metal (Ni²⁺) chromatography. There was no difference in the specific activity of the protein prepared by these two methods.

Example 1: ActE Exhibits High Cellulolytic Activity Relative to Other Cellulolytic Organisms

Prokaryotes such as *Streptomyces* are often easier to grow than eukaryotes (i.e., fungi such as *T. reesei*), and aerobes

are often easier and more energetically efficient to grow than anaerobes. *Streptomyces* may also have an advantage of producing antibiotics that limit the ability of other organisms to contaminate the culture medium during growth (Galm et al., 2011; Susi et al., 2011). This may be of advantage during large-scale culture with non-sterile biomass materials such as will be encountered in the biofuels industry.

When compared to other cellulolytic organisms (FIG. 1 and FIG. 6), ActE grows well on pure cellulose substances including amorphous cellulose (cellulose treated with phosphoric acid so as to remove all crystalline structure), filter paper (containing a mixture of amorphous and crystalline cellulose) and Sigmacell (primarily in the crystalline state as determined by X-ray powder diffraction), as well as other polysaccharides such as beta-1,3-glucan (callose), xylan, and chitin. ActE also grows well on biomass samples such as corn stover, ammonia-fiber expansion pretreated corn stover, switchgrass, ammonia-fiber expansion pretreated switchgrass, ionic liquids pretreated switchgrass, bleached spruce wood kraft pulp, and unbleached lodgepole pine kraft pulp.

FIG. 1 compares the ability of ActE, *S. coelicolor* A3(2) (NCBI taxonomy ID 100226) and *S. griseus* (NCBI CP002993.1; RefSeq: NC_015953.1) to grow in minimal medium containing filter paper as the only carbon and energy source. These images demonstrate the considerably different capabilities of the three ostensibly cellulolytic organisms. Thus ActE completely destroys the filter paper and achieves high cell density, while the two other, reputedly highly cellulolytic strains are only capable of weak colony formation attached to the filter paper. This result establishes that ActE has uniquely high cellulolytic capacity relative to other *Streptomyces* strains reported to also have this capability (Forsberg et al., 2011). In fact, the images of FIG. 1 and FIG. 6 demonstrate ActE has cellulolytic capacity rivaling that of *T. reesei* strain Rut-C30, which is widely acknowledged to be the industrial benchmark for cellulolytic capacity (Merino and Cherry, Adv. Biochem. Eng. Biotechnol. 108:95-120, 2007).

Example 2: Pretreatments Useful for Generating Fermentable Sugars

In the biofuels arena, the desired cellulose fractions of plant biomass are protected by the crystalline packing of the individual cellulose strands, and by the surrounding coating of hemicellulose and lignin. In order to most efficiently access the cellulose, chemical pretreatments are required to "loosen up" the plant cell wall structure. In this context, "loosen up" may mean removal of the lignin fraction, partial hydrolysis of feruloyl and acetyl esters present in hemicellulose, and changes in the crystallinity of the cellulose. An optimal pretreatment retains all fractions of biomass lignin, hemicellulose and cellulose in physical states that can be subsequently used by microbes and enzymes as substrates.

Ammonia-fiber expansion is a pretreatment that uses a combination of ammonia gas, low pressure, and low temperature to effect the loosening process (Balan et al., 2009; Chundawat et al., 2011; International Patent Publication No.: WO 2010/125679). It is particularly effective with grasses, and retains all fractions of the biomass for subsequent valorization without introducing water or salts into the biomass. Ionic liquids pretreatment comprises mixing a charged chemical substance (i.e., the ionic liquid) in equal mass proportions with the biomass material. Interactions between the ionic liquid substance and the biomass cause the crystalline structure of cellulose to convert to an amorphous

state (Cheng et al., 2011; Li et al., 2011) but the biomass also becomes heavily contaminated with the ionic liquid during this pretreatment, requiring extensive washing with water, a valuable resource in many localities. Kraft pulping is a method for production of paper from wood that involves treatment of the biomass material with strong alkali, sodium sulfite and moderate temperature, resulting in destruction of the lignin and hemicellulose from the desired cellulose fraction; the final biomass material is also heavily contaminated with salts that also requires extensive washing with water to remove. Acid pretreatments retain the lignin and cellulose but destroy the hemicellulose fraction, and in doing so create toxic substances derived from the decomposition of hemicellulose. Because of the need to neutralize the acid, this pretreatment generates a large contamination of salt that also requires extensive washing with water. SPORL is an acidic pretreatment that uses sulfuric acid, elevated temperature, and sodium bisulfite to effect the pretreatment (Wang et al., 2009; Tian et al., 2011). In SPORL, the lignin and hemicellulose are destroyed and cellulose is recovered, but the cellulose is again heavily contaminated with salts and toxic substances derived from chemical decomposition of hemicellulose.

ActE secretomes are highly effective for degradation of lignocellulosic material pre-treated with AFEX. ActE secretomes are also effective for degrading lignocellulosic material pretreated with ionic liquids, Kraft pulping, acid or SPORL and for degrading untreated lignocellulosic material.

Example 3: ActE Genome has High Content of Genes Encoding Carbohydrate Active Enzymes (CAZy) Relative to Other Cellulolytic Organisms

Protein-coding sequences of the ActE genome (Hyatt et al., 2010) were analyzed by BLAST comparison (Altschul et al., 1990) to the Carbohydrate Active Enzyme (CAZy) database (Cantarel et al., 2009).

Table 2 compares the genomic characteristics of ActE with well-known soil-isolated *Streptomyces* that produce antibiotics and with two model cellulolytic bacteria, *Clostridium thermocellum* and *Cellvibrio japonicas* (Lynd, Weimer, et al., 2002; Deboy, et al., 2008; Riederer, et al., 2011). Putative biomass-degrading protein-coding sequences from ActE were identified by BLAST analysis of the finished genome to the Carbohydrate Active Enzyme (CAZy) database. Among the 6357 predicted protein-coding genes, 167 have one or more domains assigned to CAZy families, including 119 glycoside hydrolases (GHs), 29 carbohydrate esterases (CEs), 6 polysaccharide lyases (PLs) and 85 carbohydrate binding modules (CBMs). ActE contains 45 different types of GH families, 4 PL families, 7 CE families, and 21 CBM families. The number of total CAZy domains and diversity of CAZy families is comparable to other highly cellulolytic organisms.

TABLE 2

Comparison of genomic composition.

	ActE	<i>S. coelicolor</i>	<i>S. griseus</i>	<i>C. thermocellum</i>	<i>C. japonicus</i>
Genome size (nt)	7414440	8667507	8545929	3843301	4576573
Proteome size	6357	8153	7136	3173	3750
Total CAZy Proteins	167	221	132	103	183

TABLE 2-continued

Comparison of genomic composition.					
	ActE	<i>S. coelicolor</i>	<i>S. griseus</i>	<i>C. thermocellum</i>	<i>C. japonicus</i>
% CAZy Proteins ^a	2.6%	2.7%	1.8%	3.2%	4.9%
Total GH ^b	119	154	80	70	124
Total PL ^c	6	11	4	6	14
Total CE ^d	29	36	23	20	28
Total CBM ^e	85	98	68	121	134
antiSMASH clusters ^f	22	24	37	3	4
Genes in clusters	620	718	1139	89	111
% antiSMASH	9.8%	8.8%	16.0%	2.8%	3.0%

^aProteins classified as Carbohydrate Active Enzymes (CAZy).

^bGH, glycoside hydrolase.

^cPL, pectate lyase.

^dCE, carbohydrate esterase.

^eCBM, carbohydrate binding module.

^fPutative antibiotic producing gene cluster.

Nearly all publically available *Streptomyces* genomes encode a relatively high percentage of genes for putative cellulolytic enzymes. Interestingly, ActE and the antibiotic producing *Streptomyces*, *S. griseus* and *S. coelicolor*, shown in Table 2 have similar numbers and compositions of CAZy families, but substantially different genome sizes. However, these antibiotic-producing *Streptomyces* are not highly cellulolytic (FIG. 1). Relative to *S. griseus* and *S. coelicolor*, the ActE genome contains two unique CAZy families but does not possess 16 CAZy families present in these species. However, ActE contains more representatives in 13 CAZy families. Enrichment of certain CAZy families was observed in other highly cellulolytic organisms. For example, *C. thermocellum* contains 16 genes in the GH9 family alone. It is interesting to consider whether the reduction in total genome size and differences in CAZy composition between ActE and other closely related soil-dwelling *Streptomyces* might have arisen from evolutionary specialization of ActE, perhaps driven by association with the *Sirex*-fungal symbiosis.

ActE contained 12 CAZy families not found in the other model cellulolytic organisms shown in FIG. 3, including GHs, CBMs, and PLs. Seven other CAZy categories, primarily hemicellulases, were shared only with *T. reesei*. ActE had 23 GH, 10 CBM and 2 PL not found in *Thermobifida fusca*, another cellulolytic Actinomycetales, which had only 1 GH and 1 CBM not found in ActE. The genome sequence revealed *C. japonicus* (strain Ueda 107) is highly enriched in GH43 enzymes required for hemicellulose utilization, but is missing a key reducing end exocellulase (bacterial GH48) required for robust growth on cellulose [e.g., see page 5459 of (DeBoy et al., 2008)]; both of these enzyme families are present in highly cellulolytic ActE. Furthermore, ActE also contained 6 genes from the CBM33 family, recently shown to catalyze oxidative cleavage of chitin (Vaaje-Kolstad et al., 2010) and cellulose (Forsberg et al., 2011). Thus, ActE has genomic composition overlapping other cellulolytic organisms, but with notable expansion in the CAZy composition for both hydrolytic and oxidative enzymes and the presence of the complete set of enzymes required for efficient cellulose deconstruction.

Example 4: Genome-Wide Gene Expression Analysis of ActE CAZy Gene

Gene expression profiles were determined for ActE grown on purified polysaccharides and plant biomass by whole

genome microarrays (FIGS. 4 and 5, FIGS. 9 to 14). Genome-wide gene expression was analyzed as a functional annotation network composed of ActE genes (circles) connected to predicted functional groups (triangles; KEGG or CAZy). In FIG. 4, the network was annotated with genome-wide microarray expression data to indicate genes that were differentially expressed when ActE was grown on either AFEX-SG or glucose, and further annotated to indicate normalized expression levels observed during growth on AFEX-SG. While many aspects of metabolism are modestly changed in response to these different carbon sources, the CAZy and ABC transporter categories were substantially enriched in differentially expressed genes (FIG. 4, green circles). Furthermore, pentose sugar metabolism, sulfur metabolism, and some amino acid biosynthesis pathways (e.g., aromatic amino acids) were also highly induced during growth on AFEX-SG relative to other carbon sources (FIGS. 9-14). In contrast, ribosomal, secondary metabolite, and DNA repair genes showed little change in expression across the conditions examined. Within the CAZy functional group, there was a large induction of genes that contained both a GH domain and a CBM2 domain. Among the 11 genes in the ActE genome that contain a CBM2 domain, 6 were induced greater than 4-fold during growth on AFEX-SG. Furthermore, 9 of the 11 CBM2 containing proteins were identified in the secreted proteome (FIG. 3).

Example 5: ActE CAZy Gene Expression is Dependent on ActE Growth Substrate

Given the large number of differentially expressed CAZy genes identified in the network analysis, Applicants analyzed the expression of this group of genes in cultures grown on different carbon sources (FIG. 5, FIG. 15 and FIG. 16). As with other cellulolytic organisms, there was strong correlation between the content of the secreted proteomes and the most highly expressed genes. Of the 167 ActE genes containing CAZy domains, 68 genes (FIG. 5, group 1) showed distinct increases in expression when grown on different polymeric substrates, 14 genes (FIG. 15, group 2) did not show any appreciable level of expression, and 85 genes (FIG. 16, group 3) showed moderate changes in expression with the different substrates. A significant frac-

tion of these genes contained translocation signals for either the Sec or twin-arginine translocation pathways, and genes encoding structural polypeptides for these translocation pathways were also highly expressed. Besides correlation with secreted proteins, the transcriptomic studies also gave insight into co-regulated gene clusters that potentially encode functional units for utilization of different polysaccharides by ActE. In the following, the 130 genes with normalized expression intensities in the top 2% of all genes are described.

During growth on cellulose, four CAZy genes (SACTE_0236, SACTE_0237, SACTE_3159, and SACTE_0482) showed >15-fold increase in transcript abundance (FIG. 5), and the corresponding proteins were highly enriched in the secreted proteome. None of these four were obviously placed in a gene cluster, and the two most highly expressed genes, SACTE_0236 and SACTE_0237, while adjacent on the chromosome, were transcribed in opposite directions. Nevertheless, these four most highly expressed genes and three others that showed >5-fold increase in transcript abundance (SACTE_3717, SACTE_6428, SACTE_2347, Table 3) were associated with a conserved 14 bp palindromic promoter sequence, TGGGAGCGCTCCA (SEQ ID NO:65) (the CebR binding element). CebR proteins are LacI/GalR-like transcriptional regulators shown to provide transcriptional control of gene expression in response to the presence of cellobiose or other small oligosaccharides in *S. griseus*, *S. reticuli*, and *Thermobifida fusca* (Marushima, Ohnishi, et al., 2009; Water and Schrempf, 1996; Deng and Fong, 2010). Likewise, the genes (SACTE_2285 to SACTE_2289) encoding a CebR regulator (SACTE_2285), a GH1 protein (β -glucosidase), a two-protein cellobiose transporter system, and an extracellular solute binding protein were associated with a CebR binding element and were also among the most highly expressed genes during growth on cellulose. These latter five genes have 75% or greater sequence identity with the cellobiose utilization operon identified in *S. griseus* and *S. reticuli* (Marushima, Ohnishi, et al., 2009; Schlosser and Schrempf, 1996). There were only 15 genes annotated as hypothetical or domain of unknown function (12%) up-regulated during growth on cellulose, a considerably smaller percentage of these than in the entire genome (27%).

TABLE 3

Analysis of upstream DNA sequence elements in ActE genes upregulated during growth on cellulose.						
Locus	Catalytic domain	CBM	Annotated function	Sequence ^a	Rank ^b	Fold change ^b
SACTE_0236	GH48	CBM2	1,4-beta cellobiohydrolase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	1	21.7
SACTE_0237	GH6	CBM2	1,4-beta cellobiohydrolase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	2	17.3
SACTE_3159	CBM33	CBM2	Cellulose-binding domain	TGGGAGCGCTC CCA (SEQ ID NO: 65)	3	16.2
SACTE_0482	GH5	CBM2	Endo-1,4-beta-glucosidase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	4	15.4
SACTE_2288			Transport systems inner membrane component	TGGGAGCGCTC CCA (SEQ ID NO: 65)	5	11.2

TABLE 3 -continued

Analysis of upstream DNA sequence elements in ActE genes upregulated during growth on cellulose.						
Locus	Catalytic domain	CBM	Annotated function	Sequence ^a	Rank ^b	Fold change ^b
SACTE_3717	GH9	CBM2	1,4-beta cellobiohydrolase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	6	9.7
SACTE_6428		CBM33	Chitin-binding, domain 3	GGGAGCGCTCC CA (SEQ ID NO: 66)	9	7.9
SACTE_2347	GH5	CBM2	Beta-mannosidase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	11	5.0
SACTE_2287			Transport systems inner membrane component	TGGGAGCGCTC CCA (SEQ ID NO: 65)	15	4.3
SACTE_2289			Family 1 extracellular solute-binding protein	TGGGAGCGCTC CCA (SEQ ID NO: 65)	19	3.9
SACTE_0352			GCN5-related N-acetyltransferase	TGGGAGCGCTC CCA (SEQ ID NO: 65)	22	3.6
SACTE_2286	GH1		Glycoside hydrolase 1	GGGAGCGCTCC CA (SEQ ID NO: 66)	27	3.4
SACTE_0483		CBM2	Cellulose-binding family protein	GGGAGCGCTCC CA (SEQ ID NO: 66)	503	1.6
SACTE_0562	GH74	CBM2	Secreted cellulase (endo)	TGGGAGCGCTC CCA (SEQ ID NO: 65)	5759	0.7
SACTE_2285			LacI family transcriptional regulator (CebR)	TGGGAGCGCTC CCA (SEQ ID NO: 65)	6229	0.6

^aPredicted binding sequence element found upstream from gene locus.^bRanking and fold change in expression intensity detected by microarray for ActE genes when grown on cellulose relative to glucose.

Several characteristics distinguished expression during growth on either xylan or chitin. First, unique sets of genes were induced, as there was only 14% and 10% overlap, respectively, when compared to cellulose. Second, ~33% of the top 2% of genes expressed during growth on either xylan or chitin were annotated as hypothetical or domain of unknown function, which greatly exceeds the unknown fraction in the cellulose secretome. During growth on xylan, two clusters of genes were up-regulated. One extended from SACTE_0357 to SACTE_0370, encoding proteins from the GH11, GH13, GH42, GH43, GH78, GH87, and CE4 families, a LacI-like transcriptional regulator, a secreted peptidase, and two sets of inner membrane transporters and associated solute binding proteins. Alternatively, during growth on chitin, three CBM33 proteins were up-regulated (SACTE_0080, SACTE_2313, SACTE_6493), and two of these had an immediately adjacent gene encoding a GH18 (SACTE_6494) or GH19 (SACTE_0081) that was up-regulated.

When ActE was grown on biomass samples, 14 additional CAZy genes were uniquely up regulated, and the corresponding proteins were identified in the proteomic analysis of biomass secretomes (FIGS. 3 and 4). A gene cluster

extending from SACTE_5858 to SACTE_5864 was uniquely up regulated during growth on biomass. Among these genes, SACTE_5860 and SACTE_5862 are annotated as a twin-arginine translocation pathway protein and an ABC transporter, respectively, while the rest are annotated either as hypothetical protein or as domain of unknown function.

Eight CAZy genes were >4-fold up-regulated during growth on cellulose, including endoglucanases, reducing and non-reducing end exoglucanases, xylanase and CBM33 proteins (FIG. 5, Table 4). During growth on xylan, eight CAZy genes were elevated >4-fold relative to glucose, including exoglucanase, xylanase, pectate lyase and other hemicellulases (Table 4). Furthermore, chitin-grown cells contained 2 up-regulated genes from CAZy families including chitinase (SACTE_4571) and a CBM33 protein [SACTE_2313, an ortholog of oxidative chitin oxidase from *S. marcescens* (Vaaje-Kolstad et al., 2010)]. Thus on a genome-wide basis ActE selectively expresses small, distinct sets of CAZy genes during growth on pure polysaccharides, which is distinct from the larger numbers of CAZy genes expressed by *T. reesei* (Herpoel-Gimbert et al., 2008), *C. thermocellum* (Raman et al., 2009; Riederer et al., 2011), and *T. fusca* (Chen and Wilson, 2007).

TABLE 4

Streptomyces sp. ActE genes with >4-fold expression increase during growth on pure polysaccharides.

Sigmacell	CAZy	Annotation	Fold increase		
			Sigmacell: glc	xylan: glc	chitin: glc
SACTE_6428	CBM33	Chitin-binding, domain 3	7.06	1.64	1.81
SACTE_3159	CBM33,2	Cellulose-binding domain, family II, bacterial type	13.03	1.90	1.29
SACTE_0358	GH11, CBM60,36	Glycoside hydrolase, family 11, active site	6.28	4.01	2.12
SACTE_0236	GH48, CBM2,37	Glycoside hydrolase, 48F	19.00	4.93	3.91
SACTE_0482	GH5, CBM2	Cellulose-binding family II/chitobiase, carbohydrate-binding domain	11.84	3.01	2.00
SACTE_2347	GH5,CE3, CBM2,37	Cellulose-binding family II/chitobiase, carbohydrate-binding domain	4.46	1.17	0.99
SACTE_0237	GH6, CBM2	1,4-beta cellobiohydrolase	15.33	1.12	0.77
SACTE_3717	GH9, CBM4,2	Carbohydrate-binding, CenC-like	8.03	2.61	1.55
SACTE_2288		Binding-protein-dependent transport systems inner membrane component	11.05	4.76	3.26
SACTE_0168		Transcription regulator LuxR, C-terminal	7.55	1.53	1.37
SACTE_0169		Glyceraldehyde 3-phosphate dehydrogenase, active site	5.01	0.75	1.08
SACTE_3594		Peptidase S1C, HrtA/DegP2/Q/S	4.52	3.36	2.70
SACTE_5228		Binding-protein-dependent transport systems inner membrane component	4.20	4.35	3.24

Xylan	CAZy	Annotation	Fold increase		
			Sigmacell: glc	xylan: glc	chitin: glc
SACTE_4029	CE4	Glycoside hydrolase/deacetylase, beta/alpha-barrel	1.07	4.35	2.22
SACTE_0358	GH11, CBM60,36	Glycoside hydrolase, family 11, active site	6.28	4.01	2.12
SACTE_0382	GH2, CBM42	Galactose-binding domain-like	1.79	4.18	2.46
SACTE_1230	GH23	Lytic transglycosylase-like, catalytic	1.29	5.64	3.70
SACTE_0816	GH31	Glycoside hydrolase, family 31	1.53	4.51	3.27
SACTE_0236	GH48, CBM2,37	Glycoside hydrolase, 48F	19.00	4.93	3.91
SACTE_1290	GH53, CBM61	Galactose-binding domain-like	1.43	4.73	2.40
SACTE_5978	PL1, CBM35	Galactose-binding domain-like	2.00	6.86	2.12
SACTE_5325		Binding-protein-dependent transport systems inner membrane component	1.78	8.26	3.76
SACTE_6023		Galactose-binding domain-like	1.92	7.84	3.34
SACTE_1834		Alkaline phosphatase D-related	1.78	7.73	3.98
SACTE_6100		Sulfate transporter	2.07	7.45	4.75
SACTE_5361		hypothetical protein	1.77	7.20	3.94
SACTE_5163		Lambda repressor-like, DNA-binding	1.47	6.89	3.29
SACTE_6365		Isocitrate lyase/phosphorylmutase	1.88	6.82	4.01
SACTE_0254		Thiolase-like	2.13	6.76	5.02
SACTE_6478		FAD-dependent pyridine nucleotide-disulfide oxidoreductase	2.00	6.72	4.46
SACTE_3570		hypothetical protein	1.61	6.71	3.72
SACTE_0590		Polyketide cyclase/dehydrase	1.55	6.67	4.42
SACTE_3152		Twin-arginine translocation pathway, signal sequence	1.41	6.60	2.98
SACTE_5285		Bacterial bifunctional deaminase-reductase, C-terminal	1.71	6.54	3.33

TABLE 4-continued

<i>Streptomyces</i> sp. ActE genes with >4-fold expression increase during growth on pure polysaccharides.				
SACTE_1383	Glycerophosphoryl diester phosphodiesterase	1.08	6.50	3.51
SACTE_4333	Binding-protein-dependent transport systems inner membrane component	1.37	6.46	3.58
SACTE_3876	hypothetical protein	1.21	6.42	2.73
SACTE_6340	Monooxygenase, FAD-binding	2.82	6.27	3.69
SACTE_4237	hypothetical protein	1.82	6.27	2.91
SACTE_5136	NAD(P)-binding domain	2.20	6.27	2.87
SACTE_6561	hypothetical protein	2.92	6.06	5.65
SACTE_0686	Transcription regulator AsnC-type	0.88	6.04	2.72
SACTE_0817	NUDIX hydrolase, conserved site	1.96	6.03	3.19
SACTE_3004	Type II secretion system F domain	1.67	6.01	4.18
SACTE_1835	DoxX	1.66	5.97	3.30
SACTE_1933	hypothetical protein	0.93	5.96	2.77
SACTE_6290	Glyoxalase/bleomycin resistance	1.86	5.95	4.10
SACTE_5583	protein/dioxygenase	1.33	5.87	4.56
SACTE_0586	hypothetical protein	1.40	5.81	2.90
SACTE_0046	NADH:flavin oxidoreductase/NADH oxidase, N-terminal	2.48	5.75	4.56
SACTE_1096	Mandelate racemase/muconate lactonizing enzyme, N-terminal	1.19	5.73	3.32
SACTE_2897	hypothetical protein	1.18	5.73	3.81
SACTE_5359	Rhs repeat-associated core	1.30	5.70	2.41
SACTE_0200	hypothetical protein	1.34	5.67	3.64
SACTE_0018	hypothetical protein	1.67	5.63	3.58
SACTE_5542	hypothetical protein	2.03	5.61	3.52
SACTE_3137	hypothetical protein	1.46	5.61	3.91
SACTE_0017	DNA helicase, UvrD/REP type	2.32	5.58	4.26
SACTE_0672	hypothetical protein	1.53	5.54	3.20
SACTE_1393	Urease, beta subunit	2.08	5.53	3.67
SACTE_0064	Transcription regulator PadR N-terminal-like	2.17	5.52	3.07
SACTE_1168	Peptidase S1/S6, chymotrypsin/Hap	0.98	5.51	3.36
SACTE_6371	hypothetical protein	1.37	5.51	3.44
SACTE_4334	Binding-protein-dependent transport systems inner membrane component	1.46	5.50	3.35
SACTE_2457	CDP-glycerol glycerophosphotransferase	1.07	5.48	3.79
SACTE_4734	Binding-protein-dependent transport systems inner membrane component	1.21	5.44	3.31
SACTE_3661	hypothetical protein	1.76	5.44	3.25
SACTE_0036	hypothetical protein	1.75	5.43	2.99
SACTE_6005	Citrate synthase-like, core	1.01	5.38	2.90
SACTE_6562	hypothetical protein	2.34	5.36	3.37
SACTE_1937	Major facilitator superfamily MFS-1	0.88	5.34	3.02
SACTE_6220	Dodecin flavoprotein	2.13	5.32	5.08
SACTE_0778	FMN-binding split barrel	1.13	5.28	2.72
SACTE_5672	Acyltransferase 3	1.33	5.28	3.09
SACTE_5989	Cysteine-rich domain	1.40	5.24	3.11
SACTE_5296	HTH transcriptional regulator, MarR	1.42	5.22	2.96
SACTE_2021	hypothetical protein	1.44	5.17	2.54
SACTE_1845	Transposase, IS4-like	1.69	5.16	3.30
SACTE_1771	Phage T4-like virus tail tube gp19	1.55	5.10	1.71
SACTE_2583	hypothetical protein	1.38	5.10	3.11
SACTE_5957	Helix-turn-helix, HxIR type	2.38	5.09	3.95
SACTE_4642	hypothetical protein	1.31	5.08	3.05
SACTE_3695	Aminoglycoside/hydroxyurea antibiotic resistance kinase	1.41	5.03	3.76
SACTE_0079	ATPase-like, ATP-binding domain	2.21	5.01	2.98

TABLE 4-continued

<i>Streptomyces</i> sp. ActE genes with >4-fold expression increase during growth on pure polysaccharides.				
SACTE_0727	hypothetical protein	2.54	5.00	3.88
SACTE_0019	hypothetical protein	1.37	5.00	2.40
SACTE_6422	<i>Streptomyces</i> cyclase/dehydrase	2.40	4.99	3.57
SACTE_4348	Bacterial extracellular solute-binding protein, family 5	1.60	4.97	3.06
SACTE_5318	Forkhead-associated (FHA) domain	1.50	4.93	2.84
SACTE_5413	Urease accessory protein UreF	1.94	4.93	2.52
SACTE_5434	Glutathione S-transferase, C-terminal-like	2.41	4.93	2.96
SACTE_6061	Glyoxalase/bleomycin resistance protein/dioxygenase	1.61	4.92	2.18
SACTE_0025	hypothetical protein	1.58	4.92	4.22
SACTE_5552	Transposase, IS4-like	1.94	4.92	3.26
SACTE_4156	HTH transcriptional regulator, LysR	1.57	4.86	2.81
SACTE_5600	hypothetical protein	1.78	4.83	2.01
SACTE_5331	Conserved hypothetical protein CHP03086	1.56	4.82	2.96
SACTE_0784	hypothetical protein	1.43	4.80	2.65
SACTE_0045	NAD(P)-binding domain	1.74	4.78	3.35
SACTE_5426	Twin-arginine translocation pathway, signal sequence	0.80	4.77	2.68
SACTE_2654	4Fe-4S ferredoxin, iron-sulfur binding domain	1.30	4.77	2.68
SACTE_2288	Binding-protein-dependent transport systems inner membrane component	11.05	4.76	3.26
SACTE_2324	Membrane insertion protein, OxaA/YidC, core	0.91	4.75	2.58
SACTE_0142	Amidohydrolase 2	1.28	4.71	2.65
SACTE_0787	hypothetical protein	1.66	4.70	2.93
SACTE_5790	hypothetical protein	1.28	4.69	2.83
SACTE_6291	hypothetical protein	1.25	4.68	3.13
SACTE_6499	hypothetical protein	1.66	4.67	3.29
SACTE_6548	Lytic transglycosylase-like, catalytic	1.97	4.66	3.20
SACTE_3087	Major facilitator superfamily MFS-1	1.30	4.66	3.26
SACTE_5512	hypothetical protein	1.79	4.64	3.48
SACTE_0491	hypothetical protein	2.44	4.63	2.71
SACTE_0312	Thiamine pyrophosphate enzyme, C-terminal TPP-binding	2.32	4.60	3.49
SACTE_6130	hypothetical protein	1.47	4.55	2.64
SACTE_3787	Helix-turn-helix type 3	1.38	4.53	2.73
SACTE_0040	hypothetical protein	1.64	4.52	4.80
SACTE_2461	Macrocin-O-methyltransferase	1.07	4.51	3.00
SACTE_5041	hypothetical protein	1.50	4.49	3.25
SACTE_5540	Transposase, IS204/IS1001/IS1096/IS1165	1.79	4.49	2.99
SACTE_0776	Protein of unknown function DUF6, transmembrane	1.34	4.48	2.52
SACTE_0785	Bacterial TniB	1.67	4.43	2.93
SACTE_0360	Binding-protein-dependent transport systems inner membrane component	1.70	4.43	2.39
SACTE_3569	Protein of unknown function DUF1023	1.00	4.42	2.78
SACTE_2986	hypothetical protein	1.62	4.42	2.96
SACTE_4732	Twin-arginine translocation pathway, signal sequence	2.08	4.41	2.72
SACTE_5228	Binding-protein-dependent transport systems inner membrane component	4.20	4.35	3.24
SACTE_0406	Binding-protein-dependent transport systems inner membrane component	1.34	4.35	2.52
SACTE_6516	Binding-protein-dependent transport systems inner membrane component	2.24	4.34	3.41
SACTE_1781	hypothetical protein	1.16	4.34	2.56
SACTE_5936	Radical SAM	1.43	4.33	2.23

TABLE 4-continued

<i>Streptomyces</i> sp. ActE genes with >4-fold expression increase during growth on pure polysaccharides.				
SACTE_0819	Protein of unknown function DUF962	1.50	4.33	2.83
SACTE_4539	NERD	1.42	4.32	3.98
SACTE_0532	Binding-protein-dependent transport systems inner membrane component	3.47	4.31	2.42
SACTE_3300	hypothetical protein	1.68	4.31	2.59
SACTE_6277	hypothetical protein	2.24	4.31	3.11
SACTE_0941	Twin-arginine translocation pathway, signal sequence	1.32	4.30	2.63
SACTE_1115	GntR, C-terminal	1.57	4.29	2.63
SACTE_6105	Fatty acid hydroxylase	1.63	4.29	2.78
SACTE_4407	Spherulation-specific family 4	1.19	4.29	4.15
SACTE_5387	hypothetical protein	1.24	4.27	3.08
SACTE_5053	NmrA-like	1.23	4.27	3.05
SACTE_5562	Amino acid ABC transporter, permease protein, 3-TM domain, His/Glu/Gln/Arg/opine family	1.37	4.26	3.75
SACTE_5522	Galactose-binding domain-like	1.82	4.26	2.62
SACTE_5484	Transcription regulator, TetR-like, DNA-binding, bacterial/archaeal	1.45	4.21	3.24
SACTE_6526	Restriction endonuclease, type IV-like, Mrf	2.31	4.20	2.40
SACTE_4164	hypothetical protein	1.06	4.19	2.48
SACTE_4979	Transcription regulator, TetR-like, DNA-binding, bacterial/archaeal	1.20	4.19	2.34
SACTE_0952	hypothetical protein	1.33	4.18	2.02
SACTE_1785	hypothetical protein	1.25	4.17	1.94
SACTE_3454	hypothetical protein	1.46	4.16	2.32
SACTE_1271	Class II aldolase/adducin, N-terminal	1.77	4.16	2.65
SACTE_1760	hypothetical protein	1.38	4.13	2.07
SACTE_0035	hypothetical protein	1.93	4.13	3.13
SACTE_0247	Protein of unknown function DUF2241	1.30	4.10	2.77
SACTE_3796	F420-dependent enzyme, PPOX class, family Rv2061, putative	1.43	4.10	3.33
SACTE_4641	hypothetical protein	1.43	4.09	2.60
SACTE_4816	Peptidase S26, conserved region	1.17	4.09	2.77
SACTE_2331	Major facilitator superfamily MFS-1	1.15	4.08	2.20
SACTE_1666	hypothetical protein	1.44	4.07	2.46
SACTE_5867	Mammalian cell entry, mce1C	1.79	4.07	2.92
SACTE_2705	AMP-binding, conserved site	1.38	4.07	2.75
SACTE_6014	Binding-protein-dependent transport systems inner membrane component	0.89	4.07	2.51
SACTE_2018	Putative DNA binding domain	1.05	4.06	2.63
SACTE_5690	Gluconate transporter	1.00	4.05	2.29
SACTE_3243	hypothetical protein	0.91	4.05	2.23
SACTE_0786	Polynucleotidyl transferase, ribonuclease H fold	1.81	4.03	2.98
SACTE_6450	Rhamnose isomerase related	2.72	4.02	2.90
SACTE_0097	Beta-lactamase-related	1.70	4.02	2.52
SACTE_6341	FMN-binding split barrel, related	1.82	4.01	2.45
SACTE_1483	hypothetical protein	0.82	4.01	2.75
SACTE_0754	Uncharacterised protein family UPF0060	1.21	4.00	2.51
SACTE_5308	Winged helix-turn-helix transcription repressor DNA-binding	1.33	4.00	1.56
SACTE_5862	ABC transporter, conserved site	1.87	4.00	3.05

TABLE 4-continued

<i>Streptomyces</i> sp. ActE genes with >4-fold expression increase during growth on pure polysaccharides.					
Chitin	CAZy	Annotation	Fold increase		
			Sigmatocell: glc	xylan: glc	chitin: glc
SACTE_2313	CBM33	Chitin-binding, domain 3	1.08	1.24	4.77
SACTE_4571	GH18, CBM57,2	EF-Hand 1, calcium-binding site	0.88	1.37	4.08
SACTE_5381		hypothetical protein	1.31	3.09	10.06
SACTE_5386		hypothetical protein	0.96	1.59	8.49
SACTE_1949		Peptidase M4, thermolysin	1.30	2.16	7.57
SACTE_6519		Binding-protein-dependent transport systems inner membrane component	2.00	3.04	7.36
SACTE_0243		Protein kinase-like domain	1.68	2.55	6.89
SACTE_6520		ABC transporter, conserved site	1.03	1.18	6.25
SACTE_5384		hypothetical protein	1.16	2.39	5.99
SACTE_6463		hypothetical protein	1.28	2.52	5.85
SACTE_6561		hypothetical protein	2.92	6.06	5.65
SACTE_5383		hypothetical protein	1.06	1.69	5.28
SACTE_6518		hypothetical protein	1.66	1.91	5.21
SACTE_4797		hypothetical protein	2.22	0.34	5.19
SACTE_6170		Domain of unknown function DUF1996	1.47	3.49	5.12
SACTE_6220		Dodecin flavoprotein	2.13	5.32	5.08
SACTE_0254		Thiolase-like	2.13	6.76	5.02
SACTE_2678		Protein of unknown function DUF397	1.40	1.13	5.02
SACTE_5968		hypothetical protein	1.58	1.31	4.90
SACTE_4757		Acetyl-coenzyme A carboxyltransferase, C-terminal	1.64	0.59	4.86
SACTE_0040		hypothetical protein	1.64	4.52	4.80
SACTE_6100		Sulfate transporter	2.07	7.45	4.75
SACTE_1833		Twin-arginine translocation pathway, signal sequence	1.64	1.56	4.64
SACTE_5583		hypothetical protein	1.33	5.87	4.56
SACTE_0046		NADH: flavin oxidoreductase/NADH oxidase, N-terminal	2.48	5.75	4.56
SACTE_5398		hypothetical protein	1.45	1.73	4.55
SACTE_6144		Twin-arginine translocation pathway, signal sequence	1.21	1.13	4.52
SACTE_6478		FAD-dependent pyridine nucleotide-disulfide oxidoreductase	2.00	6.72	4.46
SACTE_0590		Polyketide cyclase/dehydrase	1.55	6.67	4.42
SACTE_2112		Homeodomain-like	1.44	1.33	4.40
SACTE_0017		DNA helicase, UvrD/REP type	2.32	5.58	4.26
SACTE_5841		Protein of unknown function, ATP binding	1.90	3.09	4.24
SACTE_0025		hypothetical protein	1.58	4.92	4.22
SACTE_3004		Type II secretion system F domain	1.67	6.01	4.18
SACTE_4407		Spherulation-specific family 4	1.19	4.29	4.15
SACTE_0307		Protein of unknown function DUF320, <i>Streptomyces</i> species	1.13	1.79	4.15
SACTE_6290		Glyoxalase/bleomycin resistance protein/dioxygenase	1.86	5.95	4.10
SACTE_5286		hypothetical protein	1.33	3.34	4.07
SACTE_5953		Protein of unknown function, ATP binding	1.35	2.11	4.05
SACTE_6365		Isocitrate lyase/phosphorylmutase	1.88	6.82	4.01

Example 6: Composition of ActE Secretome is Dependent on ActE Growth Substrate

To identify secreted proteins, supernatants from ActE cultures grown on glucose, cellobiose, cellulose, xylan,

chitin, switchgrass, AFEX-SG, and IL-SG were analyzed by LC-MS/MS (FIG. 3 and FIG. 18). The proteins were sorted into a descending rank according to spectral counts, and sets whose spectral counts summed to 95% of the total protein in each secretome are shown. FIG. 3A summarizes the per-

centages of CAZy families in the detected proteins. The glucose secretome had a protein concentration of ~0.03 g/L of culture medium, and among the 136 proteins identified only 3% had a CAZy annotation. Indeed, the majority (>90%) likely originated from cell lysis. In contrast, the polysaccharide secretomes had a protein concentration of ~0.3 g/L of culture medium, a ~10-fold increase from the glucose secretome. Pectate lyase (SACTE_1310), chondroitin/alginate lyase (SACTE_4638), an extracellular solute binding protein (SACTE_4343), bacterioferritin (SACTE_1546), and catalase (SACTE_4439) were observed in all polysaccharide secretomes. The first two proteins, SACTE_1310 and SACTE_4638, have signal peptides and are thus secreted as part of the response needed for growth on polysaccharides.

FIG. 3 and FIG. 18 further demonstrate that 22 proteins accounted for 95% of the total spectral counts during growth on cellulose; two-thirds were from CAZy families. The five most abundant proteins, in order and representing ~85% of the total spectral counts, were reducing and non-reducing exoglucanases (SACTE_0236 and SACTE_0237), a CBM33 polysaccharide monooxygenase (SACTE_3159), an endoglucanase (SACTE_0482), and a β-mannosidase (SACTE_2347). The first four proteins encode a non-redundant set of enzymes that likely provide the essential activities required for utilization of crystalline cellulose (Deboy, et al., 2008). Among the 22 most abundant proteins, there were representatives from 9 different GH families, two CE families, two PL families, and two additional CBM33 proteins. Collectively, these secreted proteins represent ~20% of the CAZy composition in the ActE genome.

There were substantial differences in the composition of the xylan and chitin secretomes as compared to the cellulose secretome (FIG. 3 and FIG. 18). In the xylan secretome, 92 proteins comprise 95% of the detected spectral counts. Twenty GHS from 18 different CAZy families were included, along with 1 CE4 and 2 PL family proteins. Thus, growth on xylan elicits secretion of representatives from half of the total CAZy families found in the ActE genome. The broad distribution of hemicellulytic enzymes in the xylan secretome contrasts with the considerably less diverse composition of the chitin secretome, which consists of 7 representatives from GH18 (e.g., chitinase, endo beta-N-acetylglucosaminidase), 2 from GH19 (e.g., chitinase, lysozyme), and 1 chitinolytic CBM33 (FIG. 18). While chitinolytic CAZy families account for two-thirds of the proteins secreted during growth on chitin, they represent only ~6% of the diversity of CAZy families found in the genome. These results document the substantially different substrate-specific responses of ActE during growth on different polysaccharides.

The secretomes isolated from cells grown on switchgrass, AFEX-SG, and IL-SG contained the highly abundant secreted proteins identified in the purified cellulose and xylan experiments and some additional proteins. These additional proteins likely reflect cellular response to the more complex composition of polysaccharides present in the biomass samples. The increased diversity of proteins present in the biomass secretome also increased the efficiency of reaction with plant biomass (FIG. 2C). In total, the biomass secretomes contained 31 different CAZy families that contributed to the total spectral counts (~70% of the CAZy families present in the ActE genome), thus representing coordinated and extensive use of CAZyme families present in the ActE genome for biomass utilization.

The gene loci of the 117 proteins observed only in the glucose secretome are: SACTE_0494; SACTE_0514;

SACTE_0541; SACTE_0548; SACTE_0604;
SACTE_0669; SACTE_0687; SACTE_0800;
SACTE_0810; SACTE_0899; SACTE_1006;
SACTE_1045; SACTE_1068; SACTE_1069;
5 SACTE_1111; SACTE_1201; SACTE_1240;
SACTE_1285; SACTE_1328; SACTE_1344;
SACTE_1368; SACTE_1419; SACTE_1426;
SACTE_1506; SACTE_1522; SACTE_1586;
SACTE_1650; SACTE_1861; SACTE_1888;
10 SACTE_1934; SACTE_2036; SACTE_2049;
SACTE_2068; SACTE_2238; SACTE_2403;
SACTE_2431; SACTE_2468; SACTE_2558;
SACTE_2645; SACTE_2729; SACTE_2755;
SACTE_2756; SACTE_2801; SACTE_2819;
15 SACTE_3012; SACTE_3037; SACTE_3067;
SACTE_3086; SACTE_3088; SACTE_3097;
SACTE_3219; SACTE_3327; SACTE_3361;
SACTE_3371; SACTE_3385; SACTE_3389;
SACTE_3392; SACTE_3414; SACTE_3438;
20 SACTE_3511; SACTE_3604; SACTE_3716;
SACTE_3896; SACTE_3948; SACTE_3955;
SACTE_3956; SACTE_3960; SACTE_3961;
SACTE_3989; SACTE_3995; SACTE_4030;
SACTE_4031; SACTE_4038; SACTE_4039;
25 SACTE_4073; SACTE_4081; SACTE_4083;
SACTE_4145; SACTE_4191; SACTE_4194;
SACTE_4205; SACTE_4224; SACTE_4281;
SACTE_4283; SACTE_4376; SACTE_4397;
SACTE_4399; SACTE_4415; SACTE_4462;
30 SACTE_4501; SACTE_4550; SACTE_4565;
SACTE_4566; SACTE_4567; SACTE_4568;
SACTE_4591; SACTE_4610; SACTE_4616;
SACTE_4618; SACTE_4652; SACTE_4718;
SACTE_4768; SACTE_4791; SACTE_4795;
35 SACTE_4830; SACTE_4860; SACTE_4873;
SACTE_4926; SACTE_4959; SACTE_5028;
SACTE_5081; SACTE_5192; SACTE_5267;
SACTE_5482; SACTE_5519; SACTE_5983; and
SACTE_6342.
40 The gene loci of the 9 proteins observed only in the Sigmaell secretome are: SACTE_0236; SACTE_0482;
SACTE_0562; SACTE_2313; SACTE_2347;
SACTE_3590; SACTE_3717; SACTE_4571; and
SACTE_6428.
45 The gene loci of the 46 proteins observed only in the xylan secretome are: SACTE_0081; SACTE_0169;
SACTE_0365; SACTE_0379; SACTE_0383;
SACTE_0464; SACTE_0528; SACTE_0549;
SACTE_0634; SACTE_0880; SACTE_1003;
50 SACTE_1130; SACTE_1239; SACTE_1324;
SACTE_1325; SACTE_1356; SACTE_1364;
SACTE_1367; SACTE_1603; SACTE_1680;
SACTE_1858; SACTE_1949; SACTE_2768;
SACTE_3064; SACTE_4231; SACTE_4246;
55 SACTE_4363; SACTE_4459; SACTE_4483;
SACTE_4515; SACTE_4607; SACTE_4612;
SACTE_4624; SACTE_4730; SACTE_4755;
SACTE_4858; SACTE_5166; SACTE_5230;
SACTE_5231; SACTE_5418; SACTE_5457;
60 SACTE_5630; SACTE_5647; SACTE_5682;
SACTE_5751; and SACTE_6439.

In the xylan secretome, five proteins accounted for half of the total secreted protein. These were xylanases (GH10 and GH11, respectively; SACTE_0265, 9.7% and SACTE_0358, 8.1%), extracellular xylose isomerase (SACTE_5230, 12.7%), acetyl xylan esterase (CE4; SACTE_0357, 11.7%), and pectate lyase (PL1,

SACTE_5978, 6.6%). Among the remaining 98 proteins, there were numerous GH families. Given the complexity of hemicellulose, which is enriched in xylan but also contains many other sugars and many different bonding linkages between these sugars, it is noted that these additional proteins represent many GH families associated with unique hemicellulolytic activities.

Although not analyzed in FIG. 34, the chitin secretome contained ten proteins from the chitinase GH18 (49% of total protein) and GH19 (21%) families. In addition, the CBM33 protein SACTE_2313, having 50% primary sequence identity with the CBP21 chitin oxygenase from *S. marcescens*, was also detected (3.9%). Insect molt and fungal hyphae provide abundant chitin, likely accounting for the utility of these enzymes in the natural environment. There were 50 other proteins (63 total) that comprised 95% of the chitin secretome. Relative to the glucose, Sigmacell, and xylan secretomes, the following 15 proteins were observed only in the chitin secretome: SACTE_0746, SACTE_0844, SACTE_0860, SACTE_1702, SACTE_2033, SACTE_2059, SACTE_2062, SACTE_2384, SACTE_3685, SACTE_4468, SACTE_4472, SACTE_4727, SACTE_5330, SACTE_5764, and SACTE_6494.

The gene loci of the 19 proteins observed only in the switchgrass secretome are: SACTE_0642; SACTE_1130; SACTE_1250; SACTE_1858; SACTE_2033; SACTE_3012; SACTE_3777; SACTE_4198; SACTE_4571; SACTE_4624; SACTE_4669; SACTE_4676; SACTE_4718; SACTE_4738; SACTE_5220; SACTE_5418; SACTE_5685; SACTE_5751; and SACTE_5880.

The gene loci of the 8 proteins observed only in the IL-SG secretome are: SACTE_0132; SACTE_0880; SACTE_2556; SACTE_4246; SACTE_4515; SACTE_4702; SACTE_5231; and SACTE_5330.

There were no proteins observed only in the AFEX-SG secretome when compared to either the switchgrass or IL-SG secretomes.

Example 7: Minimized Size of ActE Enzymes Increases Specific Activity

When ActE is grown on Sigmacell, AFEX-SG, IL-SG, AFEX-CS, unbleached lodgepole pine kraft pulp (UBLPKP) or bleached spruce wood kraft pulp (BSKP), the characteristic secretome consists of the proteins that permit decon-

struction of these substrates into sugars that can be used for growth (FIG. 23). Interestingly, ActE is not capable of growing on lodgepole pine pretreated by SPORL, indicating this pretreatment produces toxins that inhibit the growth of highly cellulolytic microbes. When ActE is grown on cellobiose, which it does readily and rapidly, it produces a secretome that is distinct from those obtained from ActE grown on cellulose, xylan or biomass substrates, demonstrating that ActE has highly specific responses to different polymeric substances that are present in biomass. This behavior is distinct from that observed for *T. fusca*, another cellulolytic Actinomycete, and from *C. thermocellum*, where each organism produced similar sets of secreted proteins during growth on either cellulose or cellobiose (Chen and Wilson, 2007; Riederer et al., 2011). This result indicates ActE contains a unique regulatory mechanism for controlling cellulose deconstruction genes that can provide exquisite control of their production under desired circumstances.

For a single enzyme from a secretome, (Segel, Enzyme kinetics: behavior and analysis of rapid equilibrium and steady state enzyme systems. Wiley, New York, 1993) the specific activity ($\mu\text{mol}/\text{min}/\text{mg}$) is defined as mol of product formed per unit time (i.e., $\mu\text{mol}/\text{min}$) per unit mass of enzyme (i.e., mg). Specific activity is the parameter that must be used in making comparisons of catalytic properties between enzymes with different molecular masses. If two enzyme isoforms yield the same $\mu\text{mol}/\text{min}$, the isoform with the smaller molecular weight will, by definition, have the higher specific activity. In this application, it is relevant to consider the implications of a 10% or more reduction in the mass of an enzyme required to treat gigatonnes of biomass.

In the cellulose secretome, five proteins contributed ~85% of the total spectral counts. These were reducing and non-reducing end exoglucanases, endoglucanases, and CBM33 (SACTE_0237, SACTE_0236, SACTE_2347, SACTE_0482 and SACTE_3159); xylanase, another endoglucanase, and another CBM33 were also abundant (SACTE_0265, SACTE_3717 and SACTE_6428). According to the definition provided above, size minimization is a way to achieve the desired increases in specific activity. Interestingly, the set of ActE enzymes described above are on average 10% smaller in mass than their closest orthologs from *T. fusca* (Chen and Wilson, 2007), suggesting size minimization may have occurred in ActE (Table 5). These enzymes also provide all of the requisite catalytic reactions needed for the deconstruction of crystalline cellulose.

TABLE 5

ActE cellulose secretome proteins and corresponding best match
in *T. fusca*. The single protein SACTE_0237 is the best match to both Cel6A and Cel6B
suggesting one protein from ActE might replace two proteins from another organism.

ActE Gene locus	CAZy	residues	MW	identity	coverage	<i>T. fusca</i>	Protein	residues	MW
						Gene locus	name		
SACTE_0237	GH6	586		49	80	Tfu_1074	Cel6A	441	45844
SACTE_0237	GH6	586	61062	62	93	Tfu_0620	Cel6B	596	63548
SACTE_0236	GH48	954	100726	57	95	Tfu_1959	Cel48A	984	107127
SACTE_2347	GH8	562	57753	45	23	Tfu_2176	Cel9A	880	95203
SACTE_3159	CBM33	362	37787	42	71	Tfu_1665	E8	438	46808
SACTE_0482	GH5	456	47654	51	97	Tfu_0901	Cel5A	466	49807

TABLE 5-continued

ActE cellulose secretome proteins and corresponding best match in <i>T. fusca</i> . The single protein SACTE_0237 is the best match to both Cel6A and Cel6B suggesting one protein from ActE might replace two proteins from another organism.									
ActE Gene locus	CAZy	residues	MW	identity	coverage	<i>T. fusca</i> Gene locus	Protein name	residues	MW
SACTE_0265	GH10	458	47683	44	95	Tfu_2923	Xyl10A	491	53185
SACTE_3717	GH9	909	96338	61	82	Tfu_1627	Cel9B	998	107045
SACTE_6428	CBM33	222	24668	62	99	Tfu_1268	E7	222	25372
Average identity, coverage				53	82				
Sum ActE		4509	473671			with Cel6A		4920	530391
		4509	473671			with Cel6B		5075	548095
Percentage with Cel6A		92%	89%			with Cel6A,B		5516	593939
Percentage with Cel6B		89%	86%						
Percentage with Cel6A,B		82%	80%						

Example 8: ActE Secretome Specific Activity is Comparable to that of SPEZYME OPT™

The enzymatic activities of ActE secretomes were compared with a commercial secretome, SPEZYME CP. The enzyme cocktail of SPEZYME CP was prepared from *T. reesei* Rut-C30, thus providing a useful, routinely available reference point for the capabilities of other cellulolytic organisms. HPLC analysis showed that the ActE cellulose secretome released cellobiose as the primary product during reaction with cellulose (FIG. 2A, 95% of products), which is distinct from the higher proportion of glucose produced by the *T. reesei* secretome. Similarly, the primary products from xylan and mannan were xylobiose and mannobiose, respectively. Upon accounting for total glucose equivalents released, the ActE secretome obtained from growth on pure cellulose had specific activity that was about half of that provided by SPEZYME CP (FIG. 2A, inset). Interestingly, the ActE secretome obtained from growth on pure cellulose had higher specific activity for deconstruction of pure mannan than SPEZYME CP (FIG. 2B). Additionally, the ActE secretome obtained from growth on pure xylan had higher specific activity for reaction with pure xylan than SPEZYME CP. Cellulose, xylan, and mannan are all abundant in pinewood, thus accounting for the necessity of each of the major catalytic activities detected.

Anion exchange chromatography was performed to fractionate the ActE secretome obtained from cells grown on cellulose as the sole carbon source. We identified fractions that hydrolyzed pure polysaccharides by biochemical assays (FIG. 7), and confirmed the identity of the protein or proteins contained in these fractions by mass spectrometry (FIG. 17). Where multiple polypeptides were present, the identity of each was confirmed by mass spectrometry to correspond to the indicated gene locus. In several cases, these most likely arise from proteolysis of a single protein found in the secretome. Fractions containing the maximum cellulase activity were highly enriched in SACTE_0236 and SACTE_0237, reducing and non-reducing end cellobiohydrolases from the GH6 and GH48 families, respectively. SACTE_0265 and SACTE_2347 were identified as the major proteins present in fractions associated with xylan and mannan hydrolysis, respectively. A CBM33 polysaccharide monooxygenase (SACTE_3159) was also identified in the ion exchange profile. Moreover, beta-1,3 glucanase activity was identified in fractions that were enriched in SACTE_4755.

When ActE was grown on either ammonia fiber expansion-treated switchgrass (AFEX-SG) (Li, C. et al., 2011) or

ionic liquid-treated switchgrass (IL-SG), the secretomes had ~2-fold increase in specific activity relative to the cellulose secretome and were equivalent to SPEZYME CP for reaction with both the AFEX- and IL-treated biomass (FIG. 2C) (Li, C. et al., 2011). The ActE secretomes retained greater than 60% of maximal activity for the hydrolysis of AFEX- and IL-SG from 30 to 55° C. and 35 to 47° C., respectively, which is comparable to recent reports on the temperature profile of secretomes from thermophilic biomass-degrading fungi (Tolonen et al., 2011) (FIG. 8A). The secretomes showed a pH optimum of ~7 for reaction with AFEX-SG and a pH optimum of ~8 for reaction with IL-SG. Moreover, these secretomes retained greater than 60% of maximal activity over the ranges of pH 4.5 to 8.0, and pH 7.0 to 8.0, respectively (FIG. 8B). These optimal pH values are considerably higher than observed for SPEZYME CP.

Example 9: ActE Produces Cellobiose as the Primary Extracellular Product of Cellulose Utilization

The isolated ActE secretomes contained substantial ability to release reducing sugars from pure polysaccharides. Cellobiose accounted for ~95% of soluble sugar released from pure cellulose and glucose represented the remainder; cellotriose and cellobionic acid were not detected. Neither cellobiosidase nor β-glucosidase was detected in the ActE secretome. Thus ActE produces cellobiose as the primary extracellular product of cellulose utilization and also grows vigorously on this. Dominance of cellobiose may help to channel cellulolytic activity to only a subset of the *Sirex* community. Since genes encoding cellobiose oxidase and cellobiose dehydrogenase (Eastwood et al., 2011; Langston et al., 2011) were not present in ActE, biological reduction systems for the CBM33 proteins may be provided by other members of the *Sirex* community, in analogy to that described for the heterologous complex of *T. aurantiacus* GH61 and *Humicola insolens* cellobiose dehydrogenase (Langston et al., 2011).

Example 10: Enzymatic Activity of the ActE Secretome can be Improved by Adding One or More Enzymes from Other Organisms or Sources

In the ActE secretome, enzymes SACTE_0236, SACTE_0237, and SACTE_3717 (GH48, GH6, and GH9, respectively) showed decreases in content of the native forms over a 24 h period, and SACTE_0236 and SACTE_0237 were converted into ~50 kDa fragments (FIG.

24). SACTE_0359 (CBM33) also showed a time-dependent decrease. The reactions could be slowed but not eliminated by addition of phenylmethylsulfonyl fluoride (1 mM), a possible inhibitor of serine proteases (Turini et al., 1969), but not by EDTA (10 mM), a possible inhibitor of metallo-proteases (Trop and Birk, 1970).

SACTE_5668, a serine protease, was detected in all pure polysaccharide secretomes (FIG. 3), while another metallo-peptidase, SACTE_3389 annotated as peptidase M24B, X-Pro dipeptidase/aminopeptidase P, was detected in all secretomes at low level (0.026%). The protease SACTE_5530 (peptidase S1/S6, chymotrypsin/Hap, 0.1%) was also present in all polysaccharide and biomass samples. The proteases SACTE_5668 (annotated secreted peptidase, 0.3%) and SACTE_4231 (serine/cysteine peptidase, trypsin-like, 0.039%) were also detected in all pure polysaccharide secretomes, and the protease SACTE_6303 (serine/cysteine peptidase, trypsin-like, 0.039%) was also present in all biomass samples. Elimination of one or more of these proteases may impart stabilization of the enzymatic activity in the secreted proteome.

Addition of CellC_CBM3a, an engineered exoglucanase (FIG. 22) that produces cellobiose with low specific activity alone (FIG. 21), gave a synergistic increase in the activity of the ActE cellulose secretome. This result demonstrates the potential for heterologous supplementation of the ActE secretome to improve its performance by replacing an enzyme activity that is lost to proteolysis.

Example 11: ActE Cellulolytic Activity Requires a Minimal Set of Enzymes

When the ActE secretome obtained from growth on cellulose was fractionated by ion exchange chromatography (FIG. 7), several fractions were obtained that could be tested in unary, binary, ternary and quaternary combinations for reconstitution of cellulose hydrolysis and other enzymatic activities (FIG. 25). SDS PAGE and LC-MS/MS analysis showed that these fractions contained the following polypeptides in the approximate weight percentages: fraction 1, SACTE_3159 (CBM33/CBM2 oxidative endocellulase, 95%) and SACTE_4738 (GH16 β -1,3 endoglucanase, 5%); fraction 2, SACTE_0237 (GH6 non-reducing end exocellulase, 60%), SACTE_0482 (GH5 endocellulase, 25%), SACTE_4755 (GH64 β -1,3 glucanase, 10%) and SACTE_3159 (oxidative endocellulase, <5%); and fraction 3, SACTE_0236 (GH48 reducing end exocellulase, 75%), SACTE_3717 (GH9 endocellulase, 20%) and SACTE_5457 (GH46 chitinase, 5%). These results demonstrate that SACTE_3159 (oxidative endocellulase) provides a complementary activity to SACTE_0482 and SACTE_3717 (hydrolytic endocellulolytic activity). Evidently, the oxidative reaction provides breaks in the cellulose strands that can be readily used by non-reducing and reducing end exocellulases also present in the secretome to processively deconstruct the polymeric material.

According to the current understanding of reactions required for hydrolysis of crystalline cellulose, SACTE_3159 (CBM33/CBM2 oxidative endocellulase), SACTE_0482 (GH5), and SACTE_3717 provide endocellulolytic activities, while SACTE_0237 (GH6) provides non-reducing end exocellulase reaction and SACTE_0236 (GH48) provides reducing end exocellulase activity.

FIG. 16 shows that the secretome contains beta-1,3 endoglucanase activity. The majority of this activity corresponds to the fractions containing SACTE_4738 and SACTE_4755. These enzymes hydrolyze callose, a cellulose-like material

that is typically produced by plants in respond to wounding by invasive insects and other trauma.

The proteins described here constitute a naturally evolved and matched set specialized for the hydrolysis of cellulosic substrates.

Example 12: ActE Mannanase Specific Activity Increases as Mannanase Molecular Weight Decreases

FIG. 26 shows that the mannanase activity present in the ActE secretome is associated with fractions containing various naturally truncated variants of SACTE_2347 (GH5) with molecular weights of ~57, ~45, and ~37 kDa. Fractions F9 through F1 from ion exchange chromatographic separation of the ActE secretome were examined for mannan-deconstruction activity by Zymogram assay. The basis of the Zymogram assay is as follows: Congo Red stain interacts with the polysaccharide fraction (mannan) incorporated into the gel and imparts a red color. When an enzyme's activity hydrolyzes the mannan, the interaction of Congo Red with the polysaccharide is broken and the gel takes on a dark grey appearance. Of note, the strongest mannanase activity was observed in fraction F1, which primarily contains the 37 kDa truncated variant. Corresponding to the definition of specific activity given above, the 37 kDa variant has an ~35% increase in specific activity relative to the 57 kDa variant. This provides a naturally produced example of how size reduction may contribute to increased specific activity of enzymes.

Example 13: Recombination of ActE Secretome Fractions Provides Synergistic Cellulolytic Activity

FIG. 25 shows synergy of reaction obtained by recombining fractions obtained from ion exchange fractionation. In FIG. 25A, reactions were obtained from combinations of the fractions indicated by stars in FIG. 27 and FIG. 28. Combination of fractions E5 (oxidative endocellulase) and E11 (hydrolytic endo- and exocellulases) gave a ~30% increase in product yield over that expected from the arithmetic sum of reactions of E5 and E11 alone, i.e., synergy in reaction. Combination of fractions E5, E11 and F10 (hydrolytic endo- and exocellulases) gave ~60% increase in reactivity. In FIG. 25B, reactions were obtained from recombining fractions shown in FIG. 16. Titration of fraction B1 (full-length oxidative endocellulase) into D15 (hydrolytic endo- and exocellulases) shows an optimal reactivity at ~1:1 ratio of proteins from the two fractions, while an excess of B1 relative to D15 causes decrease in reaction because of depletion of required exocellulase activities. Titration of fraction C4 (truncated oxidative endocellulase and beta-1,3-endocellulase) with D15 gave maximal stimulation (62% increase) at an 80:20 proportion. These results indicate both forms of oxidative endocellulase SACTE_3159 are catalytically active, with the smaller form providing a higher synergistic response, again corresponding to a specific activity increase associated with size minimization.

Example 14: The Function of ActE Xylanases can be Assigned by Functional Assay of Proteins Produced by Using Cell-Free Translation

FIG. 29 shows that both of the xylanases identified in the fractions of ActE secretomes obtained from ion exchange chromatography can also be expressed using cell-free translation and demonstrated to be xylanases by catalytic activity

51

assays. These proteins are SACTE_0265 and SACTE_0358. Other proteins that are not secreted were successfully expressed (SACTE_2548, SACTE_2286, SACTE_437) as control proteins, and as expected from their predicted intra-cellular localization, none of these controls exhibited xylanase activity. The negative result with the control proteins also demonstrates that the wheat germ extract used for cell-free translation of novel cellulolytic enzymes does not have an endogenous xylanase activity, as established in US Patent Application Publication No.: US2010/037094 (Fox and Elsen).¹⁰

Example 15: Total Protein Secreted by ActE can be Increased

A minimal set of enzymes for biomass deconstruction can be defined by combining the additional enzymes whose expression is elicited during growth on biomass (Table 1) with enzymes uniquely expressed during growth on cellulose and xylan.¹⁵

Besides assembling the proper enzymatic constituents, the level of total protein secreted is an important biotechnological constraint for industrial enzyme production. FIG. 30 shows the non-optimized level of secreted protein obtained from growth of ActE on different biomass substrates. By use of lignocellulosic substrates for growth, secreted protein levels up to 0.25 g per liter of culture medium can be readily obtained. Growth on non-polymeric substrates such as cellobiose does not elicit a secreted protein response. FIG. 15, FIG. 16 and Table 1 indicate that the twin-arginine pathway (Tat) is used during growth, thus identifying this pathway as playing a key role in the secretion of enzymes required for extracellular deconstruction of biomass polysaccharides (Natale et al., 2008; Chater et al., 2010). Methods to increase the titer of secreted proteins are known, and have been highly effectively when applied to *Streptomyces* and other organisms (Cereghino et al., 2002; Zhang et al., 2006; Nijland and Kuipers, 2008; Chater et al., 2010; Schuster and Schmoll, 2010). These established methods can be applied to ActE to obtain more concentrated secretome preparations.³⁰

Example 16: ActE Enzymatic Activity Corresponds with Optimal Growth Conditions of Fermentation Organisms

FIG. 31 shows the temperature versus activity profile for ActE secretomes for reaction with cellulose, xylan and mannan. These profiles are well matched to the growth optima range for mesophilic fermentation organisms such as *Saccharomyces cerevisiae*, *Zymomonas mobilis*, *Escherichia coli* or others (Jarboe et al., 2010; Peralta-Yahya and Keasling, 2010), which are widely used for ethanol production from sugar hydrolysates. These hydrolysates are produced from biomass by the enzymatic action of highly cellulolytic secretomes, such as those described here from ActE. These optima are also well matched with the conditions found in the rumen, where the efficiency of conversion of animal feed, which is a biomass material, can be improved by addition of enzymes.³⁵

FIG. 32 shows the pH versus activity profiles for ActE secretomes for reaction with cellulose, xylan and mannan. These profiles are well matched to the growth optima range for fermentation organisms such as *S. cerevisiae*, *Z. mobilis*, *E. coli* or other organisms (Jarboe et al., 2010; Peralta-Yahya and Keasling, 2010) which are widely used for ethanol production from sugar hydrolysates such as might be produced from biomass by a highly cellulolytic secretome, such

52

as those described here from ActE. These optima are also well matched with the conditions found in the rumen, where the efficiency of conversion of animal feed, which is a biomass material, can be improved by addition of enzymes.⁵ The ActE secretome retains high specific activity (>80% of maximal) at pH 7, which closely approximates that of the rumen. Secretomes from fungi such as *T. reesei* are considerably less active at neutral pH, rendering them less effective at neutral pH.

The high cellulolytic capacity of ActE, and its corresponding secretomes, coupled with the temperature and pH optima described above permit assembly of two-part systems to effect the simultaneous deconstruction of biomass and fermentation to fuels.¹⁰

Example 17: ActE Induction in Medium Containing Various Percentages of Cellulose

To determine ActE's growth profile on cellulose as a carbon source ActE was grown in M63 media plus 5 g/L carbon. The carbon source ratio was adjusted from 100% cellulose to 100% glucose, total carbon in each culture was equal. Cells were grown for 6 days at 30 degrees. Supernatant was harvested, filtered, and separated by 4-20% SDS-PAGE. Results suggest that ActE is induced in media containing as little as 20% cellulose, with optimal induction in medium containing between 80%-100% cellulose (FIG. 33).²⁰

Example 18: Discussion

The work presented here provides the first genome-wide insight into how an aerobic microbe deconstructs polysaccharides. ActE achieves efficient utilization of cellulose by a simple combination of well-understood hydrolytic reactions with newly identified oxidative reactions. The two required exoglucanases are each encoded by a single gene, which also represents the only example of their respective GH families in the genome. The proteins encoded by these genes provide reactions that are complementary to the reactions of other enzymes in the secretome, and provide cellobiose as the major product of reaction. We have discovered that many of the highly abundant enzymes secreted by ActE during growth on cellulose have reduced size relative to their orthologs from closely related organisms. This novel finding suggests natural evolution to improve specific activity has already occurred in ActE in response to growth in the highly specialized insect association. Additional specializations of ActE were identified by demonstrating the secretion of a unique set of proteins in response to biomass. In addition, this work defines how simple new combinations of improved biomass deconstruction enzymes can be assembled according to the propensities of the naturally evolved system.³⁰

The present work also indicates that insect-associated microbes such as ActE are important contributors to the vigorous attack on biomass by insects. The 'highly invasive' designation given to *Sirex* has been generally attributed to the combined action of wasp and fungus (Tabata and Abe, 2000; Bergeron et al., 2011). Species convergence is now recognized in the microbial communities associated with insects (Suen et al., 2010; Hulcr et al., 2011). Given the ubiquitous presence of *Streptomyces* in these communities, the enzymatic properties described here also contribute a potential risk to pine forests, including those used for industrial purposes.³⁵

The invention has been described in connection with what are presently considered to be the most practical and preferred embodiments. However, the present invention has been presented by way of illustration and is not intended to be limited to the disclosed embodiments. Accordingly, those skilled in the art will realize that the invention is intended to encompass all modifications and alternative arrangements within the spirit and scope of the invention as set forth in the appended claims.

REFERENCES

1. Baldrian, P. & Valaskova, V. *FEMS Microbiol Rev* 32, 501-521 (2008).
2. Cantarel, B. L. et al. *Nucleic Acids Res* 37, D233-238 (2009).
3. Schuster, A. & Schmoll, M. *Appl Microbiol Biotechnol* 87, 787-799 (2010).
4. Crawford, D. L. *Appl Environ Microb* 35, 1041-1045 (1978).
5. Goodfellow, M. & Williams, S. T. *Annu Rev Microbiol* 37, 189-216 (1983).
6. McCarthy, A. J. & Williams, S. T. *Gene* 115, 189-192 (1992).
7. Schlatter, D. et al. *Microb Ecol* 57, 413-420 (2009).
8. Wachinger, G., Bronnenmeier, K., Staudenbauer, W. L. & Schrempf, H. *Appl Environ Microb* 55, 2653-2657 (1989).
9. Ishaque, M. & Kluepfel, D. *Can J Microbiol* 26, 183-189 (1980).
10. Semedo, L. T. et al. *Int J Syst Evol Microbiol* 54, 1323-1328, (2004).
11. Schlochtermeier, A., Walter, S., Schroder, J., Moorman, M. & Schrempf, H. *Mol Microbiol* 6, 3611-3621 (1992).
12. Walter, S. & Schrempf, H. *Appl Environ Microb* 62, 1065-1069 (1996).
13. Forsberg, Z. et al. *Protein Science* 20, 1479-1483 (2011).
14. Bignell, D. E., Anderson, J. M. & Crosse, R. *Fems Microbiol Ecol* 85, 151-159 (1991).
15. Pasti, M. B. & Belli, M. L. *Fems Microbiol Lett* 26, 107-112 (1985).
16. Pasti, M. B., Pometto, A. L., Nuti, M. P. & Crawford, D. L. *Appl Environ Microb* 56, 2213-2218 (1990).
17. Schafer, A. et al. *J Appl Bacteriol* 80, 471-478 (1996).
18. Adams, A. S. et al. *ISME Journal* 5, 1323-1331 (2011).
19. Bergeron, M. J. et al. *Fungal Biol* 115, 750-758 (2011).
20. Kukor, J. J. & Martin, M. M. *Science* 220, 1161-1163 (1983).
21. Lynd, L. R., Weimer, P. J., van Zyl, W. H. & Pretorius, I. S. *Microbiol Mol Biol Rev* 66, 506-577 (2002).
22. Deboy, R. T. et al. *J Bacteriol* 190, 5455-5463 (2008).
23. Riederer, A. et al. *Appl Environ Microb* 77, 1243-1253 (2011).
24. Marushima, K., Ohnishi, Y. & Horinouchi, S. *J Bacteriol* 191, 5930-5940 (2009).
25. Walter, S. & Schrempf, H. *Mol Gen Genet* 251, 186-195 (1996).
26. Deng, Y. & Fong, S. S. *Appl Environ Microb* 76, 2098-2106 (2010).
27. Li, C. et al. *Bioresour Technol* 102, 6928-6936 (2011).
28. Tolonen, A. C. et al. *Mol Syst Biol* 6, 461 (2011).
29. Hyatt, D. et al. *BMC Bioinformatics* 11, 119 (2010).
30. Rutherford, K. et al. *Bioinformatics* 16, 944-945 (2000).
31. Medema, M. H. et al. *Nucleic Acids Res* 39, W339-346 (2011).
32. Shannon, P. et al. *Genome Res* 13, 2498-2504 (2003).
33. Miller, G. L. *Anal Chem* 31, 426-428 (1959).
34. Merino and Cherry, *Adv. Biochem. Eng. Biotechnol.* 108:95-120, 2007
35. Bayer et al., *Cehm. Rec.* 8:364-377, 2008
36. Wilson, *Curr. Opin. Microbiol.* 14:259-263, 2011
37. Vuong and Wilson, *Biotechnol. Bioeng.* 107:195-205, 2010
38. Scharf et al., *PLoS One* 6:e21709, 2011
39. Luyten et al., *App. Environ. Microbiol.* 72:412-417, 2006
40. Hess et al., *Science* 331:463-467, 2011
41. Schlochtermeier et al., *App. Environ. Microbiol.* 58:3240-3248, 1992
42. Wilson, *Crit. Rev. Biotechnol.* 12:45-63; 1992
43. Langston et al., *App. Environ. Microbiol.* 77:7007-7015, 2011
44. Quinlan et al., *PNAS* 108:15079-15084, 2011
45. Vaaje-Kolstad et al., *Science* 330:219-222, 2010
46. Klepzig et al., *Environ. Entomol.* 38: 67-77, 2009
47. Teather R M, Wood P J (1982) *Appl Environ Microbiol* 43:777-780
48. Bradford, *Anal. Biochem.* 72:248-254 (1976)
49. Anne and Van Mellaert. *FEMS Microbiol. Lett.*, 114; 121-8 (1993)
50. Saloheimo and Pakula, *Microbiology*, Epub date 2011/ 11/05
51. Wood et al., *Biotechnol. Bioeng.* 55:547-55 (1997)
52. Balan et al., *Meth. Mol. Biol.* 581:61-77, 2009
53. Makino et al., *Meth. Microbiol.* 607:127-134, 2010
54. Blommel et al., *Meth. Mol. Biol.* 498:55-73, 2009
55. Galm et al., *J. Nat. Prod.* 74:526-536, 2011;
56. Susi et al. *J. Environ. Manage.* 92:1681-1689, 2011
57. Balan et al., *Meth. Mol. Biol.* 581:61-77, 2009
58. Chundawat et al., *J. Am. Chem. Soc.* 133:11163-11174, 2011
59. Cheng et al., *Biomacromol.* 12:933-941, 2011
60. Wang et al., *Biotechnol. Progress* 25:1086-1093, 2009
61. Tian et al., *Biotechnol. Progress* 27: 419-427, 2011
62. Altschul et al., *J. Mol. Biol.* 215:403-410, 1990
63. Herpoel-Gimbert et al., *Biotechnol. Biofuels* 1:18; 2008
64. Raman et al., *PLoS One* 4:e5271, 2009
65. Chen and Wilson, *J. Bacteriol.* 189:6260-6265, 2007
66. Eastwood et al., *Science* 333:762-765, 2011
67. Langston et al., *App. Environ. Microbiol.* 77:7007-7015, 2011
68. Turini et al., *J. Pharmacol. Exper. Therapeu.* 167:98-104, 1969
69. Trop and Birk, *Biochem. J.* 27:419-427, 1970
70. Natale et al., *Biochim. Biophys. Acta* 1778:1735-1756, 2008
71. Chater et al., *FEMS Microbiol. Rev.* 34:171-198, 2010
72. Cereghino et al., *Curr. Opin. Biotechnol.* 13:329-332, 2002
73. Zhang et al., *Nat. Biotechnol.* 24:100-104, 2006
74. Nijland and Kuipers, *Recent Pat. Biotechnol.* 2:79-87, 2008
75. Jarboe et al., *J. Biomed. Biotechnol.* 761042, 2010
76. Peralta-Yahya and Keasling, *Biotechnol. J.* 5:147-162, 2010
77. Tabata and Abe, *Mycoscience* 41:585-539, 2000
78. Suen et al., *PLoS Gen.* 6: e1001129, 2010
79. Hulcr et al., *Micro. Ecol.* 61:759-768, 2011
80. Kestler et al., *BMC Bioinform.* 9:67, 2008

SEQUENCE LISTING

65 This specification includes the sequence listing that is concurrently filed in computer readable form. This sequence listing is incorporated by reference herein.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 66

<210> SEQ ID NO 1

<211> LENGTH: 586

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 1

Met	Ser	Arg	Thr	Ser	Arg	Thr	Thr	Leu	Arg	Arg	Ser	Arg	Thr	Ala	Leu
1								10						15	

Met	Ala	Ala	Gly	Ala	Leu	Val	Ala	Ala	Ala	Gly	Ser	Ala	Ala	Ala
	20					25					30			

Ala	Ala	Pro	Phe	Gly	Ala	Thr	Ala	Ala	Ala	Ala	Gly	Cys	Thr	Val
	35					40					45			

Asp	Tyr	Lys	Ile	Gln	Asn	Gln	Trp	Asn	Gly	Gly	Leu	Thr	Ala	Ser	Val
	50					55					60				

Ser	Val	Thr	Asn	Asn	Gly	Asp	Ala	Ile	Ser	Gly	Trp	Gln	Leu	Gln	Trp
65						70			75		80				

Ser	Phe	Ala	Gly	Gly	Glu	Gln	Val	Ser	Gln	Gly	Trp	Asn	Ala	Thr	Val
	85					90			95						

Ser	Gln	Ser	Gly	Ser	Ala	Val	Thr	Ala	Lys	Asp	Ala	Gly	Tyr	Asn	Ala
	100					105			110						

Ala	Leu	Ala	Thr	Gly	Ala	Ser	Ala	Ser	Phe	Gly	Phe	Asn	Ala	Thr	Gly
	115					120			125						

Asn	Gly	Asn	Ser	Val	Val	Pro	Ala	Thr	Phe	Lys	Leu	Asn	Gly	Val	Thr
	130					135			140						

Cys	Asn	Gly	Gly	Thr	Thr	Gly	Pro	Thr	Asp	Pro	Thr	Asp	Pro	Thr	Asp
145						150			155		160				

Pro	Thr	Asp	Pro	Thr	Asp	Pro	Pro	Ala	Gly	Asn	Arg	Val	Asp	Asn	Pro
	165					170			175						

Tyr	Gln	Gly	Ala	Lys	Val	Tyr	Val	Asn	Pro	Glu	Trp	Ser	Ala	Asn	Ala
	180					185			190						

Ala	Ala	Glu	Pro	Gly	Gly	Asp	Arg	Ile	Ala	Asp	Gln	Pro	Thr	Gly	Val
	195					200			205						

Trp	Leu	Asp	Arg	Ile	Ala	Ala	Ile	Glu	Gly	Ala	Asn	Gly	Ser	Met	Gly
	210					215			220						

Leu	Arg	Asp	His	Leu	Asp	Glu	Ala	Leu	Thr	Gln	Lys	Gly	Ser	Gly	Glu
225						230			235		240				

Leu	Val	Val	Gln	Val	Val	Ile	Tyr	Asn	Leu	Pro	Gly	Arg	Asp	Cys	Ala
	245					250			255						

Ala	Leu	Ala	Ser	Asn	Gly	Glu	Leu	Gly	Pro	Thr	Glu	Ile	Gly	Arg	Tyr
	260					265			270						

Lys	Thr	Glu	Tyr	Ile	Asp	Pro	Ile	Ala	Glu	Ile	Leu	Gly	Asp	Pro	Lys
	275					280			285						

Tyr	Ala	Gly	Leu	Arg	Ile	Val	Thr	Val	Glu	Ile	Asp	Ser	Leu	Pro	
	290					295			300						

Asn	Leu	Val	Thr	Asn	Ala	Gly	Gly	Arg	Pro	Thr	Ala	Thr	Pro	Ala	Cys
305						310			315		320				

Asp	Val	Met	Lys	Ala	Asn	Gly	Asn	Tyr	Val	Lys	Gly	Val	Gly	Tyr	Ala
	325					330			335						

Leu	Asn	Lys	Leu	Gly	Asp	Ala	Pro	Asn	Val	Tyr	Asn	Tyr	Ile	Asp	Ala
	340					345			350						

Gly	His	His	Gly	Trp	Ile	Gly	Trp	Asp	Asp	Asn	Phe	Gly	Ala	Ser	Ala
	355					360			365						

US 10,144,941 B2

57

-continued

Glu Ile Phe His Glu Ala Ala Thr Ala Glu Gly Ala Thr Val Asn Asp
370 375 380

Val His Gly Phe Ile Thr Asn Thr Ala Asn Tyr Ser Ala Leu Lys Glu
385 390 395 400

Glu Asn Phe Ser Ile Asp Asp Ala Val Asn Gly Thr Ser Val Arg Gln
405 410 415

Ser Lys Trp Val Asp Trp Asn Arg Tyr Thr Asp Glu Leu Ser Phe Ala
420 425 430

Gln Ala Phe Arg Asn Glu Leu Val Ser Val Gly Phe Asn Ser Gly Ile
435 440 445

Gly Met Leu Ile Asp Thr Ser Arg Asn Gly Trp Gly Gly Ala Asn Arg
450 455 460

Pro Ser Gly Pro Gly Ala Asn Thr Ser Val Asp Thr Tyr Val Asp Gly
465 470 475 480

Gly Arg Tyr Asp Arg Arg Ile His Leu Gly Asn Trp Cys Asn Gln Ala
485 490 495

Gly Ala Gly Leu Gly Glu Arg Pro Gln Ala Ala Pro Glu Pro Gly Ile
500 505 510

Asp Ala Tyr Val Trp Met Lys Pro Pro Gly Glu Ser Asp Gly Ser Ser
515 520 525

Ser Glu Ile Pro Asn Asp Glu Gly Lys Gly Phe Asp Arg Met Cys Asp
530 535 540

Pro Thr Tyr Thr Gly Asn Ala Arg Asn Asn Asn Asn Met Ser Gly Ala
545 550 555 560

Leu Gly Gly Ala Pro Val Ser Gly Lys Trp Phe Ser Ala Gln Phe Gln
565 570 575

Glu Leu Met Lys Asn Ala Tyr Pro Ala Leu
580 585

58

<210> SEQ_ID NO 2
<211> LENGTH: 954
<212> TYPE: PRT
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 2

Val Ala Ala Leu Ala Leu Pro Leu Gly Met Thr Ala Ala Gly Thr
1 5 10 15

Glu Ala Gln Ala Ala Ala Val Ala Cys Ser Val Asp Tyr Thr Ser
20 25 30

Asp Trp Gly Ser Gly Phe Thr Thr Glu Leu Thr Leu Thr Asn Arg Gly
35 40 45

Ser Ala Ala Ile Asp Gly Trp Thr Leu Thr Tyr Asp Tyr Ala Gly Asn
50 55 60

Gln Gln Leu Thr Ser Gly Trp Ser Gly Thr Trp Ser Gln Ser Gly Lys
65 70 75 80

Thr Val Ser Val Lys Asn Ala Ala Trp Asn Gly Ala Ile Ala Gly
85 90 95

Ala Ala Val Thr Thr Gly Ala Gln Phe Thr Tyr Ser Gly Ala Asn Thr
100 105 110

Ala Pro Thr Thr Phe Ala Val Asn Gly Thr Val Cys Ala Gly Ala His
115 120 125

Gln Pro Pro Ile Ala Val Leu Thr Ser Pro Ala Ala Gly Ala Val Phe
130 135 140

Ser Ala Gly Asp Pro Val Pro Leu Ala Ala Thr Ala Ala Ala Asp
145 150 155 160

US 10,144,941 B2

59**60**

-continued

Gly Ala Thr Ile Ser Lys Val Glu Phe Tyr Asp Asp Thr Thr Leu Leu
165 170 175

Gly Thr Asp Thr Thr Ser Pro Tyr Ser Tyr Glu Ala Gly Gln Leu Ala
180 185 190

Ala Gly Ser His Ser Val Tyr Ala Arg Ala Tyr Asp Ser Leu Gly Ala
195 200 205

Ser Ala Asp Ser Pro Pro Ala Gly Ile Thr Val Val Thr Gly Pro Ala
210 215 220

Val Val Val Ser Pro Ala Gln Leu Gly Val Gln Gln Gly Arg Ser Gly
225 230 235 240

Thr Phe Asp Val Ser Leu Ser Thr Ala Pro Ala Ala Asp Val Thr Val
245 250 255

Thr Ala Ala Arg Ser Ala Gly Asn Thr Gly Leu Ser Val Thr Gly Gly
260 265 270

Ser Thr Leu Thr Phe Thr Pro Ala Asn Trp Ser Thr Pro Gln Lys Val
275 280 285

Thr Val Thr Ala Asp Gly Ser Gly Thr Gly Ala Ala Thr Phe Thr Val
290 295 300

Thr Ala Pro Gly His Gly Lys Ala Glu Val Thr Val Thr Gln Leu Ala
305 310 315 320

Ala Ala Lys Glu Tyr Asp Ala Arg Phe Leu Asp Leu Tyr Gly Lys Ile
325 330 335

Thr Asp Pro Ala Asn Gly Tyr Phe Ser Pro Glu Gly Ile Pro Tyr His
340 345 350

Ser Val Glu Thr Leu Ile Val Glu Ala Pro Asp His Gly His Glu Thr
355 360 365

Thr Ser Glu Ala Tyr Ser Tyr Leu Ile Trp Leu Gln Ala Met Tyr Gly
370 375 380

Lys Ile Thr Gly Asp Trp Thr Lys Phe Asn Gly Ala Trp Asp Thr Met
385 390 395 400

Glu Thr Tyr Met Ile Pro Thr His Ala Asp Gln Pro Thr Asn Ser Phe
405 410 415

Tyr Asp Ala Ser Lys Pro Ala Thr Tyr Ala Pro Glu His Asp Thr Pro
420 425 430

Asn Glu Tyr Pro Ala Val Leu Asp Gly Ser Ala Ser Ser Gly Ser Asp
435 440 445

Pro Ile Ala Ala Glu Leu Lys Ser Ala Tyr Gly Thr Asp Asp Ile Tyr
450 455 460

Gly Met His Trp Ile Gln Asp Val Asp Asn Val Tyr Gly Tyr Gly Asn
465 470 475 480

Ala Pro Gly Thr Cys Ala Ala Gly Pro Thr Gln Ala Gly Pro Ser Tyr
485 490 495

Ile Asn Thr Phe Gln Arg Gly Ser Gln Glu Ser Val Trp Glu Thr Val
500 505 510

Thr His Pro Thr Cys Asp Asn Phe Thr Tyr Gly Gly Ala Asn Gly Tyr
515 520 525

Leu Asp Leu Phe Thr Gly Asp Ser Ser Tyr Ala Lys Gln Trp Lys Phe
530 535 540

Thr Asn Ala Pro Asp Ala Asp Ala Arg Ala Val Gln Ala Ala Tyr Trp
545 550 555 560

Ala Asp Val Trp Ala Lys Glu Gln Gly Lys Ala Gly Glu Val Ala Asp
565 570 575

US 10,144,941 B2

61**62**

-continued

Thr Val Gly Lys Ala Ala Lys Met Gly Asp Tyr Leu Arg Tyr Ser Met
 580 585 590
 Phe Asp Lys Tyr Phe Lys Lys Ile Gly Asp Cys Val Gly Pro Thr Thr
 595 600 605
 Cys Pro Ala Gly Ser Gly Lys Asp Ser Ala His Tyr Leu Met Ser Trp
 610 615 620
 Tyr Tyr Ala Trp Gly Gly Ala Thr Asp Thr Ser Ala Gly Trp Ser Trp
 625 630 635 640
 Arg Ile Gly Ser Ser His Ala His Gly Gly Tyr Gln Asn Pro Met Ala
 645 650 655
 Ala Tyr Ala Leu Ser Ser Val Ala Asp Leu Lys Pro Lys Ser Ala Thr
 660 665 670
 Gly Ala Gln Asp Trp Ala Lys Ser Leu Asp Arg Gln Leu Asp Phe Tyr
 675 680 685
 Gln Trp Leu Gln Ser Asp Glu Gly Ala Ile Ala Gly Gly Ala Thr Asn
 690 695 700
 Ser Trp Lys Gly Ser Tyr Ala Gln Pro Pro Ala Gly Thr Pro Thr Phe
 705 710 715 720
 Tyr Gly Met Tyr Tyr Asp Glu Lys Pro Val Tyr His Asp Pro Pro Ser
 725 730 735
 Asn Gln Trp Phe Gly Phe Gln Ala Trp Ser Met Glu Arg Val Ala Glu
 740 745 750
 Tyr Tyr His Glu Ser Gly Asp Ala Gln Ala Lys Ala Val Leu Asp Lys
 755 760 765
 Trp Val Asp Trp Ala Leu Ser Glu Thr Thr Val Asn Pro Asp Gly Thr
 770 775 780
 Tyr Leu Met Pro Ser Thr Leu Gln Trp Ser Gly Ala Pro Asp Thr Trp
 785 790 795 800
 Asn Ala Ser Asn Pro Gly Ala Asn Ala Gln Leu His Val Thr Val Ala
 805 810 815
 Asp Tyr Thr Asp Asp Val Gly Val Ala Gly Ala Tyr Ala Arg Thr Leu
 820 825 830
 Thr Tyr Tyr Ala Ala Lys Ser Gly Asp Thr Glu Ala Glu Ala Thr Ala
 835 840 845
 Glu Ala Leu Leu Asp Gly Met Trp Gln His His Gln Asp Asp Ala Gly
 850 855 860
 Val Ala Val Pro Glu Thr Arg Ala Asp Tyr Asn Arg Phe Asp Asp Pro
 865 870 875 880
 Val Tyr Val Pro Gly Gly Trp Thr Gly Ala Met Pro Asn Gly Asp Thr
 885 890 895
 Val Asp Glu Asp Ser Thr Phe Leu Ser Ile Arg Ser Phe Tyr Lys Asp
 900 905 910
 Asp Pro Asn Trp Pro Gln Val Gln Ala Tyr Leu Asp Gly Gly Ala Ala
 915 920 925
 Pro Val Phe Thr Tyr His Arg Phe Trp Ala Gln Ala Asp Ile Ala Leu
 930 935 940
 Ala Leu Gly Ala Tyr Ala Asp Leu Leu Glu
 945 950

<210> SEQ ID NO 3

<211> LENGTH: 362

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 3

-continued

Met Ala Arg Arg Ser Arg Leu Ile Ser Leu Ala Ala Val Leu Ala Thr
 1 5 10 15
 Leu Leu Gly Ala Leu Gly Leu Thr Ala Leu Trp Pro Gly Lys Ala Glu
 20 25 30
 Ala His Gly Val Ala Met Thr Pro Gly Ser Arg Thr Tyr Leu Cys Gln
 35 40 45
 Leu Asp Ala Leu Ser Gly Thr Gly Ala Leu Asn Pro Thr Asn Pro Ala
 50 55 60
 Cys Arg Asp Ala Leu Ser Gln Ser Gly Ala Asn Ala Leu Tyr Asn Trp
 65 70 75 80
 Phe Ala Val Leu Asp Ser Asn Ala Gly Gly Arg Gly Ala Gly Tyr Val
 85 90 95
 Pro Asp Gly Ser Leu Cys Ser Ala Gly Asp Arg Ser Pro Tyr Asp Phe
 100 105 110
 Ser Ala Tyr Asn Ala Ala Arg Ala Asp Trp Pro Arg Thr His Leu Thr
 115 120 125
 Ser Gly Ala Thr Leu Lys Val Gln Tyr Ser Asn Trp Ala Ala His Pro
 130 135 140
 Gly Asp Phe Arg Val Tyr Leu Thr Lys Pro Gly Trp Ala Pro Thr Ser
 145 150 155 160
 Glu Leu Ala Trp Asp Asp Leu Gln Leu Val Gln Thr Val Ser Asn Pro
 165 170 175
 Pro Gln Gln Gly Gly Ala Gly Thr Asn Gly Gly His Tyr Tyr Trp Asp
 180 185 190
 Leu Ala Leu Pro Ser Gly Arg Ser Gly Asp Ala Leu Met Phe Ile Gln
 195 200 205
 Trp Val Arg Ser Asp Ser Gln Glu Asn Phe Phe Ser Cys Ser Asp Ile
 210 215 220
 Val Phe Asp Gly Gly Asn Gly Glu Val Thr Gly Ile Gly Gly Thr Gly
 225 230 235 240
 Thr Pro Thr Pro Thr Pro Thr Pro Thr Pro Thr Pro Thr Asp
 245 250 255
 Pro Glu His Ser Gly Ser Cys Met Ala Val Tyr Asn Val Val Ser Ser
 260 265 270
 Trp Ala Gly Gly Phe Gln Ala Ser Val Glu Val Met Asn His Gly Thr
 275 280 285
 Glu Pro Arg Asn Gly Trp Ala Val Gln Trp Lys Pro Gly Ser Gly Thr
 290 295 300
 Gln Ile Asn Ser Val Trp Asn Gly Ser Leu Ser Thr Gly Ser Asp Gly
 305 310 315 320
 Thr Val Thr Val Arg Asp Val Asp His Asn Arg Val Ile Ala Pro Asp
 325 330 335
 Gly Ser Val Thr Phe Gly Phe Thr Ala Thr Ser Thr Gly Asn Asp Tyr
 340 345 350
 Pro Ala Gly Thr Ile Gly Cys Val Thr Ser
 355 360

<210> SEQ_ID NO 4

<211> LENGTH: 456

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 4

Val Lys Arg Phe Leu Ala Leu Ala Thr Cys Ala Thr Val Leu Gly

US 10,144,941 B2

65**66**

-continued

1	5	10	15
Leu Thr Ala Leu Thr Gly Pro Gln Ala Val Ala Ala Ala Gly Cys Thr			
20	25	30	
Ala Asp Tyr Thr Ile Thr Ser Gln Trp Gln Gly Gly Phe Gln Ala Ala			
35	40	45	
Val Lys Val Thr Asn Leu Gly Thr Pro Val Thr Gly Trp Lys Leu Thr			
50	55	60	
Phe Thr Leu Pro Asp Ala Gly Gln Lys Val Val Gln Gly Trp Asn Ala			
65	70	75	80
Ala Trp Ser Gln Ser Gly Ser Ala Val Thr Ala Ala Gly Ala Asp Trp			
85	90	95	
Asn Gly Thr Leu Ala Thr Gly Ala Ser Ala Glu Ala Gly Phe Val Gly			
100	105	110	
Ser Phe Thr Gly Ala Asn Pro Pro Pro Thr Ala Phe Ala Leu Asn Gly			
115	120	125	
Val Ala Cys Thr Gly Ser Thr Gly Glu Pro Pro Ala Gly Ser Asp Gly			
130	135	140	
Gly Thr Pro Val Asp Val Asn Gly Gln Leu His Val Cys Gly Val Asn			
145	150	155	160
Leu Cys Asn Gln Tyr Asp Arg Pro Val Gln Leu Arg Gly Met Ser Thr			
165	170	175	
His Gly Ile Gln Trp Phe Asp Ala Cys Tyr Asp Ala Ala Ser Leu Asp			
180	185	190	
Ala Leu Ala Asn Asp Trp Lys Ser Asp Leu Leu Arg Ile Ala Met Tyr			
195	200	205	
Val Gln Glu Asp Gly Tyr Glu Thr Asp Pro Ala Gly Phe Thr Arg Arg			
210	215	220	
Val Asn Asp Leu Val Asp Met Ala Glu Ala Arg Gly Met Tyr Ala Leu			
225	230	235	240
Ile Asp Phe His Thr Leu Thr Pro Gly Asp Pro Asn Val Asn Leu Asp			
245	250	255	
Arg Ala Lys Thr Phe Phe Ala Ser Val Ala Ala Arg Asn Ala Gly Lys			
260	265	270	
Lys Asn Val Ile Tyr Glu Ile Ala Asn Glu Pro Asn Gly Val Thr Trp			
275	280	285	
Thr Ala Val Lys Ser Tyr Ala Glu Gln Val Ile Pro Val Ile Arg Ala			
290	295	300	
Ala Asp Pro Asp Ala Val Val Ile Val Gly Thr Arg Gly Trp Ser Ser			
305	310	315	320
Leu Gly Val Ser Asp Gly Ser Asp Glu Ser Glu Val Val Asn Ser Pro			
325	330	335	
Val Asn Ala Thr Asn Ile Met Tyr Ala Phe His Phe Tyr Ala Ala Ser			
340	345	350	
His Lys Asp Ala Tyr Arg Ser Thr Leu Ser Arg Ala Ala Arg Leu			
355	360	365	
Pro Leu Phe Val Thr Glu Phe Gly Thr Val Ser Ala Thr Gly Gly			
370	375	380	
Ala Met Asp Arg Ala Ser Thr Thr Ala Trp Leu Asp Leu Leu Asp Gln			
385	390	395	400
Leu Lys Ile Ser Tyr Ala Asn Trp Thr Tyr Ser Asp Ala Pro Glu Ser			
405	410	415	
Ser Ala Ala Phe Arg Pro Gly Thr Cys Gly Gly Asp Tyr Ser Gly			
420	425	430	

-continued

Ser Gly Val Leu Thr Glu Ser Gly Ala Leu Leu Lys Asn Arg Ile Ser
435 440 445

Thr Pro Asp Ser Phe Pro Thr Gly
450 455

<210> SEQ ID NO 5

<211> LENGTH: 458

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 5

Met Ala Lys Ile Pro Ala Arg Ala Arg Ala Leu Ser Val Leu
1 5 10 15

Thr Ala Gly Val Leu Ala Ala Ala Gly Val Val Ser Leu Ala Gly Thr
20 25 30

Ala Glu Ala Ala Gly Thr Leu Gly Asp Ala Ala Ala Lys Gly Arg
35 40 45

Tyr Phe Gly Thr Ala Val Ala Ala Asn His Leu Gly Glu Ala Pro Tyr
50 55 60

Ala Ser Thr Leu Asp Ala Gln Phe Asp Ser Val Thr Pro Glu Asn Glu
65 70 75 80

Met Lys Trp Asp Ala Val Glu Gly Ser Arg Asn Ser Phe Thr Phe Thr
85 90 95

Ala Ala Asp Gln Ile Val Ser His Ala Gln Ser Lys Gly Met Lys Val
100 105 110

Arg Gly His Thr Leu Val Trp His Ser Gln Leu Pro Gly Trp Val Gly
115 120 125

Gly Leu Gly Ala Thr Asp Leu Arg Ala Ala Met Asn Asn His Ile Thr
130 135 140

Gln Val Met Thr His Tyr Lys Gly Lys Ile His Ser Trp Asp Val Val
145 150 155 160

Asn Glu Ala Phe Gln Asp Gly Asn Ser Gly Ala Arg Arg Ser Ser Pro
165 170 175

Phe Gln Asp Lys Leu Gly Asp Gly Phe Ile Glu Glu Ala Phe Arg Thr
180 185 190

Ala Arg Thr Val Asp Pro Thr Ala Lys Leu Cys Tyr Asn Asp Tyr Asn
195 200 205

Thr Asp Gly Arg Asn Ala Lys Ser Asp Ala Val Tyr Ala Met Ala Lys
210 215 220

Asp Phe Lys Gln Arg Gly Val Pro Ile Asp Cys Val Gly Phe Gln Ser
225 230 235 240

His Phe Asn Ser Asn Ser Pro Val Pro Ser Asp Tyr Arg Ala Asn Leu
245 250 255

Gln Arg Phe Ala Asp Leu Gly Leu Asp Val Gln Ile Thr Glu Leu Asp
260 265 270

Ile Glu Gly Ser Gly Ser Ala Gln Ala Ala Asn Tyr Thr Ser Val Val
275 280 285

Asn Ala Cys Leu Ala Val Thr Arg Cys Thr Gly Leu Thr Val Trp Gly
290 295 300

Val Thr Asp Lys Tyr Ser Trp Arg Ser Ser Gly Thr Pro Leu Leu Phe
305 310 315 320

Asp Gly Asp Tyr Asn Lys Lys Pro Ala Tyr Asp Ala Val Leu Ala Ala
325 330 335

Leu Gly Gly Thr Pro Asp Gly Gly Asp Asp Gly Gly Asp Asn

-continued

340 345 350

Gly Gly Gly Asn Thr Gly Ser Cys Thr Ala Thr Tyr Thr Gln Thr Ala
 355 360 365

Thr Trp Asn Gly Gly Tyr Asn Gly Glu Val Thr Val Lys Ala Gly Ser
 370 375 380

Ser Gly Ile Thr Thr Trp Ser Val Pro Val Thr Val Pro Ser Ser Gln
 385 390 395 400

Gln Val Ser Ala Leu Trp Asn Gly Ala Pro Thr Trp Asn Ala Gly Asn
 405 410 415

Thr Val Met Thr Val Lys Pro Thr Tyr Asn Gly Thr Leu Ala Ala Gly
 420 425 430

Ala Ser Thr Ser Phe Gly Phe Thr Val Met Thr Asn Gly Asn Thr Ser
 435 440 445

Ala Pro Ala Val Gly Ala Cys Thr Ala Ser
 450 455

<210> SEQ_ID NO 6

<211> LENGTH: 562

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 6

Val Arg Thr Ala Ile Arg Thr Ala Arg Arg Pro Gln Pro Leu Ala Leu
 1 5 10 15

Leu Leu Arg Gly Leu Ala Ala Phe Leu Gly Leu Ala Leu Ala Gly Ala
 20 25 30

Leu Gly Pro Ala Thr Ala Arg Ala Ala Asp Leu Pro Gln Arg Ala Glu
 35 40 45

Ala Arg Ala Ala Gly Leu His Ile Ser Asp Gly Arg Leu Val Glu Gly
 50 55 60

Asn Gly Asn Asp Phe Val Met Arg Gly Ile Asn His Ala His Thr Trp
 65 70 75 80

Tyr Pro Gly Glu Thr Gln Ser Leu Ala Asp Ile Lys Ala Thr Gly Ala
 85 90 95

Asn Thr Val Arg Val Val Leu Ser Asp Gly Tyr Arg Trp Ser Glu Asn
 100 105 110

Ser Pro Glu Asp Val Ala Ser Ile Ile Ala Arg Cys Lys Ala Glu Arg
 115 120 125

Leu Ile Cys Val Leu Glu Val His Asp Thr Thr Gly Tyr Gly Glu Asp
 130 135 140

Ala Ala Ala Gly Thr Leu Asp His Ala Ala Asp Tyr Trp Ile Gly Leu
 145 150 155 160

Lys Asp Val Leu Asp Gly Glu Glu Asp Tyr Val Val Ile Asn Ile Gly
 165 170 175

Asn Glu Pro Trp Gly Asn Ala Asp Pro Ala Gly Trp Thr Ala Pro Thr
 180 185 190

Thr Ala Ala Ile Gln Lys Leu Arg Ala Ala Gly Phe Ala His Thr Ile
 195 200 205

Met Val Asp Ala Pro Asn Trp Gly Gln Asp Trp Glu Gly Val Met Arg
 210 215 220

Ala Asp Ala Arg Ser Val Tyr Asp Ala Asp Pro Thr Gly Asn Leu Ile
 225 230 235 240

Phe Ser Ile His Met Tyr Ser Val Tyr Asp Thr Ala Ala Lys Val Thr
 245 250 255

-continued

Asp Tyr Leu Asn Ala Phe Val Asp Ala Gly Leu Pro Leu Leu Ile Gly
260 265 270

Glu Phe Gly Gly Pro Ala Asp Gln Tyr Gly Asp Pro Asp Glu Asp Thr
275 280 285

Met Met Ala Thr Ala Glu Glu Leu Gly Leu Gly Tyr Leu Ala Trp Ser
290 295 300

Trp Ser Gly Asn Thr Asp Pro Val Leu Asp Leu Val Leu Asp Phe Asp
305 310 315 320

Pro Thr Arg Leu Ser Ser Trp Gly Glu Arg Val Leu His Gly Pro Asp
325 330 335

Gly Ile Thr Glu Thr Ser Arg Glu Ala Thr Val Phe Gly Gly Gln
340 345 350

Gly Gly Asp Thr Glu Ala Pro Thr Ala Pro Gly Thr Pro Thr Ala
355 360 365

Ser Gly Val Thr Ala Thr Ser Val Thr Leu Gly Trp Ser Ala Ala Thr
370 375 380

Asp Asp Val Gly Val Thr Ala Tyr Asp Val Val Arg Val Thr Gly Gly
385 390 395 400

Ser Glu Thr Lys Val Ala Ser Ser Ala Ala Thr Ser Val Thr Val Thr
405 410 415

Gly Leu Ser Ala Gly Thr Ala Tyr Ser Phe Ala Val Tyr Ala Arg Asp
420 425 430

Ala Ala Gly Asn Arg Ser Ala Arg Ser Gly Thr Val Ser Val Thr Thr
435 440 445

Asp Glu Gly Gly Ser Val Pro Gly Gly Ala Cys Ser Val Gly Tyr Arg
450 455 460

Val Ile Gly Glu Trp Pro Gly Gly Phe Gln Gly Glu Ile Thr Leu Arg
465 470 475 480

Asn Thr Gly Ala Ala Ala Val Asp Gly Trp Thr Leu Gly Phe Ala Phe
485 490 495

Ala Asp Gly Gln Thr Val Thr Asn Met Trp Gly Gly Thr Ala Thr Gln
500 505 510

Ser Gly Gly Ala Val Ser Val Thr Pro Ala Ser Tyr Thr Ser Thr Ile
515 520 525

Ala Ala Gly Gly Ser Val Thr Val Gly Phe Thr Gly Thr Leu Thr Gly
530 535 540

Ala Asn Ala Ala Pro Ala Ala Phe Thr Leu Asn Gly Ala Thr Cys Thr
545 550 555 560

Ala Ala

<210> SEQ ID NO 7

<211> LENGTH: 328

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 7

Met Ser Ile Thr Pro Arg Pro Ser Leu Arg Ala Met Val Thr Gly Leu
1 5 10 15

Ala Val Ala Ala Ser Ala Leu Ala Gly Gly Ala Val Thr Ala Ala Pro
20 25 30

Ala Arg Ala Ala Ala Cys Asn Gly Tyr Val Gly Leu Thr Phe Asp Asp
35 40 45

Gly Pro Ser Ala Ala Gln Thr Pro Ala Leu Leu Ser Ala Leu Lys Gln
50 55 60

-continued

Asn Gly Leu Arg Ala Thr Met Phe Asn Gln Gly Asn Tyr Ala Ala Ser
 65 70 75 80
 Asn Pro Ala Gln Val Lys Ala Gln Val Asp Ala Gly Met Trp Val Gly
 85 90 95
 Asn His Ser Tyr Ser His Pro His Leu Thr Gln Gln Ser Gln Ala Gln
 100 105 110
 Met Asp Ser Glu Ile Ser Arg Thr Gln Gln Ala Ile Ala Ala Gly Gly
 115 120 125
 Gly Gly Thr Pro Lys Leu Phe Arg Pro Pro Tyr Gly Glu Thr Asn Ala
 130 135 140
 Thr Leu Arg Ser Val Glu Ala Lys Tyr Gly Leu Thr Glu Val Ile Trp
 145 150 155 160
 Asp Val Asp Ser Gln Asp Trp Asn Gly Ala Ser Thr Asp Ala Ile Val
 165 170 175
 Gln Ala Val Ser Arg Leu Thr Ala Gly Gln Val Ile Leu Met His Glu
 180 185 190
 Trp Pro Ala Asn Thr Leu Ala Ala Ile Pro Arg Ile Ala Gln Thr Leu
 195 200 205
 Ser Ala Lys Gly Leu Cys Ser Gly Met Ile Ser Pro Gln Thr Gly Arg
 210 215 220
 Ala Val Ala Pro Asp Gly Gly Asn Gly Gly Gly Gly Gly Gly Gly
 225 230 235 240
 Gly Gly Cys Thr Ala Thr Leu Ser Ala Gly Glu Lys Trp Gly Asp Arg
 245 250 255
 Tyr Asn Leu Asn Val Ala Val Ser Gly Ser Ser Asn Trp Thr Val Thr
 260 265 270
 Met Asn Val Pro Ser Gly Glu Arg Val Met Thr Thr Trp Asn Val Ser
 275 280 285
 Ala Ser Tyr Pro Ser Ala Gln Val Leu Val Ala Lys Pro Asn Gly Ser
 290 295 300
 Gly Asn Asn Trp Gly Ala Thr Ile Gln Ala Asn Gly Asn Trp Thr Trp
 305 310 315 320
 Pro Thr Val Ser Cys Thr Thr Ser
 325

<210> SEQ ID NO 8
 <211> LENGTH: 335
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 8

Met Asn Pro Leu Val Tyr Thr Glu Arg Arg Arg Arg Gly Arg Leu Thr
 1 5 10 15
 Ser Leu Ala Gly Ser Val Cys Ala Leu Val Leu Ala Ala Ala Ala
 20 25 30
 Met Leu Leu Pro Gly Thr Ala Ser Ala Asp Thr Val Val Thr Thr Asn
 35 40 45
 Gln Thr Gly Asn Asn Asn Gly Tyr Tyr Tyr Ser Phe Trp Thr Asp Gly
 50 55 60
 Gly Gly Gln Val Ser Met Asn Leu Ala Ser Gly Gly Ser Tyr Ser Thr
 65 70 75 80
 Ser Trp Thr Asn Thr Gly Asn Phe Val Ala Gly Lys Gly Trp Ser Thr
 85 90 95
 Gly Gly Arg Lys Ser Val Thr Tyr Ser Gly Thr Phe Asn Pro Ser Gly
 100 105 110

-continued

Asn Ala Tyr Leu Thr Leu Tyr Gly Trp Ser Thr Asn Pro Leu Val Glu
 115 120 125
 Tyr Tyr Ile Val Asp Asn Trp Gly Thr Tyr Arg Pro Thr Gly Thr Phe
 130 135 140
 Lys Gly Thr Val Ser Ser Asp Gly Gly Thr Tyr Asp Ile Tyr Glu Thr
 145 150 155 160
 Thr Arg Thr Asn Ala Pro Ser Ile Glu Gly Thr Lys Thr Phe Lys Gln
 165 170 175
 Phe Trp Ser Val Arg Gln Ser Lys Arg Thr Gly Gly Thr Ile Thr Thr
 180 185 190
 Gly Asn His Phe Asp Ala Trp Ala Arg Asn Gly Met Asn Leu Gly Thr
 195 200 205
 Met Asn Tyr Met Ile Leu Ala Thr Glu Gly Tyr Gln Ser Ser Gly Ser
 210 215 220
 Ser Asn Ile Thr Val Ser Glu Gly Gly Ser Gly Gly Gly Asp Asn
 225 230 235 240
 Gly Gly Gly Gly Gly Gly Cys Thr Ala Thr Leu Ser Ala
 245 250 255
 Gly Glu Lys Trp Gly Asp Arg Tyr Asn Leu Asn Val Ala Val Ser Gly
 260 265 270
 Ser Ser Asn Trp Thr Val Thr Met Asn Val Pro Ser Ala Glu Lys Val
 275 280 285
 Leu Ser Thr Trp Asn Ile Ser Ala Ser Tyr Pro Ser Ser Gln Val Leu
 290 295 300
 Val Ala Lys Pro Asn Gly Ser Gly Asn Asn Trp Gly Ala Thr Ile Gln
 305 310 315 320
 Ala Asn Gly Asn Trp Thr Trp Pro Thr Val Ser Cys Thr Thr Ser
 325 330 335

<210> SEQ ID NO 9
 <211> LENGTH: 280
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 9

Met Ser Glu Arg Ala Ala Ser Pro Arg Thr His Arg Arg Arg Pro Gly
 1 5 10 15
 Arg Arg Arg Ile Ala Thr Ala Leu Thr Ala Ala Leu Gly Leu Thr Gly
 20 25 30
 Ala Ala Leu Ala Thr Gly Val Met Leu Gln Pro Ala Gly Ala Ala Thr
 35 40 45
 Thr Ala Ile Pro Ala Trp Pro Ser Ala Thr Gly Ser Gln Ser Val Ser
 50 55 60
 Lys Thr Ile Glu Val Ser Gly Thr Tyr Asp Gly Gly Leu Lys Arg Phe
 65 70 75 80
 Thr Gly Ser Gly Asp Leu Gly Asp Gly Gly Gln Asp Glu Gly Gln Asp
 85 90 95
 Pro Ile Phe Lys Leu Lys Asp Gly Ala Thr Ile Lys Asn Val Ile Leu
 100 105 110
 Gly Thr Pro Ala Ala Asp Gly Ile His Cys Ser Gly Ser Cys Thr Ile
 115 120 125
 Gln Asn Val Trp Trp Glu Asp Val Gly Glu Asp Ala Ala Ser Phe Lys
 130 135 140
 Gly Thr Ser Thr Ser Ser Val Tyr Thr Val Tyr Gly Gly Ala Lys

US 10,144,941 B2

77

-continued

78

145	150	155	160
Lys Ala Ser Asp Lys Val Phe Gln Phe Asn Gly Ala Gly Lys Leu Val			
165	170	175	
Val Thr Lys Phe Gln Val Ala Asp Phe Gly Lys Leu Val Arg Ser Cys			
180	185	190	
Gly Asn Cys Ser Lys Gln Tyr Lys Arg Glu Ile Ile Val Asn Asp Val			
195	200	205	
Asp Val Thr Ala Pro Gly Lys Ser Leu Val Gly Ile Asn Thr Asn Tyr			
210	215	220	
Gly Asp Thr Ala Ala Leu Arg Ser Val Arg Val His Gly Asp Ser Ser			
225	230	235	240
Lys Lys Ile Lys Pro Cys Val Arg Tyr Thr Gly Asn Ser Thr Gly Ala			
245	250	255	
Glu Pro Lys Glu Thr Gly Ser Gly Pro Asp Gly Thr Tyr Cys Lys Tyr			
260	265	270	
Thr Ala Ser Asp Leu Ser Tyr Asp			
275	280		

<210> SEQ_ID NO 10

<211> LENGTH: 909

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 10

Met Trp Cys His Pro Tyr Leu Arg Leu Arg Thr Ser Gly Arg Lys Val			
1	5	10	15
Ser Ser Val Asn Ala Leu Pro Pro Pro Ala Arg Pro Ala Pro Val Arg			
20	25	30	
Pro Arg Ser Arg Tyr Gly Arg Arg Val Leu Gly Met Ser Ala Ala Ala			
35	40	45	
Leu Leu Cys Ala Gly Ala Leu Ala Val Pro Gly Thr Ala Met Ala Asp			
50	55	60	
Asp Ala Glu Pro Gly Pro Gly Pro Glu Gln Ile Thr Asn Gly Asp Phe			
65	70	75	80
Ala Thr Gly Thr Ser Ala Pro Trp Trp Trp Thr Pro Asn Ala Ser Ala			
85	90	95	
Ala Val Ser Glu Gly Arg Leu Cys Val Glu Val Pro Ala Gly Thr Ala			
100	105	110	
Asn Ala Trp Asp Val Ile Val Gly Gln Asn Asp Val Pro Ile Val Ala			
115	120	125	
Gly Glu Ser Tyr Glu Leu Ser Tyr Thr Ala Arg Ser Thr Val Pro Leu			
130	135	140	
Thr Val Gln Thr Arg Val Gln Glu Ala Val Glu Pro Tyr Thr Thr Val			
145	150	155	160
Leu Ala Thr Ala Asp Pro Val Gly Ala Glu Asp Thr Arg Val Ala Arg			
165	170	175	
Thr Phe Thr Ala Ser Val Asp Gln Pro Ala Ala Ser Val Gln Leu Gln			
180	185	190	
Ile Gly Gly Gly Glu Arg Ala Thr Thr Phe Cys Leu Asp Asp Val Ser			
195	200	205	
Leu Arg Gly Gly Ala Glu Pro Pro Val Tyr Val Pro Asp Thr Gly Ser			
210	215	220	
Pro Val Arg Val Asn Gln Val Gly Tyr Leu Pro Arg Gly Pro Lys Ser			
225	230	235	240

US 10,144,941 B2

79

80

-continued

Gly Thr Val Val Thr Asp Ala Glu Ala Pro Leu Thr Trp Thr Val Lys
245 250 255

Ala Glu Asp Gly Ser Thr Ala Ala Thr Gly Thr Thr Val Pro Arg Gly
260 265 270

Glu Asp Pro Ser Ser Arg Arg Arg Val His Thr Phe Asp Phe Gly Asp
275 280 285

Leu Thr Thr Ala Gly Asp Gly Tyr Thr Val Glu Val Asp Gly Glu Val
290 295 300

Ser Glu Pro Phe Ser Ile Arg Gly Asp Leu Tyr Asp Ser Leu Arg Ser
305 310 315 320

Asp Ala Leu Ala Tyr Phe Tyr His Asn Arg Ser Gly Ile Glu Ile Asp
325 330 335

Ala Asp Leu Val Gly Glu Gln Tyr Ala Arg Pro Ala Gly His Ile Gly
340 345 350

Val Ala Pro Asn Lys Gly Asp Thr Asp Val Pro Cys Arg Pro Gly Val
355 360 365

Cys Asp Tyr Arg Leu Asp Val Ser Gly Gly Trp Tyr Asp Ala Gly Asp
370 375 380

His Gly Lys Tyr Val Val Asn Gly Gly Ile Ser Val Ala Gln Leu Met
385 390 395 400

Ala Thr Tyr Glu Arg Thr Leu Thr Ala Pro Asp Ala Glu Ser Ala Glu
405 410 415

Leu Gly Asp Gly Ala Leu Arg Val Pro Glu Arg Asp Asn Gly Val Pro
420 425 430

Asp Ile Leu Asp Glu Ala Arg Trp Glu Met Asp Phe Leu Ile Lys Met
435 440 445

Gln Val Pro Ala Gly Glu Gln Leu Ala Gly Met Val His His Lys Met
450 455 460

His Asp Ala Glu Trp Thr Gly Leu Pro Met Lys Pro His Leu Asp Pro
465 470 475 480

Gln Gln Arg Glu Leu His Pro Pro Ser Thr Ala Ala Thr Leu Asn Leu
485 490 495

Ala Ala Thr Ala Ala Gln Cys Ala Arg Leu Tyr Ala Pro Phe Asp Ala
500 505 510

Asp Phe Ala Asp Arg Cys Leu Arg Ala Ala Glu Thr Ala Trp Asp Ala
515 520 525

Ala Lys Arg His Pro Asp Val Leu Ala Asp Pro Asn Asp Gly Ile Gly
530 535 540

Gly Gly Ala Tyr Asn Asp Asp Asp Val Ser Asp Glu Phe Tyr Trp Ala
545 550 555 560

Ala Ala Glu Leu Phe Thr Thr Gly Lys Asp Ile Tyr Arg Gln Ala
565 570 575

Val Leu Ser Ser Ala Trp His Gly Asp Ala Gly Ala Val Phe Pro Ala
580 585 590

Gly Gly Ile Ser Trp Gly Ser Thr Ala Gly Leu Gly Val Leu Thr
595 600 605

Leu Ala Thr Val Pro Asn Ala Leu Thr Ser Asp Gln Leu Ala Gln Val
610 615 620

Arg Thr Val Val Thr Glu Gly Ala Asp Arg Tyr Ala Ala Gln Ser Arg
625 630 635 640

Glu Gln Ala Tyr Gly Leu Pro Tyr Ala Pro Arg Gly Glu Asp Tyr Val
645 650 655

Trp Gly Ser Asn Ser Gln Val Leu Asn Asn Met Val Val Leu Ala Thr

US 10,144,941 B2

81**82**

-continued

660	665	670
Ala His Asp Leu Thr Gly Asp Ala Ala Tyr Gln Asp Ala Val Leu Arg		
675	680	685
Gly Ala Asp Tyr Leu Leu Gly Arg Asn Pro Leu Asn Gln Ser Tyr Val		
690	695	700
Thr Gly Tyr Gly Glu Arg Asp Ser His Asn Gln His His Arg Phe Trp		
705	710	715
720		
Ala His Gln Asn Asp Pro Ser Leu Pro Asn Pro Ala Pro Gly Ser Ile		
725	730	735
Ala Gly Gly Pro Asn Leu Thr Ala Ile Ala Ser Gly Asp Pro Val Ala		
740	745	750
Ala Glu Lys Leu Ser Gly Cys Ala Pro Ala Met Cys Tyr Val Asp Asp		
755	760	765
Ile Gly Ser Trp Ala Thr Asn Glu Ile Thr Ile Asn Trp Asn Ala Pro		
770	775	780
Leu Ala Phe Ile Ala Ser Tyr Leu Asp Asp Ala Gly Glu Gly Gly Gln		
785	790	795
800		
Thr Ala Ala Ala Arg Thr Cys Gln Val Thr Tyr Ser Ser His Pro Trp		
805	810	815
Asn Ser Gly Ser Thr Val Thr Val Arg Val Glu Asn Thr Gly Ser Asp		
820	825	830
Pro Val Ser Pro Trp Ala Leu Thr Trp Leu Leu Pro Gly Glu Gln Arg		
835	840	845
Leu Ser His Thr Trp Ser Ala Glu Phe Asp Gln His Gly Arg Thr Val		
850	855	860
Ser Ala Arg Pro Leu Ser Trp Asn Arg Thr Leu Ala Pro Gly Ala Ala		
865	870	875
880		
Val Asp Phe Gly Phe Asn Thr Ser Ala Ala Gly Ser Ser Pro Glu Pro		
885	890	895
Gly Ala Phe Lys Leu Asn Gly Arg Ala Cys Ser Ala Gly		
900	905	

<210> SEQ ID NO 11

<211> LENGTH: 405

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 11

Met Arg Thr Gly Ser Ile Ala Arg Val Leu Gly Leu Ala Ala Ala Leu		
1	5	10
15		
Ala Ala Leu Leu Thr Thr Ala Phe Met Ala Pro Ala Met Ala Gly Lys		
20	25	30
His Asp Ala Thr Asp Ser Pro Ser Ala Ala Ala Pro Ala Ser Phe		
35	40	45
Thr His Pro Gly Val Leu Val Ser Arg Pro Gln Leu Asp Phe Val Arg		
50	55	60
Gly Lys Val Gln Ala Gly Ala Gln Pro Trp Lys Gly Ala Tyr Asp Gln		
65	70	75
80		
Met Leu Ala Ser Pro Tyr Ala Ser Leu Ser Arg Thr Ala Lys Pro Arg		
85	90	95
Ala Val Val Glu Cys Gly Ser Tyr Ser Asn Pro Asn Asn Gly Cys Thr		
100	105	110
Asp Glu Arg Glu Asp Ala Leu Ala Ala Tyr Thr Leu Ser Leu Ala Trp		
115	120	125

-continued

Tyr Ile Ser Gln Asp Gly Arg Tyr Ala Gln Lys Ala Ile Gln Ile Met
130 135 140

Asp Ala Trp Ser Gly Val Ile Lys Asp His Thr Asn Ser Asn Ala Pro
145 150 155 160

Leu Gln Thr Gly Trp Ala Gly Ser Ser Trp Pro Arg Ala Ala Glu Ile
165 170 175

Ile Lys Tyr Thr Tyr Gly Asn Trp Pro Ala Ser Gly Arg Phe Gly Thr
180 185 190

Met Leu Arg Asp Val Tyr Leu Pro Lys Val Ala Asn Gly Ser Asn Ser
195 200 205

Asn Gly Asn Trp Glu Leu Ser Met Thr Glu Ala Ala Ile Gly Ile Ala
210 215 220

Val Phe Leu Glu Asp Arg Gly Ala Tyr Asp Arg Ala Val Ala Lys Phe
225 230 235 240

Arg Gly Arg Val Pro Ala Tyr Ile Tyr Val Thr Ala Asp Gly Ser Leu
245 250 255

Pro Lys Ala Ala Pro Gly Ser Gly Leu Asp Thr Arg Glu Lys Ile Ile
260 265 270

Asn Tyr Trp Gln Gly Gln Ser Thr Phe Val Asp Gly Leu Ser Gln Glu
275 280 285

Thr Cys Arg Asp Leu Thr His Thr Gly Tyr Gly Leu Ser Ala Ile Ser
290 295 300

His Ile Ala Glu Thr Ser Arg Ile Gln Gly Gln Asp Leu Tyr Pro Glu
305 310 315 320

Val Ala Asp Arg Leu Arg His Ala Leu Gly Leu His Ala Lys Tyr Gln
325 330 335

Leu Gly Glu Lys Val Pro Ser Ser Leu Cys Gly Gly Ser Leu Lys Asp
340 345 350

Ser Leu Gly Pro Val Thr Glu Val Gly Phe Asn Ala Leu His Asn Arg
355 360 365

Met Gly Tyr Ala Met Thr Asn Thr Gln Thr Leu Thr Glu Arg Gln Arg
370 375 380

Pro Ala Ala Ser Asn Asn Leu Phe Val Ala Trp Glu Thr Leu Thr His
385 390 395 400

Ala Asp Asn Pro Asn
405

<210> SEQ ID NO 12

<211> LENGTH: 626

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 12

Met Pro Ser Arg Thr Thr Leu Ile Ala Thr Thr Ala Ala Leu Val Ala
1 5 10 15

Leu Ala Ala Pro Met Ala Phe Ala Ala Pro Ala Pro Ala Pro Asp Pro
20 25 30

Ala Val Glu Ala Ala Ala Ala Ala Trp Asp Thr Asp Arg Ala Ala Ser
35 40 45

Ala Tyr Ala Ala Asn Pro Ala Ala Val Thr Ala Ser Gly Ser Glu Asn
50 55 60

Pro Ala Ser Gly Pro Gly Ala Ala Thr Asp Gly Asp Ala Thr Thr Arg
65 70 75 80

Trp Ser Ser Asp Phe Ala Asp Asn Ala Trp Ile Arg Val Asp Leu Gly
85 90 95

-continued

Ser Thr Ile Arg Ile Asn Gln Val Lys Leu Glu Trp Glu Ala Ala Tyr
 100 105 110
 Gly Lys Lys Tyr Val Leu Glu Val Ser Lys Asp Gly Thr Asn Trp Thr
 115 120 125
 Pro Phe Tyr Thr Glu Asp Ala Gly Thr Gly Gly Thr Val Thr Ala His
 130 135 140
 Thr Tyr Pro Gln Glu Val Thr Gly Arg Tyr Val Arg Met Arg Gly Val
 145 150 155 160
 Glu Arg Ala Thr Ala Trp Gly Tyr Ser Leu Phe Ser Phe Gln Val Tyr
 165 170 175
 Gly Gly Glu Pro Ala Pro Ala Ser Thr Thr Arg Ser Asn Leu Ala Leu
 180 185 190
 Asn His Pro Ala Tyr Gly Asp Leu Tyr Gln His Ala Gly Asn Ser Pro
 195 200 205
 Ala Phe Val Thr Asp Gly Gly Trp Pro Ala Asp Leu Lys Ala Asp Arg
 210 215 220
 Ser Arg Trp Ser Ser Asp Trp Asn Ala Asp Arg Trp Val Gly Val Asp
 225 230 235 240
 Leu Gly Ala Thr Ser Thr Ile Asn Ser Val Asp Leu Tyr Trp Glu Ala
 245 250 255
 Ala Tyr Ala Val Asp Tyr Glu Ile Gln Val Ser Asp Asp Asn Arg Thr
 260 265 270
 Trp Arg Thr Val His Arg Pro Ser Ala Ala Glu Val Ala Ala Arg Arg
 275 280 285
 Ala Asp Val Lys Ala Pro Ala Glu Ala Val Gly His Asp Thr Ile
 290 295 300
 Asn Leu Pro Thr Pro Ala Thr Gly Arg Tyr Val Arg Met Leu Gly Lys
 305 310 315 320
 Glu Arg Arg Ser Phe Tyr Asn Pro Ala Pro Ser Thr Ala Gln Phe Gly
 325 330 335
 Tyr Ser Leu Tyr Glu Phe Gln Val Trp Gly Thr Gly Gly Ser Ala Asp
 340 345 350
 Ala Ala Tyr Pro Ala Leu Pro Lys Asn Pro Gly Gly Ala Tyr Arg Thr
 355 360 365
 Thr Phe Phe Asp Asp Phe Thr Gly Ser Gly Leu Asp Arg Ser Lys Trp
 370 375 380
 Arg Val Val Arg Thr Gly Thr Glu Met Gly Pro Val Asn Gly Glu Ser
 385 390 395 400
 Gln Ala Tyr Val Asp Ser Pro Asp Asn Ile Arg Thr Glu Asn Gly Ala
 405 410 415
 Leu Val Leu Glu Ser Lys Tyr Cys Lys Gly Cys Thr Pro Thr Pro Asn
 420 425 430
 Gly Thr Phe Asp Phe Thr Ser Gly Arg Val Asp Thr Asn Thr Lys Phe
 435 440 445
 Asp Phe Thr Tyr Gly Lys Val Ser Ala Arg Met Lys Leu Pro Val Gly
 450 455 460
 Asp Gly Phe Trp Pro Ala Phe Trp Leu Leu Gly Ser Asp Val Asp Asp
 465 470 475 480
 Pro Ala Val Ser Trp Pro Gly Ser Gly Glu Thr Asp Ile Met Glu Asn
 485 490 495
 Ile Gly Tyr Gly Asp Trp Thr Ser Ser Gly Leu His Gly Pro Gly Tyr
 500 505 510

US 10,144,941 B2

87

88

-continued

Ser Ala Asp Gly Asn Ile Gly Ala Ser Gln Thr Tyr Pro Asn Gly Gly
515 520 525

Arg Ala Asp Glu Trp His Thr Tyr Gly Val Glu Trp Thr Pro Glu Gly
530 535 540

Met Thr Phe Thr Val Asp Asp Arg Val Val Gln Gln Thr Ser Arg Gln
545 550 555 560

Lys Leu Glu Ser Thr Arg Gly Lys Trp Val Phe Asp His Asn Gln Tyr
565 570 575

Val Ile Leu Asn Leu Ala Leu Gly Gly Ala Tyr Pro Gly Gly Tyr Asn
580 585 590

Gln Val Thr Gln Pro Tyr Trp Gly Leu Pro Gln Ser Ser Val Asp Arg
595 600 605

Ile Ala Gln Gly Gly Ile Lys Ala Glu Ile Asp Trp Val Arg Val Glu
610 615 620

Gln Lys
625

<210> SEQ ID NO 13

<211> LENGTH: 408

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 13

Val Ile Ser Arg Arg Met Phe Leu Thr Gly Ala Ala Ala Ser Ala Thr
1 5 10 15

Ala Leu Thr Tyr Pro Leu Trp Gly Thr Ala Leu Ser Pro Arg Thr Ser
20 25 30

Ala Ala Ala Ala Thr Cys Glu Leu Ala Leu Glu Asn Arg Ser Leu Pro
35 40 45

Gly Thr Val His Ala Tyr Val Thr Gly His Glu Gln Gly Thr Asp Ser
50 55 60

Trp Val Leu Leu Arg Ala Asp Gly Ser Val Tyr Arg Pro Glu Ser Pro
65 70 75 80

Gly Ala Pro Gln Thr Pro Leu Pro Val Asp Cys Ala Ile Pro Leu Asn
85 90 95

Gly Ala Gly Ala Gly Pro Val Val Leu Thr Leu Pro Gln Met Tyr Gly
100 105 110

Ala Arg Val Tyr Phe Val Arg Asp Asp Lys Leu Asp Phe Tyr Leu Asn
115 120 125

Pro Gly Pro Ser Leu Val Glu Pro Ala Phe Ala Thr Pro Thr Asp Pro
130 135 140

Asn Tyr Gly Arg Thr Trp Ser Phe Cys Glu Phe Thr Phe Asn Pro Gln
145 150 155 160

Gln Leu Tyr Ala Asn Ile Ser Tyr Val Asp Leu Val Thr Ala Leu Pro
165 170 175

Ile Gly Leu Thr Leu Glu Gly Asp Ser Thr His Thr Val Ala Pro Leu
180 185 190

Pro Asp Gly Ala Val Gln Arg Ile Ala Asp Asp Leu Thr Ala Gln Ala
195 200 205

Ala Ala Asp Gly Gln Pro Trp Asp Lys Leu Val Thr Arg Gly Ser Asp
210 215 220

Gly Gln Val Leu Arg Val Val Ser Pro Gln Asn Leu Met Ala Pro Tyr
225 230 235 240

Phe Asp Arg Pro Asp Glu Met Pro Phe Arg Asp Leu Phe Ala Ala Gln
245 250 255

US 10,144,941 B2

89

90

-continued

Ile Asp Glu Val Trp Glu Lys Tyr Arg Ser Thr Asp Leu Arg Ile Asp
260 265 270

Leu Gln Gly Gly Arg Gly Thr Leu Ala Gly Arg Val Ser Gly Asp Thr
275 280 285

Leu Thr Phe Glu Gly Gly His Thr Phe Ser Lys Pro Thr Ser Lys Asp
290 295 300

Ile Phe Thr Cys Asn His Gly Pro Phe Thr Asn Asn Pro Ser Asp Ser
305 310 315 320

Asp Asp Lys Lys Ala Leu Leu Ala Arg Ile Ala Ala Gly Phe Asn Arg
325 330 335

Ser Ile Met Leu Ser His Pro Ser Gln Pro Asn Gly Thr Ser Val Ala
340 345 350

Asp Tyr Tyr Gln Asp Ala Val Thr Asn His Trp Ser Arg Val Val His
355 360 365

Ala Asn Ser Pro Ile Gly Tyr Ala Phe Pro Tyr Asp Asp Val Arg Pro
370 375 380

Asp Gly Glu Pro Asp Val Ser Gly Ala Ala Asn Asp Gly Asn Pro Arg
385 390 395 400

Arg Phe Thr Val Ser Val Gly Ser
405

<210> SEQ ID NO 14

<211> LENGTH: 289

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 14

Val Leu His Pro His Asn Arg Thr Ala Arg Arg Thr Thr Arg Leu Thr
1 5 10 15

Arg Thr Gly Gly Leu Ala Ala Ala Leu Gly Leu Ala Leu Met Ala
20 25 30

Leu Pro Val Thr Ala His Ala Gly Ala Pro Thr Gln Pro Ala Ala His
35 40 45

His Leu Glu Ala Ala Ala Thr Gly Leu Asp Asp Pro Ala Lys Lys Asp
50 55 60

Ile Ala Met Gln Leu Val Ser Ser Ala Glu Asn Ser Thr Leu Asp Trp
65 70 75 80

Lys Ala Gln Tyr Gly Tyr Ile Glu Asp Ile Gly Asp Gly Arg Gly Tyr
85 90 95

Thr Ala Gly Ile Ile Gly Phe Cys Ser Gly Thr Gly Asp Met Leu Ala
100 105 110

Leu Val Glu Arg Tyr Thr Asp Arg Ser Pro Gly Asn Val Leu Ala Ser
115 120 125

Tyr Leu Pro Ala Leu Arg Glu Val Asp Gly Thr Asp Ser His Asp Gly
130 135 140

Leu Asp Pro Gly Phe Pro Arg Asp Trp Ala Glu Ala Ala Lys Asp Pro
145 150 155 160

Val Phe Gln Gln Ala Gln Asn Asp Glu Arg Asp Arg Val Tyr Phe Asp
165 170 175

Pro Ala Val Arg Gln Ala Lys Asp Asp Gly Leu Gly Thr Leu Gly Gln
180 185 190

Phe Ala Tyr Tyr Asp Ala Ile Val Met His Gly Gly Gly Asp Ser
195 200 205

Thr Ser Phe Gly Ser Ile Arg Gln Arg Ala Leu Ala Glu Ala Glu Pro

-continued

210 215 220

Pro Ser Arg Gly Gly Asp Glu Val Ala Tyr Leu Asp Ala Phe Leu Asp
 225 230 235 240

Ala Arg Val Trp Ala Met Arg Gln Glu Glu Ala His Ser Asp Thr Ser
 245 250 255

Arg Val Asp Thr Ala Gln Arg Val Phe Leu Arg Asp Gly Asn Leu Asn
 260 265 270

Leu Asp Pro Pro Leu Asp Trp Gln Val Tyr Gly Asp Ser Phe His Ile
 275 280 285

Gly

<210> SEQ ID NO 15

<211> LENGTH: 790

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 15

Met Thr Pro Pro His Arg His Arg Leu Phe Arg Arg Ser Val Ser Ala
 1 5 10 15

Ser Leu Ser Leu Ala Leu Thr Ala Val Gly Thr Ala Ala Ala Val Val
 20 25 30

Leu Ala Gly Ala Pro Ala Ala Gln Ala Ala Ala Val Pro Ala Pro Ser
 35 40 45

Pro Val Gly Ile Ser Gly Arg Gly Ala Ala Val Pro Phe Thr Glu Gln
 50 55 60

Glu Ala Glu Tyr Ala Ala Thr Asn Gly Thr Leu Ile Gly Pro Asp Arg
 65 70 75 80

Arg Tyr Gly Ser Leu Pro Ser Glu Ala Ser Gly Arg Gln Ala Val Thr
 85 90 95

Leu Asp Ala Ala Gly Glu Tyr Val Glu Phe Thr Leu Thr Ala Pro Ala
 100 105 110

Asn Ala Met Thr Phe Arg Tyr Ser Leu Pro Asp Asn Ala Ala Gly Thr
 115 120 125

Gly Arg Asp Ala Ser Leu Asp Leu Arg Val Asn Gly Ser Val Leu Lys
 130 135 140

Ser Val Pro Val Thr Ser Lys Tyr Gly Trp Tyr Tyr Gly Gly Tyr Pro
 145 150 155 160

Phe Asn Asn Pro Gly Asp Thr Asn Pro His His Phe Tyr Asp Glu
 165 170 175

Thr Arg Thr Met Phe Gly Ser Thr Leu Pro Ala Gly Thr Lys Val Arg
 180 185 190

Leu Gln Val Ala Ser Thr Ala Gly Ser Pro Ser Phe Thr Val Asp Leu
 195 200 205

Ala Asp Phe Glu Gln Val Ala Ala Pro Val Gly Lys Pro Ser Gly Ala
 210 215 220

Leu Asp Val Val Ser Asp Phe Gly Ala Asp Pro Thr Gly Ala Ala Asp
 225 230 235 240

Ser Thr Ala Lys Ile Gln Ala Ala Val Asp Ala Gly Arg Thr Gln Gly
 245 250 255

Lys Val Val Tyr Ile Pro Gln Gly Thr Phe Gln Val Arg Asp His Ile
 260 265 270

Val Val Asp Gln Val Thr Leu Arg Gly Ala Gly Pro Trp Tyr Ser Val
 275 280 285

Leu Thr Gly Arg His Pro Thr Asp Arg Ser Lys Ala Val Gly Val Tyr

US 10,144,941 B2

93**94**

-continued

290	295	300
Gly Lys Tyr Ser Ala Gln Gly Gly Ser Arg Asn Val Thr Leu Lys Asp		
305	310	315
320		
Phe Ala Ile Ile Gly Asp Ile Gln Glu Arg Val Asp Asn Asp Gln Val		
325	330	335
Asn Ala Ile Gly Gly Ala Met Ser Asp Ser Val Val Asp Asn Val Trp		
340	345	350
Met Gln His Thr Lys Cys Gly Ala Trp Met Asp Gly Pro Met Asp Asn		
355	360	365
Phe Thr Ile Lys Asn Ser Arg Ile Leu Asp Gln Thr Ala Asp Gly Val		
370	375	380
Asn Phe His Tyr Gly Val Thr Asn Ser Thr Val Thr Asn Thr Phe Val		
385	390	395
400		
Arg Asn Thr Gly Asp Asp Gly Leu Ala Met Trp Ala Glu Asn Val Pro		
405	410	415
Asn Val Lys Asn Lys Phe Thr Phe Asn Thr Val Ile Leu Pro Ile Leu		
420	425	430
Ala Asn Asn Ile Val Thr Tyr Gly Gly Lys Asp Ile Thr Ile Ser Asp		
435	440	445
Asn Val Met Ala Asp Thr Ile Thr Asn Gly Gly Leu His Ile Ala		
450	455	460
Asn Arg Tyr Pro Gly Val Asn Ser Gly Gln Gly Thr Ala Val Ala Gly		
465	470	475
480		
Thr His Thr Ala Ala Arg Asn Thr Leu Ile Arg Thr Gly Asn Ser Asp		
485	490	495
Phe Asn Trp Asn Phe Gly Val Gly Ala Ile Trp Phe Ser Gly Leu Asn		
500	505	510
Glu Pro Ile Ser Asn Ala Thr Ile Asn Ile Thr Asp Ser Glu Val Leu		
515	520	525
Asp Ser Ser Tyr Ala Ala Ile His Leu Ile Glu Gly Ala Ser Asn Gly		
530	535	540
Leu His Phe Lys Asn Val Lys Ile Asp Gly Ala Gly Thr Tyr Ala Leu		
545	550	555
560		
Gln Ile Gln Ala Pro Gly Thr Ala Thr Phe Glu Asn Val Val Ala Thr		
565	570	575
His Ile Ala Gln Ser Asn Pro Ile His Asn Cys Val Gly Ser Gly Phe		
580	585	590
Gln Ile Thr Arg Gly Ser Gly Asn Ser Gly Trp Tyr Ala Asp Pro Pro		
595	600	605
Ala Cys Thr Gly Val Trp Pro Asp Pro Val Trp Thr Asn Gly Gly Val		
610	615	620
Pro Gly Gly Gly Pro Thr Asn Pro Thr Asp Pro Thr Asp Pro Thr		
625	630	635
640		
Asp Pro Thr Asp Pro Thr Asp Pro Pro Glu Glu Thr Gly Asn Leu Ala		
645	650	655
Arg Gly Arg Thr Val Thr Glu Thr Ser His Thr Asp Val Tyr Gly Ala		
660	665	670
Ala Asn Thr Val Asp Gly Asn Ala Asp Thr Tyr Trp Glu Ser Arg Asn		
675	680	685
Asn Ala Phe Pro Gln Ser Val Thr Val Asp Leu Gly Ala Ala Lys Ala		
690	695	700
Val Lys Arg Val Val Leu Lys Leu Pro Pro Ala Ala Ala Trp Ala Thr		
705	710	715
720		

-continued

Arg Thr Gln Thr Leu Ser Val Ser Gly Ser Thr Asp Asn Gly Thr Tyr
 725 730 735

Asn Ser Leu Lys Ala Ser Ala Gly Tyr Thr Phe Asn Pro Ser Ser Gly
 740 745 750

Asn Thr Ala Thr Val Ser Leu Pro Gly Thr Pro Val Arg Tyr Leu Arg
 755 760 765

Leu Thr Phe Thr Gln Asn Thr Gly Trp Pro Ala Ala Gln Leu Ser Glu
 770 775 780

Leu Glu Ala Tyr Thr Ser
 785 790

<210> SEQ ID NO 16
 <211> LENGTH: 514
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 16

Met Arg Arg Pro Val Ala Leu Arg Leu Ser Ala Ala Gly Ala Thr Leu
 1 5 10 15

Ala Leu Ala Ala Ala Thr Gly Ala Leu Met Ala Met Pro Glu Ala Ala
 20 25 30

Ser Ala Ala Thr Gly Gly Val Thr Gly Tyr Ala Thr Gln Asn Gly Gly
 35 40 45

Thr Thr Gly Gly Ala Gly Gly Gln Thr Val Arg Ala Thr Thr Gly Thr
 50 55 60

Ala Ile His Ala Ala Leu Cys Gly Arg Ala Ser Ser Ser Thr Pro Leu
 65 70 75 80

Thr Ile Gln Val Glu Gly Thr Ile Asn His Gly Asn Thr Asp Lys Val
 85 90 95

Ser Gly Ser Ser Cys Asn Thr Ala Ala Gly Val Ile Glu Leu Lys Gln
 100 105 110

Ile Ser Asn Val Thr Ile Val Gly Val Gly Gly Ala Val Phe Asp
 115 120 125

Gln Val Gly Ile His Val Arg Glu Ser Ser Asn Ile Ile Ile Gln Asn
 130 135 140

Val Thr Val Lys Asn Val Lys Lys Ser Gly Ser Pro Thr Ser Asn Gly
 145 150 155 160

Gly Asp Ala Ile Gly Met Glu Lys Asp Val Arg Asn Val Trp Val Asp
 165 170 175

His Thr Thr Leu Glu Ala Ser Gly Glu Ser Glu Gly Phe Asp Gly
 180 185 190

Leu Phe Asp Met Lys Ala Gly Thr Gln Tyr Val Thr Leu Ser Tyr Ser
 195 200 205

Ile Leu Arg Asn Ser Gly Arg Gly Leu Val Gly Ser Ser Glu Ser
 210 215 220

Asp Leu Ser Asn Gly Phe Ile Thr Tyr His His Asn Leu Tyr Glu Asn
 225 230 235 240

Ile Asp Ser Arg Ala Pro Leu Leu Arg Gly Gly Val Ala His Ile Tyr
 245 250 255

Asn Asn His Tyr Val Gly Leu Ser Lys Ser Gly Ile Asn Ser Arg Ala
 260 265 270

Gly Ala Arg Ala Lys Val Asp Asn Asn Tyr Phe Glu Asp Ser Lys Asp
 275 280 285

Val Leu Gly Thr Phe Tyr Thr Asp Ala Ala Gly Tyr Trp Gln Val Ser

US 10,144,941 B2

97**98**

-continued

290	295	300
Gly Asn Val Phe Asp Asn Val Thr Trp Ser Gly Arg Ser Ser Asp Asn		
305	310	315
320		
Asn Pro Ala Gly Pro Asp Pro Gln Ser Asn Thr Ser Val Ser Ile Pro		
325	330	335
Tyr Ala Tyr Thr Leu Asp Gly Ala Asn Cys Val Pro Ser Val Val Ser		
340	345	350
Arg Thr Ala Gly Ala Asn Thr Gly Leu Lys Val Ser Asp Gly Ser Cys		
355	360	365
Ser Pro Gln Thr Pro Asp Pro Thr Asp Pro Thr Pro Asp Pro Thr Pro		
370	375	380
Asp Pro Thr Asp Pro Thr Pro Pro Thr Gly Thr Asn Leu Ser Leu Gly		
385	390	395
400		
Ala Gly Ser Asp Gly Ser Ser Lys Ala Ser Gly Thr Ser Tyr Gly Asp		
405	410	415
Val Arg Asp Gly Asp Met Ser Thr Tyr Trp Ser Pro Ser Gly Ser Thr		
420	425	430
Gly Ser Val Ser Ile Lys Trp Ser Ser Ala Thr Thr Val Ser Lys Ile		
435	440	445
Asn Val Arg Glu Ala Ala Gly Ser Thr Gly Ser Ile Thr Ser Trp Lys		
450	455	460
Val Gly Asn Ala Asp Thr Gly Ala Val Leu Ala Ser Gly Ser Gly Ala		
465	470	475
480		
Gly Val Ile Thr Phe Pro Gln Thr Ser Leu Arg Lys Ile Thr Phe Glu		
485	490	495
Ile Thr Gly Ser Thr Gly Thr Pro Lys Val Ala Glu Phe Glu Thr Tyr		
500	505	510
Ala Gly		

<210> SEQ ID NO 17
<211> LENGTH: 1761
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 17

atgagccgca	cgagccgcac	caccctgccc	cgatccccaa	cagcactcat	ggcgccgggc	60
gccctcgatcg	ccgcagccgc	gggtccgcgc	gcagccgcgg	cacccttcgg	tgcacccgccc	120
cccgccggcg	ccggctgcac	cgtcgactac	aagatccaga	accagtggaa	ccggccggctc	180
accgccttcgg	tgagcgtcac	caacaacggg	gacgccatct	ccggctggca	gttccagttgg	240
agcttcgccc	gcggcgagca	ggtcagccag	gggttggaaacg	ccaccgtctc	tcagagccgc	300
tccggccgtca	ccgccaaggaa	cgccggctac	aacgcccggcc	tggccacccgg	ggcatcgccc	360
tccttcgggtt	tcaacgcgtac	ggcaacggc	aacagcgtcg	tcccgccgac	gttcaagctg	420
aacggcgtca	cctgcaacgg	cgccaccacgg	ggcccgaccgg	atcccacggaa	ccccacggac	480
ccgacggacc	cgaccgaccc	gccccggggc	aaccgtgtgg	acaaccccta	ccagggagcc	540
aagggtctatg	tgaacccgga	gtggtcggcg	aacgcccggc	cogagccggg	cgccgacaga	600
atcgccgacc	agccccacgg	cgtctggctg	gaccgcatacg	ccgcgatcga	gggcgcgaac	660
gttccatgg	gtctcgccga	ccatctcgac	gaggccctga	cgcagaaggg	ctccggcgaa	720
ctcgtcgatcc	aggcgtcat	ctacaacctg	cccgccggag	actgcgcggc	gctggcctcc	780
aacggtgagc	tcggaccgac	cgagatccgc	cgctacaaga	ccgagttacat	cgacccgatc	840

US 10,144,941 B2

99**100**

-continued

gccccatcc tcggcgaccc gaagtacgag ggctcgccca tcgtcaccac ggtcgagatc	900
gactcgctgc cgaaccttgt caccaacgcc ggccggccgc ccacggccac tccggccctgt	960
gacgtcatga aggccaacgg caactacgtc aaggcgctcg gctacgctgc caacaagctc	1020
ggcgacgcgc ccaacgtcta caactacatc gacgcccccc accacggctg gatcggtgg	1080
gacgacaact tcggcgccctc cgccggagatc ttccacgagg ccgcgcaccgc cgaggcgccg	1140
accgtcaacg acgtgcacgg ctcatcacc aacacccgcca actacagcgc gctgaaggag	1200
gagaacttct ccategacga cgccgtgaac ggacacgtcg tccggcagtc gaagtgggtc	1260
gactggaaacc gctacacggg cgagctgtcc ttccgcgcagg cttccgcacaa cgagctggtc	1320
tccgtcggtct tcaactccgg catcgccatg ctcatcgaca cctccgcacaa cggctggggc	1380
ggcgacgaaacc ggcccgagccg accggggccgc aacaccagccg tccgcacccata tggacccggc	1440
gggcgcgtacg accgcggcat ccacctgggc aactggtgca accaggcagg agcgggtctc	1500
ggcgaacggc cgcaggccgc ccccgagccg gggatcgacg cgtacgtctg gatgaagccc	1560
cggggggagt cccgacggttc cagctcgagg atcccgaaacg acgaggggcaa gggattcgac	1620
cggatgtgcg accccgaccta cacgggttaac gcccgttaaca acaacaacat gtcggggcg	1680
ctgggtggcg ccccggtctc cgggaagtgg ttctcgccca agttccagga gctcatgaag	1740
aacgcctacc cggcgctcta g	1761

<210> SEQ ID NO 18

<211> LENGTH: 2865

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 18

gtggccgccc tcggccatccc cttggaaatg accgcagccg ccggcacggg ggcccgaggcc	60
gccgcgcgtcg cgtcgacgtcg cgactacacg accagtgact gggatcgccc gttcaccacc	120
gaactcaccc tgaccaaccc gggctccgcg gcatcgacg gctggaccct gacgtacgac	180
tacgcggga accagcagct cacgagccgc tggagccgc cctggccca gtcaggcaag	240
accgtcagcg tgaagaacgc agcctggaaac ggtgcgtatcg ccgcgggtgc cgccgtcact	300
accggccgcg agttcaccta cagccggccg aacacccgac cgaccaccc tcgcgtcaac	360
ggcacggctc gcgcgggggc ccaccagccg ccgcgtccgc tcctcacctc cccggccgcg	420
ggcgccgtct tctccgcgg ggacccgggtt ccgcgtgggg cgaccggccg ggccggggac	480
ggggcgacga tcagcaaggt cgaggcttac gacgacacga ccctccctgg caccgacacc	540
acctccccgt acagctacga ggccgggcaa ctggccggccg gcagccactc cgtgtacgcc	600
aggggctacg acagectcgcc cgcctccgcg gattccccgc ccgcggccat caccgtcgcc	660
accggccccg cggtcgtcg ctcccccgat caactcgccg tccagcaggg caggtcgaaa	720
accttcgacg tctcgatgtc caccggcccc gggccggacg tcaccgtcac ggccggccgg	780
tccgcgggta acacccggct gagegtcacc ggccgggtcg ccctcacctt caccggccgc	840
aactggtcca cacccagaa ggtgaccgtc acggccgacg gctccggcac cggggccgcg	900
accttcacccg tcacggcccc cggccacggc aaggccgagg tcaccgtcac ccagctggcg	960
gccccgaaagg agtacgacgc cggttccctc gacccatcg ggaagatcac cgatcccg	1020
aacggctact tctcgccggaa gggatcccc taccactccg tccggacgtc gatcgatcg	1080
ggccccgacc acggggcacga gaccacctcg gaggcctaca gctaccgtat ctggctgcag	1140
cgatgtacg gcaagatcac cggcgactgg accaagttca acgggtcgat ggacaccatg	1200

US 10,144,941 B2

101

102

-continued

gagacgtaca tgatccccac ccacgcccac cagcccacga actccttcta cgacgcgtcc	1260
aagcccgcca octacgcgcc cgagcacacg accccgaaacg agtaccccgc ggtgtcgac	1320
ggctccgect cctccggctc cgaccggata gcggcagagc tgaagagcgc gtacggcacc	1380
gacgacatct acggcatgca ctggatccag gacgtcgaca acgtctacgg atacggcaac	1440
gcgcggcga cgtgcgcggc cggcccccacc caggccggtc cgtectacat caacaccc	1500
cagcggggct cgcaggagtc ggtctggag accgtcaccc acccgacctg cgacaacttc	1560
acgtacggcg gcgcacaacgg ctacactcgac ctgttacccg gggactcctc gtacgccaag	1620
cagtggaaat tcaccaacgc ccccgacgcc gacgccccgc cctgtcagggc cgcctactgg	1680
gcgcacgtct gggcgaagga gcaggggaag gcggggcgaag tccggcggacac cgtcgccaa	1740
gcggcgaaga tgggtacta cctgcgtac tccatgttc acaagtactt caagaagatc	1800
ggcgactgcg tcggcccgac cacctgccccg gccggctccg gcaaggacag cgcgcactac	1860
ctgatgtctt ggtactacgc ctggggcgcc gccaccgaca ctcggccgg ctggccctgg	1920
cggatcggtt ccacggggga taccagaacc cgatggggc ctacgcgtt	1980
agctccgtgg cgcacctcaa gcccaagtcg gccaccggag cgcaggactg ggcaagagc	2040
ctggaccggcc aactggactt ctaccagtcg ctccagtcg acgagggtgc catcgccggc	2100
ggtgtcgacca acagctggaa gggcagctac gcccagcccc cggccggcac gccgaccttc	2160
tacggcatgt actacgacga gaagccctgt taccacgacc cgccgtccaa ccagtggttc	2220
ggcttccagg cgtggtccat ggagcgcgtc gcccggactt accacggatc gggtgacgccc	2280
caggcgaagg cctgtctcgaa caagtgggtc gactggggcc tggccgagac gaccgtcaac	2340
ccggacggca cctatctgtat gcccctacc ctccagtggtt cggggcgcgc ggacacctgg	2400
aacgcctcgaa accccgggtgc caacgcccacg ctccacgtca cggcgcgcga ctacaccgac	2460
gacgtcgccg tggccggccgc gtaegcccg acactgacact actacgcccgc caagtccgg	2520
gacacggagg cgcggccac cggcgaggccg ctgctcgacg gcatgtggca gcaccacca	2580
gacgacggccg cggtggccgt gcccggagacc cgcggccact acaaccgggtt cgacgacccg	2640
gtcttaegtcc cccgtggctg gacggggcgc atgcccacg gtgacaccgt cgacgaggac	2700
tgcacgttcc tctccatccg ctcccttctac aaggacgacc cgaactggcc ccaggtgcag	2760
gcgtacctgg acgggggtgc cgcggccgtc ttacacccacc accgggttctg ggcgcaggcc	2820
gacatcgac tggccctggg ggcgtacgac gacccctctgg agtga	2865

<210> SEQ ID NO 19

<211> LENGTH: 1089

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 19

atggcttagac gcagcagact catctccctg gcagcgggtgc tggccaccct gctcgccggc	60
ctcgccctca ccgcactctg gccggggcaag gcggaggccgc acgggtgtcgc gatgaccc	120
ggatcgcgtt octatctgtt ccacgtcgac gcccgtccg gacccggccgc gctgaaccc	180
acgaacccgg cctgcccggc cgcgtcgacg cagagcggccg cgaacgcgtt gtacaactgg	240
tgcggccgtc tcgactccaa cgcggggccgc cgcggccgcg gatatgtgcc ggacggcagc	300
ctgtgcgttcc cccgtgtaccg ctccccgtac gacttctccg cctacaacgc cgcggccgc	360
gactggccccc ggacacatct gacccctccgtt ggcgtacgac aggtgcagta cagcaactgg	420

US 10,144,941 B2

103

104

-continued

gcccggccacc	ccgggtgactt	ccgggtctac	ctgacccaagc	cgggctgggc	acccacgtcc	480
gaactcgctt	gggacgaccc	tcaagtggta	cagaccgtaa	gcaacccggc	gcagcaggcc	540
ggggcgggca	ccaaacggcgg	gcactactac	tgggacctgg	cgctgecgtc	ggcccggtcc	600
ggtgacgcgc	tgtatgttcat	ccagtggtgt	cgttcggaca	gtcaggagaa	cttcttctcc	660
tgtctggaca	tcgtcttcga	cggcggcaac	ggcgaggtga	cgggaatcgg	cggcacggc	720
accccccaccc	ccactccgac	ccccgactccg	accccgaccc	cgacggaccc	ggageactcc	780
ggttcttgca	tggccgtcta	caacgtcgtc	agtccttggg	ccggtggctt	ccaggccctcc	840
gtcgagggtga	tgaaccacgg	tacggAACCG	cgcaacggct	ggggcgtgca	gtggaaagccc	900
ggttccggga	cgcagatcaa	cagcgtgtgg	aacggctccc	tctccaccgg	gtccgacggc	960
accgtgacgg	tgcgcgacgt	ggaccacaac	cgtgtcatcg	ccccggacgg	cagtgtgacc	1020
ttcgggttca	ccggcacctc	cacgggcaac	gactaccccg	ccgggacgat	cgggtgtgtg	1080
acgtccttag						1089

<210> SEQ ID NO 20

<211> LENGTH: 1371

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 20

gtgaaaacgct	ttctggcctt	actggccacc	tgcgcgacgg	tcctgggcct	cacggcactg	60
accggccccc	aggcggtggc	cggcgccggc	tgcacggccc	actacacgt	caccagccag	120
tggcaggggcg	gcttccaggc	cgcgggtgaag	gtcaccaacc	tgggaacccc	cgtgaccggg	180
tggaagctca	cgttcacccct	gccggacgac	ggacagaagg	tgcgtccagg	ctggaacgccc	240
gcctggcgc	agtcgggttc	cgcggtcacc	gccggccggc	ccgactggaa	cggcacactg	300
gccacccggcg	cgtcgccgc	ggcgggcttc	gtgggctct	tcacgggccc	caacccgcct	360
cccacggcgt	tgcgcgtcaa	cgggtgtcgc	tgtacgggt	ccacccggaga	accccccggcc	420
ggctccgacg	ggggaccccc	cgtggacgac	aacgggcage	tccacgtctg	cgggggtgaac	480
ctctgcaacc	agtacgaccg	gcccgtgcag	ctgggggtta	tgagcacgca	cggcatccag	540
tggttcgacg	cctgtacga	cggcgctcc	ctggacgacg	tggcgaacga	ctggaagtgc	600
gacctgtgc	gcatcgccat	gtacgtgcag	gaggacgggt	acgagaccga	cccgccgggc	660
ttcacccggc	gcgtgaacga	cctcgac	atggccgagg	cccgccggat	gtacgcgttgc	720
atcgacttcc	acacccgtac	cccgccggc	ccgaacgtca	acctcgaccc	cggcaagacg	780
ttcttcgcgt	ccgtcgccgc	gcgcaacgc	ggcaagaaga	acgtgatcta	cgagatcgcc	840
aacgagccca	acggcggtgac	ctggacggcc	gtcaagagct	acgcccggac	ggtcatcccg	900
gtgatccggg	ccgcccaccc	ggacggccgc	gtcatcgctg	gcacccgggg	ctggtcctcg	960
ctggggcgct	oggacggctc	cgacgagac	gagggtgtca	acagccccgt	caatggccacc	1020
acacatcatgt	acgcgttcca	cttctacga	gcgagccaca	aggacgcata	ccgctccacg	1080
ctgagccggg	cgccggccgc	gctccgc	ttcgttaccc	agttcggcac	ggtgagcgcc	1140
accggccggcg	gggcgtatgga	ccggccgcgc	accacggcc	ggctggaccc	gctcgaccag	1200
ctgaagatca	gctatgcgaa	ctggacctat	tccgacgcgc	ccgagagcag	cgcggcggtc	1260
cgccggggca	cctgccccgg	cgccggactac	agccggcagcc	gggtgtgtgac	cgagtccggg	1320
ggctgtgtca	agaaccggat	cagcaccccc	gattccctcc	ccacccggctg	a	1371

US 10,144,941 B2

105**106**

-continued

<210> SEQ ID NO 21
<211> LENGTH: 1377
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 21

atggccaaga aaatccccgc ccgtgccaga cgggcactct ccgtcctgac ggccggcggt	60
ctcgccgccc cggcgctgt ctgcgtcgcc ggcacggccg aggacgagg caccctgggt	120
gacgcggccg cggcgaaggg cgggtacttc ggcacccggg tcgccccgaa ccacccggc	180
gaggcaccgt acgcgtccac gctggacgac cagttcgact cggtaacccc ggagaacgag	240
atgaagtggg acgcggcgtca gggcagccgc aactccttca ctttcacggc cgccgaccag	300
atcgtagtc acgcccagag caaggaaatg aagggtgcgcg ggcacacccct ggtgtggcac	360
tgcgacgtgc ogggctgggt cggcgccctg ggccgcaccc acctccggc ggcgtatgaa	420
aaccacatca cccaggtgat gacgcactac aagggaaga tccattccgt ggacgtggtg	480
aacgaggccct tccaggacgg caacagcggt gccccggcga gctctccctt ccaggacaag	540
ctgggtgacg gtttcatcga ggaggcggtt cgcacccggcc gtacggcgtca tccgaccggc	600
aagctctgtt acaacgacta caaacccgac ggccggaaacg cgaagagcga cgcggcttac	660
ccatggcga aggacttcaa gcagcgcgtt gtggcgatcg actgcgtggg ttccagttcc	720
cacttcaaca gcaactcccc cgtggccctcc gactaccggg ccaatctcca gcgcttgc	780
gacctcggtc tcgaegtcca gatcaccgaa ctggacatcg agggttccgg ctggcccgag	840
gcccgcgaact acacgagcgt cgtgaacgcg tgccctggcc tgaccccgctg caccggcc	900
accgtctggg gtgttaccgaa caagtactcc tggcgacgca gggcacggc gctgttcc	960
gacggcgact acaacaagaa gcccgggtac gacgcgggtc tcgccccgct cggccggcacc	1020
cccgacgggt gcggtgacga cggggggggc gacaacggcg gggggaaacac cggcagctc	1080
acggcgacgt acacgcagac cggccacgtgg aacggcggtt acaacgggtga ggtgacggc	1140
aaggcaggct cctccggcat caccacctgg tcgggtggcc tgaccgtgcc ctgcgtcc	1200
cagggtctccg ccctctggaa cggggcccc acgtggaaacg cggcaacac cgtgtatgac	1260
gtgaagccca cctacaacgg gaccctggcg gccgggtgcct cgacgagctt cgggttcc	1320
gtcatgacga acggcaacac ctggcgcccc gccgtcgccg cctgcaccgc ctcc	1377

<210> SEQ ID NO 22
<211> LENGTH: 1689
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 22

gtgagaacag cgatacgac acgacgacga ccacagcccc tggcccttct gctgagagg	60
ctggccgcct tcctggggct cgcgcgtcgcc ggagccctcg gcccggccac cgccggcc	120
gccccacgtgc cccagggggc ggaggcgccgg gccggccggcc tccacatcg cgcggccgc	180
ctgggtgaaag gcaacggcaa cgacttcgtc atgcgcggca tcaaccacgc ccacac	240
tatccggccg agacccagtc cctcgccgac atcaaggcgaa cggcgccgaa cacggccgc	300
gtgggtctgt cccacggcta cccgtggac gagaacacgc cccggacgtt cgcctcgatc	360
atcgccccgtt gcaaggccga gccggctatc tgcgttctgg aggtccacga caccacccgg	420
tacggggagg acggccgcgc cggaaaccctc gaccacgcgg ccgactactg gatggcc	480
aaggacgtac tcgaacggcga ggaggactac gtcgtcatca acatcgccaa cgagcc	540

US 10,144,941 B2

107

108

-continued

ggcaacgccc atccggcgaa ctggaccgccc cccacgacgg ccgcgatcca gaagctgcgc 600
 gcccgggtt tcgcccacac gatcatggtg gacgcgcccc actggggcca ggactggag 660
 ggcgcatgc gggccgacgc ccggagcgtg tacgacgccc acccgaccgg caatctgatc 720
 ttctcgatcc acatgtacag cgtctacgac accgcccggaa aggtcaccga ctacctaacc 780
 gccttcgtcg acgccggact tccctgtc atcggcgagt tcggcgcccc cgccggacag 840
 tacgggacc cggacgagga cacgtatgt gecacccggc aggagttggg gctcggtac 900
 ctggcctggt cctggagcgg caacacggat cccgttctcg acctggctct cgacttcgac 960
 cccacccggc ttagctcgta gggcgagcgc gtccctccacg gccccgacgg catcaccgg 1020
 acgtcccgta agggcacggt cttcgccggc gggcaggcgg ggggcgacac cgaggcccg 1080
 accgcacccggc gcaccccgac ggccctccggg gtacggcggc cttccgtcac cttcggtgg 1140
 agtgccgcca cggacgacgt cggcgtcacc gcttacgacg tggtccggcgt gaccggccgc 1200
 tccgagacga aggtcgccctc ctccgccggc acctcggtca ccgtgaccgg tctgagcgcc 1260
 ggaccccgct acagttcgcc cgtctacgccc cgggacgccc ccggcaacccg ttccggcgcc 1320
 tccggcaccggc tgtcggtcac caccgacgag ggcggcagggc tgccgggggg cgcctgtcc 1380
 gtgggctacc gggtgatcggt cgagtggcgg ggcggcttcc agggggagat caccctccgg 1440
 aacacccggc cccggccgtt cggacggctgg acgtgggttcc tcggccttcgc cgacggggcag 1500
 accgtcaca gcatgtgggg cggcaccggc acgcagagcg ggggcgcccgtt gagcgtcacc 1560
 cccggctcgta acacctccac gatcgccggc gggggctcggt tcaccgtgg cttcaccggc 1620
 accctgactg gcgcaacgc cggccggcggc gccttacgc tcaacggcgc cacctgcacc 1680
 cgccctgaa 1689

<210> SEQ ID NO 23

<211> LENGTH: 987

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 23

atgagcatca caccgggtcc tccctgcgc gcatggtca ccgggttcgc cgtcgccgg 60
 tccggccctgg cggccggcgc cgttacccgc gacccggccc gggccggccgc ttgcaacggc 120
 tacgtcgccggc tcacccgtca cggacggacgg tcggcgcccc agacccggc cctgtgtcc 180
 gcgctcaagc agaacggcct gggggccacc atgttcaacc agggcaacta cgccgcctcc 240
 aacccggccc aggtcaaggc ccagggtcgac gccggcatgt gggtcggcaa ccacagctac 300
 agccacccgc acctgaccca gcagagccag ggcgcagatgg actccgagat ctccggacc 360
 cagcaggcaca tcggccggcgg agggggggcgc acaccggaaac tggccggccc gccgtacggc 420
 gagaccaacg ccacgtcgat gtcgggtcgag gcaaggatcg gtttcaccga ggtcatctgg 480
 gacgtcgact cggcggactg gaacggccgcg agcaccggacgc cgatcggtca ggcgggttcc 540
 cggcttacccg cccggctaggat cttccgtatg caccgtggc ccggcaacac cctcgccgg 600
 atcccgccca tcggcccgac cctgtccggc aagggttgtt gttccggcat gatctcccg 660
 cagacccggcc gcgccgtcgcc tcccgacggc ggcggcaacg gtggaggggg cgggtggcggt 720
 ggcgggtgca cccgcacgtt gtcggcggtt gagaagggtt gttccggat caacctgaac 780
 gtggcggtgaa gggggccatg caactggacg gtcggatgatc acgtgcggc gggcgagagg 840
 gtcatgacga cctggaaacgt cggcgcgatg tatccgacggc cggcggatctt ggtcgccaaag 900
 ccgaacggga gcgaaaacaa ctgggggtcgac acgtccagg ccaacggcaa ctggaccctgg 960

US 10,144,941 B2

109**110**

-continued

ccgaccgtct cctgcaccac gagctga	987
-------------------------------	-----

<210> SEQ ID NO 24
<211> LENGTH: 1008
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 24

atgaaccac tcgtgtacac ggagcgccgc agacgcggcc ggctcaccc gctggccggc	60
agcgttgcg ccctggtaact ggcgcgcggc gcccgcgtgc tgctgccccg cacggccagt	120
gcccacacgg tcgtcacgac gaaccagacc ggcaacaaca acggctacta ctactcggttc	180
tggaccgacg gggggggcca ggttccatg aacctggcct cggcgccag ctacagcacc	240
tctgtggacga acacccggcaa ctgcgtcgcc ggcaaggcggt ggagcacggg cggccgtaa	300
agcgtcacct actcgggcac cttcaaccccg tccggcaacg cctacctgac gctgtacgga	360
tggtcgacga accccgtcggt cgagtactac atcgtggaca actggggcac ctacggccc	420
accggtaacgt tcaagggcac ggttccacgc gacggcgccca cgtacgacat ctacgagacc	480
acccgcacca acggcccccgc categagggt acgaagaccc tcaagcagtt ctggagcgtc	540
cggcagtcga agcgggaccgg cggcaccatc accacccggca accacttgcg cgcctggcc	600
cgcaacggca tgaacctcggt caccatgaac tacatgatcc tggccaccga gggctaccag	660
agcagcggca gctccaacat cacggtgacg gagggcggtt cgggtgggtgg cggcgacaac	720
ggtgtgggggg gcggtggcggt tggtgggtgc accggccacgt tgctggcggt tgagaagtgg	780
ggtgaccgggt acaacctgaa cgtggcggtg agcggctcca gcaactggac ggtgacgatg	840
aacgtggcggtt cggcgagaaa ggtgtgtcg acctggaaaca tcaagcgcgag ttatccgagc	900
tcccagggtcc tggtcgcca gccgaacggg agcgggaaca actgggggtgc gacgatccag	960
gccaacggca actggacgtg gcccaccgtc tcctgcacca cgagctga	1008

<210> SEQ ID NO 25
<211> LENGTH: 843
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 25

atgagtgaaa gagccgcattt cccacgtacc caccggcgcc gccccggcccg ccggcgcatc	60
gccaccggcgc tgacggcgcc actggggcctc accggcgccgg cactggccac cggcggtatg	120
ctccagccgg cccggcgccgc caccacccggc atccccgcct ggccctccgc cacgggcagc	180
cagtccgtct cgaagaccat cgaggcttcc gggacgtacg acggcggtct gaagcgcttc	240
accggcagcg gtgacctggg cgacgggtgc caggacgagg gccaggaccc gatcttcaag	300
ctgaaggacg gggcgacgtt caagaacgtc atcctggca cttccggccgc cgacggcatc	360
cactgtcccg gcagctgcac gatccagaac gtctgggtgg aggacgtcggtt cgaggacgcc	420
gcgtccttca agggcacctc cacgtcgccgtt gtgtacacgg tgtaacggcgcc gggcgcaag	480
aggcctccg acaaggcttt ccgttcaac ggccggccca agctgggtgtt gacgaaatcc	540
cagggtcgccg acttcggcaaa gctggtccgc tcgtgcggca actgtccaa gcaactaa	600
cgcgagatca tcgtcaacga cgtcgacgtc acggcgccgg gcaagtccct ggtcgccatc	660
aacaccaact acggggacac cggcgccgtg cgctcggtgc gcttccacgg cgacagcagc	720
aagaagatca agccctgcgtt ccgttacacc ggcaacagca cggcgccggaa accgaaggag	780

-continued

acgggcagcg	gtccggacgg	cacgtactgc	aagtacaccg	cctcggacct	gagctacgac	840
tag						843
<210> SEQ ID NO 26						
<211> LENGTH: 2730						
<212> TYPE: DNA						
<213> ORGANISM: Streptomyces sp. ACTE						
<400> SEQUENCE: 26						
atgtggtgtc	acccgtaccc	ccgcctccgc	acgtccggac	gaaaggtttc	ctcggtgaac	60
gcccttccac	ccccggcccg	gcccgcaccc	gtccgaccac	ggtcccggtt	cgggcgccgc	120
gtgctcggtt	tgctggccgc	cgcctgtgt	tgcgcaagggg	ccctggccgt	gcccgtacg	180
gcatatggccg	acgacgcccga	acccggaccc	ggcccccggac	agatcaccaa	cgccgacttc	240
gccacccggta	cctcagcccc	gtggtgggtgg	acgcccgaacg	cctcgccgc	cgtgtccgag	300
ggccggctct	gcgtggaggt	gcccggccgc	acggccaacg	cctggacgt	catacgtcggc	360
cagaacgacg	taccgtacgt	cgcggggcgg	agctacgacg	tgtcctacac	ggcgcttcg	420
accgtgtcccc	tgaccgttca	gaccggggtc	caggaggcgg	tggagcccta	cacgacgggt	480
ctggcgacgg	cggtatccgg	gggcgcggag	gacacgcggg	tgcggccac	gttcacggcc	540
tcgggtggacc	agcccgccgc	gtcggtgcag	ttgcagatcg	gtggcgggg	gcggggcag	600
acgttctgc	tggacgacgt	gtcgctgcgg	ggcggggccgt	agccggccgt	gtacgtacccg	660
gacacccggct	cgcgggtccg	cgtcaaccag	gtcggtatc	tgcggccgg	tcccaagagc	720
ggcacccgtgg	tcacccgtac	cgaggcgcgg	ctgacccgtt	cggtcaaa	cgaggacgg	780
tcgacggccg	ccacccgtac	gaccgttccg	cgagggtgagg	accccagctc	gcgcgcacgg	840
gtccacaccc	tgcacttcgg	cgcacccatcc	acggcggggg	acggctacac	cgtggaggt	900
gacggtgagg	tgagcgagcc	gttctcgatc	cgcggggacc	tgtacgactc	cctgcgtc	960
gacgcgttgg	cgtacttcata	ccacaaccgc	acggccatcg	agatcgacgc	ggacccgtc	1020
ggtgagcagt	acgcgcgc	ggccgggtac	atcgccgtcg	cgcccaacaa	gggcgcacacg	1080
gacgtggcgt	gcccacccgtt	ggtctcgac	taccggctgg	acgtgtcg	cggtgttac	1140
gacgcggccg	accacggcaa	gtacgtggtc	aacggcgaaa	tctcggtgg	ccagctgtat	1200
gcccacgtac	acggccaccc	caccggcccg	gacgcggag	cggccgagct	cggcgcacggc	1260
gacgtggcgg	tgcccgagcg	cgacaacggg	gtgcggaca	tctggacga	ggcgccgttgg	1320
gagatggact	tcctcatcaa	gatcgaggtc	ccggcgccgg	agcagctggc	ggggatggtc	1380
caccacaaga	tgcacgtacgc	cgagtggtacc	gggctggcga	tgaagccgca	cctggacccg	1440
cagcagcgcg	agctgcaccc	gcccgtcgac	gcccacac	tcaacccgtc	cgccacggcc	1500
gcccagtgcg	cccggtctta	cgcgccttc	gacgcggact	tgcggacccg	ctgcgtcg	1560
gcccggaga	ccgcgtggga	cgcggcgaag	cgccacccgg	acgtgtcg	cgacccgaac	1620
gacggcatcg	gcggcggtgc	gtacaacgac	gacgacgtct	cgacggagg	ctactggcg	1680
gcccggcgc	tcttcaccac	gacggcaag	gacatctacc	ggcaggcggt	gtctccctcc	1740
gcatggcactg	gtgacgcggg	cgcggcttc	ccggcgccgg	gcggaatctc	ctggggctcc	1800
acggccggac	tcggcggtct	caccctggcc	accgtgccc	acgcctgac	gtccgatcg	1860
ctcgcccaagg	tgcgacacgg	ggtgacccgg	ggcgccgacc	gotacgcgc	cgagtcccg	1920
gagcaggcgt	acgggctgcc	gtacgcgc	cgggggggagg	actacgtctg	gggtccaa	1980
agtcagggtgc	tcaacaacat	ggtcgtctgt	gcccacccccc	acgacccgtac	cggtgacgccc	2040

-continued

gcctaccagg acgcccgtgct	gccccggcgcc	gactatctgc	tggggcgc当地	cccgcgtgaac	2100
cagtcgtacg tcacccggcta	cggcgagcg	gactcgcaca	accagcacca	ccgcttctgg	2160
gcccaccaga acgacccca	cctgccc当地	ccggcgccc当地	gttcgatcgc	ggggcgccc当地	2220
aacctccacgg cgatcgcctc	cggtgaccgg	gtggcgccgg	agaagctcag	cggctgc当地	2280
ccggccatgt gctacgtcga	cgacatcggc	tcctggc当地	ccaacgagat	caccatcaac	2340
tggAACGCAc	cgatcgc当地	catcgcc当地	tacctggacg	acggcgccg	2400
acggccgc当地	ccggcacctg	ccaggtc当地	tactcctc当地	acccgtggaa	2460
acgggtacgg tacgggtcga	gaacacccgc	tcggatccc当地	tctcgeccctg	ggcgctgacc	2520
tggctgtcc	ccggcgagca	ggggctgagc	cacacgtgga	gcccggagtt	2580
ggccgtacgg tca	ccggccccc当地	ccggctgtcg	tggaaacccg	ccctggacc	2640
gtcgacttcg	gcttcaacac	ctcgccggcg	ggctcctc当地	ccgagccogg	2700
ctgaacggcc	gggcctgctc	agcgggctg	ta		2730

<210> SEQ ID NO 27
<211> LENGTH: 1218
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 27

atgcgtaccc	gatccatcgc	gcgcgtcctg	ggcctcgccg	ccgcctggc	cgcactgctc	60
accacggcc	tcatggccccc	ggccatggcc	ggcaaaca	acgc当地	ccgc当地	120
gccgc当地	ccccggcg	cttacccac	ccggcg	tggc当地	gccg	180
gacttcgtac	gcggcaaggt	ccaggc当地	ccaggc当地	ggaagggg	gtacgaccag	240
atgctggcc	gtccctacgc	ctcgctctcg	cgaccgc当地	agcc	ccgc当地	300
tgccgc当地	actccaaccc	caacaacggc	tgcaccgc当地	agcgc当地	cgcg	360
gcttgc当地	tctcgctggc	ctggtacatc	agccaggac	gccgctacgc	ccagaagg	420
atccagatc	tggacgc当地	gtcggg	atcaaggacc	acaccaac	caacgccc当地	480
ctgc当地	gttggccgg	ctcc	ccggccgg	ccgagatcat	caagtacacg	540
tacggcaact	ggccggcg	cgccgc当地	ggcaccatgc	tgcgtacgt	ctac	600
aagg	ccgc当地	actggaa	actggaa	tctccatgac	cgaggccg	660
atcg	cggttctc	ggaggac	ggcgctac	acagg	cccaagttc	720
cgccgc当地	tccccc当地	catatacgt	accgc当地	gatcg	gtccgc当地	780
ccggc当地	gtctcg	acac	gccc当地	atcatcaact	actggcagg	840
ttcgtggac	ggctctcg	ggac	ccgc当地	cccacccg	ctacgg	900
tccgc当地	cccacatcgc	cgagacc	cgatcc	agg	ctacccg	960
gtcgccgacc	ggctccgt	ca	cg	gttgc当地	agtacc	1020
gtccgc当地	ccctgtcg	cg	gg	ccgttcc	gggg	1080
ggcttcaac	ccctgc当地	ccgc当地	ccgc当地	atgc当地	gaga	1140
gagccgc当地	ggccgc当地	ctcg	aa	ccgc当地	ccgtgac	1200
ccgc当地	acaacc	cgactg	aa	ccgc当地	ccgtgac	1218

<210> SEQ ID NO 28
<211> LENGTH: 1881
<212> TYPE: DNA

-continued

<213> ORGANISM: *Streptomyces* sp. ACTE

<400> SEQUENCE: 28

atgcctcc	gtacgacgtt	gategccacc	accggggccc	tggtcgcct	cggcgc	cccc	60
atggcattcg	cggtcccg	ccccggcccc	gaccccgccc	tgcaggccgc	cggcg	cgcc	120
tgggacacccg	accggcg	gtccgcctac	ggggcgaacc	ccggccgcgt	caccgcgtcc		180
ggcagegaga	accccgctc	cggaccgggc	gcccaccc	acggcgacgc	caccacccgc		240
tggtccagcg	acttcgcga	caacgcctgg	atacgcgtcg	acctcggtc	caccatccgg		300
atcaaccagg	tgaagcttga	gtgggaggcc	gcctacggca	agaagtacgt	ccttggaa	gtc	360
tccaaggacg	gcaccaactg	gaccccttc	tacacggagg	acgcgggcac	cggcg	ggcacc	420
gtcaccggcc	acacctaccc	cgaggagg	acccggccgt	acgtgcggat	gcgcgg	cggtc	480
gaacgcgcca	cggtctgggg	ctactccctc	tttctccttcc	aggctacgg	gggcg	agccgg	540
gccccggcct	cgaccacccg	cagcaaccc	gcccctaacc	accccgccct	cggcg	acactc	600
taccagcacg	cgggcaactc	gcccgcattc	gtcaccgcacg	ggggctggcc	cggcg	acactg	660
aaggcggacc	gtccccgtg	gtcctccgac	tggAACGCG	accgcgtgggt	cggcgtc	gac	720
ctcggcgcga	cctccacca	caacagcg	tcacgcgtc	gacctctact	gggaggccgc	ctacgcgtc	780
gactacgaga	tccagggtgc	cgacgacaac	cggaacctggc	ggaccgtcca	ccggcc	ctcc	840
gcccggcagg	tcggcccg	acgcgcgcac	gtcaaggccc	cgcccgaggc	cgtcg	gacgc	900
cacgacacca	tcaacctg	ccacccggcc	accggccgt	acgtccggat	gttggca	aa	960
gagcgcgtt	ccttctacaa	cccgccaccc	tccaccgccc	agttcggtct	ctcgct	tac	1020
gagttccagg	tgtggggc	ccggccgc	gcccgc	cctacccgc	cctgccc	aa	1080
aaccccgccg	gcccgc	caccaccc	tgcacgcact	tcacccgc	cggc	ctgg	1140
cgcgtca	gtgcgttgtt	gcccgcgt	acggagatgg	gcccggtaa	cgggg	agg	1200
caggcctacg	tgcactcg	ggacaacatc	cgtaccgaga	acggcgcc	gttctgg	gag	1260
tccaagtact	gcaagggt	cacccca	cccaacggca	cttcgactt	cac	tcgg	1320
cgcgtcaca	ccaacacaa	gttcgact	acca	tcacggc	cgttat	gaa	1380
c	ccccgg	gtggcgg	tttctgg	tggcagc	cgtcg	ac	1440
ccggcgtct	cctggcc	ctccggc	acggacat	tggagaacat	cggctac	ggc	1500
gactggacca	gtccggc	gcacggac	ggactact	cagacggca	catcg	cgcc	1560
tcccagac	acccgaa	cgccgg	gacgagt	ccgg	acac	ctcg	1620
accccgaa	gcatgac	caccgt	gaccgc	tgcagc	actt	ccgc	1680
aagctggagt	ccacccg	caagtgg	tgcacc	accagtac	gttct	caac	1740
ctggccctcg	gcccgc	cccg	tacaacc	tcaccc	ctact	gggc	1800
cttccgc	caacgcgt	ccgc	tcacgg	tcacgg	gatcg	acttg	1860
gtacgggtcg	agcaga	agta	a				1881

<210> SEQ ID NO 29

<211> LENGTH: 1227

<212> TYPE: DNA

<213> ORGANISM: *Streptomyces* sp. ACTE

<400> SEQUENCE: 29

gtgatttcgc	gcagaatgtt	cctgaccggc	gcccgcgt	ccgcgaccgc	gttcac	ctat	60
cgcgtctggg	gcacccgcct	gagcccgcgc	acgtcgccgc	cgcccgccac	gtgcg	actg	120

-continued

gcccctcgaga accgttcggtt gcccggtacg gtgcacgcct acgtcaccgg tcacgagcag 180
 ggcaaccgaca gctgggtgct gctggggcc gacggcagcg tgtaccgc cggatcgccg 240
 ggcgcctccgc agacccctct gccgggtggac tgcgcctatcc cgctgaacgg cgccggcgcc 300
 ggcccggtcg tcctgacgct gccccagatg tacggcgcgcg gggctactt cgtccgtgac 360
 gacaagctgg acttctacctt gaacccgggc ccctcgctgg tgcgatcgcc cttcgcgacg 420
 cccacccgacc cgaactacgg ggcacccctgg tgcgtctgcg agttcacctt caaccccgacg 480
 cagctgtacg cgaacatcag ctacgtcgac ctggtaaccg ccctgcgat cggcctgacc 540
 ctggaggcg actccaccca caccgtcgcc cgcgtcccg gacggcgcgt gcagcgatc 600
 gccgacgacc tgacggccca ggcggccgcg gacggcagc cgtggacaa gctggtcacc 660
 cgtggctcgg acggccaggt gctggggtc gtctcgccg agaacctgtat ggcgcgtac 720
 ttcgaccggc ccgacgagat gccgttcgg gacctgttcg cggcccgat cgacgaggc 780
 tgggagaagt accgctccac cgcacgtcgat atcgacccatc agggcggccg gggcacccctg 840
 gggggccggg tcaacggggaa cacgctgacc ttcaacggcg gacacaccc ttccaagccc 900
 acctcgaagg acatcttcac ctgcacccac ggtccgttca cgaacaaccc gagegactcg 960
 gacgacaaga aggcgtgtt ggcacggatc gggccgggtc tcaacccgtc gatcatgtc 1020
 agccacccca gccagccaa cggcacctcg gtggcggact actaccagga cgcggcgacc 1080
 aaccacttgtt cgcgggtcgat ccacgcgaaat tccccatcg ggtacgcgtt cccgtacgac 1140
 gacgtacgcc cgcacggta gccggacgtc tccggcgcgg cgaacgacgg caaccccg 1200
 cgcttcacgg tgacgtggg ttccgtga 1227

<210> SEQ ID NO 30

<211> LENGTH: 870

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 30

gtgcgttcacc cccacaaccc cacggcaacgt cgcaccactc ggctcaccccg caccggcggt 60
 ctgcgcgcg cggccctcggt gctcgcgctc atggcgctcc ccgtcacccgc tcacgcccgc 120
 gccccccacgc agccggccgc tcatacatcg gaggccgcgc cgacccggact ggacgatccc 180
 gcaagaagg acatcgccat gcagttggtc tccagcgcgg agaactccac gctggactgg 240
 aaggcgcagt acggctacat cgaggacatc ggacgcggac ggcgcgtacac cggccggatc 300
 atcggcttct gctccggac cggagacatg ctgcgcctgg tgcgatcgatc cacggaccgc 360
 tcaccggcga acgtactggc gtgcgtacatcg cccgcctgc gcgaggtcga cgggaccgc 420
 tccgcacgacg ggctcgaccc cggctccccc cggactggg cggaggccgc gaaggacccg 480
 gtgttccagc aggccgcagaa cgacgcgcgg gacccgggtt acttcgaccc ggcggcgcc 540
 caggccaaagg acgacgggtt ggggacgcgc ggcgcgttccg cgtactacga cgcgcgttc 600
 atgcacggag gggggggggaa cagcacgacg ttccgggttca tccggcagcg cgcgcgtcg 660
 gaggcggaaac cggccctcggt gggcggtgac gaggtcgatc acctcgacgc gttccgtggac 720
 ggcgcgggtct gggcgatcg cgcggaggag gcccactcg gacccagccg ggtcgacacc 780
 ggcgcggcgcc tcttcgtcg cgcacggaaat ctgaacccgtt atccgcgcgtt ggactggcag 840
 gtgtacggcg acagttcca catcggtcgat 870

<210> SEQ ID NO 31

US 10,144,941 B2

119**120**

-continued

```

<211> LENGTH: 2373
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE
<400> SEQUENCE: 31

atgaccccac cgacacagaca ccgcctgttc aggcgctcggt tgcggcttc cctctcgctg      60
gcccttacccg ccgtcggcac cgccgcccgg gtcgtctgg ccgggtcccc ggccggccagg      120
ggcgccgggg tccccggacc ctccccgggtc ggcatatccg gccggggccgc cgccgtcccg      180
ttcacggagc aggaggccga gtacgcccgg accaacggca cgctcatcggt cccggacccgg      240
cgctacggct cactgcccgg ggaggcggtcc ggccggcagg ccgtcacgct cgacgcggcc      300
ggtgagtacg tggagttcac cctcaccggcc cccgccaacg cgatgacctt ccgttatccg      360
ctgcccggaca acggccgggg gacggggccgg gacggctctc tgcacctcggt ggtgaacccgg      420
tcgggtctca agagcgtgcc ggtgacctcg aagtacggct ggtactacgg ggttacccccc      480
ttcaacaaca accccggggaa caccaacccgg caccatttc acgacgagac ccggaccatg      540
ttcggtctcgaa ccctgcccgc cggtacgaaag gtccggctgc aggtggcgcc caccggccgc      600
tcggccctcgat tcaccgtcgaa cctggccgac ttcgagcagg tggccggccgc cgccggcaag      660
ccgtccggcgcc cactggacgt ggtgagcgtac ttccggggccgg accccgggggg ggccggccgac      720
tccaccgcga agatccaggc ggccgtcgac gcccggggccaa cccagggccaa ggtcgctac      780
atccccggcagg ggaccttcca ggtgcgtgac cacatcggtcg tggaccaggta gacgtgcgc      840
ggccggccggcc cctggtagcag cggtgtcgac gggccgtcacc ccacggacccgg gagcaaggcg      900
gtcggtgtct acgggaagta ctcggcgcag ggccggcagca ggaacgtcac cctcaaggac      960
ttcgccatca tcggcgacat ccaggagcgt gtggacaacg accaggtcaa cgccatcgcc      1020
ggggccatgt ccgactcggt cggtcgacaaac gtctggatgc agcacaacccaa gtgcggccgc      1080
tggatggacg gcccgtatggaa caatttcacc atcaagaaca gtgcgtatccct ggaccagacc      1140
ggccggccggcc tgaacttcca ctacggggcc acgaactcgaa ccgtcacgaa cacccgtcgcc      1200
cgcaacacccg gtgacgacgg cctggccatg tggccggaga acgtccggaa cgtgaagaac      1260
aagttcacgt tcaacacgggt gatctggccg atccctggccca acaacatcggt gacgtacggc      1320
ggcaaggaca tcaacatcgat cgcacacccg atggccggaca ccatcacccaa cggccggccgg      1380
ctgcacatcg ccaacccgcta cccggggccgc aactcggggcc aggggacggc cgccggccgggg      1440
acgcacacccg ccgcgcgcac cacccgtatc cgtaacggca acagcgactt caactggaaac      1500
ttcggtgtcg gggcgatctg gttcagcggtt ctcacacgaa acgtcgacaa cgccaccatc      1560
aacatcaccgg acagcgaggt cctggacagc tcctacggccg cgatccacccgt gatcgagggt      1620
ggcgacacccg ggctgcactt caagaacgtc aagatcgaccc gggccgggtac ctacggccctg      1680
cagatccagg cacccggccac ggccacccatc gagaacgtcg tggccacccaa catcgcccg      1740
tccaacccgaa tccacaactg tgcggcagc gggttccaga tcacccgggg cagccggcaac      1800
tccggctgggt acggccaccc gccccctgc accgggggtct ggccccgaccc ggtgtggacc      1860
aacggccggcc tgccgggggg cggccggccccc accaaccggc ccgaccccccac cgaccccccacc      1920
gaccccgacgg accccccggc cccggccctgag gagacggggca acctcgcccg gggacgcacc      1980
gtcaccgaga ccagccacac ggacgtgtac ggccggccca acaccgtcgaa cggcaacccg      2040
gacacgtact gggagagccg caacaacgc ttcggccagtc ccgtcacccgt cgacccgtcc      2100
gtcgccaaagg cggtgaagcg ggtgggtctg aagctccgc cggccggccgc gtggggccacc      2160
cgcaacgcaga cgctctccgt gtccggcagc accgacaacg ggacgtacaa ctcgtgaag      2220

```

-continued

gcgtcgccgg gttacacctt caaccgcgtc agcggcaaca ccgcgacggt ctccctcccg 2280
 gggacgcggg tccggcacct gcggctgacc ttccacccaga acaccgggtg gcccggcc 2340
 cagctgtccg aactggaggc ctacaccaggc tga 2373

<210> SEQ ID NO 32
 <211> LENGTH: 1545
 <212> TYPE: DNA
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 32

atgaggagac cagtcgcctt gcgactcagg gggcggggg ccaccctggc cctggctgcc 60
 gcgaccggcg cactgatggc gatgcccgg gggcggtgg cagcgaccgg cgccgtcacc 120
 ggatacgcga occagaacgg cgccaccacc ggccggcgccg gggccggac ggtggggcc 180
 accacccggga cccgcgttcca cggcccttg tggggggggg ccagcagctc caccggctc 240
 accatccagg tcgagggggac catcaaccac ggcaacaccc acaaggcttc gggcagcagc 300
 tgcaacaccc cccggggagt catcgagctg aagcagatca gcaacgtcac gatcgctggc 360
 gtggggggcg gggccgttcc cgaccaggta ggcattccacg tccggcgagtc cagcaacatc 420
 atcatccaga acgttaccgt caagaacgtc aagaaggccg gtcgcacac gtccaaacggc 480
 ggtgacgcca tggcatggaa gaaggacgtc cgcaacgtct gggggacca caccacccctg 540
 gaggcctcg gggggggcgg gggggcttc gacggccctt tccggatgaa gggggccacc 600
 cagtacgta cgctgtccca cagcatccgt cgcaactccg gccccggggg cctcgctggc 660
 tccagegaga gcaaccttcc gaaaggcttc atcaccatcc accacaacct gtacgagaac 720
 atcgactccc gggccctct gtcggggggc ggcgtcgccc acatctacaa caaccactac 780
 gtggggactca gcaagtcggg catcaactcc cggggccggcg cccggcccaa ggtggacaac 840
 aactactcg aggactccaa ggacgttccg ggcacccctt acaccgacgc gggggctac 900
 tggcagggtca gggcaacgt ctggacaaac gtgacgttgt cggccggcag cagcgacaac 960
 aacccggcg gcccggaccc gcaacttccac acctcggtca gcatccctca cgcctacacc 1020
 ctcgaegggg cgaactcggt accgtccgtc gtgagccggc gggggggcgc gAACACGGG 1080
 ctgaagggtgt cggacggcag ctgtcgccg cagacggccg accccgaccga ccccccccc 1140
 gacccgacgc cggacccgac cgacccact cggccaccc ggaccaacct cagctcggg 1200
 gccggctcg acggctccag caaggcgacg gggaccagct acggcgacgt gggggacgg 1260
 gacatgagca cctactggc accgtccggc tccggatgggt ccgtctcgat caagtggagc 1320
 tccggccacca cggctccaa gatcaacgtg cggcaggccg cggccctccac gggctccatc 1380
 acctcttggg aggtcgccaa cggcggaccc ggcggccgtcc tggcctccgg cagccccgg 1440
 ggcgtcatca cttcccgca gacctcggtc cgcaagatca cttcgagat cacggctcg 1500
 acggggcaccg cgaagggtcgc cggatcgag acgtacggccg gctga 1545

<210> SEQ ID NO 33
 <211> LENGTH: 389
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 33

Met Pro Glu Arg Phe Thr Pro Thr Pro Glu Asp Lys Phe Thr Phe Gly
 1 5 10 15

Leu Trp Thr Val Gly Trp Arg Gly Asn Asp Pro Phe Gly Glu Pro Thr

US 10,144,941 B2

123

124

-continued

20	25	30	
Arg Pro Val Leu Asp Pro Val Glu Ser Val Glu Arg	Leu Ala Glu Leu		
35	40	45	
Gly Ala His Gly Val Thr Phe His Asp Asp Asp	Leu Ile Pro Phe Gly		
50	55	60	
Ser Asp Asp Arg Glu Arg Ala Arg	Leu Val Gly Arg Phe Arg Glu Ala		
65	70	75	80
Leu Glu Arg Thr Gly Leu Lys Val Pro Met Ala Thr	Thr Asn Leu Phe		
85	90	95	
Thr His Pro Val Phe Lys Asp Gly Gly Phe Thr Ser	Asn Asp Arg Asp		
100	105	110	
Val Arg Arg Phe Ala Leu Arg Lys Val Ile Arg Asn	Ile Asp Leu Ala		
115	120	125	
Val Glu Leu Gly Ala Gln Thr Tyr Val Ala Trp Gly	Gly Arg Glu Gly		
130	135	140	
Ala Glu Ser Gly Ala Ala Lys Asp Val Arg Ser Ala	Leu Asp Arg Met		
145	150	155	160
Lys Glu Ala Phe Asp Leu Leu Gly Asp Tyr Val Thr	Glu Gln Gly Tyr		
165	170	175	
Asp Leu Arg Phe Ala Ile Glu Pro Lys Pro Asn Glu	Pro Arg Gly Asp		
180	185	190	
Ile Leu Leu Pro Thr Ile Gly His Ala Leu Ala Phe	Ile Glu Arg Leu		
195	200	205	
Glu Arg Pro Glu Leu Val Gly Val Asn Pro Glu Thr	Gly His Glu Gln		
210	215	220	
Met Ala Gly Leu Asn Phe Pro His Gly Ile Ala Gln	Ala Leu Trp Ala		
225	230	235	240
Gly Lys Leu Phe His Ile Asp Leu Asn Gln Ser Gly	Ile Lys Tyr		
245	250	255	
Asp Gln Asp Phe Arg Phe Gly Ala Gly Asp Leu Arg	Gln Ala Phe Trp		
260	265	270	
Leu Val Asp Leu Leu Glu Thr Ala Gly Trp Asp Gly	Ser Arg His Phe		
275	280	285	
Asp Phe Lys Pro Val Arg Thr Asp Gly Ile Asp Gly	Val Trp Glu Ser		
290	295	300	
Ala Lys Asn Cys Met Arg Asn Tyr Leu Ile Leu Lys	Glu Arg Ala Ala		
305	310	315	320
Ala Phe Arg Ala Asp Pro Ala Val Gln Glu Ala Leu	Thr Ala Ser Arg		
325	330	335	
Leu Asp Glu Leu Ala Arg Pro Thr Ala Asp Asp Gly	Leu Lys Ala Leu		
340	345	350	
Leu Ala Asp Arg Thr Ala Tyr Glu Asp Phe Asp Ala	Thr Ala Ala Ala		
355	360	365	
Glu Arg Ser Met Ala Phe Glu Ala Leu Asp Gln Leu	Ala Met Asp His		
370	375	380	
Leu Leu Asn Val Arg			
385			

<210> SEQ ID NO 34

<211> LENGTH: 655

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 34

-continued

Met Thr Ser Ala Leu Arg Ala Thr Gln Gly Leu Gln Ser Thr Asn His
 1 5 10 15

Pro Arg Leu Ser Asp Leu Thr Arg Gly Ala Pro Leu Ser Thr Glu Ser
 20 25 30

Pro Arg Arg Ser Ser Arg Leu Arg Trp Arg Leu Gly Pro Gly Arg Ala
 35 40 45

Thr Arg Ala Lys Ala Val Ala Gly Phe Thr Ala Leu Leu Leu Pro Leu
 50 55 60

Ala Ala Met Val Gly Leu Ala Ser Pro Ala Gln Ala Ala Thr Ser Ala
 65 70 75 80

Thr Ala Thr Tyr Leu Lys Lys Ser Asp Trp Gly Ser Gly Phe Glu Gly
 85 90 95

Gln Trp Thr Val Lys Asn Thr Gly Thr Thr Ala Leu Ser Ser Trp Thr
 100 105 110

Ile Glu Trp Asp Phe Pro Ser Gly Thr Ala Val Gly Ser Ala Trp Asp
 115 120 125

Ala Ser Val Thr Ser Ser Gly Thr His Trp Thr Ala Lys Asn Leu Gly
 130 135 140

Trp Asn Gly Thr Val Ala Pro Gly Ala Ser Ile Ser Phe Gly Phe Asn
 145 150 155 160

Gly Thr Gly Ser Gly Ser Pro Thr Gly Cys Lys Leu Asn Gly Ala Ser
 165 170 175

Cys Asp Gly Gly Thr Val Pro Gly Asp Ser Ala Pro Ser Lys Pro
 180 185 190

Gly Thr Pro Thr Ala Ser Gly Ile Thr Asp Thr Ser Val Lys Leu Ser
 195 200 205

Trp Ser Ala Ala Thr Asp Asp Lys Gly Ile Lys Asn Tyr Asp Val Leu
 210 215 220

Arg Asp Gly Ala Lys Val Ala Thr Val Thr Thr Thr Tyr Thr Asp
 225 230 235 240

Thr Gly Leu Thr Lys Gly Thr Asp Tyr Ser Tyr Ser Val Gln Ala Arg
 245 250 255

Asp Thr Ala Asp Gln Thr Gly Pro Val Ser Gly Ala Val Ala Val Arg
 260 265 270

Thr Thr Gly Gly Asn Asp Asn Pro Gly Pro Gly Thr Gly Ser Lys Val
 275 280 285

Asn Leu Gly Tyr Phe Thr Asn Trp Gly Val Tyr Gly Arg Asn Tyr His
 290 295 300

Val Lys Asn Leu Val Thr Ser Gly Ser Ala Glu Lys Ile Thr His Ile
 305 310 315 320

Asn Tyr Ala Phe Gly Asn Val Gln Gly Gly Lys Cys Thr Ile Gly Asp
 325 330 335

Ser Tyr Ala Asp Tyr Asp Lys Ala Tyr Thr Ala Asp Gln Ser Val Asp
 340 345 350

Gly Val Ala Asp Thr Trp Asp Gln Pro Leu Arg Gly Asn Phe Asn Gln
 355 360 365

Leu Arg Lys Leu Lys Ala Lys Tyr Pro His Ile Lys Val Ile Trp Ser
 370 375 380

Phe Gly Gly Trp Thr Trp Ser Gly Gly Phe Gly Ala Ala Gln Asn
 385 390 395 400

Pro Ala Ala Phe Ala Gln Ser Cys Tyr Asp Leu Val Glu Asp Pro Arg
 405 410 415

Trp Ala Asp Val Phe Asp Gly Ile Asp Ile Asp Trp Glu Tyr Pro Asn

US 10,144,941 B2

127

-continued

128

420	425	430
Ala Cys Gly Leu Thr Cys Asp Thr Ser Gly Pro Ala Ala Leu Lys Asn		
435	440	445
Leu Ser Ser Ala Leu Arg Ala Lys Phe Gly Ala Lys Asn Leu Val Thr		
450	455	460
Ala Ala Ile Thr Ala Asp Gly Ser Asp Gly Gly Lys Ile Asp Ala Ala		
465	470	475
Asp Tyr Ala Gly Ala Ala Gln Ser Phe Asp Trp Tyr Asn Val Met Thr		
485	490	495
Tyr Asp Phe Phe Gly Ala Trp Glu Ala Lys Gly Pro Thr Ala Pro His		
500	505	510
Ser Pro Leu Asn Ala Tyr Ala Gly Ile Pro Gln Asp Gly Phe Asn Ser		
515	520	525
Ala Ala Ala Ile Ala Lys Leu Lys Ala Lys Gly Val Pro Ala Ser Lys		
530	535	540
Leu Leu Leu Gly Ile Gly Phe Tyr Gly Arg Gly Trp Thr Gly Val Thr		
545	550	555
Gln Ala Ala Pro Gly Gly Thr Ala Thr Gly Ala Ala Pro Gly Thr Tyr		
565	570	575
Glu Ala Gly Ile Glu Asp Tyr Lys Val Leu Lys Thr Ser Cys Pro Ala		
580	585	590
Thr Gly Thr Ile Ala Gly Thr Ala Tyr Ala His Cys Gly Thr Asn Trp		
595	600	605
Trp Ser Tyr Asp Thr Pro Ala Thr Ile Thr Ser Lys Met Ala Trp Ala		
610	615	620
Asn Ser Gln Gly Leu Gly Gly Ala Phe Phe Trp Glu Phe Ser Gly Asp		
625	630	635
Thr Ala Asn Gly Glu Leu Val Ser Ala Met Asp Ser Gly Leu Asn		
645	650	655

<210> SEQ ID NO 35

<211> LENGTH: 196

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 35

Met Arg Lys Arg Ala Ser Ala Ala Val Ile Gly Leu Ala Ile Ala Gly		
1	5	10
Val Ser Met Phe Ala Thr Ser Ser Ala Ser Ser His Gly Tyr Thr Asp		
	20	25
Ser Pro Ile Ser Arg Gln Lys Leu Cys Ala Asn Gly Thr Val Thr Gly		
	35	40
Cys Gly Asn Ile Gln Trp Glu Pro Gln Ser Val Glu Gly Pro Lys Gly		
	50	55
Phe Pro Ala Ala Gly Pro Ala Asp Gly Lys Ile Cys Ala Gly Gly Asn		
65	70	75
Ser Ser Phe Ala Ala Leu Asp Asp Pro Arg Gly Gly Asn Trp Pro Ala		
85	90	95
Thr Gln Val Thr Gly Gly Gln Gly Tyr Asn Phe Arg Trp Gln Phe Thr		
100	105	110
Ala Arg His Ala Thr Thr Asp Phe Arg Tyr Tyr Ile Thr Lys Asp Gly		
115	120	125
Trp Asp Ser Thr Lys Pro Leu Thr Arg Ala Ala Leu Glu Ser Gln Pro		
130	135	140

US 10,144,941 B2

129**130**

-continued

Phe Met Thr Val Pro Tyr Gly Asn Gln Gln Pro Pro Ala Thr Leu Thr
 145 150 155 160

His Gln Gly Thr Ile Pro Thr Gln Lys Ser Gly Lys His Ile Ile Leu
 165 170 175

Ala Val Trp Asn Val Ala Asp Thr Ala Asn Ala Phe Tyr Ala Cys Ser
 180 185 190

Asp Val Lys Phe
 195

<210> SEQ ID NO 36
 <211> LENGTH: 556
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 36

Val Ala Ala Leu Ala Ala Gly Ala Leu Thr Val Thr Gly Leu Val Gly
 1 5 10 15

Thr Ala Gln Ala Ala Asp Ile Asn Val Ala Lys Asn Ala Gly Phe Glu
 20 25 30

Ser Gly Leu Ser Gly Trp Thr Cys Thr Gly Gly Ser Gly Ala Thr Val
 35 40 45

Ser Ser Pro Val His Gly Gly Ser Ala Ala Leu Lys Ala Thr Pro Ser
 50 55 60

Gly Gln Asp Asn Ala Lys Cys Thr Gln Thr Val Ala Val Lys Pro Asn
 65 70 75 80

Ser Thr Tyr Ala Leu Ser Ser Trp Val Gln Gly Gly Tyr Ala Tyr Leu
 85 90 95

Gly Ala Ser Gly Thr Gly Thr Asp Val Ser Thr Trp Thr Pro Gly
 100 105 110

Ser Thr Gly Trp Thr Gln Leu Arg Thr Ser Phe Thr Thr Gly Pro Ser
 115 120 125

Thr Thr Ser Val Gln Val Tyr Thr His Gly Trp Tyr Gly Gln Ala Ala
 130 135 140

Tyr Tyr Ala Asp Asp Val Ala Val Thr Gly Pro Asp Gly Gly Gly
 145 150 155 160

Thr Glu Glu Pro Gly Pro Ala Ile Pro Gly Ala Pro Ala Gly Leu Ala
 165 170 175

Val Gly Thr Thr Ser Ser Val Ala Leu Ser Trp Asn Ala Val
 180 185 190

Ser Gly Ala Thr Gly Tyr Thr Val Tyr Arg Asp Gly Thr Lys Ala Thr
 195 200 205

Thr Thr Thr Gly Thr Ser Ala Thr Val Ser Gly Leu Ala Ala Asp Thr
 210 215 220

Ala Tyr Gln Phe Ser Val Ser Ala Thr Asn Ala Ala Gly Glu Ser Val
 225 230 235 240

Arg Ser Ala Thr Val Ser Gly Arg Thr Ala Lys Lys Asp Glu Thr Gly
 245 250 255

Pro Gly Pro Ser Thr Ser Val Pro Lys His Ala Val Thr Gly Tyr Trp
 260 265 270

Gln Asn Phe Asn Asn Gly Ala Ala Val Gln Lys Leu Ser Asp Val Pro
 275 280 285

Ala Asn Tyr Asp Ile Ile Ala Val Ser Phe Ala Asp Ala Ala Gly Thr
 290 295 300

Pro Gly Ala Val Thr Phe Asn Leu Asp Ser Ala Gly Leu Asn Gly Tyr
 305 310 315 320

-continued

Thr Val Ala Gln Phe Lys Ala Asp Ile Lys Ala Lys Gln Ala Ala Gly
 325 330 335
 Lys Asn Val Ile Ile Ser Val Gly Gly Glu Lys Gly Thr Val Ser Val
 340 345 350
 Asn Ser Asp Ala Ser Ala Asn Ala Phe Ala Asp Ser Leu Tyr Thr Leu
 355 360 365
 Ile Gln Glu Tyr Gly Phe Asn Gly Val Asp Ile Asp Leu Glu Asn Gly
 370 375 380
 Leu Asn Ser Thr Tyr Met Thr Lys Ala Leu Arg Ser Leu Ser Ser Lys
 385 390 395 400
 Val Gly Ser Gly Leu Val Ile Thr Met Ala Pro Gln Thr Ile Asp Met
 405 410 415
 Gln Ser Thr Ser Gly Glu Tyr Phe Lys Thr Ala Leu Asn Ile Lys Asp
 420 425 430
 Ile Leu Thr Val Val Asn Met Gln Tyr Tyr Asn Ser Gly Ser Met Leu
 435 440 445
 Gly Cys Asp Gly Lys Val Tyr Ser Gln Gly Ser Val Asp Phe Leu Thr
 450 455 460
 Ala Leu Ala Cys Ile Gln Leu Glu Gly Leu Ala Pro Ser Gln Val
 465 470 475 480
 Gly Leu Gly Val Pro Ala Ser Thr Arg Gly Ala Gly Ser Gly Tyr Val
 485 490 495
 Ala Pro Ser Val Val Asn Ala Ala Leu Asp Cys Leu Ala Lys Gly Thr
 500 505 510
 Gly Cys Gly Ser Phe Lys Pro Ser Arg Thr Tyr Pro Asp Ile Arg Gly
 515 520 525
 Ala Met Thr Trp Ser Thr Asn Trp Asp Ala Thr Ala Gly Asn Ala Trp
 530 535 540
 Ser Asn Ala Val Gly Pro His Val His Gly Leu Pro
 545 550 555

<210> SEQ_ID NO 37
 <211> LENGTH: 295
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 37

Val Ile Arg Arg Val Met Gly Leu Leu Thr Ala Leu Ala Ala Val Val
 1 5 10 15
 Ala Thr Leu Val Phe Leu Pro Ala Ala Thr Ala Ser Ala Ala Thr Cys
 20 25 30
 Ala Pro Ala Trp Asn Ala Ser Ser Val Tyr Thr Gly Gly Ser Ala
 35 40 45
 Ser Tyr Asn Gly His Asn Trp Ser Ala Lys Trp Trp Thr Gln Asn Glu
 50 55 60
 Arg Pro Gly Thr Ser Asp Val Trp Ala Asp Gln Gly Ala Cys Gly Ser
 65 70 75 80
 Gly Gly Gly Thr Asp Pro Asn Pro Ser Gly Phe Val Val Ser Glu
 85 90 95
 Ala Gln Phe Asn Gln Met Phe Pro Ser Arg Asn Ser Phe Tyr Thr Tyr
 100 105 110
 Ser Gly Leu Thr Ala Ala Leu Ser Ala Tyr Pro Ala Phe Ala Asn Thr
 115 120 125
 Gly Ser Asp Thr Val Lys Lys Gln Glu Ala Ala Phe Leu Ala Asn

US 10,144,941 B2

133

134

-continued

130	135	140
Val Ser His Glu Thr Gly	Gly Leu Val His Ile Val Glu Gln Asn Thr	
145	150	155
160		
Ala Asn Tyr Pro His Tyr Cys Asp Thr Ser Gln Ser Tyr Gly Cys Pro		
165	170	175
Ala Gly Gln Ala Ala Tyr Tyr Gly Arg Gly Pro Ile Gln Leu Ser Trp		
180	185	190
Asn Phe Asn Tyr Lys Ala Ala Gly Asp Ala Leu Gly Ile Asp Leu Leu		
195	200	205
Gly Asn Pro Trp Gln Val Glu Gln Asn Ala Ser Val Ala Trp Lys Thr		
210	215	220
Gly Leu Trp Tyr Trp Asn Thr Gln Ser Gly Pro Gly Thr Met Thr Pro		
225	230	235
240		
His Asn Ala Ile Val Asn Gly Ser Gly Phe Gly Glu Thr Ile Arg Ser		
245	250	255
Ile Asn Gly Ser Ile Glu Cys Asn Gly Gly Asn Pro Gly Gln Val Gln		
260	265	270
Ser Arg Val Asn Thr Tyr Gln Ser Phe Val Gln Ile Leu Gly Thr Thr		
275	280	285
Pro Gly Ser Asn Leu Ser Cys		
290	295	

<210> SEQ ID NO 38

<211> LENGTH: 507

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 38

Met Arg Arg Ser Arg Ser Val Arg Ala Leu Val Thr Ala Ala Val Thr		
1	5	10
15		
Thr Val Ala Ala Ala Gly Met Ala Val Leu Gly Ser Gly Thr Ala Gln		
20	25	30
Ala Ala Thr Pro Leu Pro Asp His Val Phe Ala Pro Tyr Phe Glu Ser		
35	40	45
Trp Thr Gly Glu Ser Pro Ala Ala Met Ala Ala Glu Ser Gly Ala Lys		
50	55	60
His Leu Thr Met Ala Phe Leu Gln Thr Thr Ala Lys Gly Ser Cys Thr		
65	70	75
80		
Pro Tyr Trp Asn Gly Asp Thr Gly Leu Pro Ile Ala Gln Ala Ser Phe		
85	90	95
Gly Ala Asp Ile Asp Thr Ile Gln Ala Gly Gly Asp Val Ile Pro		
100	105	110
Ser Phe Gly Gly Tyr Thr Ala Asp Thr Thr Gly Thr Glu Ile Ala Asp		
115	120	125
Ser Cys Thr Asp Val Asp Gln Ile Ala Ala Tyr Gln Lys Val Val		
130	135	140
Thr Thr Tyr Asp Val Ser Arg Leu Asp Met Asp Ile Glu Val Asp Ser		
145	150	155
160		
Leu Asp Asp Thr Ala Gly Ile Asp Arg Arg Asn Lys Ala Ile Lys Lys		
165	170	175
Leu Gln Asp Trp Ala Asp Ala Asn Gly Arg Asp Leu Glu Ile Ser Tyr		
180	185	190
Thr Leu Pro Thr Thr Arg Gly Leu Ala Ser Ser Gly Leu Ala Val		
195	200	205

US 10,144,941 B2

135

136

-continued

Leu Arg Asn Ala Val Thr Asn Gly Ala Arg Val Asp Val Val Asn Leu
 210 215 220
 Met Thr Phe Asp Tyr Tyr Asp Asn Ala Ser His Asp Met Ala Ala Asp
 225 230 235 240
 Thr Glu Thr Ala Ala Gln Gly Leu Tyr Asp Gln Leu Ala Lys Leu Tyr
 245 250 255
 Pro Gly Arg Thr Ala Thr Gln Leu Trp Ser Met Val Gly Val Thr Glu
 260 265 270
 Met Pro Gly Val Asp Asp Phe Gly Pro Ala Glu Thr Phe Thr Leu Ala
 275 280 285
 Asn Ala Ala Arg Val Tyr Asp Trp Ala Val Ala Lys Gly Ile Asn Thr
 290 295 300
 Leu Ser Phe Trp Ala Leu Gln Arg Asp Asn Gly Gly Cys Pro Gly Gly
 305 310 315 320
 Pro Ala Ala Asp Asp Cys Ser Gly Ile Gln Gln Asn Thr Trp Asp Phe
 325 330 335
 Thr Arg Val Phe Ala Pro Phe Thr Ser Gly Thr Thr Ala Pro Asp Asp
 340 345 350
 Asp Phe Ser Val Thr Ala Thr Pro Ala Ser Gly Thr Val Thr Ala Gly
 355 360 365
 Gly Ser Ala Thr Thr Thr Val Lys Thr Ala Val Thr Lys Gly Ala Ala
 370 375 380
 Gln Gln Val Gly Leu Thr Val Ser Gly Val Pro Ala Gly Val Thr Ala
 385 390 395 400
 Ser Leu Ser Pro Ser Ser Val Thr Ala Gly Gly Arg Ser Thr Leu Thr
 405 410 415
 Leu Ala Thr Thr Gln Ala Ala Val Ser Gly Thr Tyr Arg Ile Ser Val
 420 425 430
 Thr Gly Thr Ser Pro Ser Gly Ser His Ala Thr Ala Tyr Thr Leu Thr
 435 440 445
 Val Thr Gly Gly Thr Gly Ser Gln Cys Thr Ala Gly Pro Trp Ala Gly
 450 455 460
 Gly Thr Val Tyr Thr Gly Gly Gln Gln Val Ser Tyr Lys Gly His Thr
 465 470 475 480
 Trp Lys Ala Lys Trp Trp Thr Thr Gly Glu Glu Pro Gly Thr Thr Gly
 485 490 495
 Glu Trp Gly Val Trp Gln Asp Leu Gly Ala Cys
 500 505

<210> SEQ ID NO 39

<211> LENGTH: 483

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 39

Val Thr Gln Gly Pro Leu Thr Thr Glu Ala Gly Ala Pro Val Ala Asp
 1 5 10 15
 Asn Gln Asn Ser Glu Thr Ala Gly Pro Gly Gly Pro Val Leu Val Gln
 20 25 30
 Asp Gln Ala Leu Leu Glu Lys Leu Ala His Phe Asn Arg Glu Arg Ile
 35 40 45
 Pro Glu Arg Val Val His Ala Arg Gly Ala Gly Ala Tyr Gly Thr Phe
 50 55 60
 Thr Leu Thr Arg Asp Val Ser Gln Trp Thr Arg Ala Lys Phe Leu Ser
 65 70 75 80

-continued

Glu Val Gly Lys Glu Thr Glu Thr Phe Leu Arg Phe Ser Thr Val Ala
 85 90 95
 Gly Asn Leu Gly Ser Ala Asp Ala Ala Arg Asp Pro Arg Gly Trp Ala
 100 105 110
 Leu Lys Phe Tyr Thr Glu Glu Gly Asn Tyr Asp Leu Val Gly Asn Asn
 115 120 125
 Thr Pro Val Phe Phe Ile Lys Asp Ala Ile Lys Phe Pro Asp Phe Ile
 130 135 140
 His Thr Gln Lys Arg Asp Pro Tyr Thr Gly Ser Gln Glu Ala Asp Asn
 145 150 155 160
 Val Trp Asp Phe Trp Gly Leu Ser Pro Glu Ser Thr His Gln Val Thr
 165 170 175
 Trp Leu Phe Gly Asp Arg Gly Ile Pro Ala Ser Phe Arg His Met Asn
 180 185 190
 Gly Tyr Gly Ser His Thr Phe Gln Trp Asn Asn Glu Ala Gly Glu Val
 195 200 205
 Phe Trp Val Lys Tyr His Phe Lys Thr Asp Gln Gly Ile Lys Asn Leu
 210 215 220
 Thr Thr Glu Glu Ala Val Arg Leu Ser Gly Val Asp Pro Asp Ser His
 225 230 235 240
 Gln Arg Asp Leu Arg Glu Ser Ile Glu Arg Gly Asp Phe Pro Thr Trp
 245 250 255
 Thr Val Gln Val Gln Ile Met Pro Ala Ala Glu Ala Ala Thr Tyr Arg
 260 265 270
 Phe Asn Pro Phe Asp Leu Thr Lys Val Trp Pro His Glu Asp Tyr Pro
 275 280 285
 Pro Ile Glu Ile Gly Lys Leu Glu Leu Asn Arg Asn Pro Glu Asn Ile
 290 295 300
 Phe Ala Glu Val Glu Gln Ser Ile Phe Ser Pro Ala His Phe Val Pro
 305 310 315 320
 Gly Ile Gly Pro Ser Pro Asp Lys Met Leu Gln Gly Arg Leu Phe Ala
 325 330 335
 Tyr Gly Asp Ala His Arg Tyr Arg Val Gly Ile Asn Ala Asp His Leu
 340 345 350
 Pro Val Asn Arg Pro His Ala Thr Glu Ala Arg Thr Asn Ser Arg Asp
 355 360 365
 Gly Tyr Leu Tyr Asp Gly Arg His Lys Gly Thr Lys Asn Tyr Glu Pro
 370 375 380
 Asn Ser Phe Gly Gly Pro Val Gln Thr Asp Arg Pro Leu Trp Gln Pro
 385 390 395 400
 Val Ser Val Thr Gly Gly Thr Gly Asn His Glu Ala Ala Val His Ala
 405 410 415
 Glu Asp Asn Asp Phe Val Gln Ala Gly Asn Leu Tyr Arg Leu Met Ser
 420 425 430
 Glu Asp Glu Lys Gly Arg Leu Ile Asp Asn Leu Ala Gly Phe Ile Ala
 435 440 445
 Lys Val Ser Arg Asp Asp Ile Ala Asp Arg Ala Ile Asn Asn Phe Arg
 450 455 460
 Gln Ala Asp Ala Asp Phe Gly Lys Arg Leu Glu Val Ala Val Gln Ala
 465 470 475 480
 Leu Arg Gly

US 10,144,941 B2

139**140**

-continued

<210> SEQ ID NO 40

<211> LENGTH: 926

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 40

Val Tyr Ala Met Pro Ser Thr Ala Pro Ala Ala Val Gln Ser Gly Glu
 1 5 10 15

Asp Ala Pro Val Arg Ser Ser Pro Arg Pro Phe Ala Ala Leu Leu Ala
 20 25 30

Ala Leu Ala Leu Thr Ala Gly Leu Ser Leu Ile Gly Thr Pro Ala Val
 35 40 45

Ala Arg Ser Asp Glu Ala Pro Ala Ala Thr Glu Ala Ser Asp Val Ser
 50 55 60

Ile Ala Ala Asp Thr Tyr Thr Trp Lys Asn Ala Arg Ile Asp Gly Gly
 65 70 75 80

Gly Phe Val Pro Gly Ile Val Phe Asn Arg Ser Glu Lys Asn Leu Ala
 85 90 95

Tyr Ala Arg Thr Asp Ile Gly Gly Ala Tyr Arg Trp Asp Gln Ser Gly
 100 105 110

Lys Gln Trp Lys Pro Leu Leu Asp Trp Val Asp Trp Asp Arg Trp Gly
 115 120 125

Trp Thr Gly Val Val Ser Leu Ala Ser Asp Thr Val Asp Pro Asp Asn
 130 135 140

Val Tyr Ala Ala Val Gly Thr Tyr Thr Asn Ser Trp Asp Pro Thr Asp
 145 150 155 160

Gly Ala Val Leu Arg Ser Ser Asp Arg Gly Ala Ser Trp Lys Ala Ala
 165 170 175

Thr Leu Pro Phe Lys Leu Gly Gly Asn Met Pro Gly Arg Gly Met Gly
 180 185 190

Glu Arg Leu Ala Val Asp Pro Asn Lys Asn Ser Val Leu Tyr Leu Gly
 195 200 205

Ala Pro Ser Gly Asn Gly Leu Trp Arg Ser Thr Asp Ala Gly Val Ser
 210 215 220

Trp Ser Glu Val Thr Ala Phe Pro Asn Pro Gly Asn Tyr Ala Gln Asp
 225 230 235 240

Pro Ser Asp Thr Ser Gly Tyr Gly Asn Asp Asn Gln Gly Ile Val Trp
 245 250 255

Val Thr Phe Asp Glu Arg Ser Gly Ser Ala Gly Ser Ala Thr Gln Asp
 260 265 270

Ile Tyr Val Gly Val Ala Asp Lys Glu Asn Thr Val Tyr Arg Ser Thr
 275 280 285

Asp Gly Gly Ala Thr Trp Ser Arg Ile Pro Gly Gln Pro Thr Gly Tyr
 290 295 300

Leu Ala His Lys Gly Val Leu Asp Ser Ala Thr Gly His Leu Tyr Leu
 305 310 315 320

Thr Leu Ser Asp Thr Gly Gly Pro Tyr Asp Gly Gly Lys Gly Arg Ile
 325 330 335

Trp Arg Tyr Asp Thr Ala Ser Gly Ala Trp Gln Asp Val Ser Pro Val
 340 345 350

Ala Glu Ala Asp Ala Tyr Tyr Gly Phe Ser Gly Leu Ser Val Asp Arg
 355 360 365

Gln Lys Pro Gly Thr Leu Met Ala Thr Ala Tyr Ser Ser Trp Trp Pro
 370 375 380

-continued

Asp Thr Gln Ile Phe Arg Ser Thr Asp Ser Gly Ala Thr Trp Thr Gln
 385 390 395 400

 Ala Trp Asp Tyr Thr Gly Tyr Pro Asn Arg Ser Asn Arg Tyr Thr Leu
 405 410 415

 Asp Val Ser Ser Val Pro Trp Leu Ser Trp Gly Ala Ser Pro Ala Pro
 420 425 430

 Pro Glu Thr Ala Pro Lys Leu Gly Trp Met Thr Glu Ala Leu Glu Ile
 435 440 445

 Asp Pro Phe Asp Ser Asp Arg Met Met Tyr Gly Thr Gly Ala Thr Val
 450 455 460

 Tyr Gly Thr Glu Asp Leu Thr Ser Trp Asp Ser Gly Gly Thr Phe Arg
 465 470 475 480

 Ile Thr Pro Met Val Lys Gly Ile Glu Glu Thr Ala Val Asn Asp Leu
 485 490 495

 Ala Ser Pro Pro Ser Gly Ala Pro Leu Leu Ser Ala Leu Gly Asp Ile
 500 505 510

 Gly Gly Phe Arg His Thr Asp Leu Asp Ala Val Pro Asp Leu Met Tyr
 515 520 525

 Thr Ser Pro Asn Leu Asp Ser Thr Thr Ser Leu Asp Phe Ala Glu Ser
 530 535 540

 Ser Pro Gly Thr Val Val Arg Val Gly Asn Ser Asp Ala Ala Pro His
 545 550 555 560

 Ile Gly Phe Ser Thr Asp Asn Gly Ala Asn Trp Phe Gln Gly Ser Glu
 565 570 575

 Pro Ser Gly Val Thr Gly Gly Thr Val Ala Ala Ala Ala Asp Gly
 580 585 590

 Ser Gly Phe Val Trp Ser Pro Glu Gly Ala Gly Val His His Thr Thr
 595 600 605

 Gly Phe Gly Thr Ser Trp Thr Ala Ser Thr Gly Ile Pro Ala Gly Ala
 610 615 620

 Thr Val Glu Ser Asp Arg Lys Asn Pro Glu Lys Phe Tyr Gly Phe Glu
 625 630 635 640

 Ala Gly Thr Phe Tyr Val Ser Thr Asp Gly Gly Ala Thr Phe Thr Ala
 645 650 655

 Glu Ala Thr Gly Leu Pro Ala Glu Gly Asn Val Arg Phe Gln Ala Leu
 660 665 670

 Pro Gly Thr Glu Gly Asp Ile Trp Leu Ala Gly Gly Ser Asp Thr Gly
 675 680 685

 Ala Tyr Gly Leu Trp Arg Ser Thr Asp Ser Gly Ala Thr Phe Thr Lys
 690 695 700

 Ser Ala Gly Val Glu Gln Ala Asp Ser Val Gly Phe Gly Lys Ala Ala
 705 710 715 720

 Pro Gly Ala Ser Tyr Arg Thr Val Phe Val Ser Ala Lys Ile Gly Gly
 725 730 735

 Val Arg Gly Ile Phe Arg Ser Thr Asp Ala Gly Ala Ser Trp Thr Arg
 740 745 750

 Ile Asn Asp Asp Ala His Gln Trp Gly Trp Thr Gly Ala Ala Ile Thr
 755 760 765

 Gly Asp Pro Arg Val Tyr Gly Arg Val Tyr Val Ser Thr Asn Gly Arg
 770 775 780

 Gly Ile Gln Val Gly Glu Thr Ser Asp Ser Gly Gly Gly Thr Asp
 785 790 795 800

 Pro Gly Thr Asp Pro Gly Thr Asp Pro Gly Thr Asp Pro Gly Pro Glu

US 10,144,941 B2

143

144

-continued

805	810	815
-----	-----	-----

Gln Pro Ala Asp Ala Ala Cys Ala Val Thr Tyr Ala Val Thr Asn Gln
820 825 830

Trp Pro Gly Gly Phe Gln Ala Asp Val Thr Val Thr Asn Thr Gly Asp
835 840 845

Ala Ala Tyr Asn Gly Trp Lys Leu Gly Trp Ser Phe Pro Gly Gly Gln
850 855 860

Gln Ile Ser Gln Ile Trp Asn Ala Ser His Arg Gln Asp Gly Val Lys
865 870 875 880

Val Thr Val Thr Asp Ala Gly Trp Asn Gly Thr Val Ala Pro Gly Ser
885 890 895

Ser Ala Gly Phe Gly Phe Thr Gly Ser Trp Ala Gly Ser Asn Ala Glu
900 905 910

Pro Ala Ala Phe Thr Leu Asp Gly Gln Ala Cys Thr Val Gly
915 920 925

<210> SEQ_ID NO 41

<211> LENGTH: 543

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 41

Met Arg Gly Ala Lys Ser Ala Lys Trp Val Ala Gly Ala Ala Ile Ile
1 5 10 15

Ala Leu Ala Ala Thr Ala Cys Gly Gly Asp Ser Asp Ser Asp Asn
20 25 30

Gly Ala Lys Gly Ala Val Asp Ala Asp Gly Ile Phe Ser Val Glu Val
35 40 45

Gly Glu Pro Gln Asn Pro Leu Gln Pro Ala Asn Thr Met Glu Ser Asn
50 55 60

Gly Ser Ile Val Thr Asp Ala Ile Phe Ser Gln Leu Val Asp Tyr Asp
65 70 75 80

Pro Asp Gly Lys Leu Glu Met Ile Asn Ala Glu Ser Val Glu Thr Thr
85 90 95

Asp Ser Lys Leu Trp Thr Val Lys Leu Lys Asp Trp Lys Phe His
100 105 110

Asp Gly Thr Pro Val Thr Ala Asp Ser Tyr Val Lys Ala Trp Asn Trp
115 120 125

Ala Ala Asn Ile Glu Asn Ala Gln Thr Asn Ala Ser Trp Phe Ala Asp
130 135 140

Ile Lys Gly Tyr Ala Asp Val His Pro Asp Gly Glu Gly Ala Lys Pro
145 150 155 160

Lys Ser Asp Ala Met Ser Gly Leu Lys Lys Val Asp Asp Tyr Thr Phe
165 170 175

Thr Ile Glu Leu Asn Ser Ala Val Pro Tyr Phe Ser Tyr Lys Leu Gly
180 185 190

Tyr Thr Val Phe Ser Pro Leu Pro Glu Ser Phe Tyr Ala Asp Pro Lys
195 200 205

Ala Ala Gly Glu Lys Pro Val Gly Asn Gly Ala Tyr Lys Phe Val Ser
210 215 220

Trp Asp His Lys Lys Gln Ile Lys Val Val Arg Asn Asp Asp Tyr Lys
225 230 235 240

Gly Pro Asp Lys Ala Lys Asn Gly Gly Val Ile Phe Lys Asn Tyr Thr
245 250 255

US 10,144,941 B2

145**146**

-continued

Thr Leu Glu Thr Ala Tyr Glu Asp Leu Lys Ser Gly Asn Val Asp Val
 260 265 270
 Leu Arg Gln Ile Gly Pro Lys Asp Leu Pro Val Tyr Arg Ala Asp Leu
 275 280 285
 Glu Asp Arg Ala Val Asp Lys Ala Tyr Ser Ala Val Gln Thr Leu Gly
 290 295 300
 Val Ala Met Tyr Thr Asp Gln Trp Lys Asn Thr Asp Pro Lys Val Leu
 305 310 315 320
 Gln Gly Leu Ser Met Ala Ile Asp Arg Asp Thr Ile Thr Lys Thr Val
 325 330 335
 Leu Gln Gly Thr Arg Glu Pro Ala Thr Gly Trp Val Ala Lys Gly Val
 340 345 350
 Leu Gly Tyr Gln Glu Asn Val Ala Gly Asp Val Thr Lys Tyr Asp Pro
 355 360 365
 Ala Lys Ala Lys Ala Leu Ile Lys Glu Gly Gly Val Pro Gly Asn
 370 375 380
 Glu Ile Phe Ile Gln Phe Asn Ala Asp Gly Gly His Lys Glu Trp Ile
 385 390 395 400
 Glu Ala Val Cys Asn Ser Ile Thr Gln Ala Thr Gly Val Lys Cys Thr
 405 410 415
 Gly Asp Ser Lys Ala Asp Phe Gln Ala Asp Leu Asn Ala Arg Asp Ala
 420 425 430
 Lys Gln Val Lys Ser Phe Tyr Arg Ser Gly Trp Val Leu Asp Tyr Pro
 435 440 445
 Val Asn Ala Asn Phe Ile Ser Asp Leu Phe Arg Thr Gly Ala Ala Gly
 450 455 460
 Asn Asn Gly Phe Ser Asn Lys Asp Leu Asp Ala Lys Ile Lys Ala
 465 470 475 480
 Ala Asp Ser Ala Ala Ser Leu Asp Asp Ser Val Lys Ala Tyr Gln Glu
 485 490 495
 Ile Glu Lys Glu Leu Val Asn Tyr Met Pro Ser Ile Pro Leu Trp Tyr
 500 505 510
 Tyr Lys Val Asn Ala Gly Tyr Ser Glu Asn Val Lys Asn Val Asp Tyr
 515 520 525
 Ala Gln Asp Gly Asp Pro Ile Leu Thr Glu Val Gln Val Ile Lys
 530 535 540

<210> SEQ ID NO 42
 <211> LENGTH: 159
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 42

Met Gln Gly Asp Pro Glu Val Leu Glu Phe Leu Asn Glu Gln Leu Thr
 1 5 10 15
 Ala Glu Leu Thr Ala Ile Asn Gln Tyr Phe Leu His Ala Lys Met Gln
 20 25 30
 Asp His Arg Gly Trp Thr Lys Leu Ala Lys His Thr Arg Ala Glu Ser
 35 40 45
 Phe Asp Glu Met Lys His Ala Glu Ile Leu Thr Asp Arg Ile Leu Leu
 50 55 60
 Leu Asp Gly Leu Pro Asn Tyr Gln Arg Leu Phe His Val Arg Val Gly
 65 70 75 80
 Gln Thr Val Thr Glu Met Phe Gln Ala Asp Arg Gln Val Glu Val Glu
 85 90 95

-continued

Ala Ile Asp Arg Leu Arg Arg Gly Val Asp Leu Met Arg Ala Lys Ser
 100 105 110

Asp Ile Thr Ser Ala Asn Ile Phe Glu Arg Ile Leu Glu Asp Glu Glu
 115 120 125

His His Ile Asp Tyr Leu Asp Thr Gln Leu Glu Leu Ile Glu Lys Leu
 130 135 140

Gly Glu Pro Leu Tyr Leu Ala Gln Val Ile Glu Gln Val Glu Leu
 145 150 155

<210> SEQ ID NO 43
 <211> LENGTH: 297
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 43

Met Ser Pro Tyr Thr Ala Thr Arg Arg Thr Phe Leu Thr Gly Ala Leu
 1 5 10 15

Ala Ala Ala Thr Gly Val Val Leu Gly Gly Thr Pro Ala Leu Ala Ala
 20 25 30

Pro Ala Arg Val Leu Gly Thr Gln Asp Trp Met Gly Ala Leu Ala Asp
 35 40 45

Ser Thr Pro Leu Arg Arg Leu Thr Ile Pro Gly Thr His Asn Ala Gly
 50 55 60

Ala Arg Tyr Gly Gly Pro Trp Thr Glu Cys Gln Asn Thr Thr Val Ala
 65 70 75 80

Glu Gln Leu Gly Ser Gly Ile Arg Phe Leu Asp Val Arg Cys Arg Ile
 85 90 95

Thr Gly Asp Ala Phe Ala Ile His His Gly Ala Ser Tyr Gln Asn Leu
 100 105 110

Met Phe Gly Asp Val Leu Ile Ala Cys Arg Asp Phe Leu Ala Ala His
 115 120 125

Pro Ser Glu Thr Val Leu Met Arg Val Lys Gln Glu Tyr Ser Glu Glu
 130 135 140

Ser Asp Ala Ala Phe Arg Gln Ile Phe Asp Leu Tyr Leu Asp Gly Lys
 145 150 155 160

Gly Trp Arg Pro Leu Phe Arg Leu Asp Pro Thr Leu Pro Asp Leu Gly
 165 170 175

Gly Ala Arg Gly Lys Val Val Leu Leu Ala Asp Asn Gly Gly Leu Pro
 180 185 190

Gly Val Arg Tyr Ala Asp Pro Ala Val Phe Asp Ile Gln Asp Asp Tyr
 195 200 205

Met Ala Glu Pro Phe Gly Lys Tyr Pro Lys Ile Glu Ala Gln Phe Arg
 210 215 220

Lys Ala Ala Gln Gln Pro Gly Lys Leu Phe Met Asn Tyr Val Ser Thr
 225 230 235 240

Ala Ala Leu Leu Pro Pro Arg Ser Asn Ala Asp Arg Leu Asn Pro Gln
 245 250 255

Val His Thr Phe Leu Asp Gly Ser Glu Ala Ala Gly Trp Thr Gly Leu
 260 265 270

Gly Ile Val Pro Leu Asp Tyr Pro Ala Thr Arg Pro Gly Leu Val Glu
 275 280 285

Ser Leu Ile Arg His Asn Pro Val Ala
 290 295

US 10,144,941 B2

149**150**

-continued

<210> SEQ_ID NO 44
 <211> LENGTH: 432
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 44

Val	Ser	Glu	His	Thr	Asn	Asn	Ala	Val	Val	Leu	Arg	Tyr	Gly	Asp	Asp
1								5	10					15	
Glu	Tyr	Thr	Tyr	Pro	Val	Ile	Asp	Ser	Thr	Val	Gly	Asp	Lys	Gly	Phe
	20				25					30					
Asp	Ile	Gly	Lys	Leu	Arg	Ala	Asn	Thr	Gly	Leu	Val	Thr	Leu	Asp	Ser
	35				40				45						
Gly	Tyr	Gly	Asn	Thr	Ala	Ala	Tyr	Lys	Ser	Ala	Ile	Thr	Tyr	Leu	Asp
	50				55			55	60						
Gly	Glu	Gln	Gly	Ile	Leu	Arg	Tyr	Arg	Gly	Tyr	Pro	Ile	Glu	Gln	Leu
	65				70			75				80			
Ala	Glu	Ser	Ser	Thr	Phe	Leu	Glu	Val	Ala	Tyr	Thr	Leu	Ile	Asn	Gly
	85				90			90			95				
Asp	Leu	Pro	Lys	Val	Asp	Glu	Leu	Ser	Ala	Phe	Lys	Asn	Glu	Ile	Thr
	100				105			105			110				
Gln	His	Thr	Leu	Leu	His	Glu	Asp	Val	Lys	Arg	Phe	Asp	Gly	Phe	
	115				120			120			125				
Pro	Arg	Asp	Ala	His	Pro	Met	Ala	Met	Leu	Ser	Ser	Val	Val	Ser	Ala
	130				135			135			140				
Leu	Ser	Thr	Phe	Tyr	Gln	Asp	Ser	His	Asn	Pro	Phe	Asp	Glu	Gln	
	145				150			150			155			160	
Arg	His	Leu	Ser	Thr	Ile	Arg	Leu	Leu	Ala	Lys	Leu	Pro	Thr	Ile	Ala
	165				170			170			175				
Ala	Tyr	Ala	Tyr	Lys	Lys	Ser	Ile	Gly	His	Pro	Phe	Val	Tyr	Pro	Arg
	180				185			185			190				
Asn	Asp	Leu	Gly	Tyr	Val	Glu	Asn	Phe	Leu	Arg	Met	Thr	Phe	Ser	Val
	195				200			200			205				
Pro	Ala	Gln	Glu	Tyr	Val	Pro	Asp	Pro	Ile	Val	Val	Ser	Ala	Leu	Glu
	210				215			215			220				
Lys	Leu	Leu	Ile	Leu	His	Ala	Asp	His	Glu	Gln	Asn	Cys	Ser	Thr	Ser
	225				230			230			235			240	
Thr	Val	Arg	Leu	Val	Gly	Ser	Ser	Gln	Ala	Asn	Met	Phe	Ala	Ser	Ile
	245				250			250			255				
Ser	Ala	Gly	Ile	Ser	Ala	Leu	Trp	Gly	Pro	Leu	His	Gly	Gly	Ala	Asn
	260				265			265			270				
Gln	Ser	Val	Leu	Glu	Met	Leu	Glu	Gly	Ile	Gln	Ala	Asn	Gly	Gly	Asp
	275				280			280			285				
Val	Asp	Ser	Phe	Ile	Gln	Lys	Val	Lys	Asn	Lys	Glu	Asp	Gly	Val	Arg
	290				295			295			300				
Leu	Met	Gly	Phe	Gly	His	Arg	Val	Tyr	Lys	Ser	Phe	Asp	Pro	Arg	Ala
	305				310			310			315			320	
Lys	Ile	Ile	Lys	Ala	Ala	Ala	His	Asp	Val	Leu	Ser	Ser	Leu	Gly	Lys
	325				330			330			335				
Ser	Asp	Glu	Leu	Leu	Asp	Ile	Ala	Leu	Lys	Leu	Glu	Glu	His	Ala	Leu
	340				345			345			350				
Ser	Asp	Asp	Tyr	Phe	Val	Ser	Arg	Asn	Leu	Tyr	Pro	Asn	Val	Asp	Phe
	355				360			360			365				
Tyr	Thr	Gly	Leu	Ile	Tyr	Arg	Ala	Met	Gly	Phe	Pro	Thr	Glu	Met	Phe
	370				375			375			380				

US 10,144,941 B2

151

152

-continued

Thr Val Leu Phe Ala Leu Gly Arg Leu Pro Gly Trp Ile Ala Gln Trp
 385 390 395 400
 His Glu Met Ile Lys Glu Pro Gly Ser Arg Ile Gly Arg Pro Arg Gln
 405 410 415
 Ile Tyr Thr Gly Glu Val Leu Arg Asp Phe Val Pro Val Glu Ser Arg
 420 425 430

<210> SEQ ID NO 45
 <211> LENGTH: 527
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 45

Met	Thr	Lys	Arg	Ala	Gly	Ile	Leu	Val	Ala	Val	Gly	Ala	Thr	Val	Ala
1				5					10						15
Gly	Leu	Val	Thr	Ala	Val	Pro	Ser	Ala	Ala	Ser	Thr	Ala	Pro	Gly	Ala
			20				25							30	
Pro	Gly	Ala	Ala	Ala	Pro	Leu	Lys	Trp	Thr	Ala	Cys	Gly	Thr	Lys	Ala
	35					40						45			
Tyr	Pro	Thr	Gln	Gln	Cys	Ala	Thr	Val	Arg	Ala	Pro	Leu	Asp	His	Asp
	50				55				60						
Arg	Pro	Ser	Gly	Arg	Gln	Val	Thr	Leu	Ala	Leu	Ala	Arg	Ile	Pro	His
	65				70				75					80	
Thr	Ala	Lys	Thr	Ser	Gln	Gly	Pro	Leu	Leu	Val	Asn	Pro	Gly	Gly	Pro
		85						90						95	
Gly	Gly	Ser	Gly	Leu	Ser	Met	Ala	Gly	Phe	Val	Ala	Ser	Ser	Leu	Pro
	100					105						110			
Ala	Lys	Leu	Ala	Ala	Gln	Tyr	Asp	Val	Ile	Gly	Phe	Asp	Pro	Arg	Gly
	115						120					125			
Val	Gly	Arg	Ser	Ser	Pro	Ala	Leu	Asp	Cys	Val	Pro	Lys	His	Phe	Asp
	130					135					140				
Pro	Val	Arg	Pro	Asp	Thr	Val	Pro	Gly	Ser	Pro	Arg	Asp	Glu	Arg	Thr
	145					150				155				160	
Asn	Arg	Glu	Arg	Ala	Ala	Ser	Phe	Ala	Asp	Ala	Cys	Gly	Glu	Lys	His
		165						170					175		
Gly	Asp	Leu	Leu	Pro	Phe	Met	Asp	Thr	Val	Ser	Thr	Ala	Lys	Asp	Leu
		180					185					190			
Asp	Val	Ile	Arg	Arg	Ala	Leu	Gly	Ala	Arg	Gln	Ile	Asn	Tyr	Phe	Gly
	195					200					205				
Tyr	Ser	Tyr	Gly	Thr	Tyr	Leu	Gly	Ala	Val	Tyr	Ala	Lys	Leu	Phe	Pro
	210					215					220				
Glu	Arg	Val	Arg	Arg	Leu	Val	Leu	Asp	Ser	Ile	Val	Asp	Pro	Asp	Gly
	225				230				235				240		
Val	Trp	Tyr	Glu	Asp	Asn	Leu	Gly	Gln	Asp	Tyr	Ala	Phe	Asp	Ala	Arg
		245						250					255		
His	Lys	Ala	Phe	Ala	Ala	Trp	Val	Ala	Lys	Asn	Asp	Ala	Thr	Tyr	Arg
		260					265					270			
Leu	Gly	Thr	Asp	Pro	Ala	Lys	Val	Glu	Ala	Ala	Trp	Tyr	Arg	Met	Arg
	275					280					285				
Ala	Ala	Val	Lys	Lys	His	Pro	Ala	Ala	Gly	Lys	Val	Gly	Pro	Ser	Glu
	290				295					300					
Leu	Glu	Asp	Thr	Phe	Leu	Pro	Gly	Gly	Tyr	Tyr	Asn	Gly	Tyr	Trp	Pro
	305				310				315				320		
Gln	Leu	Ala	Glu	Ala	Phe	Ala	Ala	Tyr	Val	Asn	Asp	Lys	Asp	Glu	Asp
		325					330					335			

Ala Leu Ala Thr Ala Tyr Asp Asp Phe Ala Ala Val Asp Ala Ser Gly
 340 345 350
 Asp Asn Gly Tyr Ser Val Tyr Thr Ala Val Gln Cys Arg Asp Thr Gly
 355 360 365
 Trp Pro Lys Ser Trp Thr Thr Trp Arg Asn Asp Thr Trp Gln Ala His
 370 375 380
 Arg Lys Ala Pro Phe Met Ser Trp Asn Asn Thr Trp Tyr Asn Ala Pro
 385 390 395 400
 Cys Ala Thr Trp Pro Val Ala Pro Leu Arg Pro Val Arg Val Thr Asn
 405 410 415
 Arg Glu Ile Pro Pro Ala Leu Leu Phe Gln Ala Thr Asp Asp Ala Ala
 420 425 430
 Thr Pro Tyr Glu Gly Gly Leu Ser Met His Arg Lys Leu Lys Gly Ser
 435 440 445
 Arg Leu Val Val Glu Glu Gly Gly Asn His Gly Ile Ser Leu Ser
 450 455 460
 Gly Asn Asp Cys Leu Asp Ala His Leu Ile Ala Tyr Leu Thr Asp Gly
 465 470 475 480
 Thr Leu Pro Arg Ser Gly Gly Ser Gly Ala Asp Ala Val Cys Asp Ala
 485 490 495
 Leu Pro Glu Pro Glu Ala Ala Ala Thr Ala Lys Ala Lys Ala Ala Thr
 500 505 510
 Gly Gln Lys Gly Ser Thr Leu His Ser Leu Leu Gly Phe Arg Gly
 515 520 525

<210> SEQ ID NO 46
 <211> LENGTH: 222
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 46

Met	Asn	Cys	His	Asp	Arg	Ile	Asn	Leu	Arg	Gly	Trp	Thr	Thr	Arg	Leu
1						5		10						15	

Ser	Gly	Leu	Phe	Val	Ala	Ala	Val	Leu	Cys	Leu	Leu	Pro	Trp	Thr	Gly
				20				25					30		

Thr	Ala	Glu	Ala	His	Gly	Ser	Val	Val	Asp	Pro	Ala	Ser	Arg	Asn	Tyr
						35		40				45			

Gly	Cys	Trp	Leu	Arg	Trp	Gly	Ser	Asp	Phe	Gln	Asn	Pro	Ala	Met	Ala
						50		55			60				

Gln	Glu	Asp	Pro	Met	Cys	Trp	Gln	Ala	Trp	Gln	Ala	Asp	Pro	Asn	Ala
						65		70		75		80			

Met	Trp	Asn	Trp	Asn	Gly	Leu	Tyr	Arg	Asn	Glu	Ser	Ala	Gly	Asn	Phe
						85		90				95			

Pro	Ala	Val	Ile	Pro	Asp	Gly	Gln	Leu	Cys	Ser	Gly	Gly	Arg	Thr	Glu
						100		105				110			

Gly	Gly	Arg	Tyr	Asn	Ala	Leu	Asp	Thr	Val	Gly	Ala	Trp	Gln	Ala	Thr
						115		120				125			

Asp	Ile	Thr	Asp	Asp	Phe	Thr	Val	Arg	Leu	Glu	Asp	Gln	Ala	Ser	His
						130		135			140				

Gly	Ala	Asp	Tyr	Phe	Arg	Val	Tyr	Val	Thr	Glu	Gln	Gly	Phe	Asp	Pro
						145		150		155		160			

Thr	Ala	Gln	Pro	Leu	Thr	Trp	Gly	Ala	Leu	Asp	Leu	Val	Ala	Glu	Thr
						165		170				175			

Gly Arg Tyr Gly Pro Ser Thr Ser Tyr Glu Ile Pro Val Ser Thr Ser

US 10,144,941 B2

155**156**

-continued

180

185

190

Gly Tyr Thr Gly Arg His Val Val Tyr Thr Ile Trp Gln Ala Ser His
 195 200 205

Met Asp Gln Thr Tyr Phe Leu Cys Ser Asp Val Asn Phe Gly
 210 215 220

<210> SEQ ID NO 47

<211> LENGTH: 1065

<212> TYPE: PRT

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 47

Val Ile Ser Arg Arg Arg Leu Leu Ser Thr Thr Ala Ala Thr Ala Ala
 1 5 10 15

Leu Ala Ala Val Ser Ser Pro Ala Ala Arg Ala Ala Ala Pro Ala Asp
 20 25 30

Thr Ala Ala Gly Arg Leu Arg Val Thr Gly Pro Thr Val Glu Tyr Val
 35 40 45

Arg Arg Pro Leu Gly Leu Asp Val Ser Arg Pro Arg Leu Ser Trp Pro
 50 55 60

Leu Ala Ser Asp His Pro Asp His Gly Gln Ser Ala Tyr Gln Val Arg
 65 70 75 80

Val Ala Thr Ser Pro Asp Arg Leu Ala Arg Pro Asp Val Trp Asp Ser
 85 90 95

Gly Lys Val Val Ser Pro Thr Ser Val Leu Val Pro Tyr Ala Gly Pro
 100 105 110

Ala Leu Val Ser Arg Thr Arg Tyr His Trp Ser Val Arg Val Trp Asp
 115 120 125

Gln Asp Gly Arg Val Ser Ala Trp Ser Glu Pro Ser Trp Trp Glu Thr
 130 135 140

Gly Leu Leu Asp Glu Ala Asp Trp Ser Ala Gly Trp Ile Gly Ala Pro
 145 150 155 160

Ala Ala Leu Thr Ser Ser Pro Ser Leu Glu Ala Ala Ser Trp Ile Trp
 165 170 175

Phe Pro Glu Gly Asp Pro Ala Val Gly Ala Pro Ala Ala Thr Arg Trp
 180 185 190

Phe Arg Gly Arg Val Glu Ile Pro Glu Gly Val Thr Arg Ala Arg Leu
 195 200 205

Val Met Thr Ala Asp Asp Gly Phe Thr Ala Leu Val Asp Gly Val Gln
 210 215 220

Val Ala Arg Thr Glu Pro Asp Gly Pro Ala Glu Asn Trp Arg Arg Pro
 225 230 235 240

Val Val Val Asp Val Thr Ala His Leu Ser Pro Gly Ser Arg Val Val
 245 250 255

Ala Val Thr Ala Thr Asn Ala Val Asp Gly Pro Ala Gly Leu Leu Gly
 260 265 270

Ala Leu Glu Leu Thr Thr Ala Asp Gly Ala Val Thr Leu Ala Thr Gly
 275 280 285

Thr Gly Trp Arg Ala Thr Asp Arg Glu Pro Asp Gly Asp Trp Ala Ser
 290 295 300

Gly Gly Tyr Asp Asp Thr Gly Trp Pro Ala Ala Val Leu Ala Pro
 305 310 315 320

Trp Gly Ser Gly Pro Trp Gly Glu Val Arg Ala Ala Leu Ser Pro Ala
 325 330 335

US 10,144,941 B2

157

-continued

Thr Gln Leu Arg Thr Glu Phe Arg Leu Gly Arg Lys Arg Val Ala Arg
 340 345 350
 Ala Arg Leu Tyr Ser Thr Ala Leu Gly Leu Tyr Glu Val Phe Leu Asn
 355 360 365
 Gly Ala Arg Val Gly Glu Asp Arg Leu Ala Pro Gly Trp Thr Asp Tyr
 370 375 380
 Arg Lys Arg Val Gln Tyr Gln Thr Tyr Asp Val Thr Ala Leu Leu Arg
 385 390 395 400
 Ser Gly Gly Asn Ala Leu Gly Val Thr Leu Ala Pro Gly Trp Tyr Ala
 405 410 415
 Gly Asn Ile Ala Trp Phe Gly Pro His Gln Tyr Gly Glu Arg Pro Ala
 420 425 430
 Val Leu Ala Gln Leu Glu Val Thr Phe Thr Asp Gly Ser Ile Glu Arg
 435 440 445
 Val Leu Ser Gly Thr Gly Trp Ala Ala Ala Thr Gly Pro Val Thr Ala
 450 455 460
 Thr Asp Leu Met Ala Gly Glu Glu Tyr Asp Ala Arg Leu Glu Thr Asp
 465 470 475 480
 Gly Trp Ser Arg Ala Gly Phe Asp Ala Ser Gly Trp Leu Ala Ala Glu
 485 490 495
 Ala Val Glu Gly Val Thr Ala Val Pro Val Ala Ala Val Asp Gly Ala
 500 505 510
 Cys Arg Val Glu Arg Glu Leu Thr Ala Arg Glu Val Thr Glu Pro Glu
 515 520 525
 Pro Gly Val Tyr Val Phe Asp Leu Gly Gln Asn Met Val Gly Thr Val
 530 535 540
 Arg Leu Leu Val Ser Gly Pro Ala Gly Thr Thr Val Arg Leu Arg His
 545 550 555 560
 Ala Glu Val Leu Asn Pro Asp Gly Thr Leu Tyr Thr Ala Asn Leu Arg
 565 570 575
 Thr Ala Arg Ala Thr Asp Thr Tyr Thr Leu Arg Gly Gly Pro Glu
 580 585 590
 Thr Tyr Glu Pro Arg Phe Thr Phe His Gly Phe Arg Tyr Val Glu Val
 595 600 605
 Thr Gly Phe Pro Gly Arg Pro Gly Pro Asp Ala Val Val Gly Arg Val
 610 615 620
 Ile His Thr Ser Ala Pro Phe Thr Met Ala Phe Ser Thr Asp Val Pro
 625 630 635 640
 Met Leu Asp Arg Leu His Ser Asn Ile Thr Trp Gly Gln Arg Gly Asn
 645 650 655
 Phe Leu Ser Val Pro Thr Asp Thr Pro Ala Arg Asp Glu Arg Leu Gly
 660 665 670
 Trp Thr Gly Asp Ile Asn Val Phe Ala Pro Thr Ala Ala Tyr Thr Met
 675 680 685
 Glu Ser Ala Arg Phe Leu Gly Lys Trp Leu Gln Asp Leu Arg Asp Asp
 690 695 700
 Gln Leu Ala Asp Gly Ala Phe Pro Asn Val Ala Pro Asp Leu Pro Gly
 705 710 715 720
 Val Gly Ser Gly Ala Ala Gly Trp Gly Asp Ala Gly Val Thr Val Pro
 725 730 735
 Trp Ala Leu Tyr Gln Ala Tyr Gly Asp Val Arg Val Leu Glu Gln Ser
 740 745 750
 Trp Ser Ser Met Val Ala Trp Leu Glu Tyr Leu Gln Ala His Ser Asp

158

US 10,144,941 B2

159**160**

-continued

755	760	765
Gly Leu Leu Arg Pro Ala Asp Gly Tyr Gly Asp Trp Leu Asn Ile Glu		
770	775	780
Asp Glu Thr Pro Lys Asp Val Ile Gly Thr Ala Tyr Phe Ala His Ser		
785	790	795 800
Ala Asp Leu Thr Ala Arg Thr Ala Glu Val Leu Gly Lys Asp Pro Gly		
805	810	815
Pro Tyr Arg Thr Leu Ser Gly Arg Val Arg Asp Ala Phe Arg Ala Ala		
820	825	830
Tyr Val Gly Asp Gly Gly Arg Val Lys Gly Asp Thr Gln Thr Ala Tyr		
835	840	845
Val Leu Ala Leu Ser Met Asp Leu Leu Glu Pro Gly Asp Arg Ala Pro		
850	855	860
Ala Ala Asp Arg Leu Val Ala Leu Ile Glu Ala Lys Asp Trp His Leu		
865	870	875 880
Ser Thr Gly Phe Leu Gly Thr Pro Arg Leu Leu Pro Val Leu Thr Asp		
885	890	895
Thr Gly His Thr Asp Val Ala Tyr Arg Leu Leu Thr Arg Arg Thr Phe		
900	905	910
Pro Ser Trp Gly Tyr Gln Ile Asp Arg Gly Ala Thr Thr Met Trp Glu		
915	920	925
Arg Trp Asp Ser Val Arg Pro Asp Gly Gly Phe Gln Asp Ala Gly Met		
930	935	940
Asn Ser Phe Asn His Tyr Ala Tyr Gly Ser Val Gly Glu Trp Met Tyr		
945	950	955 960
Ala Asn Ile Ala Gly Ile Ala Pro Ala Ala Pro Gly Phe Arg Glu Ile		
965	970	975
Arg Val Arg Pro Arg Pro Gly Gly Val His Arg Ala Glu Ala Arg		
980	985	990
Phe Asp Ser Leu Tyr Gly Pro Val Thr Thr Arg Trp Thr Ser Asp Gly		
995	1000	1005
Gly Gly Phe Ala Leu Arg Val Val Leu Pro Ala Asn Thr Thr Ala		
1010	1015	1020
Glu Val Trp Val Pro Gly Gly Asp Gly Arg Ser Ser Val Arg Gly		
1025	1030	1035
Thr Ala Val Phe Leu Arg Arg Glu Asp Gly Cys Ala Val Phe Ala		
1040	1045	1050
Ala Gly Ser Gly Ile His Arg Phe Thr Ala Pro Ala		
1055	1060	1065

<210> SEQ ID NO 48

<211> LENGTH: 1170

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 48

atgccggagc	gtttcactcc	cactcctgag	gacaaggta	cgttcggtct	gtggaccgtg	60
ggctggcgaa	gcaacgaccc	gttcggtag	ccgacgcgtc	cggtgtggaa	cccggtggag	120
tccgtcgagc	ggctggcgaa	gttcggtag	cacgggtga	cgttccatga	cgacgactg	180
attccgttgc	ggtcggacga	ccgtgagcgg	gcgcggctgg	tccggcggtt	cagggaggcg	240
ctggagcgta	ccgggctcaa	ggtgccgtat	gcgacgcgat	acctgttcac	gcacccggtg	300
ttcaaggacg	gcgggttac	ctccaacgac	cgtgacgtgc	ggcggttgc	gctgacgcaag	360

US 10,144,941 B2

161

162

-continued

gtgatccgca acatcgatct cgccgggtggag	ctcggcgccgc agacgtatgt ggcctgggc	420
gggcgtgagg ggcggagtc cggtgccggcc aaggacgtgc	ggtcggccct ggacggatg	480
aaggaggcct tcgacctgct gggegactac	gtcaccgagc agggctacga cctgeggttc	540
gcgatcgagc ccaagccaa cgagccccgc	ggtgacatcc tgcgtgeccac gatcgggcac	600
gcgctggccct tcatcgagcg	cctggagcgc cccgagctgg tcggggtgaa cccggagacc	660
gggcacgagc agatggccgg	gctgaacttc ccccacggca tgcgcaggg cctgtggcg	720
ggcaagctct tccacatcga	cctcaacggc cagtccggga tcaagtacga ccaggactc	780
cgcttcggcg cccggtagcc	gcccggatcg ttctggctcg tggacccctt ggagacggcc	840
ggctgggacg gtcacgcca	cttcgacttc aagccggatc gcacccggacg catcgacggg	900
gtgtgggagt cccgcaagaa	ctgcatcgcc aactaccta tccctcaagga ggcgcggcc	960
gccttcggcg cccggccggc	cgtccaggag gcccctcaccc cctccggcct cgacgaaactc	1020
gcccggccca cccggacgca	cggcctcaag gcactcctcg cccggccac cgcctacgg	1080
gacttcgacg ccacccggc	cggcggaaacgc tccatggccct tcgaaggccct cgaccagctc	1140
gccccatggacc	accttcctcaa cgtccgctga	1170

<210> SEQ_ID NO 49

<211> LENGTH: 1968

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 49

atgacaacgcg cgctcagggc gacgcagggt	ttgcagtcca cgaaccaccc ccgtttgtcg	60
gacctcaccc gaggagcacc gttgagcact	aatcccccc gaagaagttc ccgtctcaga	120
tggagactcg gcccggggcg ggccacccgg	gccaaggcgg tcgcggggctt caccgcactg	180
ctgctgccgc tcgcccgcgt	ggtcggccctg gctccccgg cccaggccgc gacctcgccg	240
accggccacct acctaagaa	gtcggactgg ggcagcggct tcgaggggca gtggacggtg	300
aagaacacccg	gcaccacccgc cctgtccctcc tggacgatcg agtgggactt cccctccggc	360
accgcggctcg	gtccgcctcg ggacgcctcc gtgaccagct cccggccacca ctggaccggcc	420
aagaacacccg	gtctggaaacgg taaggctgccc cccgggtgcca gcatcagctt cggcttcaac	480
ggcacccggat	ccggctccccc caccggctgc aagctgaacg gtgcctccctg tgacggccgc	540
ggcacggtcc	ccggcggacag cgcggcgatc aagccggca ccccccacccgc gagcggcatc	600
accgcacccct	cggtgaagct ctccctggacgc gcagccaccc acgacaaggg catcaagaac	660
tacgacgtcc	tgcgcgacgg cgccaagggtc ggcacgggtca ccacgcacgc gtacaccgc	720
accggccctca	ccaaggccac ggactactcc tactccgtgc aggccccgca caccggccac	780
cagacccggac	cggtcagcgg cgccggggcc gtgcgcacca cggggcggaa cgacaacccg	840
ggccccggca	ccggcggacaa ggtcaaccc tcggacttca ccaactgggg cgtctacggg	900
cgcaactacc	acgtcaagaa cctgggtgacc tcgggctcg cccgagaagat cacgcacatc	960
aactacgcct	tccggcaacgt ccaggccggc aagtgcacca tcggcgactc ctacgcccac	1020
tacgacaagg	cctacaccgc cgaccagtc gtcgacggcg tcggccacac gtgggaccag	1080
ccgctgcgcg	gcaacttcaa ccagctgcgc aagctcaagg cgaagtaccc gcacatcaag	1140
gtgatctggt	cggtcggccgg ctggacccctgg tcggggccgt tcgggtgcgc ggcgcagaac	1200
ccggccggcgt	tcgcccagtc ctgctacgac ctgggtggagg accccccctg ggccgatgtc	1260
ttcgacggca	tcgacatcga ctgggagttac cccaaacgcct gccggccgtac ctgtgacacc	1320

-continued

```

agcggccccg ccgcgtgaa gaacctgtcc tccgcgtcc ggcgaagtt cgccgcaag 1380
aacctggta cgcgcgcgtat caccggggac ggctcgacg gggcaagat cgacggcc 1440
gactacgggg ggcgcgcgtca gtccctcgac tggtacaacg tggatgacgta cgacttctc 1500
ggcgcctggg aggccaagggt tccgacggcc ccgcactccc cgctgaacgc gtacggcc 1560
atccccgggg acggcttcaa ctccggccgc gccatcgccca agctgaaggc caagggcg 1620
ccggcctcgaa agctgtgtt cggcatcggt ttctacggcc gggctggac gggcggtgacc 1680
caggcggcac cggggggcac ccgcacccggc gggggggggc gcacgtacga ggccggcatc 1740
gaggactaca aggttcctcaa gaccagctgc ccggccaccc gcacgtatgc cgccacccgc 1800
tacgcgact gcccaccaa ctggtgagc tacgacaccc cggcgaccat cacctccaa 1860
atggcctggg cgaacagccca gggctcggtc ggtgcgttct tctgggagtt cagcgccgac 1920
acgcacaacg gcgagctgt gagegccatg gacagcgcc tcaactag 1968

```

<210> SEQ ID NO 50
<211> LENGTH: 591
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 50

```

atgcggaaaa gggcaagcgc ggcgtcata ggcctggcga tggccggcgt ctgcgttgc 60
gccaccagca gtgccagcag ccacggctac accgattccc ccatcagcag acagaagctg 120
tgtgccaacg gcacccgtac cggctcgcc aacatccagt gggagccgc gaggcgtcgag 180
ggcccgaaagg gcttcccgcc ggcagggtccg gggacggca agatctgcgc cggcggaaac 240
agctccctcg ccgcgtcga cggccgcgc gggggcaact gggccgcac ccagggtcacc 300
ggcggccagg gctacaactt ccgtggcag ttcaccgc ggcacgcac gaccgacttc 360
cggtactaca tcaccaagga cggctggac tccaccaagc cgctcaccag ggccgcctg 420
gagtgcagc cttcatgac ggtgcgtac gggaccgcg agcccccggc gaccctgacc 480
caccaggcga ccateccccac ccagaagtcc ggcaagcaca tcatcctggc cgtctggAAC 540
gtggctgaca cggccaaacgc gttctacggt tgctcgacg tgaagttctg a 591

```

<210> SEQ ID NO 51
<211> LENGTH: 1671
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 51

```

gtggccgccc tggggccgg cgccctgacc gtgaccggtc tggtcggcac cgcacaggcg 60
ccgcacatca acgtcgccaa gaacgcggg ttcgagacg ggcgtcgcc ctggacctgt 120
accggccggca cggccgcac cgtctccctcc cccgtgcacg gggctccgc cggccctcaag 180
gccaccccgaa gggccaggaa caacgcgaag tgcacccaga ccgtggccgt gaagcccaac 240
tccacctatg cgctcagttc ctgggtgcag ggcgggtacg cctacctcg ggcgagccgc 300
acggccacca cggacgttcc cacctggacc cccggcagca cggcgacgac ccagctgcgc 360
acgagcttca ccacccggcc gtccacccacc tgggtgcagg tctacacccca cggctggtag 420
ggccaggccgg cctactacgc ggacgacgtc ggggtcaccg gacccgacgg cggccggcggt 480
acggaggagc cggccggccgc gatccccggc gccccggcg gttctggccgt cggccaccacc 540
acgtccctct cggtgccct gtcgtggaaac ggggtctccg gggccacccgg ctacaccgtc 600

```

US 10,144,941 B2

165

-continued

166

tacggggacg	gcaccaaggc	gaccaccacc	accggcacct	ccgcgcacgg	gagcggcctg	660
gcccggaca	ccgcgtacca	gttctcggtg	agcgccacca	acgcccgg	tgagtccgtc	720
aggtcggcga	ccgtgagcgg	acgtacggcc	aagaaggacg	agaccggccc	gggcccctcg	780
acctccgtc	ccaagcacgc	cgtgacccgc	tactggcaga	acttcaacaa	cggcgcggcc	840
gtccagaagc	ttagcgacgt	gcccgcaac	tacgacatca	tcgcccgtctc	cttcgcggac	900
gcccgggta	ccccgggtgc	cgtcaccttc	aacctcgact	eggggggcct	gaacggctac	960
accgtcgccc	agttcaaggc	cgacatcaag	gccaagcagg	ccgggggcaa	gaacgtcatc	1020
atctccgtcg	gcggcgagaa	gggcacccgtc	tcggtaaca	gcgacgcctc	ggcgaacgcg	1080
ttcgcggact	cgctgtacac	gctgatccag	gagtacggct	tcaacggcgt	cgacatcgac	1140
ctggagaacg	gcctcaactc	cacctacatg	acgaaggccc	tcgggtcgct	gtcctcgaa	1200
gtgggctccg	gtctcgcat	cacgatggcg	ccgcagacga	tcgacatgca	gtcgacgtcg	1260
ggtagtact	tcaagacggc	gctcaacatc	aaggacatcc	tgaccgtcgt	caacatcgag	1320
tactacaaca	gcgggttcgtat	gctgggctgc	gacggcaagg	tctactcgca	gggctcggtg	1380
gacttctca	ccgcgtcgc	ctgcattccag	ttggagggcg	gcctcgcccc	gtcccaggc	1440
ggcctcggtg	tgccccctc	caccccgccgc	gcggggcagcg	gctacgtcgc	cccgtcggtc	1500
gtgaacgcgg	ccctggactg	cctggccaag	ggcacccgg	gcgggtccctt	caagccgtcc	1560
aggacgtacc	cggacatccg	tggtgcgtat	acctggtcga	cgaactggga	cgccacggcg	1620
ggcaacgcct	ggtccaaacgc	ggtcggcccg	cacgtccacg	gccttcggta	a	1671

<210> SEQ ID NO 52

<211> LENGTH: 888

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 52

tgatcgac	gcgtcatggg	cctgctcacc	gcgcgtggccg	cggtcgctgc	gacgtcg	60
ttctccccg	ccgcacggc	ctcgccggcc	acctgcgc	cggcctggaa	cgccctcg	120
gtgtacacgg	gcgggggctc	cgcctcgat	aacgggcaca	actggtcggc	gaagtgg	180
acgcagaacg	gcgtccggg	cacctcgac	gtctggccg	accaggcgc	ctgcgg	240
ggcggcggcg	gcacccgaccc	gaacccctcg	ggcttcgtcg	tcaagcgg	gcagtca	300
cagatgttcc	cgagccggaa	ctccttctac	acctacagcg	ggctcaccgc	cgcg	360
gcctacc	ccttcgccaa	cacccggcgc	gacaccgtga	agaaggcagg	ggcggcggcg	420
ttcctcgcca	acgtcagcca	tgagacccgc	ggcttgtcc	acatcg	tgga	480
gcacactacc	cgcactactg	cgacaccgc	cagtccatcg	gtgc	ccggc	540
gcctactacg	gcgcggccccc	catccagctc	agctgaaact	tcaactaca	ggcggccgg	600
gacgcctcg	gcatcgaccc	gctggcaac	ccctggcagg	tggagcaga	ccgc	660
gccttggaa	ccggcctctg	gtactggaa	accaggcc	gccccggc	acatgac	720
cacaacgc	tcgtcaacgg	ctccggattc	ggtagacca	tccggccat	caacggc	780
atcgagtgc	acggcggcaa	ccccggccag	gtccagagcc	gcgtcaacac	ctacc	840
ttcgtccaga	tcctcggtac	cacggccggc	tcgaacctga	gctgctga		888

<210> SEQ ID NO 53

<211> LENGTH: 1524

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

US 10,144,941 B2

167

-continued

<400> SEQUENCE: 53

atgagacgt cactatccgt ccgcgcgtg gtgacggcg gggttcaccac ggtggcccg	60
gcaggcatgg ccgtgtggg ctccggcacc gcccaggcg cgaccccgct gcccgaccac	120
gtcttcggcc cctacttcga gtctgtggacc ggagagagcc cggggccat ggccggccgag	180
tccggggcga aacacctgac catggcggtt ctccagacga cggccaaggg ctctgcacg	240
cctgactgga acggcgacac cggcgtcgat atcgcccagg cgtcttcgg cggcgacatc	300
gacacgtacc agggccggagg cggcgacgtc atccccgtgt tccggggcta caccggcgac	360
accacccggca cggagatcgc cggacgtgc acccgacgtc accagatcgc cggggctac	420
cagaagggtcg tcacgacgta cggacgtctcg cggctcgaca tggacatcga ggtcgactcc	480
ctcgacgaca cccggggat cggccgggg aacaaggccca tcaagaagct ccaggactgg	540
ggggacgsga acggccgtga cctggagatc ttctacacgc ttccgacgac caccggcgga	600
ctggcctcca gggccgtcgc cgtgtcgcc aacggccgtga ccaacggggc acgggtcgac	660
gtcgtgaacc tgatgacgtt cgactactac gacaacgcgt cccacgacat ggccggcgac	720
accgagacgg cccggccaggg cctgtacgac cagctcgca agctgtaccc gggcaggacc	780
gcacacccagc tgggtccat ggtggcggtc accgagatgc cggcgctgaa cggactcgcc	840
ccggccgaga ctttcacgct cggcaacgcg gcccgggtgt acgactgggc ggtggccaag	900
ggcatcaaca ccctgtcctt ctggcgctc cagcgacaca acggccggctg cccggggc	960
ccggccgccc acgactgtc cggcatccag cagaacacccat gggacttac ccgcgtttc	1020
ggcccttca ccageggcac cacggcgccg gacgacgact tctcggtgac ggccacgccc	1080
gcctccggga cggtgaccgc ggggggttcg gccaccacca cggtaagac cggcggtgacc	1140
aaggggcgccg cacagcaggt cggcgtcaeg gtcagcgaaa tccggccgg tggcaccgcc	1200
tccctcagcc ctttcgtgtt gacggcgccc ggccggtaa cgttcaccc cggccacgacc	1260
caggccgccc ttcgggcac gtacggatc agcgttaccc gtagcgccccc gtcggccagc	1320
cacgcgtacgg ctttacacgct gacggtcacc ggccggccacg gcagccatgt cacggccggg	1380
ccgtggccgg gggggacggc ctacacccgc ggccagcagg tctcgtaaa gggccacacc	1440
tggaaaggccca agtgggtggac gacggggcgag gagcccgccaa ccacccgtga gtggggcgtc	1500
tggcaggacc tggggcgctg ctga	1524

<210> SEQ ID NO 54

<211> LENGTH: 1452

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 54

gtgacgcagg gaccgtcac cacggaggcc ggcgcgcgg tagccgacaa ccagaacagt	60
gagaccgcag gccccgggtgg accgggttctc gttcaggacc aggcgtttct ggagaagctg	120
gcccacttca accgggagcg catccggag cgcgtcgatc atgccccggg agccggcg	180
tacggcgttgc ttcacgttgc cggacgttc tccgtgtggaa cggcgatggaa gttctctcg	240
gaggtcgca aggagacccga gacccgttccg cgttctccaa cggcgccggg caaccctccg	300
tccggccgacg cggcggtgtga cccggcgccg tggcgctgaa agttctacac cgaagaggc	360
aactacgacc tccgtcgccaa caacaccccg gtgttcttca tcaaggacgc catcaagttc	420
cccgacttca tccacacccca gaagcgccac cggtaacacgg gttcccgagga ggcggacaac	480

168

US 10,144,941 B2

169

170

-continued

gtctggact tctggggct gtcgccgaa tccacccacc aggtgacctg gctttcggt	540
gaccgcggca tccgggcctc gttecgtaatc atgaacggct acggctcgca cacgttccag	600
tggaacaacg aggccggcga ggtttctgg gtcaagtacc acttcaagac cgaccaggc	660
atcaagaacc tcaccaccga ggaggccgta cgctctccg cggtcgaccc ggacagccac	720
cagcgcgatc tgcgtgagtc catcgagcgc ggtgacttcc cgacctggac ggtcaggtc	780
cagatcatgc cggcgccgaa ggcggccaaatc taccgcttca acccggtcga cctgaccaag	840
gtgtggccgc acgaggacta cccgcccata gagatggca agctggagct caaccgcaac	900
ccggagaaca tcttcggcga ggtcgagcag tcatcttca gcccggcga ctctcgatcc	960
ggcatacgcc cgtccccggaa caagatgtc caggccgccc tgttcgcttca cggcgcaccc	1020
caccgctacc gcgtcgccat caacgcccac cacctggccg tgaaccgtcc gcacgcccacc	1080
gaggcgcgtt ccaacagccg tgacggctac ctgtacgacg gcccgcacaa gggcacgaaag	1140
aactacgacg cgaacagctt cggcgccccg gtccagaccc acaggccgct ctggcagccc	1200
gtctccgtca cccggcggtac gggcaaccac gaggccgccc tccacgcccga ggacaacgac	1260
ttcgtgcagg cccgcaatct ctaccggctg atgtcgagg acgagaaggg ccggctgtatc	1320
gacaacctgg cccgggttcat cgcgaagggtg tgcgcgcacg acatcgccga tgcgcgcac	1380
aacaacttcc gtcaggccga cgcggacttc ggcaagcggc tggaggtcgc ggtccaggcc	1440
ctgcgcggct ga	1452

<210> SEQ ID NO 55

<211> LENGTH: 2781

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 55

gtgtatgcca tgccctccac cggccctcgcg ggggtccagt cggagagga cgctcccg	60
cgttcaagcc ccagaccctt cggccctcgctg ctggccggcgc tccacgtac cgcagggtt	120
tcactcatcg gaacccctgc cgtggcgccgc tccgacgagg cacctgtgc gacagaagca	180
tcggatgtgt ccatagccgc ggacacctac acctggaaaga acgccccgat cgacggccgc	240
ggcttcgtcc cccggatctgt cttcaacccgg tccgagaaga acctcgccata cggccggacc	300
gacatcgccgc ggcctaccgc ctgggaccag tccggcaagc agtggaaagcc cctgtggac	360
tgggtggact gggaccgctg gggctggacg ggcgtggta gcctcgccctc cgacacggc	420
gaccccgaca acgtgtacgc cggcggtgggg acgtacacca acagctggga cccgaccgac	480
ggcgccgtcc tgcgtccctc ggacggggccgc ctctctggaa aggccggccac cctcccg	540
aagctcgccgc gcaacatgcc cggacgcccgc atggggggacg ggctcgccgtt cgacccgaa	600
aagaactccg tgctctaccc gggcgccccc agcggcaacg gcctctggcg gtccaccgac	660
gggggagtcg gctggccgaa ggtacggccctt ttcggccaaatc cggggaaacta cgcgcaggac	720
cgcgtcgacca ccagcggtca cggcaacgcac aaccaggccatc tgcgtcggtt gacccgtac	780
gagcgttccg gcagcgccgg cagcgccacc caggacatct acgtcggtt cggccacaag	840
gagaacaccgc tctaccgctc cacggacggc ggcggccaccc ggtcgccgtt cccggccag	900
ccacccggctt acctcgccca caagggcgta ctgcactccg cgaccggccca cctctatct	960
acgctgacgcg acacggccgg cccctacgc ggcggcaagg ggcggatctg ggcgtacgac	1020
acggcgcccg ggcgtggca ggacgtcagc cccgtggggaggccgc ctactacggc	1080
ttcagcgccgc tctccgtggaa cccggcagaag cccggccaccc tgcgtggccac cgcctacac	1140

-continued

tcctggtggc ccgacaccca gatctccgc tccacggaca gcggtgccac ctggaccag 1200
 gcctggact acacccgcta cccgaaccgc tccaaaccgt acacgctgga cgtctctcc 1260
 gtgccgtggc ttcctgggg cgcttcccc gcaccgccc agaccgccc gaagctggc 1320
 tggatgacgg aggccgtgga gatcgaccgg ttcgactcg accggatgtat gtacggcacc 1380
 ggagcgacgg tctacggcac cgaggaccc acgtcctggg actccggcgg cacgttcagg 1440
 atcacccca tggtaaggg gatcgaggag acggccgtca acgacctggc cagccggccc 1500
 tccggggcac cgttgttag cgcaactggt gacatgggg gttccggca caccgaccc 1560
 gacgcgtgc cggacctgtat gtacacctcc cccgaacctcg actcgaccac cagccgtggac 1620
 ttccggaga gtcgccccgg cacggctgac cgggtcggca actccgacgc cgcgeccac 1680
 atcggttctt ccacccgacaa cggggccaaac tggttccagg gtcggagcc ttccggcgtc 1740
 accggggcg gCACGGTGC GGCAGGCGCG gacggcagcg gttcgtgt gaggccggag 1800
 ggccggggcg tccaccacac cacggccttc ggcacccctc ggaccggcc caccggcatc 1860
 cggccggtg ccacggtcga gtccgaccgg aagaaccccg agaagttcta cggattcgag 1920
 gggggcacct ttcacgttcc gacggacggc gggggcgaccc tcaaccggca ggccaccggg 1980
 ctggccggcg agggcaacgt ccgttccag gcaactggcc ggacggaggg cgacatctgg 2040
 ctccggggcg gtcggacac cggggcgtac ggtctgtggc gtcaccggca ctccggggcg 2100
 acgttacga agtccggcg cgtegagcag gggacacgc tggttccgg caaggccgcc 2160
 cggggccct cgtacggac ggttccgtc agcgcgaaga tggccgggg ggcggccatc 2220
 ttccggtcca ccgaegccgg ggcgagctgg accaggatca acgacgacgc ccaccagtgg 2280
 ggctggaccg ggcggccgtat cacggccgac cccagggtct acggggcgct ctacgttcc 2340
 accaaacgggc gcgggatcca ggtggggcag acctccgaca gggccggccg aggacggac 2400
 cccggccaccc atccggcac cgtacccggc accgatcccg gtccggagca gcccggccac 2460
 gccgcctgtg cgggtacgta cgcggtcacc aaccagtggc cggggccgtt ccaggccgat 2520
 gtgacggta ccaacacggg tgaecggcgg tacaacggc ggaagetcgg ctggtcgttc 2580
 cccggggggc agcagatctgg ccagatctgg aacgcctcgc accggcagga cgggggtgaag 2640
 gtcacccgtca cggacggccgg ctggAACGGC acgggtggccg cggcgtcgac ggccggcttc 2700
 ggcttcaccg gcaagttgggc gggagcaac gccaacccgg ccgccttac cctggacggc 2760
 caggccgtca ccgtggctg a 2781

<210> SEQ ID NO 56

<211> LENGTH: 1632

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 56

atgcgcggtg ccaagagcgc caagtgggtc gggggagcgg caatcatcgc cctggccgc 60
 accgcctgtg gtggggcga cagcgacagc gacaacggtg ccaagggcgc cgtcgacgc 120
 gagggcatat ttcctgtca ggtcggttag cccgagaacc cgctgcagcc ggccaacacg 180
 atggagtcga acggcagcat cgtacccgac gccatcttc cgcagctcg cgtactacgc 240
 cccgacggca agctcgagat gatcaacggc gagtccgtcg agacgaccga cagcaagctg 300
 tggacggta agctcaagaa ggactggaa ttccacgacg gcaccccccgt caccggcacc 360
 tcctacgtca aggccctggaa ctggccgcg aacatcgaga acgcgcagac gaacgcctcc 420

-continued

tggttcgccg acatcaaggg ctacgcccac gtccaccccg acggcgaggg cgccaagccg	480
aagtccgacg ccatgtccgg cctgaagaag gtggacact acacacctac catcgagctc	540
aactcggccg tccccgtactt ctcgtacaag ctccgtaca cggtcttctc gccgtcgccc	600
gagtccttct acgcggaccc gaaggccgcc ggtgagaagc cggtcggcaa cggcgctac	660
aagttcgta gctgggacca caagaaggcag atcaaggctcg tccgcaacgc cgactacaag	720
ggcccccaca aggcgaagaa cggtggtgtg atcttcaaga actacaccac cctcgagacc	780
gcctacgagg acctaagtc cggcaacgtc gacgtgtcc gccagatcg cccgaaggac	840
ctcccggtct accgtgcccga cctcgaggac cgcgcgtgg acaaggccctt ctcccggtt	900
cagacgctcg gtgtcgccat gtacaccgc cagtggaaa acacggaccc gaaggctc	960
cagggccgtg cgtatggccat cgaccggac acgtacccca agacgggtct ccaggccacc	1020
cgcgagccgg ccacgggctg ggtcgccaaag ggcgtctcg gttaccagga gaacgtcgcc	1080
ggtgacgtca ccaagtacga cccggcgaag gccaaggccc tcatcaagga gggtggcggt	1140
gttccgggca acgagatctt catccagttc aacgcccacgc gcggccacaa ggagtggatc	1200
gaggccgtct gcaacagcat cacgcaggcc accggcgtca agtgcaccgg cgactcgaaag	1260
gcccacttcc aggccgaccc gaacgcccgc gacgccaacgc aggtgaagtc gttctaccgc	1320
agtggctggg tcctcgacta cccggtaaac gccaacttca tcagcgaccc gttccgcacc	1380
gggtcgcccg gcaacaacgg cttdttctcc aacaaggacc tcgacgcgaa gatcaaggcc	1440
gcggactccg cccggagccct cgacgattcg gtcaaggccct accaggagat cgagaaggag	1500
ctggtaact acatgcccag catcccgctc tggtaactaca aggtcaacgc cggctactcg	1560
gagaacgtca agaacgtgga ctacgcgcag gacggcgacc cgatcctgac cgaagtccag	1620
gtcatcaagt aa	1632

<210> SEQ ID NO 57

<211> LENGTH: 480

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 57

atgcagggcg accccgaggt cctcgagttc ctgaacgaac agctgaccgc cgaattgact	60
cccatcaatc agtacttcct gcacgcgaag atgcaggatc accgcggctg gaccaagctc	120
gccaaacaca cccggccga gtcgttcac gagatgaagc acgcggagat cctgaccgc	180
cggatctgc tgctggacgg cctgccaac tatcagcggc tggccacgt gcgggtggc	240
cagaccgtca cggagatgtt ccaggccac cggcaggctcg aggtcgaggc gatcgaccga	300
ctgcggcgcg gtgtcgatct gatgcgcgc aagagcaca tcacgtccgc caacatctc	360
gaacggatcc tggaggacga ggagcaccac atcgactatc tcgacacccca gctggagctg	420
atcgagaagc tcggggagcc gctctaccc gcccaggta tcgagcagggt cgagcttga	480

<210> SEQ ID NO 58

<211> LENGTH: 894

<212> TYPE: DNA

<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 58

atgagcccgat acacccgcac gcgcggacc ttctcacccg ggcctggc cgccgcacc	60
ggagtcgtcc tcgggtgtac gcccgcctc gccgcccccg cgagactct gggaccac	120
gactggatgg gggccctcgcc gactccacc cccgtcgac gctcacgat cccggcacc	180

-continued

cacaacgcgg gggcccgcta cggcggaccc tggaccgagt gccagaacac cacggtgcc	240
gagcagctcg gcageggcat ccgcttcgt gacgtgcgtt gcccgtcac cggcgcacgc	300
ttcgcgtatcc accacggcgc ctcgttccag aacctgtatg tccggggacgt cctcatcgcc	360
tgccggactt ccctggccgc gcacccgtcc gagacgggtc tgatgggggtt caaggcaggag	420
tactcggagg agagcgcacgc cgcgttccgg cagatctcg acctgttaccc cgacggcaag	480
ggctggccgc cgctttcccg cctcgaccgc accctggggg acctcgccgg cgccgggggc	540
aaggctgtgc tcctcgcgga caacggggc ctggccgggg tccggtagc cgacccggcg	600
gtcttcgaca tccaggacga ctatcgcc gagcccttcg gcaagtaccc caagatcgag	660
gcccggatcc gcaaggccgc ccagcagccc ggcaagctct tcatgaacta cgtgtccacc	720
gtgtccctgc tgccggccgc ctcgaacgcg gacgggtcata accccgcagggtt ccacacgttc	780
ctcgacggctt ccggaggccgc gggctggacc ggctcgaa tcttcccgctt ggactatccg	840
ggacccggcc ccggccctggt cgactcgctg atcaggcaca accccgggtggc ctga	894

<210> SEQ ID NO 59
<211> LENGTH: 1299
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 59

gtgagcgcgc acaccaacaa cgctgttagta ctgcggtagc ggcgtacgcgtt gtacaccatc	60
ccgggtatcg acagcaccgtt cgccgcacaa ggcttcgaca tcggggatctt ccggggccat	120
acggggctgg tcacgttggc cagggatac ggcaacaccgc cggcctataa atccggccatc	180
acctatctcg acggcgaaca gggcatctcg cgctaccgcg gctaccggat cgacgcgc	240
ggggagagctt cgacgttccctt cgaggtagc tacacgttgc tcaacggcgtt cttcccaag	300
gtcgacgcgc tgccggccctt caagaacggat atcaccggatc acacgttgcgtt gcacggaggac	360
gtcaaggcgctt tcttcgacgg cttccgcgc gacggccaccat cgatggccat gctgtccctcg	420
gtcgacgcgc cgctgtccac gttctaccag gacagccaca acccggttgcg cggaggac	480
cgtcacccctt cgacgttcccg gctgttggcc aagctcccgat cgatcgccgc gtacgttgc	540
aagaagtcga tcgggttccacc gttctgttac ccggcgtacgc acctcggttgc cgtcgagaac	600
ttccctgtcgca tgaccccttc ggtccggccat caggatgttgc tgccggaccat gategttgc	660
tcggcgctcg agaagctgttgc catctgttgc acggaccacgc agcagaactgtt tcggacccatc	720
accgtcgcttc tggtcggcttc ctcgcaggcc aacatgttgc cttccatctc cggccggatc	780
tcggcgctgtt gggggccgtt gcacgggtggc gccaaccatgtt gatgttggaa	840
ggcatccagg ccaacggccgg cgacgttgcac ttccatcc agaaggatcaa gaacaaggag	900
gacggcgtcc gcctgtatggg cttccggccat cgggtgtaca agtcccttgcg cccggccgc	960
aagatcatca aggccggccgc ccacgttgc cttcccttcgc tcggcaagtc cgacgttgc	1020
ctggacatcg cgctcaagctt ggaggaggac ggcgttcccg acgactactt cgtctcgcc	1080
accccttaccat ccaacgttggc cttctacacgc ggcgtatctt accggggccat gggcttcccg	1140
accggatgtt tcaccgttgc cttccgttcc ggcgttcccg cgggttggat cgttgcgtt	1200
cacggatgtt tcaaggagcc ggggtcccgatc atcggccgcgc cggccagat ctacaccggc	1260
gaggttccctgc ggcgttccgtt cccggccatc agccgttgc	1299

<210> SEQ ID NO 60

US 10,144,941 B2

177

178

-continued

<211> LENGTH: 1584
 <212> TYPE: DNA
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 60

atgacgaaac gtgcaggcat tctggtcga gtcggcgcca cggtcgcgg gctggtcacc	60
gccccgttccgtt cccggcgccc acccgccccc gggggccctgg gggccgcgc gccgcgtgaag	120
tggacccgtt gcggggacgaa ggccgtatccg acccagcagt ggcgcacccgt tcgcgcgc	180
ctggaccatg acaggccgtc aggacggcag gtcacgcctcg ccctcgcccg gatccgcac	240
acggcgaaga cctcgcaggg tccgcgtcg gtcacccccc gggggccccc cggcagcggg	300
ctctcgatgg ccgggttcgtt ggccgtcccg ctgcggcgca agctcgccgc ccagtacgac	360
gtgatcggtt tcgaccccgcg cggggtcggc aggagcagcc cggcgcttggc ctgcgttaccg	420
aagcacttcg accccggtacg ccccgacacc gtgcggcgat ccccgccggc cgagcggacc	480
aaccgggaac gcgcggcgcc ttccgcgcac ggcgtccggc agaagcacgg ggacctgtcg	540
ccgttcatgg acacgggttccg caccgcgaag gacccgtggc tgatccgcg ggcgcgttcc	600
gcacggcaga tcaactactt cggctactcc tacggcacctt acctggcgcc cgtctacgac	660
aagctgttcc cggagcggcgtt gcccggccgtt gtgcgtcgactt cgatcggttcc cccggacggc	720
gtctggtacg aggacaacctt cggccaggac tacggcttcg acggccgtca caaggcggttcc	780
gccccctgggg tggcgaagaa cggccgcacc tacggctcg gcacccgaccc ggcgaagggtc	840
gaagccgcctt ggtaccggat gccccggcggtt gtgaagaaggc accccgcggc gggcaagggtc	900
ggcccgagcg agctggagga cacccctcg cccggccgtt actacaacgg ctactggccg	960
caactggccg aggccgttccg cgcgtacgtt aacgacaagg acgaggacgc gctggccacg	1020
gcgtacgacg acttcgcggc ggtcgacgcg agcggggaca acggctactc cgtctacacg	1080
gccgtccagt gccgcgcacac gggctggccg aagtccgttcc acacccgttcc caacgcacacc	1140
tggcaggccg accgcgaaggc gcgcgttccatg tccctggaaaca acacccgttcc caacgcgc	1200
tgcgcacccgtt ggcccggttcc acccgctggc cccggctgggg tccaccaaccc cgagatcccg	1260
ccggcgctcc tttcccgatgc caccgcacgc gccggcgaccc cgtacgggg cggccgttcc	1320
atgcacccgc accgtcaaggc ctgcgcctcg gtcgtcgagg agggccggcc caaccacggc	1380
atcagccgttgc gcccggcaacgc ctgcgcgttgc ggcgcacccgtt tccctggaaaca acacccgttcc	1440
accctggcccg gctccggccg cggccggccgac gacccgttcc ggcgcgttcc ccccgagccg	1500
gaggccggccg cggccggccgaa ggcgaaggcc gctacggccg agaaggccgac caccctgcac	1560
aggccgttccg gcttccgggg ctgttcc	1584

<210> SEQ ID NO 61
 <211> LENGTH: 669
 <212> TYPE: DNA
 <213> ORGANISM: Streptomyces sp. ACTE
 <400> SEQUENCE: 61

atgaattgttc atgatcgcat caacttacgc ggctggacga cacggcttgcg cggctgttcc	60
gtcgcccgccg tgctctgtct gctcccgatgc acggggcaccgg ccgaggccca cggctcggttcc	120
gtcgaccccg cgtcccgcaa ctacggctgc tggctccgtt gggccaggcga cttccagaac	180
cccgccatgg cgcaggaaaga ccccatgtgc tggcaggcat ggcaggccga cccgaacgc	240
atgttggaaact ggaacggccct gtaccgcac gatccgcggc gcaacttccc ggcagtgtatc	300
cccgacgggc agctgtgttccg cggccggccgg accgaggccg gccggatccaa cggctggac	360

-continued

accgtggcg cctggcaggc cacggacatc acggacgact tcaccgtgag gctggaggac	420
caggccagcc acggcgccga ctacttccgg gtgtacgtca ccgagcaggg cttcgacccc	480
actgctcagc ccctgacactg gggcgactc gacctggtgg cggagacccg acgttacggt	540
cccaggtacga gctacgagat cccctgtgagt acgtcggggt acacccggcg ccatgtcgtc	600
tacacgatct ggcaggcctc gcacatggac cagacgtact tccgtgcag tgacgtgaac	660
ttcggctga	669

<210> SEQ ID NO 62
<211> LENGTH: 3198
<212> TYPE: DNA
<213> ORGANISM: Streptomyces sp. ACTE

<400> SEQUENCE: 62	
gtgatcagca gaagacgact gctcagcacc accggccgcca cccggccacct cgccgggtc	60
tcctcgcccg cccggccgccc cggccggcccg gccgacaccc cggccgggtcg gtcggcggtc	120
accggggcga cccgtggagta cgtacgccc cccgtcggtcc tccgttcccg cccggcccg	180
ctgagctggc ccctcgccctc ggaccacccg gaccacggc agtccgoccta ccagggtgcgg	240
gtcgccacct cccgggacccg cccggcccgccc cccgacgtct gggacacggg caagggtcg	300
tccccgacgt cgggtgttgtt cccgtacggc gggccggccg tgggttcccg tacggcgat	360
cactggtcgg tgcgtgtgtt ggaccaggac ggacgggtct cggccgtggag cgagccgtcc	420
tgggtggaga cccgggttccct ggacggggcc gactgggtgg cgggggtggat cggccggccc	480
ggccgcgtga cccctcacc cccctggag gggccctctt ggttcccg cccggaggcc	540
gatccggccg tggggcgccacc gggggccacc cgggtggttcc gggggccgggt ggagatcccc	600
gaggggcgta cccggggcccg cccgggttccctt accggccacg acgggttccac cggccgttcc	660
gacgggttcc accggggcccg tccggggccg gacggggcccg cggagaactg ggttcc	720
gtgggtgtgg acgtgacggc gcaacttccccc cccgggttccctt ggggtgtcg cgtgacggcc	780
accaacggc tggacggccc ggggggttccctt cccgggttccctt ggggtgttccctt cccggggcc	840
gggtgcgttca cactggccac gggaaaccggta tggggggccca cccggggggaa gccggacggg	900
gactggggcgta cccggggcccg cccgggttccctt ggggtgttccctt cccggggcccg	960
tgggggttccctt ggggggttccctt cccggggcccg cccggggcccg cccggggcccg	1020
acggaaattcc gggtggggccg caacggccgtc gggggggccg ggggttccctt cccggggcc	1080
ggcctgttccctt ggggggttccctt cccggggcccg cccggggcccg cccggggcccg	1140
tggaccgact accggcaagcg cgtccagtttccctt cccggggcccg cccggggcccg	1200
tccggggccca acgggttccctt cccggggcccg cccggggcccg cccggggcccg	1260
tgggttccctt cccggggcccg cccggggcccg cccggggcccg cccggggcccg	1320
ttcacccgacg ggttccctt cccggggcccg cccggggcccg cccggggcccg	1380
cccggttccctt cccggggcccg cccggggcccg cccggggcccg cccggggcccg	1440
gggtggggccg cccggggcccg cccggggcccg cccggggcccg cccggggcccg	1500
gtcaccggccg tgggggttccctt cccggggcccg cccggggcccg cccggggcccg	1560
ggccggggccg tgggggttccctt cccggggcccg cccggggcccg cccggggcccg	1620
gtggggcccg tgggggttccctt cccggggcccg cccggggcccg cccggggcccg	1680
ggcgaggtgc tgggggttccctt cccggggcccg cccggggcccg cccggggcccg	1740

US 10,144,941 B2

181

182

-continued

accgacacct acacgctca gggggcgga ccggagacgt acgagccccg cttcaccc	1800
cacggttcc gtaacgtcga ggtgacgggc ttccggggc gccccgggc ggacgcggtg	1860
gtggggcggg tcatecacac ctcggcgccg ttcaccatgg ctttctcgac cgacgtcccc	1920
atgctcgacc ggctccacag caacatcacc tgggggcagc gggcaactt cctctccgtc	1980
cgcaccaca cggccgcgcg cgacgaacgc ctggctgga cggcgacat caacgtctc	2040
gcccaccgc cgcgtacac gatggagtgc gcccgttcc tcggcaagtgc gtcaggac	2100
ctgcgcgacg accagctggc cgacggcgc ttcggaaacg tggccggga cttccggc	2160
gtcggcagcg gggggccgg ctggggcgac gccgggtga cgggtccgtg gggctgtac	2220
caggcgtacg gggacgtgcg ggtgctggag cagtcctggt cgtcgtatggt ggcctggctg	2280
gagtacctcc aggacgacag cgacggctc ctggggccgg cggatggta cggggactgg	2340
ctcaacatcg aggacgagac acccaaggac gtcatcgca cggcgtactt cgccacagc	2400
gccgacctca cggccggac cgccgagggtg ctggcaagg accccgggc ctaccgcacg	2460
ctgtccggcc gggtgccgcg cgggtccgg gccggcgtacg tggcgcacgg cgggggggtg	2520
aaggcgcaca cgcagaccgc gtacgtcctg gcccgtcga tggacctgtc ggagccggc	2580
gaccgcgcac cggctgcggg caggctggc gcgctgtacg aggcgaagga ctggcacctg	2640
tgcacggggg tccctggcac accgcgcctg ctggcggtgc tgaccgacac cgggcacacg	2700
gacgtcgctt accgggtgct gacgcggggg acgttcccgaa gctgggggtt ccagatcgac	2760
cgggggtgcac ccacgtgtg ggagcgtgg gactccgtgc ggccggacgg cgggtttcag	2820
gacgcggggg tgaactcctt caaccactac gcctacgggtt cgggtggcga gtggatgtac	2880
gcgaacatcg cggcatcgc cccggcgccg cccggcttcc gcgagatccg ggtgcgtccg	2940
cgtccggggg ggggggtgca cggggccgag gcccgggtcg actccctgtt cggggcggtc	3000
accacccgtt ggacctcgga cggggggggc ttgcgcgttc ggggtggctt gcccggcaac	3060
acgacggccg aggtgtgggt gccggggggt gacggggagga gtcgggtccg gggcacccgc	3120
gtgttctgc ggcgggagga cgggtgcgcg gtcttcggg cgggtcggtt catccacccgc	3180
ttcaccgcgc cggcctga	3198

<210> SEQ ID NO 63

<211> LENGTH: 1872

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: CelLcc_CBM3a DNA

<400> SEQUENCE: 63

atgggacatc accatcatca ccatcaccat gcatccgaaa acctgtactt ccaggcgatc	60
gcatggatc cgaacaatga cgactggctg catgttgaag gtaacaaaat agtggacatg	120
tacggtaatc aggtctggct gacggcgtgc aactgggtt gattcaatac cggtaaccaat	180
gtgttgtacg gagtatggag ctgcaatattt agagaagccc tcaagggtat ggcggacaga	240
ggaataaaatt ttttggaaat acctatttca acagaatttc tttatcaatg gtctcaagga	300
atatatccca aagcaaattttaatgt taatgatttt gtaaatccgg agctgaaagg aaagaacacgc	360
cttgagcttt ttgactttgc cggtcagtgc tgcaaaagaat tcggaaataaa gataatggtg	420
gtatatacaca gtccggcaac agatgccatg gggcatatgt atcctttatg gtatgacggt	480
caatttacaa cagagatatg gattcaact ttggagtggt tgacggaaag atataaaaat	540
gatgacacaa ttcttgcaact ggaccttaaa aatgagcctc acggcacccc gggcagcga	600

-continued

ttaatggcca aatgggatgg ttccacggat ttgaacaact ggaagcatgc tgctgaaaca	660
tgcgcaaga gaatccgtc aataaatccg aatatttcta ttgtggtaga aggagtggaa	720
gtttatccaa agcctggcta tgattatacc gcagtggacg aatggggaaa agagagtaaa	780
tatttctata actgggtgggg agggaaattta agaggagtca gggattatcc cattgacctt	840
ggcaaggcatc agaaggcagct tgtatactca cctcacgatt acggtcccct cgtagataaa	900
caaccttgggt tcttatgagg cttaacaaa gaaactttgt ataatgattt ctggagagat	960
aactgggcat acatacacga gggaaacatc gtccttgc tagtgggtga atggggaggt	1020
ttcatggacc gcggagacaa cgagaaatgg atgaaagcgc tgagagatata tatttttttt	1080
aataaaatat cccacactt ttggtgctat aatgcaaatt ccggtgatac cgaggactt	1140
gtataactatg atttttattac ctgggacgaa gaaaaatatg ctcttgcggatgcata	1200
tggcagacag aggacggaaa gttttaggc ctgaccatc agataccctt tggttcaat	1260
ggagggtttaa acgcgactcc cactaaaggt gccactccta ccaatacggc gactccgact	1320
aagtccggcaa cggcaacgcc cactcgcccc agcgtaccga ccaatactcc gactaatacc	1380
ccggcgaaca ccccgatcaag cggtaacctg aagggttgaat ttataactc caacccaaagc	1440
gacacaacga atagcatcaa tcggcagttc aaagtacgaa acactggcag ttcagctatc	1500
gatctgtcga aactgaccct tcggtactac tatacggttt atggccaaa agatcagacc	1560
ttttggtgcg accatgcage aatcatcggt agcaatgggtt cttataacgg cattacttct	1620
aatgtaaaag gcacccctgt gaagatgtca agtagccacca acaatgctga tacctactg	1680
gaaattagct tcacgggtgg cacacttggaa ccaggagccc acgtccagat ccagggccgt	1740
tttgcggaaa acgattggag caactatacg caatcaaacc attatagtt caaaagcgcg	1800
tctcaattcg tagaatggga tcaggtgacc gcatatttga acggagtgtt gggttgggg	1860
aaagaaccag ga	1872

<210> SEQ ID NO 64

<211> LENGTH: 624

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: CellLcc_CBM3a Amino acids

<400> SEQUENCE: 64

Met	Gly	His	His	His	His	His	His	Ala	Ser	Glu	Asn	Lle	Tyr
1													
								10				15	

Phe	Gln	Ala	Ile	Ala	Met	Asp	Pro	Asn	Asn	Asp	Asp	Trp	Lle	His	Val
					20		25					30			

Glu	Gly	Asn	Lys	Ile	Val	Asp	Met	Tyr	Gly	Asn	Gln	Val	Trp	Lle	Thr
					35		40					45			

Gly	Cys	Asn	Trp	Phe	Gly	Phe	Asn	Thr	Gly	Thr	Asn	Val	Phe	Asp	Gly
					50		55					60			

Val	Trp	Ser	Cys	Asn	Met	Arg	Glu	Ala	Lle	Lys	Gly	Met	Ala	Asp	Arg
					65		70		75			80			

Gly	Ile	Asn	Phe	Lle	Arg	Ile	Pro	Ile	Ser	Thr	Glu	Lle	Lle	Tyr	Gln
					85		90					95			

Trp	Ser	Gln	Gly	Ile	Tyr	Pro	Lys	Ala	Asn	Val	Asn	Phe	Val	Asn
					100		105					110		

Pro	Glu	Lle	Lys	Gly	Lys	Asn	Ser	Lle	Glu	Lle	Phe	Asp	Phe	Ala	Val
					115		120					125			

US 10,144,941 B2

185**186**

-continued

Gln Cys Cys Lys Glu Phe Gly Ile Lys Ile Met Val Asp Ile His Ser
130 135 140

Pro Ala Thr Asp Ala Met Gly His Met Tyr Pro Leu Trp Tyr Asp Gly
145 150 155 160

Gln Phe Thr Thr Glu Ile Trp Ile Ser Thr Leu Glu Trp Leu Thr Glu
165 170 175

Arg Tyr Lys Asn Asp Asp Thr Ile Leu Ala Leu Asp Leu Lys Asn Glu
180 185 190

Pro His Gly Thr Pro Gly Ser Glu Leu Met Ala Lys Trp Asp Gly Ser
195 200 205

Thr Asp Leu Asn Asn Trp Lys His Ala Ala Glu Thr Cys Ala Lys Arg
210 215 220

Ile Leu Ala Ile Asn Pro Asn Ile Leu Ile Val Val Glu Gly Val Glu
225 230 235 240

Val Tyr Pro Lys Pro Gly Tyr Asp Tyr Thr Ala Val Asp Glu Trp Gly
245 250 255

Lys Glu Ser Lys Tyr Phe Tyr Asn Trp Trp Gly Gly Asn Leu Arg Gly
260 265 270

Val Arg Asp Tyr Pro Ile Asp Leu Gly Lys His Gln Lys Gln Leu Val
275 280 285

Tyr Ser Pro His Asp Tyr Gly Pro Leu Val His Lys Gln Pro Trp Phe
290 295 300

Tyr Glu Gly Phe Asn Lys Glu Thr Leu Tyr Asn Asp Cys Trp Arg Asp
305 310 315 320

Asn Trp Ala Tyr Ile His Glu Glu Asn Ile Ala Pro Leu Ile Val Gly
325 330 335

Glu Trp Gly Gly Phe Met Asp Arg Gly Asp Asn Glu Lys Trp Met Lys
340 345 350

Ala Leu Arg Asp Tyr Met Ile Glu Asn Lys Ile Ser His Thr Phe Trp
355 360 365

Cys Tyr Asn Ala Asn Ser Gly Asp Thr Gly Gly Leu Val Tyr Tyr Asp
370 375 380

Phe Ile Thr Trp Asp Glu Glu Lys Tyr Ala Leu Leu Lys Pro Ala Leu
385 390 395 400

Trp Gln Thr Glu Asp Gly Lys Phe Ile Gly Leu Asp His Gln Ile Pro
405 410 415

Leu Gly Ser Asn Gly Gly Leu Asn Ala Thr Pro Thr Lys Gly Ala Thr
420 425 430

Pro Thr Asn Thr Ala Thr Pro Thr Lys Ser Ala Thr Ala Thr Pro Thr
435 440 445

Arg Pro Ser Val Pro Thr Asn Thr Pro Thr Asn Thr Pro Ala Asn Thr
450 455 460

Pro Val Ser Gly Asn Leu Lys Val Glu Phe Tyr Asn Ser Asn Pro Ser
465 470 475 480

Asp Thr Thr Asn Ser Ile Asn Pro Gln Phe Lys Val Thr Asn Thr Gly
485 490 495

Ser Ser Ala Ile Asp Leu Ser Lys Leu Thr Leu Arg Tyr Tyr Tyr Thr
500 505 510

Val Asp Gly Gln Lys Asp Gln Thr Phe Trp Cys Asp His Ala Ala Ile
515 520 525

Ile Gly Ser Asn Gly Ser Tyr Asn Gly Ile Thr Ser Asn Val Lys Gly
530 535 540

Thr Phe Val Lys Met Ser Ser Ser Thr Asn Asn Ala Asp Thr Tyr Leu

-continued

545	550	555	560
Glu Ile Ser Phe Thr Gly Gly Thr Leu Glu Pro Gly Ala His Val Gln			
565		570	575
Ile Gln Gly Arg Phe Ala Lys Asn Asp Trp Ser Asn Tyr Thr Gln Ser			
580		585	590
Asn Asp Tyr Ser Phe Lys Ser Ala Ser Gln Phe Val Glu Trp Asp Gln			
595		600	605
Val Thr Ala Tyr Leu Asn Gly Val Leu Val Trp Gly Lys Glu Pro Gly			
610		615	620

<210> SEQ ID NO 65
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic - 14 bp palindromic promoter sequence

<400> SEQUENCE: 65

tgggagcgct cccca

14

<210> SEQ ID NO 66
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic

<400> SEQUENCE: 66

gggagcgctc cca

13

We claim:

1. A microbial host cell comprising at least one exogenous nucleic acid molecule encoding a *Streptomyces* sp ActE (SActE) enzyme, wherein said enzyme is selected from the group consisting of:

SActE_0237 (GH6),
SActE_0236 (GH48),
SActE_3159 (CMB33),
SActE_0482 (GH5),
SActE_0265 (GH10),
SActE_2347 (GH5),
SActE_0357 (CE4),
SActE_0358 (GH11),
SActE_1310 (PL3),
SActE_3717 (GH9),
SActE_4638,

and wherein the enzyme is expressed in the microbial host cell.

2. The microbial host cell of claim 1, wherein enzyme: SActE_0237 (GH6) comprises the amino acid sequence SEQ ID NO: 1,

SActE_0236 (GH48) comprises the amino acid sequence SEQ ID NO: 2,

SActE_3159 (CMB33) comprises the amino acid sequence SEQ ID NO: 3,

SActE_0482 (GH5) comprises the amino acid sequence SEQ ID NO: 4,

35 SActE_0265 (GH10) comprises the amino acid sequence SEQ ID NO: 5,
SActE_2347 (GH5) comprises the amino acid sequence SEQ ID NO: 6,
SActE_0357 (CE4) comprises the amino acid sequence SEQ ID NO: 7,
40 SActE_0358 (GH11) comprises the amino acid sequence SEQ ID NO: 8,
SActE_1310 (PL3) comprises the amino acid sequence SEQ ID NO: 9,
SActE_3717 (GH9) comprises the amino acid sequence SEQ ID NO: 10,
45 SActE_4638 comprises the amino acid sequence SEQ ID NO: 11,
SActE_4738 (GH16) comprises the amino acid sequence SEQ ID NO: 12,
50 SActE_4755 (GH64) comprises the amino acid sequence SEQ ID NO: 13,
SActE_5457 (GH46) comprises the amino acid sequence SEQ ID NO: 14,
55 SActE_5647 (GH87) comprises the amino acid sequence SEQ ID NO: 15, and
SActE_5978 (PL1) comprises the amino acid sequence SEQ ID NO: 16.

3. The microbial host cell of claim 1, wherein enzymes

SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159

(CMB33), SActE_0482 (GH5), and SActE_3717 (GH9) are

expressed in the microbial host cell from the at least one

exogenous nucleic acid molecule.

4. The microbial host cell of claim 1, wherein enzymes SActE_0265 (GH10), SActE_0357 (CE4), SActE_0358

(GH11), and SActE_5978 (PL1) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

189

5. The microbial host cell of claim 4, said microbial host cell further comprising an exogenous nucleic acid molecule encoding xylose isomerase SActE_5230.

6. The microbial host cell of claim 1, wherein enzyme SActE_2347 (GH5) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

7. The microbial host cell of claim 1, wherein enzyme SActE_1310 (PL3) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

8. The microbial host cell of claim 1, wherein chondroitin/alginate lyase SActE_4638 is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

9. The microbial host cell of claim 1, wherein enzyme SActE_5647 (GH87) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

10. The microbial host cell of claim 1, wherein enzymes SActE_4738 (GH16) and SActE_4755 (GH64) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

11. The microbial host cell of claim 1, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), SActE_0265 (GH10), SActE_2347 (GH5), SActE_0358 (GH11), SActE_1310 (PL3), and SActE_3717 (GH9) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

12. The microbial host cell of claim 1, wherein said microbial host cell is selected from the group consisting of *Streptomyces lividans*, *Trichoderma reesei*, *Saccharomyces cerevisiae*, and *Escherichia coli*.

13. A *Streptomyces* sp. ActE strain host cell comprising at least one exogenous nucleic acid molecule encoding a *Streptomyces* sp. ActE (SActE) enzyme, wherein said enzyme is selected from the group consisting of:

SActE_0237 (GH6),
 SActE_0236 (GH48),
 SActE_3159 (CBM33),
 SActE_0482 (GH5),
 SActE_0265 (GH10),
 SActE_2347 (GH5),
 SActE_0357 (CE4),
 SActE_0358 (GH11),
 SActE_1310 (PL3),
 SActE_3717 (GH9),
 SActE_4638,
 SActE_4738 (GH16),
 SActE_4755 (GH64),
 SActE_5457 (GH46),
 SActE_5647 (GH87), and
 SActE_5978 (PL1),
 and wherein the enzyme is expressed in the *Streptomyces* sp. ActE strain host cell.

14. The *Streptomyces* sp. ActE strain host cell of claim 13, wherein enzyme:

SActE_0237 (GH6) comprises the amino acid sequence SEQ ID NO: 1,
 SActE_0236 (GH48) comprises the amino acid sequence SEQ ID NO: 2,
 SActE_3159 (CBM33) comprises the amino acid sequence SEQ ID NO: 3,
 SActE_0482 (GH5) comprises the amino acid sequence SEQ ID NO: 4,
 SActE_0265 (GH10) comprises the amino acid sequence SEQ ID NO: 5,
 SActE_2347 (GH5) comprises the amino acid sequence SEQ ID NO: 6,

190

SActE_0357 (CE4) comprises the amino acid sequence SEQ ID NO: 7,
 SActE_0358 (GH11) comprises the amino acid sequence SEQ ID NO: 8,

5 SActE_1310 (PL3) comprises the amino acid sequence SEQ ID NO: 9,
 SActE_3717 (GH9) comprises the amino acid sequence SEQ ID NO: 10,

SActE_4638 comprises the amino acid sequence SEQ ID NO: 11,
 SActE_4738 (GH16) comprises the amino acid sequence SEQ ID NO: 12,

10 SActE_4755 (GH64) comprises the amino acid sequence SEQ ID NO: 13,
 SActE_5457 (GH46) comprises the amino acid sequence SEQ ID NO: 14,

SActE_5647 (GH87) comprises the amino acid sequence SEQ ID NO: 15, and
 SActE_5978 (PL1) comprises the amino acid sequence SEQ ID NO: 16.

15 15. The *Streptomyces* sp. ActE strain host cell of claim 13, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), and SActE_3717 (GH9) are expressed in the *Streptomyces* sp.

20 SActE strain host cell from the at least one exogenous nucleic acid molecule.

16. The *Streptomyces* sp. ActE strain host cell of claim 13, wherein enzymes SActE_0265 (GH10), SActE_0357 (CE4), SActE_0358 (GH11), and SActE_5978 (PL1) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

25 17. The *Streptomyces* sp. ActE strain host cell of claim 16, said *Streptomyces* sp. ActE strain host cell further comprising an exogenous nucleic acid molecule encoding xylose isomerase SActE_5230.

30 18. The *Streptomyces* sp. ActE strain host cell of claim 13, wherein enzyme SActE_2347 (GH5) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

35 19. The *Streptomyces* sp. ActE strain host cell of claim 13, wherein enzyme SActE_1310 (PL3) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

20. The *Streptomyces* sp. ActE strain host cell of claim 13, 40 wherein chondroitin/alginate lyase SActE_4638 is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

21. The *Streptomyces* sp. ActE strain host cell of claim 13, 45 wherein enzyme SActE_5647 (GH87) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

22. The *Streptomyces* sp. ActE strain host cell of claim 13, 50 wherein enzymes SActE_4738 (GH16) and SActE_4755 (GH64) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

23. The *Streptomyces* sp. ActE strain host cell of claim 13, 55 wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), SActE_0265 (GH10), SActE_2347 (GH5), SActE_0358 (GH11), SActE_1310 (PL3), and SActE_3717 (GH9) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

24. An animal feed comprising the microbial host cell of 60 claim 1.

25. An animal feed comprising the *Streptomyces* sp. ActE strain host cell of claim 13.

191

26. A method for digesting a lignocellulosic material, said method comprising exposing the lignocellulosic material to the microbial host cell of claim 1, wherein the exposed lignocellulosic material is at least partially digested by the microbial host cell.

27. The method of claim 26, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), and SActE_3717 (GH9) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

28. The method of claim 26, wherein enzymes SActE_0265 (GH10), SActE_0357 (CE4), SActE_0358 (GH11), and SActE_5978 (PL1) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

29. The method of claim 28, wherein said microbial host cell further comprises an exogenous nucleic acid molecule encoding xylose isomerase SActE_5230.

30. The method of claim 26, wherein enzyme SActE_2347 (GH5) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

31. The method of claim 26, wherein enzyme SActE_1310 (PL3) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

32. The method of claim 26, wherein chondroitin/alginate lyase SActE_4638 is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

33. The method of claim 26, wherein enzyme SActE_5647 (GH87) is expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

34. The method of claim 26, wherein enzymes SActE_4738 (GH16) and SActE_4755 (GH64) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

35. The method of claim 26, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), SActE_0265 (GH10), SActE_2347 (GH5), SActE_0358 (GH11), SActE_1310 (PL3), and SActE_3717 (GH9) are expressed in the microbial host cell from the at least one exogenous nucleic acid molecule.

36. A method for digesting a lignocellulosic material, said method comprising exposing the lignocellulosic material to the *Streptomyces* sp. ActE strain host cell of claim 13,

192

wherein the exposed lignocellulosic material is at least partially digested by the microbial host cell.

37. The method of claim 36, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), and SActE_3717 (GH9) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

38. The method of claim 36, wherein enzymes SActE_0265 (GH10), SActE_0357 (CE4), SActE_0358 (GH11), and SActE_5978 (PL1) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

39. The method of claim 38, wherein said *Streptomyces* sp. ActE strain host cell further comprises an exogenous nucleic acid molecule encoding xylose isomerase SActE_5230.

40. The method of claim 36, wherein enzyme SActE_2347 (GH5) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

41. The method of claim 36, wherein enzyme SActE_1310 (PL3) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

42. The method of claim 36, wherein chondroitin/alginate lyase SActE_4638 is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

43. The method of claim 36, wherein enzyme SActE_5647 (GH87) is expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

44. The method of claim 36, wherein enzymes SActE_4738 (GH16) and SActE_4755 (GH64) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

45. The method of claim 36, wherein enzymes SActE_0237 (GH6), SActE_0236 (GH48), SActE_3159 (CBM33), SActE_0482 (GH5), SActE_0265 (GH10), SActE_2347 (GH5), SActE_0358 (GH11), SActE_1310 (PL3), and SActE_3717 (GH9) are expressed in the *Streptomyces* sp. ActE strain host cell from the at least one exogenous nucleic acid molecule.

* * * * *