



US008790875B2

(12) **United States Patent**
Khatib

(10) **Patent No.:** **US 8,790,875 B2**
(45) **Date of Patent:** **Jul. 29, 2014**

(54) **METHODS AND COMPOSITIONS FOR
IMPROVED FERTILIZATION AND
EMBRYONIC SURVIVAL**

(75) Inventor: **Hasan Khatib**, Fitchburg, WI (US)

(73) Assignee: **Wisconsin Alumni Research
Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/563,750**

(22) Filed: **Aug. 1, 2012**

(65) **Prior Publication Data**

US 2013/0041209 A1 Feb. 14, 2013

Related U.S. Application Data

(60) Provisional application No. 61/514,030, filed on Aug.
2, 2011.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
A61B 17/43 (2006.01)

(52) **U.S. Cl.**
USPC **435/6.1**; 435/6.11; 435/6.12; 600/33;
600/34

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Zhang B. et al. Journal of Dairy Science vol. 94 No. 8, 2011, pp.
4174-4182.*

GenBank locus NM_001050332 (Sep. 12, 2013) "Bos taurus DnaJ
(Hsp40) homolog, subfamily C, member 15 (DNAJC15), mRNA."
from www.ncbi.nlm.nih.gov, printed pp. 1-2.*

GenBank locus NW_003104311 (Jun. 3, 2010) "Bos taurus breed
Hereford chromosome 12 genomic scaffold, alternate assembly
Bos_taurus_UMD_3.1, whole genome shotgun sequence.", and
BLAST alignment output, from www.ncbi.nlm.nih.gov and blast.
ncbi.nlm.nih.gov, printed pp. 1-7.*

* cited by examiner

Primary Examiner — Stephen Kapushoc

(74) *Attorney, Agent, or Firm* — Kening Li; Pinsent Masons
LLP

(57) **ABSTRACT**

Single nucleotide polymorphic sites of the bovine HSP genes
are associated with improved fertilization rate and/or
improved embryo survival rate in cattle. Nucleic acid mol-
ecules, kits, methods of genotyping and marker assisted
bovine breeding methods based on these SNPs are disclosed.

16 Claims, 7 Drawing Sheets

83821 tagcaattca gatccttgatg tgaaaatgtc agtgcctccc tcaagggtgg aaagctaaact
83881 atagctcctt gaggccacc cttccattgc tgattcattt cctgcttccc ttttccttct
83941 caaatgggtg gtgtgatato agtaaaaggc ccagataaag tcaggacagt gcatacgcca
84001 ctgctgtcta gatgttttgt gtgtgtgtgt ggtttgttgt gagccactca ggtcctaaga
84061 tgttttcaac agctgtggat catttgtact caagggggag aataaaaaat aaaaaacaaa
84121 cacagagact ggtgagctgg actgttagct tgagtgcata gcatgaaagt gttcaggact
84181 agaagttaga aggtgtatga tgcttaataa tttttagtgat gacagtttgg ggccagtatt
84241 tttatatgta agggagctcc tgtttgtcat ttttaacgat tctgaatttg aaattcacct
84301 cagttttcct ctggttcttt acgttagaaa taattcattg tgaagaatta tgaaaactga
84361 cttcatttta ttttaaggcg agtgacgctg tctgtgctcc ataagtattc tgatggcact
84421 gtgtcagaga gcagggggca taccctttta cgttattttt attatcactg attttttatc
84481 atattattta ctaatattga ccaatattag tgaaatgctt ttagagctga gtagtatata
84541 gttgcctttt taacttattt ttatgaaaat gtatggtttg ataaaacttt gaaggcacc
84601 ttggattgct tcacctcctt cagaaaagata acttccttca tagaagaatg
(DNAJC15-6F)
84661 aaaaaaaca agaaaaaatt acgtcttggg gaaggactgt tattttattt acctaa
84721 tctg gtcagatatt tggcatttga aaggtaatgt ctgaccagtg
84781 aaatgtttga gtgtgtttta aattactcct gatcataagt ttttttttct cagagattat
84841 ccttgactgt taggtatctg tagtgcttca ttgaacgagt ttaacttata ctgttgtgat
84901 ttctagagag cttataaaat aatttgtcag ggcatgaatg tgtgggaaaa tatcagagac
84961 atctaactga gtgtttttct ttatttgaga ctttaaaaaa tgtcagtagt gaagtgtcct
85021 ttaatccatt ttcttttctt tgtcctctca ggtgggtctc cttacttagc aacaaaaatt
85081 aatgaagcaa aagacctgct agaagcaacc accaaacact gatggacgt caaggaccac
85141 tctgaaggag aagaagggg gctttataaa agctgtcctg cacattagt c taaaccgtgg
85201 cattcacaaat ttcatatgtg caccgaccac agccatttct tccgtgtctg agctatgtga
85261 taataaaaaga tgaacagtct agctttgtat tgcctttcag aggagcacat ggaa
(DNAJC15-6R)
85321 ctca ttttcttctt ctctctgata gtcatacatt tgaagacttc
85381 tactactaag ttcccccgac cccagccctt gcagcagtat cattatcagt aatataattg
85441 gaattttgga gcaagagtgt agaatttttt gctgcttata tttatgctcg cattgtccct
85501 tgtttcttca aaggaattta gcactctgcc tatgtgaagt gctttaaatg ttatgagacg
85561 aaaccctgac tgccaccoca ggaagtgaa ctgaagcttg atggaattta ctaaagggtg
85621 ttttgtaaaag cagtggtag gaaagaaagc tgctctccca gagatcccaa ggctgtagca
85681 tgagaacttg gagagctgct aaaatgatta aaggtttaca gaatgagccc tgtgaggaaa
85741 gcctggaat tgctgggaga agtctgaggg attcgtactt aattatatag tcttctggtg
85801 ctgtgtactt attcaattat gcattctaca aatattttatt taccacctgt agcgggtgag
85861 gtactgtagg ggtcctcagg agttccagg gtgaaccagg cacggatcgt gacccaaagg
85921 aacttgaatt tagaggaaag ataaaccaca ggagtactat tagagaggca cagatgatag
85981 gcaaaagagt tcaaaataaa agactatttt acactgggca tcaggggggc gtgctataga
86041 atcagttagc tagagctggg ctttaaaaaa aatgagatat accatatgat gggtagtca
86101 gaattagcag aaatagggtg tattagtcca ttggagtga gggtttctcc aaagtaacgg
86181 atgaacacac agtctggaag ggctggccct gtaatgtgca gtgcattcaa tttttttttt
86221 tttccccctc aacacagccc ctggttatca tcaattttct aaaaaggaga aaagagagag
86281 tcagaatacc agcatatttc ttgtactgct ttacgacgt gggctcatcc caggtaaaca
86341 gaactaccac gtttcatctg tcttaaacgc tcttgtaata gtctagaggc atctttcatt
86401 ttacgacatc ctgtagttag gagttctcag cattggtgat actgactcat tgagctgctg
86461 attcctgctt gtggggggct ctctgtgtga tggtaggatg attagcaaca tccctgctg
86521 cacctacgaa aggccaggag cgccccctct gttgtgacaa ccgaaaatgt atctagatgt
86581 tgccaaacgt tcccttgagg gcaaaattgc cccaggtaga gagccactgc tctaaatggt
86641 aaccgtgttc tcccttggtg catgaaataa tggatggcct gacttgaaat tgattcttag
86701 gttcactgaa atacggctct cgttttaaaa tcccctgtgt gtcacagttt tttgtgcttt
86761 agtatgtttt ctggccact gctgcccgg taccttaagt gcctctctga taccagatc
86821 tccacgtgtt caaagcctgg ggatagttga aggccaagct gaaaagtctg tctctccttt
86881 tctggtgctt ccagcaaaag taatcccatg aattgtcagc tctcagcaaa caaggattgt
86941 atctgtcttg gtcactgcta cattgacact caataaatat ttgttaaatg aacgaatata
87001 ttcccgtagc actttattct tcacatttgg taacttgtga aacaactctt tgtgtgacat
87061 gacttatttc ttctacaaaa gtacaagaac ctttagagca agaattctgt catttgtctt
87121 tgtaacctcc ttaagagtgg agcacttggg actttgtaga catcataaag ttgtaagtac
87181 taaaaaaaat gttttttgaa tactatgctt caaaaaatgt aaccactagt tttgaaagtt

Figure 1


87241 atgttatgca atattgggggt accctcagaa catacctggt agttgacatc attttctagt
87301 ccacactgtg ttcataaaaa tttaagtcct cagcactagc ttgaaatttc cttcattcta
87361 ttctcattta tctgttgagt gtgtgcccac aaatcagaca cctggtttta agtactatcc
87421 tcctatcaca taatctgggg caattaatct gcaaggcttt aattgcctca tctataaaat
87481 agagaaaata ataaccctat ctaaaaaggt ccatctagtc aaggctatgg tttttccagt
87541 ggtcatgtat ggatgtgaga gttggactgt gaagaaggct gagcgccgaa gaattgatgc
87601 ttttgaactg tgggtgttga gaagactcct gagagtcct tggactgcaa ggacatccaa
87661 ccagtccatc ctaaaggaga tcagtcctgg gtgttcattg gagggactga tgctgaagct
87721 gaaactccaa tactttggcc acctcatgcg aagagctgac tcattggaaa agacctgat
87781 gctgggaggg gttgggggca ggaggagaag gggacgacag aggatgagat ggctggatgg
87841 catcaccaac tcgatggatg tgagtttggg taagctccgg gagtgggtgc tggacaggga
87901 ggctggcat gctgcgattc atggggttgc aaagagtcgg acacgactga gtgactgaac
87961 tgaactgaaa aaggttgtt

Figure 1 (continued)

34081 ttgctcagtt gtgcccgaact ctttgcaacc ccatggactg tagcctacta ggttccttgc
34141 tccatgggat tctccaggca agagtaccag agtgggttgc catttccttc tccagaggat
34201 cttcctgacc cagggatcga acccaggtct cccgcattcc aggcagacac tttaacctct
34261 gagccactag ggatgggaag agggagacag ttatttggtca cccttgattt gacaagcgtt
34321 actgaggatg ctttaaacac ctgacgaaat tcagatgacc ttgtagcgt aagttaattg
34381 gtggctttttg gggttccca ggtggcacag tggtaaagaa tctgcctgcc catgcaagag
34441 atgcaagaga cttgggttcg atccctgggt ggggaagatc ctctgaagaa gggaatggcc
34501 acccactcca gtattcttgc ctggagaatt ccatggacag aggagcctgg tgggctacag
34561 gccatagggt tgcaaagagt cgggcacaac tgagcaacta atattacttt tactattgta
34621 tatcaaattc agcttttttg ttgttttgc ctttccctta tggaccctgc tccctggctg
34681 actttgtccg ctgtttgctt ctggaactac atgccgtgtc tctgttccct cctggcctc
34741 tgccattttct tccctgcgg ctctgtgttt cccatcaccc gatcaattcc tgaaccccat
34801 cttcttgcag agtctctctc cagggggggcc ttgcctgat ggctcttct cctcatcgct
34861 tccatcagca attatttgcg gagtgcctaa tgcctgtgag gcgctaacca ctaattttca
34921 gttgattatg ttgtataact cactttgtat tgtctgtttt tctacttgaa tgttacatat
34981 tatttgtcca ttgtttgttt cccagtcctt tagtctaacc tttgagaaag ggtcacgata
35041 atttcagata tgcttaactg ~~ttgtttgttt cca~~ ggcagctgaa atgcagtga
35101 cagagggttg ggctgacctc cctactgaca tagagctctg ctttttcttt gcagggatga
35161 agtcaataaa gcatatcgga agcttgctgt gctgcttcac cctgacaaat gtgtagcacc
35221 tggtagtgaa gatgccttca aagcctgtgt gaatgcccg acagccctcc tgaaaaaacat
35281 caagtagaaa ataggaaaaa aagacaggtg taggcctcga gtccaaacag actttctctg
35341 gaagtaaaat agccaacatg ggtttttcct ccacaaaatt cttacagccc ttttactca
35401 cgtgctgtca tttgtgaatc agtaatttgg ctgcctgttt ccatgtcaca gacattttgc
35461 agagagacaa aatgaagaga accgagtgcc acttatacac tgtcgttcat tttctatttc
35521 tgaatgatga tagatttttt tttttttctc atgagttggt aagctgtcgg tatagcatca
35581 cttgggcact ttgctcagga aagtcacaaa agtttctgga agaaggatag gaggatcctt
35641 ttgctttttc tttattaaaa aaacattcat catcagagga aaaaaaaca aaaatggaag
35701 ~~cagatccttg agatccttg aagtgagggt~~ gactttacaa gcaagttgcc cttttacaat
35761 gagttttctaa ggtgatattt taagcatcag tctgcttaat taatgaatga actcttggca
35821 gcccttcgga agtgaaacaa ggtgatgcta tcttgagctg agtggccctt ttttgacggt
35881 cgtctctgtc ccttctctca tcaccggggg agagaccagc cagtccctgag agtagagggtg
35941 gcggtgaccg caggttctgg tagactggct gctgctagac ctgcccctgg gtctgaggcc
36001 tttccctggc cgtgggagtg ctggtagctg taggggcac ttcacctag agtctggagt
36061 cagatcccaa gacacaaatg cctgcaggtc aaactggatt gtgtacacat tcccagaggg
36121 cctgaccctg ctggttcggg cgtgtctgat gccctaggag agcccccata ggaggcttat
36181 ~~gcccttccttg ccttccttg ccttccttg ccttccttg~~ gcaggg ttgctctca
36241 acaggggggc tcccttcgct tctctgctcag tgtgagctcc ccttccctct tctgttctctg
36301 ccttgtctct caaggcaccc catctgccac cagctctctc ttctcgaatc cccaccttcc
36361 ggctattgcc tctctgtctc tgtgcagcct ttgctggggc ctggtagacc agagacaagt
36421 atagatcctt ggggtggttg atggagagat ctttaaggcct gtgtgggaag aatttgttac
36481 ttttgtttag gttccaaatt tctgatcagt aatactttta agctgcagat tatgaaaatt
36541 aatttcacat agagaaatcc attctcttgc tatcgtctgt ctctcagtat ccaagagatg
36601 gatgaagatc aagtgtcagg accggccttc cttccagagc tcacagaata gagtaccacac
36661 gtggcgtggt acaccatcac tctcacgtta ctacagattgt tagctgtgac caagctgcac
36721 taattggggc tttaaagtat atgaagcttt gccatcagtg aattttaagc agctatccca
36781 tattgattgc caagtag ~~ga~~ agaaaaacca ctttcattta
36841 attcaaaccc actgcagctt actgaacct tcccttctcc agagcagctt cgctttgtgt
36901 ggctactctt gggcagcact gagctggttg tcaggaaacc agagcattgt tttctagtaa
36961 gagttttcat ttgatatta aaaatgcttt ttttcttag taaaattgag atatgagata
37021 tgattttttg tgggattttc ctcttgccct ttaattttaa ctgactctg ggccgtgaaa
37081 actcaggcca aagtctctca gggttttaat ttgtctgttg gccttcacatc tggccaaagt
37141 aacaaacata gatctgttct tgtggcagaa ctttaagatc acatgagtga ggaactgtgc
37201 atgggtggtg cagccacat tgacatgaca gctcacttgt gtatgtggcc ctgtgtaaaa
37261 actggtgcca gttttcatca ctgtcattaa acactagatg ggtcacctca ttcagctaca
37321 ttgtgcagat gageagttat agcagataga gttctctagt tcttccattc acatccttat
37381 gtaatgagtg acttggtgac atttagcctg gtcttgatt ~~tttctctctg~~
37441 ~~tttctctctg~~ cacttga ttactaatgt taaggatggc catgtactac ttcatttatg gaaaagaaag
37501 aaaagaaaga taaaggttca gggctcctaga attttctga tttccataat cctgagtaga
37561 atgtcaagaa gtaatgaaat gaagcctgat ctgcaatgaa aagtaacttc atataaaatt

Figure 2

Figure 2 (continued)

**Figure 3**




Figure 4




Figure 5

1

METHODS AND COMPOSITIONS FOR IMPROVED FERTILIZATION AND EMBRYONIC SURVIVAL

STATEMENT OF GOVERNMENT INTEREST

This invention was made with government support under 09-CRHF-0-6055 awarded by the USDA/NIFA. The government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to a method of genetic testing for improved fertilization rate and embryonic survival rate in animals, especially dairy cattle, based on single nucleotide polymorphisms (SNPs) in genes encoding the heat shock proteins (HSP).

BACKGROUND OF THE INVENTION

Poor reproductive performance, in particular in high-producing dairy cows, is a major problem on dairy farms throughout the world and has been identified as the single most important problem in dairy herd management in many countries (Royal et al., 2000; Dobson et al., 2008). In addition to direct financial losses, infertility can result in increased management complexity, for example, an inability to achieve a compact calving pattern, which is of critical importance in maximizing milk production from grazed grass in seasonal production systems.

This is of particular interest in cattle because of declining fertility over the past few decades (Dobson et al., 2007; Leroy et al., 2008). Furthermore, recent studies have shown that low fertilization rates and embryonic loss seem to be the main factors in dairy cattle infertility (Santos et al., 2004; Morris and Diskin, 2008).

While reproductive performance is influenced by a large number of factors, low fertilization rate and early embryonic loss are the primary factors contributing to poor reproductive performance in dairy cattle (Santos et al., 2004; Morris and Diskin, 2008). Enormous efforts, such as animal breeding and artificial insemination, have been and continue to be invested in ensuring adequate fertility in the cattle herd. Typically, artificial insemination in dairy cattle is successful only 30-35% of the time. The reasons for this are not clear. However, it is understood that both biological and environmental factors affect fertility rate. Some environmental factors such as high temperature, and lack of precipitation can cause stress in cattle and can drop the fertility rate to 10-15%. Commercial artificial insemination operations often shut down in July and August due to the drop in fertility caused by the hot, dry weather.

Genetics is also a prominent factor in fertility, and accounts for about one-third of the decline in pregnancy rate of dairy cows (Shook, 2006). In particular, identifying highly fertile bulls has been a time-consuming and expensive process. It can take 5-10 years of tracking the attempts of artificial insemination using semen from a bull before it can be certified as a quality bull.

Marker-assisted selection, on the other hand, can lower the high cost and reduce the extended time commitment of progeny testing currently used to improve sires, since young bull progeny could be evaluated immediately after birth or even prior to birth for the presence/absence of the marker, and young bulls that are determined by genetic testing to have undesirable markers would never be progeny tested.

2

There is thus a need for a method of genetically evaluating the bulls, as well as the cows, e.g., by genetic testing, to enable a quick and accurate evaluation of its fertility as well as the survival rate of embryos conceived therefrom.

Heat shock proteins (HSPs) are among the most highly conserved proteins in nature and have been found in all organisms studied from bacteria to humans (Becker and Craig, 1994). The structure and roles of HSPs as molecular chaperones in the folding, transport, and assembly of proteins as well as in protecting the cell under different stress conditions have been extensively studied and reported in the scientific literature (Becker and Craig, 1994; Nollen and Morimoto, 2002; Qiu et al., 2006). The fact that HSPs are essential in the folding, stability, and cellular localization of newly synthesized proteins implies key roles of these proteins in apoptosis, cell differentiation, and regulation of the embryo cell cycle (Luft and Dix, 1999; Lanneau et al., 2007). Strong evidence which has accumulated on the expression of HSPs during spermatogenesis, oogenesis, and embryogenesis suggests they have important functions in fertilization and during the pre-implantation period (Neuer et al., 1999). Mouse embryos cultured with monoclonal antibodies to HSPs have been found to display a significantly reduced blastocyst rate (Neuer et al., 1998). Al-Katanani and Hansen (2002) reported that the addition of antibodies for the induced form of HSP70 reduced blastocyst rate of cattle embryos, suggesting that HSP70 is involved in proper embryonic development. Matwee and colleagues (2001) showed that fertilization—measured as the number of spermatozoa tightly bound to a zona pellucida—and embryo development in cattle were significantly affected by the presence of different concentrations of anti-HSP70.

Although there is strong evidence in the literature on the roles of HSPs in early embryonic development, most of the studies have focused on the mouse and only a few HSP genes have been studied in cattle embryos. No evidence existed that any polymorphism in HSPs, if existed, is related to dairy cattle fertility in any way.

SUMMARY OF THE INVENTION

The present inventor investigated the expression profiles of HSPs and their splice variants in bovine embryos (degenerates vs. blastocysts), and also analyzed the association of these profiles with fertility traits. Some splice variants showed differential expression between degenerates and blastocysts while others were not expressed at all in embryos, implying different functions of these transcripts in embryonic development. Among the HSPs investigated, DNAJC15, DNAJC19, DNAJC24 and DNAJC27, all of the HSP40 family, had the most significant expression differential. They were further investigated for association with fertility and development traits. Single nucleotide polymorphisms (SNP) in DNAJC15 and DNAJC27 were found to be associated with blastocyst rate and fertilization rate, respectively.

Accordingly, the present invention provides an isolated nucleic acid molecule comprising at least one polymorphic site selected from the group consisting of positions 85146, 85161, 85216, 85292, and 85300 of the nucleic acid sequence shown in FIG. 1 (SEQ ID NO: 1) (part of the DNAJC15 gene), and at least 10 contiguous nucleotides of the sequence shown in FIG. 1, wherein position 85146 is guanine, position 85161 is guanine, position 85216 is adenine, position 85292 is cytosine, or position 85300 is guanine; or an isolated nucleic acid molecule comprising a polymorphic position selected from the group consisting of positions 35728, 36016, and 38867 of the nucleic acid sequence shown in FIG. 2 (SEQ ID

3

NO: 2) (part of the DNAJC27 gene), and at least 10 contiguous nucleotides of the sequence shown in FIG. 2, wherein position 35728 is guanine, position 36016 is guanine, or position 38867 is guanine. It is recognized that SEQ ID NO: 1 is already known, and the nucleic acid molecule therefore does not encompass one that consists of SEQ ID NO: 1.

Preferably, the nucleic acid molecule which comprises at least 15, more preferably at least 20, still more preferably at least 25, contiguous bases of SEQ ID NO: 1 or SEQ ID NO: 2 adjacent to the polymorphic site. In one embodiment, the isolated nucleic acid molecule comprises not more than 1,500 nt, preferably not more than 1000 nt, more preferably not more than 900 nt, more preferably not more than 800 nt, more preferably not more than 700 nt, preferably not more than 600 nt, more preferably not more than 500 nt, preferably not more than 400 nt, more preferably not more than 300 nt, more preferably not more than 150 nt, preferably not more than 100 nt, still more preferably not more than 50 nt.

The nucleic acid molecule preferably contains the polymorphic site which is within 4 nucleotides of the center of the nucleic acid molecule. Preferably, the polymorphic site is at the center of the nucleic acid molecule.

In another embodiment, the nucleic acid molecule contains the polymorphic site which is at the 3'-end of the nucleic acid molecule.

In another embodiment, the nucleic acid molecule contains the polymorphic site which is at the 5'-end of the nucleic acid molecule.

The present invention also provides an array of nucleic acid molecules comprising at least two nucleic acid molecules described above.

The present invention further provides a kit comprising a nucleic acid molecule described above, and a suitable container.

Also provided is a method for detecting single nucleotide polymorphism (SNP) in the bovine DNAJC15 or DNAJC27 genes, wherein the DNAJC15 gene is partially shown in FIG. 1, and the DNAJC27 gene is partially shown in FIG. 2, the method comprising optionally isolating a nucleic acid sample from the bovine animal or a tissue sample therefrom, determining the identity of a nucleotide of at least one position selected from the group consisting of a first polymorphic position comprising positions 85146, 85161, 85216, 85292, and 85300 of SEQ ID NO: 1, and a second polymorphic position comprising positions 35728, 36016, and 38867 of SEQ ID NO: 2, and comparing the nucleotide identity of the position respectively to the nucleotide identity at a corresponding position of SEQ ID NO: 1 or SEQ ID NO: 2.

Also provided is a method for genotyping a bovine cell, comprising obtaining a nucleic acid sample from said cell and determining the identity of a nucleotide of at least one position selected from the group consisting of positions 85146, 85161, 85216, 85292, and 85300 of SEQ ID NO: 1, and positions 35728, 36016, and 38867 of SEQ ID NO: 2. In one embodiment, the bovine cell is an adult cell, an embryo cell, a sperm, an egg, a fertilized egg, or a zygote. In one embodiment, both copies of the gene in the cell are genotyped.

The present invention further provides a method for progeny testing of cattle, the method comprising collecting a nucleic acid sample from said progeny, and genotyping said nucleic sample.

In another embodiment, a method is provided for selectively breeding cattle using a multiple ovulation and embryo transfer procedure (MOET), the method comprising superovulating a female animal, collecting eggs from said superovulated female, in vitro fertilizing said eggs from a suitable male animal, implanting said fertilized eggs into other

4

females allowing for an embryo to develop, and genotyping said developing embryo, and terminating pregnancy if the developing embryo does not have at least one polymorph selected from the group consisting of a first polymorph selected from the group consisting of guanine at position 85146, guanine at position 85161, adenosine at position 85216, cytosine at position 85292, guanine at position 85300 of SEQ ID NO: 1; and a second polymorph selected from the group consisting of guanine at position 35728, guanine at position 36016, and guanine at position 38867 of SEQ ID NO: 2.

In one embodiment, pregnancy is terminated if the developing embryo does not have either the first polymorph or the second polymorph.

In another embodiment, pregnancy is terminated if the developing embryo is not homozygous in the first polymorph and the second polymorph.

Further provided is a method for selecting a cattle as a breeder, wherein the cattle is genotyped according to the present invention, and the animal is only selected for breeding purpose if it comprises at least a first polymorph or a second polymorph, wherein the first polymorph is selected from guanine at position 85146, guanine at position 85161, adenosine at position 85216, cytosine at position 85292, guanine at position 85300 of SEQ ID NO: 1; and the second polymorph is selected from the group consisting of guanine at position 35728, guanine at position 36016, and guanine at position 38867 of SEQ ID NO: 2. In one embodiment, the cattle animal is selected only if it comprises both the first and second polymorphs. In another embodiment, the cattle animal is selected only if it is homozygous with regard to both the first and second polymorphs.

DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a portion of the DNAJC15 gene sequence (SEQ ID NO: 1) where the polymorphic sites and some of the PCR primers are shown. The five polymorphic sites are highlighted.

FIG. 2 shows a portion of the DNAJC27 gene sequence (SEQ ID NO: 2) where the polymorphic sites and some of the PCR primers are shown. The three polymorphic sites are highlighted, bold faced and underlined.

FIG. 3 shows the changes in expression levels of heat shock protein genes in bovine embryos using qRT-PCR. Data are shown as mean+/-maximum and minimum fold changes. Upregulation in degenerate or blastocyst embryos is represented by bars above or below, respectively, the x axis. qRT-PCR was performed in 4 sets of biological replicates of blastocysts and degenerate embryos.

FIG. 4 shows the splice variants of DNAJC5, DNAJC19, DNAJC24, and DNAJB12 genes. Positions of start and stop codons are indicated by vertical arrows and positions of primers used in the qRT-PCR are indicated by horizontal arrows. Black boxes represent coding sequences and white boxes represent untranslated regions. Due to overlap for DNAJB12, only transcript DNAB12-1220 could be amplified using transcript-specific primers.

FIG. 5 is a heat map of linkage disequilibrium (r^2) between SNPs in DNAJC15 and DNAJC27. SNPs 85146, 85161, 85216, 85292 and 85300 are located in DNAJC15 while SNPs 35728, 36016 and 38867 are located in DNAJC27.

DETAILED DESCRIPTION OF THE INVENTION

The present inventor has constructed an in-vitro fertilization (IVF) system which has the advantages of a unified

5

environment and well-isolated components of the embryonic development process, and used this system to characterize genetic factors involved in embryonic loss, and hence infertility, and to associate candidate genes and pathways with fertilization rate and embryonic survival at both the genomic and gene expression levels (Khatib et al., 2008a,b; Khatib et al., 2009; Huang et al., 2010a,b).

As indicated above, HSPs are among the first proteins produced during embryonic development (Neuer et al., 1999). The present inventor hypothesized that determining expression patterns of these genes in developed vs. arrested embryos could lead to the identification of specific genes involved in reproductive success. The high structural and functional conservation of HSPs during evolution suggests crucial roles of these proteins in fertilization, embryo development, and thus fertility in cattle. Also, some HSPs are considered housekeeping genes that are essential for many cell functions.

In efforts leading to the present invention, the expression levels of HSP genes in biological replicate bovine embryo pools that differ in their morphology and developmental statuses were evaluated, and associations of polymorphisms with fertility traits were examined. The HSP genes were found to be differentially expressed between blastocysts and degenerate embryos.

SNPs in two HSP genes were found to be associated with fertility traits. Specifically, a total of 17 candidate genes were chosen based on differential expression results observed in a previous microarray analysis of bovine embryos. To test whether or not these genes have roles in early development of cattle embryos, their expression levels were quantified and compared between blastocysts and degenerates. All 17 genes showed expression differences that ranged between 1.5- and 7.6-fold between the embryo groups. Two of these bovine HSP genes, DNAJC15 and DNAJC27, both of the HSP40 gene family, have several polymorphic positions, and these polymorphisms are associated with fertilization and blastocyst rates. Furthermore, all SNPs in each of DNAJC15 and DNAJC27 genes showed high linkage disequilibrium with each other.

The term “polymorphic” or “polymorphism” as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or individuals. “Polymorphic” refers to the condition in which two or more variants of a specific genomic sequence can be found in a population. A “polymorphic site” is the locus at which the variation occurs. Polymorphisms generally have at least two alleles, each occurring at a significant frequency in a selected population. A polymorphic locus may be as small as one base pair. The first identified allelic form is arbitrarily designated as the reference form, and other allelic forms are designated as alternative or variant alleles. The allelic form occurring most frequently in a selected population is sometimes referred to as the wild type form. Diploid organisms may be homozygous or heterozygous for allelic forms. A biallelic polymorphism has two forms, and a triallelic polymorphism has three forms, and so on.

Polymorphisms may provide functional differences in the genetic sequence, through changes in the encoded polypeptide, changes in mRNA stability, binding of transcriptional and translation factors to the DNA or RNA, and the like. Polymorphisms are also used to detect genetic linkage to phenotypic variation.

One type of polymorphism, single nucleotide polymorphisms (SNPs), has gained wide use for the detection of genetic linkage recently. SNPs are generally biallelic systems, that is, there are two alleles that an individual may have

6

for any particular SNP marker. In the instant case, the SNPs are used for determining the genotypes of the two HSP genes, which are found to have strong correlation to fertilization rate and embryonic survival.

The provided sequences also encompass the complementary sequence corresponding to any of the provided polymorphisms. In order to provide an unambiguous identification of the specific site of a polymorphism, the numbering of the original sequence in the GenBank is shown in FIG. 1 and is used throughout this disclosure.

The present invention provides nucleic acid based genetic markers for identifying bovine animals with superior breeding (such as fertility and embryo survival rates) traits. In general, for use as markers, nucleic acid fragments, preferably DNA fragments, may be as short as 7 nucleotides (nt), but may preferably be at least 12 nt, 15 nt, usually at least 20 nt, often at least 50 nt. Such small DNA fragments are useful as primers for the polymerase chain reaction (PCR), and/or probes for hybridization screening, etc.

The term primer refers to a single-stranded oligonucleotide capable of acting as a point of initiation of template-directed DNA synthesis under appropriate conditions (i.e., in the presence of four different nucleoside triphosphates and an agent for polymerization, such as, DNA or RNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 15 to 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to hybridize with the template. The term primer site, or priming site, refers to the area of the target DNA to which a primer hybridizes. The term primer pair means a set of primers including a 5' upstream primer that hybridizes with the 5' end of the DNA sequence to be amplified and a 3', downstream primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

The term “probe” or “hybridization probe” denotes a defined nucleic acid segment (or nucleotide analog segment) which can be used to identify by hybridizing to a specific polynucleotide sequence present in samples, said nucleic acid segment comprising a nucleotide sequence complementary of the specific polynucleotide sequence to be identified. “Probes” or “hybridization probes” are nucleic acids capable of binding in a base-specific manner to a complementary strand of nucleic acid.

An objective of the present invention is SNP genotyping, that is, to determine which embodiment of the SNP polymorphisms a specific sample of DNA has. For example, it is desirable to determine whether the nucleotide at a particular position is A or C. Many references describe genotyping methods, such as Chen et al., “Single nucleotide polymorphism genotyping: biochemistry, protocol, cost and throughput”, *Pharmacogenomics J.* 2003; 3(2):77-96; Kwok et al., “Detection of single nucleotide polymorphisms”, *Curr Issues Mol Biol.* April 2003; 5(2):43-60; Shi, “Technologies for individual genotyping: detection of genetic polymorphisms in drug targets and disease genes”, *Am J Pharmacogenomics.* 2002; 2(3):197-205; and Kwok, “Methods for genotyping single nucleotide polymorphisms”, *Annu Rev Genomics Hum Genet* 2001; 2:235-58. Exemplary techniques for high-throughput SNP genotyping are described in Marnellos, “High-throughput SNP analysis for genetic association studies”, *Curr Opin Drug Discov Devel.* May 2003; 6(3):317-21. Common SNP genotyping methods include, but are not limited to, TaqMan assays and modifications thereof such as

Molecular Beacon assays, SNPlex platforms, Bio-Plex system, CEQ and SNPstream systems, Molecular Inversion Probe array technology, BeadArray Technologies (e.g., Illumina GoldenGate and Infinium assays), single stranded conformation polymorphism assays (SSCP), molecular beacon assays, nucleic acid arrays, allele-specific primer extension, allele-specific PCR, arrayed primer extension, homogeneous primer extension assays, primer extension with detection by mass spectrometry, pyrosequencing, multiplex primer extension sorted on genetic arrays, ligation with rolling circle amplification, homogeneous ligation, OLA (U.S. Pat. No. 4,988,167), multiplex ligation reaction sorted on genetic arrays, restriction-fragment length polymorphism, single base extension-tag assays, and the Invader assay. Such methods may be used in combination with detection mechanisms such as, for example, luminescence or chemiluminescence detection, fluorescence detection, time-resolved fluorescence detection, fluorescence resonance energy transfer, fluorescence polarization, mass spectrometry, and electrical detection.

The sequence neighboring the SNP site can be used to design SNP detection reagents such as oligonucleotide probes, which may optionally be implemented in a kit format. Preferably, the oligonucleotide probe will have a detectable label, and contains for example an A at the corresponding position. Experimental conditions can be chosen such that if the sample DNA contains an A, a hybridization signal can be detected because the probe hybridizes to the corresponding complementary DNA strand in the sample, while if the sample DNA contains a G, no hybridization signal is detected.

Similarly, PCR primers and conditions can be devised, whereby the oligonucleotide is used as one of the PCR primers, for analyzing nucleic acids for the presence of a specific sequence. These may be direct amplification of the genomic DNA, or RT-PCR amplification of the mRNA transcript of the target gene. The use of the polymerase chain reaction is described in Saiki et al. (1985) *Science* 230:1350-1354. Amplification may be used to determine whether a polymorphism is present, by using a primer that is specific for the polymorphism. Alternatively, various methods are known in the art that utilize oligonucleotide ligation as a means of detecting polymorphisms, for examples see Riley et al (1990) *Nucleic Acids Res.* 18:2887-2890; and Delahunty et al (1996) *Am. J. Hum. Genet.* 58:1239-1246. The detection method may also be based on direct DNA sequencing, or hybridization, or a combination thereof. Where large amounts of DNA are available, genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis. The nucleic acid may be amplified by PCR, to provide sufficient amounts for analysis.

Hybridization may be performed in solution, or such hybridization may be performed when either the oligonucleotide probe or the target polynucleotide is covalently or non-covalently affixed to a solid support. Attachment may be mediated, for example, by antibody-antigen interactions, poly-L-Lys, streptavidin or avidin-biotin, salt bridges, hydrophobic interactions, chemical linkages, UV cross-linking baking, etc. Oligonucleotides may be synthesized directly on the solid support or attached to the solid support subsequent to synthesis. Solid-supports suitable for use in detection methods of the invention include substrates made of silicon, glass, plastic, paper and the like, which may be formed, for example, into wells (as in 96-well plates), slides, sheets, membranes, fibers, chips, dishes, and beads. The solid support may be treated, coated or derivatized to facilitate the

immobilization of the allele-specific oligonucleotide or target nucleic acid. For screening purposes, hybridization probes of the polymorphic sequences may be used where both forms are present, either in separate reactions, spatially separated on a solid phase matrix, or labeled such that they can be distinguished from each other.

Hybridization may also be performed with nucleic acid arrays and subarrays such as described in WO 95/11995. The arrays would contain a battery of allele-specific oligonucleotides representing each of the polymorphic sites. One or both polymorphic forms may be present in the array, for example the polymorphism of a SNP position may be represented by either, or both, of the listed nucleotides. Usually such an array will include at least 2 different polymorphic sequences, i.e. polymorphisms located at unique positions within the locus, and may include all of the provided polymorphisms. Arrays of interest may further comprise sequences, including polymorphisms, of other genetic sequences, particularly other sequences of interest. The oligonucleotide sequence on the array will usually be at least about 12 nt in length, may be the length of the provided polymorphic sequences, or may extend into the flanking regions to generate fragments of 100 to 200 nt in length. For examples of arrays, see Ramsay (1998) *Nat. Biotech.* 16:4044; Hacia et al. (1996) *Nature Genetics* 14:441-447; Lockhart et al. (1996) *Nature Biotechnol.* 14:1675-1680; and De Risi et al. (1996) *Nature Genetics* 14:457-460.

The identity of polymorphisms may also be determined using a mismatch detection technique, including but not limited to the RNase protection method using riboprobes (Winter et al., *Proc. Natl. Acad. Sci. USA* 82:7575, 1985; Meyers et al., *Science* 230:1242, 1985) and proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein (Modrich, *P. Ann. Rev. Genet.* 25:229-253, 1991). Alternatively, variant alleles can be identified by single strand conformation polymorphism (SSCP) analysis discussed above (Orita et al., *Genomics* 5:874-879, 1989; Humphries et al., in *Molecular Diagnosis of Genetic Diseases*, R. Elles, ed., pp. 321-340, 1996) or denaturing gradient gel electrophoresis (DGGE) (Wartell et al., *Nucl. Acids Res.* 18:2699-2706, 1990; Sheffield et al., *Proc. Natl. Acad. Sci. USA* 86:232-236, 1989).

A polymerase-mediated primer extension method may also be used to identify the polymorphism(s). Several such methods have been described in the patent and scientific literature and include the "Genetic Bit Analysis" method (WO92/15712) and the ligase/polymerase mediated genetic bit analysis (U.S. Pat. No. 5,679,524). Related methods are disclosed in WO91/02087, WO90/09455, WO95/17676, U.S. Pat. Nos. 5,302,509, and 5,945,283. Extended primers containing a polymorphism may be detected by mass spectrometry as described in U.S. Pat. No. 5,605,798. Another primer extension method is allele-specific PCR (Ruao et al., *Nucl. Acids Res.* 17:8392, 1989; Ruao et al., *Nucl. Acids Res.* 19, 6877-6882, 1991; WO 93/22456; Turki et al., *J. Clin. Invest.* 95:1635-1641, 1995). In addition, multiple polymorphic sites may be investigated by simultaneously amplifying multiple regions of the nucleic acid using sets of allele-specific primers as described in Wallace et al. (WO 89/10414).

A detectable label may be included in an amplification reaction. Suitable labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine

(TAMRA), radioactive labels, e.g. ^{32}P , ^{35}S , ^3H ; etc. The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

It is readily recognized by those ordinarily skilled in the art that in order to maximize the signal to noise ratio, in probe hybridization detection procedure, the polymorphic site should be at the center of the probe fragment used, whereby a mismatch has a maximum effect on destabilizing the hybrid molecule; and in a PCR detection procedure, the polymorphic site should be placed at the very 3'-end of the primer, whereby a mismatch has the maximum effect on preventing a chain elongation reaction by the DNA polymerase. The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the polynucleotide is considered to be "at the center" of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is considered to be "within 1 nucleotide of the center." With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be "within 1 nucleotide of the center" and any of the four nucleotides in the middle of the polynucleotide would be considered to be "within 2 nucleotides of the center," and so on.

In some embodiments, a composition contains two or more differently labeled oligonucleotides for simultaneously probing the identity of nucleotides or nucleotide pairs at two or more polymorphic sites. It is also contemplated that primer compositions may contain two or more sets of allele-specific primer pairs to allow simultaneous targeting and amplification of two or more regions containing a polymorphic site.

Alternatively, the relevant portion of the target gene of the sample of interest may be amplified via PCR and directly sequenced, and the sequence be compared to the wild type sequence. It is readily recognized that, other than those specifically disclosed herein, numerous primers can be devised to achieve the objectives. PCR and sequencing techniques are well known in the art and reagents and equipments are readily available commercially.

The so-called next generation sequencing, and high-throughput sequencing methods, may also be used. For example, Massively Parallel Signature Sequencing (MPSS); Polony sequencing, pyro sequencing, SOLiD sequencing; ion semiconductor sequencing, and DNA nanoball sequencing systems are well-known to those skilled in the art.

DNA markers have several advantages; segregation is easy to measure and is unambiguous, and DNA markers are co-dominant, i.e., heterozygous and homozygous animals can be distinctively identified. Once a marker system is established selection decisions could be made very easily, since DNA markers can be assayed any time after a sample is collected from the individual animal, or even earlier by testing embryos in vitro if very early embryos are collected. The use of marker assisted genetic selection will greatly facilitate and speed up cattle breeding problems. For example, a modification of the

multiple ovulation and embryo transfer (MOET) procedure can be used with genetic marker technology. Specifically, females are superovulated, eggs are collected, in vitro fertilized using semen from superior males and implanted into other females allowing for use of the superior genetics of the female (as well as the male) without having to wait for her to give birth to one calf at a time. Developing blastomeres at the 4-8 cell stage may be assayed for presence of the marker, and selection decisions made accordingly.

In one embodiment of the invention an assay is provided for detection of presence of a desirable genotype using the markers.

The term "genotype" as used herein refers to the identity of the alleles present in an individual or a sample. In the context of the present invention a genotype preferably refers to the description of the polymorphic alleles present in an individual or a sample. The term "genotyping" a sample or an individual for a polymorphic marker refers to determining the specific allele or the specific nucleotide carried by an individual at a polymorphic marker.

The present invention is suitable for identifying a bovine, including a young or adult bovine animal, an embryo, a semen sample, an egg, a fertilized egg, or a zygote, or other cell or tissue sample therefrom, to determine whether said bovine possesses the desired genotypes of the present invention, some of which are indicative of improved fertilization rate and embryonic survival.

Further provided is a method for genotyping the bovine HSP gene, comprising determining for the two copies of the HSP gene present the identity of the nucleotide pair at positions 25402 and 19069.

One embodiment of a genotyping method of the invention involves examining both copies of the relevant HSP gene, or a fragment thereof, to identify the nucleotide pair at the polymorphic site in the two copies to assign a genotype to the individual. In some embodiments, "examining a gene" may include examining one or more of: DNA containing the gene, mRNA transcripts thereof, or cDNA copies thereof. As will be readily understood by the skilled artisan, the two "copies" of a gene, mRNA or cDNA, or fragment thereof in an individual may be the same allele or may be different alleles. In another embodiment, a genotyping method of the invention comprises determining the identity of the nucleotide pair at the polymorphic site.

The present invention further provides a kit for genotyping a bovine sample, the kit comprising in a container a nucleic acid molecule, as described above, designed for detecting the polymorphism, and optionally at least another component for carrying out such detection. Preferably, a kit comprises at least two oligonucleotides packaged in the same or separate containers. The kit may also contain other components such as hybridization buffer (where the oligonucleotides are to be used as a probe) packaged in a separate container. Alternatively, where the oligonucleotides are to be used to amplify a target region, the kit may contain, preferably packaged in separate containers, a polymerase and a reaction buffer optimized for primer extension mediated by the polymerase, such as PCR.

In one embodiment the present invention provides a breeding method whereby genotyping as described above is conducted on bovine embryos, and based on the results, certain cattle are either selected or dropped out of the breeding program.

Through use of the linked marker loci, procedures termed "marker assisted selection" (MAS) may be used for genetic improvement within a breeding nucleus; or "marker assisted

introgression” for transferring useful alleles from a resource population to a breeding nucleus (Soller 1990; Soller 1994).

Testing of Roles of 17 HSP Genes

The roles of 17 HSP genes in early embryonic development and fertility in cattle at both genomic and gene expression levels were investigated. These genes include 8 HSP40 genes, six HSP70 genes, one HSPP, and 2 HSP10 genes. (see Table 1) These genes were chosen based on differential expression results observed in a previous microarray analysis of bovine embryos; and their expression levels were quantified and compared between blastocysts and degenerates to test whether or not these genes have roles in early development of cattle embryos.

All 17 genes showed expression differences that ranged between 1.5- and 7.6-fold between the embryo groups. All HSP40 family genes were found to be upregulated in degenerate embryos compared to blastocysts. Although the specific functions of HSP40s have not been reported in cattle, studies from other species have shown that these genes play important roles in protecting cells under stress conditions (Lanneau et al., 2007). It is well established that ATP hydrolysis is necessary for protein folding activity of HSP70s and that HSP40s stimulate ATPase activity and stabilize the interaction of HSP70s with their substrates (Qiu et al., 2006). Gotoh and colleagues (2004) reported that HSP40 combines with HSP70 to act as chaperones to protect cells from apoptosis. Degenerate embryos in the IVF system are growth-arrested embryos and seem to undergo partial apoptosis. Research has shown that HSPs are upregulated to prevent apoptosis triggered by different stimuli by interacting with key factors of the apoptotic signaling pathways (Lanneau et al., 2007). Therefore, it is feasible that upregulation of HSP40 genes observed in degenerates is a response to stress on these embryos so they can maintain protein homeostasis.

The qRT-PCR analysis revealed moderate fold change expression of HSPE1, HSPH1, HSB1, and six HSP70 genes (HSPA2, HSPA4, HSPA5, HSPA8, HSPA9 and HSPA14). HSPA2, HSPA4, HSPA5, and HSPA8 showed higher expression in blastocysts than degenerates, whereas expression levels of HSPA9 and HSPA14 were higher in degenerate embryos. Although the specific functions of most HSP70s have not been reported in cattle, studies in human and mouse have indicated important roles for these genes in early embryogenesis. It has been reported that Hspa2 is essential for normal spermatogenesis and for growth and survival of cancer cells and that Hspa8 knockout mice are not viable because of the housekeeping functions of this gene (Daugaard et al., 2007). Human HSPA5 is involved in the folding and transport of proteins into the endoplasmic reticulum and Hspa5 knockout mice embryos die at day 3.5, and therefore it, too, is considered a housekeeping gene (Daugaard et al., 2007). Furthermore, Matwee et al. (2001) have reported a reduced blastocyst rate and an increased apoptosis of embryos cultured in the presence of anti-HSP70. Thus, upregulation and downregulation of HSPs in bovine embryos observed in the present study imply that these genes play vital roles in early embryo development.

Given that alternative splicing is a major source of phenotypic complexity in mammals (Wang et al., 2008), a search was conducted of the Ensembl annotation of the bovine genome and found that only 4 out of 17 HSP candidate genes have annotated splice variants. In contrast, for the 17 human and mouse counterparts, 15 and 13 genes, respectively, have 2 or more splice variants. The low number of alternative transcripts identified in cattle is presumably due to the incomplete annotation of the bovine genome. Indeed, in a previous study, thousands of alternative splicing events were charac-

terized in blastocysts and degenerative embryos, and some of those events were found to be associated with the developmental status of embryos (Huang and Khatib, 2010).

In order to explore the effects of the different splice variants of HSPs on embryo development, expression levels of the identified variants were estimated and compared between blastocysts and degenerate embryos. The results clearly show that different splice variants display different expression levels which imply different functions in embryo development. For example, while the long variant of DNAJC24 was highly expressed in degenerate embryos compared to blastocysts, the short variant of this gene was found to be lowly expressed in both embryo groups. Also, for DNAJC5, one transcript could not be detected using qRT-PCR. These results are consistent with human studies in which only some of the splice variants of genes were found to be associated with a disease state (Wang and Cooper, 2007). Collectively, these results testify to the importance of alternative splicing as a key regulator of phenotypic variation in mammals.

To better understand the involvement of the HSP genes in fertility traits, 4 genes that showed the most significant expression differences between embryo groups were tested for SNP association.

Recently, an IVF experimental system in cattle has been developed in our laboratory (see Methods) aimed at identifying genetic factors affecting fertilization and embryo development. The effectiveness of this system in identifying genes and pathways associated with developmental and fertility traits has been demonstrated in several studies (Khatib et al., 2008a,b; Khatib et al., 2009; Driver et al., 2009; Wang et al., 2009; Huang et al., 2010a).

Analysis of DNAJC15 and DNAJC27 SNP revealed significant associations with blastocyst and fertilization rates, respectively. These results are of particular significance because these 2 genes showed a remarkable difference in expression between blastocysts and degenerate embryos, providing strong evidence for the involvement of these genes in embryo production and development in cattle. The results also provide further evidence for the roles of HSPs in the fertilization process and blastocyst rates observed in mice and cattle in previous studies. Neuer et al. (1998) found that the presence of anti-HSP60 monoclonal antibodies had significant effects on blastocyst rates of mouse embryos. Matwee et al. (2001) reported that the addition of anti-HSP70 monoclonal antibodies to the culture medium reduced the mean number of spermatozoa bound to zona pellucida by about 50% to that of controls. Also, the authors reported that anti-HSP70 significantly reduced the number of bovine embryos that reached the blastocyst stage (Matwee et al., 2001). A recent study by Rosenkrans et al. (2010) reported an association between SNP in the promoter region of HSP70 and calving percentages in Brahman cows. Although a small sample size was used in the association analysis, the results point to an important role of HSP70 in cow fertility.

The identification of specific HSP genes contributing to embryo survival can be a unique opportunity to improve protection of IVF embryos from different toxic conditions and to enhance pregnancy rates in cattle (Hansen, 2007). HSP genes investigated in this study were found to be differentially expressed between blastocysts and degenerate embryos and SNP in 2 genes were associated with fertility traits. Blastocysts and degenerate embryos have distinct morphological and developmental features. As such, the present study provides a set of candidate genetic markers for pre-implantation embryo development. However, it has been reported that relatively small changes (decrease or increase) in expression levels of heat shock proteins can result in growth abnormali-

ties and cell death (Nollen and Morimoto, 2002). Also, the assessment of the expression of alternative transcript isoforms in the embryos examined in this study testifies to the importance of inclusion of this approach in studying gene expression.

The present invention accordingly provides novel cattle genotyping, selective cattle breeding and related methods, based on the discovery that certain HSP40 polymorphism confers improved fertilization and embryonic survival rate to cattle.

The following examples are intended to illustrate preferred embodiments of the invention and should not be interpreted to limit the scope of the invention as defined in the claims.

EXAMPLES

Materials and Methods

In-Vitro Maturation, Fertilization, and Embryo Culture

In this study, fertilization and embryo production were performed in 2 different experiments. The first experiment is a comparison of expression profiles of 17 HSPs between 2 populations of embryos differing in their morphology, and the second experiment is a genomic association analysis of genes—found to be differentially-expressed in the first experiment—with fertilization and blastocyst rates.

Experiment I: Expression Analysis of HSP Genes in Cattle Embryos

Ovaries were obtained from a local abattoir and processed for IVF using standard protocol (Huang et al., 2010b; Khatib et al., 2008a). Briefly, oocytes were aspirated from 2-6 mm follicles and washed in Tyrode's albumin lactate pyruvate (TALP)-Hepes after which they underwent maturation in supplemented M199 media (Khatib et al., 2008a). Incubation was completed at 5% CO₂ in air at 39° C. and high humidity for 22-24 hours at which time oocytes were then washed in TALP-Hepes and moved into fresh IVF-TALP (Biowhittaker, Walkersburg, Md.). Semen samples from 2 different bulls were used for fertilization; sperm underwent Percoll separation (45-90% gradient) and were adjusted to a final concentration of 1×10⁶/ml (Parrish et al., 1995). Fertilization was marked as day 0 of development and completed by combining sperm, heparin, and PHE with the oocytes in 44 µl drops of media (Khatib et al., 2008a). Incubation was then continued for 22-24 hours after which the putative zygotes were washed in TALP-Hepes, denuded of remaining cumulus cell complexes, and placed into supplemented synthetic oviductal fluid (Biowhittaker) and returned to incubation (Khatib et al., 2008a).

Morphological Assessment

Putative zygotes were cultured over an 8-day period during which assessments of their developmental progress were made. By day 5 of development, a bovine embryo should attain approximately 16-32 cells and show signs of cellular compaction deeming it as morula. Embryos failing to show these characteristics were assumed to have been arrested in development and were excluded from analysis. Morulas were then continued in culture until day 8. By this time an embryo should show evidence of a fluid filled cavity (blastocoele), which gives rise to the differentiation of the inner cell mass and the trophectoderm, qualifying it as a blastocyst. For expression analysis, there were 2 populations of embryos collected. The first consisted of embryos that attained a compacted morula status by day 5 but failed to form a blastocoele by day 8, referred as degenerate embryos. The second consisted of embryos that developed into blastocysts by day 8. Embryos from each morphological group were collected in

pools of 20 and preserved in RNALater (Ambion, Austin, Tex.). Two bulls were used with 2 sets of biological replicate pools (total of 4 biological replicates) from each to prevent maternal crossover.

Quantitative Real-Time PCR (qRT-PCR)

A previous study in our laboratory using microarray expression analysis has revealed that many genes are differentially expressed between blastocysts and degenerate embryos (Huang et al., 2010b). Seventeen HSP genes that showed 1.5-fold or higher differences in expression between the embryo groups described in the study of Huang et al. (2010b) were chosen for validation of differential expression and further investigation in new sets of blastocysts and degenerate embryo pools using qRT-PCR analysis. Primers (Table 1) were designed to amplify fragments spanning more than one exon to exclude the possibility of genomic DNA contamination in the qRT-PCR reactions using the Beacon Designer software (Premier Biosoft International, Palo Alto, Calif.). Total RNA was extracted from pools of embryos using RNeasy Micro (Ambion) and quality controlled using a RNeasy6000 PicoChip (Agilent Technologies, Calif.). Messenger RNA was amplified using MessageAmp II (Ambion, Austin, Tex.), followed by cDNA synthesis using iScript (Bio-Rad, Hercules, Calif.) according to manufacturers' instructions. Reactions of qRT-PCR were run on a DNA Engine-Opticon 2 Detection System (MJ Research, Watertown, Mass.) using iQ SYBR Green Supermix kit (Bio-Rad Laboratories, Calif.). Each sample was run in quadruplicate and all 4 expression results were averaged. Analysis of gene expression levels was conducted using the 2^{-ΔΔC_t} method (Livak and Schmittgen, 2001). The selection of the house-keeping gene GAPDH as an endogenous control was as described in Huang et al. (2010b).

Experiment II: Association Analysis of Differentially-Expressed HSPs with Fertilization and Blastocyst Rates

In order to further investigate the involvement in fertility of genes (DNAJC15, DNAJC19, DNAJC24, and DNAJC27) that showed the highest fold change in expression in degenerate embryos vs. blastocysts, an association analysis between SNPs in these genes and 2 main fertility traits: fertilization rate and blastocyst rate was performed. Using the IVF experimental system, several genes were found to be significantly associated with variation in fertilization and embryonic survival rates (Khatib et al., 2008a,b; Khatib et al., 2009).

Phenotypic Data.

In order to generate phenotypic data for association analysis, a total of 6,893 in-vitro fertilizations were performed using oocytes from 399 ovaries (obtained from 399 Holstein cows) and semen samples from 12 Holstein bulls. For 92 ovaries, oocytes were fertilized by two different bulls each. Fertilization rate was calculated as the number of cleaved embryos 48 h postfertilization out of the total number of oocytes exposed to sperm. Blastocyst rate was calculated as the number of embryos that reached blastocyst stage out of the total embryos cultured by day 8. asdf

Genotyping.

DNA was extracted from ovaries (n=399) using standard phenol/chloroform protocols. The DNA concentrations were measured using a spectrophotometer (Ultraspec 2100; Amersham Biosciences). For polymorphism identification, 3 DNA pools were constructed from 20 different ovary/cow samples to contain 25 ng of DNA from each sample. DNA Pools were amplified with different sets of primers designed from the coding and 5' and 3' UTR regions of the 4 candidate genes (Table 3). Amplification, sequencing of PCR products, and SNP identification were as described in Khatib et al. (2008a,

b). SNP identified in DNAJC15, DNAJC19, DNAJC24, and DNAJC27 were genotyped for the 399 ovary/cow samples at GeneSeek (Lincoln, Nebr.).

Statistical Analysis

For expression analysis, normalized gene expression values (ΔCt) were analyzed using a general linear model including the fixed effects of the bull, the type of embryo (blastocyst or degenerate), and the random effect of the pool. Association between the normalized gene expression and the type of embryo was tested using a likelihood ratio test by comparing this model to a reduced model without the embryo effect. The mean and the range of the fold change for each gene were calculated as $2^{-\Delta\Delta Ct}$ using the estimated $\Delta\Delta Ct$ value $\pm$ standard error.

For genomic association analysis between SNP and fertilization and blastocyst rates, the following mixed linear model was used,

$$y_{ijk} = \mu + o_i + b_j + \text{SNP}_{ijk} + e_{ijk}$$

where y_{ijk} represents in turn, the fertilization or survival rate of oocytes k from ovary i fertilized with semen from bull j ; μ represents a general mean for the trait considered, o_i represents the random effect of the individual ovary from which oocytes were harvested; b_j represents the random effect of the sire used in the fertilization; SNP_{ijk} represents the fixed effect of the genotypic class for the SNP considered; and e_{ijk} represents the residuals, assumed normal, independent and identically distributed with mean 0 and variance σ_e^2 . Ovaries and bulls were assumed uncorrelated with variance structures σ_o^2 and σ_b^2 , respectively. Association between the SNP and fertilization or blastocyst rates was tested using a likelihood ratio test by comparing the full model to a reduced model without the SNP effect. All analyses were performed using the lme4 package of R language/environment (R Development Core Team, 2009).

Results

Expression Profiling of HSP Genes in Embryos

Expression differences between degenerate embryos and blastocysts were estimated for 17 HSP genes in 4 sets of biological replicates using qRT-PCR (FIG. 1). A range of 1.5- to 7.6-fold difference in expression was observed between embryo groups. Interestingly, all HSP40 gene family members were found to be upregulated in degenerate embryos compared to blastocysts (FIG. 1). For example, DNAJC15 ($P < 0.0001$), DNAJC19 ($P < 0.0001$), DNAJC24 ($P = 0.002$), and DNAJC27 ($P = 0.0098$) showed an average of 7.6-, 4.8-, 3.3-, 4.33-fold differences in expression, respectively. In contrast, only 2 members of HSP70 family (HSPA14 and HSPA9) showed upregulation in degenerates, whereas 4 HSP70 genes (HSPA2, HSPA4, HSPA5, and HSPA8) showed higher expression levels in blastocysts than in degenerates (FIG. 1).

Differential Expression of Alternatively-Spliced Transcripts

Splice variants of HSP genes were identified using the genebuild procedure of Ensembl (<http://uswest.ensembl.org/>). In this procedure, annotation of transcripts is based on mRNA and protein sequences deposited in public scientific databases. Ensembl search revealed one transcript per gene for 13 genes, 2 transcripts each for DNAJB12 and DNAJC19, and 3 transcripts each for DNAJC5 and DNAJC24. Expression levels of splice variants of DNAJC5, DNAJC19, DNAJC24, and DNAJB12 (FIG. 2) were estimated in 4 sets of biological replicates of blastocysts and degenerate embryos using qRT-PCR. Transcripts DNAJC5-1049 and DNAJC5-667 (FIG. 2) showed 2.53- and 1.67-fold higher expression, respectively, in degenerate embryos com-

pared to blastocysts while the DNAJC5-725 transcript was not detected in any of the embryo groups. For DNAJC19, only transcript DNAJC19-536 was detected in embryos. For DNAJC24, DNAJC24-2172 transcript showed a 3.22-fold higher expression in degenerate embryos compared to blastocysts, whereas transcript DNAJC24-628 was lowly expressed in all embryo samples. DNAJC24-1893 transcript could not be detected in embryos. The sequences of the 2 DNAJB12 transcripts were overlapping so that only transcript DNAB12-1220 could be amplified using transcript-specific primers. DNAB12-1220 showed a 1.95-fold higher expression in degenerate embryos vs. blastocysts.

Association of HSP Polymorphisms with Fertilization and Blastocyst Rates

SNP in genes DNAJC15, DNAJC19, DNAJC24, and DNAJC27 were tested for association with fertilization and blastocyst rates. These genes showed the highest fold differences in expression between embryo groups (FIG. 1) and the highest statistical significance in the general linear model analysis ($P < 0.01$). Using the pooled DNA sequencing approach, 5 SNP were identified in the 3'UTR of DNAJC15, one SNP in exon 5 of DNAJC19, one SNP and one 4-bp deletion in the 3' UTR of DNAJC24, and 3 SNP in the 3' UTR of DNAJC27. All SNP in each of DNAJC15 and DNAJC27 genes showed high linkage disequilibrium with each other (FIG. 3). Polymorphisms in DNAJC19 and DNAJC24 did not show significant associations with fertility traits. Table 2 shows fertilization and blastocysts rates for the genotypic classes of the DNAJC15 and DNAJC27 SNP. For DNAJC27 SNP36016, oocytes collected from genotype GG ovaries showed a 69.3% fertilization rate vs. 62.2% for oocytes collected from CC ovaries ($P = 0.034$). For DNAJC15 SNP85146, the blastocyst rate of embryos produced from GG dams was 40.1% vs. 31.0% and 28.1% for embryos produced from AA and AG dams, respectively (Table 2).

While the invention has been described in connection with one or more embodiments, it should be understood that the invention is not limited to those embodiments, and the description is intended to cover all alternatives, modifications, and equivalents, as may be included within the spirit and scope of the appended claims. All referenced cited herein are further incorporated by reference in their entirety.

REFERENCES CITED

- Becker, J., and E. A. Craig. 1994. Heat-shock proteins as molecular chaperones. *Eur. J. Biochem.* 219:11-23.
- Carlborg, O., and C. S. Haley. 2004. Epistasis: too often neglected in complex trait studies? *Nat. Rev. Genet.* 5:618-625.
- Cobanoglu, O., I. Zaitoun, Y. M. Chang, G. E. Shook, and H. Khatib. 2006. Effects of the signal transducer and activator of transcription 1 (STAT1) gene on milk production traits in Holstein dairy cattle. *J. Dairy Sci.* 89:4433-4437.
- Darnell, J. E. 1997. STATs and gene regulation. *Science* 277:1630-1635.
- Daugaard, M., M. Rohde, and M. Jäättelä. 2007. The heat shock protein 70 family: Highly homologous proteins with overlapping and distinct functions. *FEBS Lett.* 581:3702-3710.
- Dobson, H., R. Smith, M. Royal, Ch. Knight, and I. Sheldon. 2007. The high-producing dairy cow and its reproductive performance. *Reprod Domest Anim.* 42 Suppl 2:17-23.
- Dobson, H., S. L. Walker, M. J. Morris, J. E. Routly, and R. F. Smith. 2008. Why is it getting more difficult to successfully artificially inseminate dairy cows? *Animal* 2:1104-1111.

Driver, A. M., W. Huang, S. Gajic, R. L. Monson, G. J. Rosa, and H. Khatib. 2009. Effects of the progesterone receptor variants on fertility traits in cattle. *J. Dairy Sci.* 92:4082-4085.

Hansen, P. J. 2007. To be or not to be—determinants of embryonic survival following heat shock. *Theriogenology* 68 Suppl 1:S40-8.

Hombria, J. C., and S. Brown. 2002. The fertile field of *Drosophila* Jak/STAT signalling. *Curr. Biol.* 12:R569-R575.

Huang, W., B. W. Kirkpatrick, G. J. Rosa, and H. Khatib. 2010a. A genome-wide association study using selective DNA pooling identifies candidate markers for fertility in Holstein cattle. *Anim. Genet.* 41:570-578.

Huang, W., B. S. Yandell, and H. Khatib. 2010b. Transcriptomic profiling of bovine IVF embryos revealed candidate genes and pathways involved in early embryonic development. *BMC Genomics* 11:23.

Huang, W., and H. Khatib. 2010. Comparison of transcriptomic landscapes of bovine embryos using RNA-Seq. *BMC Genomics* 11:711.

Kisseleva, T., S. Bhattacharya, J. Braunstein and C. W. Schindler. 2002. Signaling through the JAK/STAT pathway, recent advances and future challenges. *Gene* 285:1-24.

Khatib, H., R. L. Monson, V. Schutzkus, D. M. Kohl, G. J. Rosa, and J. J. Rutledge. 2008a. Mutations in the STATSA gene are associated with embryonic survival and milk composition in cattle. *J. Dairy Sci.* 91:784-793.

Khatib, H., C. Maltecca, R. L. Monson, V. Schutzkus, X. Wang, and J. J. Rutledge. 2008b. The fibroblast growth factor 2 gene is associated with embryonic mortality in cattle. *J. Anim. Sci.* 86:2063-2067.

Khatib, H., W. Huang, X. Wang, A. H. Tran, A. B. Bindrim, V. Schutzkus, R. L. Monson, and B. S. Yandell. 2009. Single gene and gene interaction effects on fertilization and embryonic survival rates in cattle. *J. Dairy Sci.* 92:2238-2247.

Kodama, H., K. Fukuda, J. Pan, S. Makino, A. Baba, S. Hori, and S. Ogawa. 1997. Leukemia inhibitory factor, a potent cardiac hypertrophic cytokine, activates the JAK/STAT pathway in rat cardiomyocytes. *Circ. Res.* 81:656-663.

Lanneau, D., A. de Thonel, S. Maurel, C. Didelot, and C. Garrido. 2007. Apoptosis versus cell differentiation: role of heat shock proteins HSP90, HSP70 and HSP27. *Prion* 1:53-60.

Leroy, J. L., G. Opsomer, A. Van Soom, I. G. Goovaerts, and P. E. Bols. 2008. Reduced fertility in high-yielding dairy cows: are the oocyte and embryo in danger? Part I. The importance of negative energy balance and altered corpus luteum function to the reduction of oocyte and embryo quality in high-yielding dairy cows. *Reprod Domest Anim.* 43:612-622.

Livak, K. J., and T. D. Schmittgen. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25:402-408.

Luft, J. C., and D. J. Dix. 1999. Hsp70 expression and function during embryogenesis. *Cell Stress Chaperones* 4:162-170.

Maj, T., and A. Chelmonska-Soyta. 2007. Pleiotropy and redundancy of STAT proteins in early pregnancy. *Reprod. Domest. Anim.* 42:343-353.

McCullagh, P., and J. A. Nelder. 1989. Generalized Linear Models, 2nd edn. Chapman and Hall, London.

Morris, D., and M. Diskin. 2008. Effect of progesterone on embryo survival. *Animal* 2:1112-1119.

Neuer, A., C. Mele, H. C. Liu, Z. Rosenwaks, and S. S. Witkin. 1998. Monoclonal antibodies to mammalian heat shock proteins impair mouse embryo development in vitro. *Hum. Reprod.* 13:987-990.

Neuer, A., S. D. Spandorfer, P. Giraldo, J. Jeremias, S. Dietlerle, I. Korneeva, H. C. Liu, Z. Rosenwaks, and S. S. Witkin. 1999. Heat shock protein expression during gametogenesis and embryogenesis. *Infect. Dis. Obstet. Gynecol.* 7:10-6.

Nollen, E. A., and R. I. Morimoto. 2002. Chaperoning signaling pathways: molecular chaperones as stress-sensing 'heat shock' proteins. *J. Cell Sci.* 115:2809-2816.

Parrish, J. J., A. Krogenaes, and J. L. Susko-Parrish. 1995. Effect of bovine sperm separation by either swim-up or Percoll method on success of in vitro fertilization and early embryonic development. *Theriogenology* 44:859-869.

Qiu, X. B., Y. M. Shao, S. Miao, and L. Wang. 2006. The diversity of the DnaJ/Hsp40 family, the crucial partners for Hsp70 chaperones. *Cell Mol. Life Sci.* 63:2560-2570.

R Development Core Team. 2008. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. (www.R-project.org).

Royal, M., G. E. Mann, and A. P. Flint. 2000. Strategies for reversing the trend towards subfertility in dairy cattle. *Vet. J.* 160:53-60.

Santos, J. E. P., W. W. Thatcher, R. C. Chebel, R. L. A. Cerri, and K. N. Galvao. 2004. The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. *Anim. Reprod. Sci.* 83:513-535.

Shook G. E. 2006. Major advances in determining appropriate selection goals. *J. Dairy Sci.* 89:1349-1361.

Soller, M. (1990) Genetic mapping of the bovine genome using DNA-level markers with particular attention to loci affecting quantitative traits of economic importance. *J. Dairy Sci.* 73:2628-2646.

Soller, M. (1994) Marker-assisted selection, an overview. *Anim. Biotech.* 5:193-208.

Takeda, K., K. Noguchi, W. Shi, T. Tanaka, M. Matsumoto, N. Yoshida, T. Kishimoto, and S. Akira. 1997. Targeted disruption of the mouse Stat3 gene leads to early embryonic lethality. *Proc. Natl. Acad. Sci. USA.* 94:3801-3804.

Teglund, S., C. McKay, E. Schuetz, J. M. van Deursen, D. Stravopodis, D. Wang, M. Brown, S. Bodner, G. Grosveld and J. N. Ihle. 1998. StatSa and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell* 93:841-850.

Truchet, S., M. Chebrou, C. Djedati, J. Wietzerbin, and P. Debey. 2004. Presence of permanently activated signal transducers and activators of transcription in nuclear interchromatin granules of unstimulated mouse oocytes and preimplantation embryos. *Biol. Reprod.* 71:1330-1339.

VanRaden, P. M., A. H. Sanders, M. E. Tooker, R. H. Miller, H. D. Norman, M. T. Kuhn, and G. R. Wiggans. 2004. Development of a national genetic evaluation for cow fertility. *J. Dairy Sci.* 87:2285-2292.

Wang, G. S., and T. A. Cooper. 2007. Splicing in disease: disruption of the splicing code and the decoding machinery. *Nat. Rev. Genet.* 8:749-761.

Wang, X., V. Schutzkus, W. Huang, G. J. Rosa, and H. Khatib. 2009. Analysis of segregation distortion and association of the bovine FGF2 with fertilization rate and early embryonic survival. *Anim. Genet.* 40:722-728.

Wang, E. T., R. Sandberg, S. Luo, I. Khrebukova, L. Zhang, C. Mayr, S. F. Kingsmore, G. P. Schroth, and C. B. Burge. 2008. Alternative isoform regulation in human tissue transcriptomes. *Nature* 456:470-476.

TABLE 1

Primers for total expression and alternative splicing analyses of HSP genes				
Gene		Family forward primer 5'-3'	reverse primer 5'-3'	amplicon size (bp)
GAPDH		TGCCCGAATATCATCCC (SEQ ID NO: 3)	AGGTCAGATCCACAACAG (SEQ ID NO: 4)	134
DNAJC5	HSP40	CTACGACAAGTACGGCTCAC (SEQ ID NO: 5)	GGCAGCAGCAACAGTAGC (SEQ ID NO: 6)	149
DNAJC15	HSP40	AGGTCGCTACGCATTTCAG (SEQ ID NO: 7)	GACTTGCTTCTCGCCTACTC (SEQ ID NO: 8)	137
DNAJC19	HSP40	GGACTGACCATTGCTGCTG (SEQ ID NO: 9)	CAAAACCCACCTCTGTAATAGC (SEQ ID NO: 10)	140
DNAJC24	HSP40	GAAATATGGGACCAGTAGATGC (SEQ ID NO: 11)	TGTAACCTTCTTCTGCTTCATCC (SEQ ID NO: 12)	134
DNAJC27	HSP40	AACAAGCGGACACCATTCG (SEQ ID NO: 13)	TGAAGCAGCACAGCAAGC (SEQ ID NO: 14)	124
HSPA5	HSP70	CAACCAACTGTTACCATCAAGG (SEQ ID NO: 15)	AAAGGTGACTTCAATCTGTGG (SEQ ID NO: 16)	133
HSPA9	HSP70	GACCAACTGCCTGCTGATG (SEQ ID NO: 17)	GATGCCCGCTGCCTTATG (SEQ ID NO: 18)	113
HSPA8	HSC70	CGCAGAAGCCTACCTTGG (SEQ ID NO: 19)	GTTGAGACCAGCAATAGTTCC (SEQ ID NO: 20)	115
HSPA14	HSP70	AACCTTAGCACAGTACCTAGC (SEQ ID NO: 21)	TGTCAGCACCGTTCATCAG (SEQ ID NO: 22)	101
HSBP1	HSBP	CATGTCCGACCAGATCATTG (SEQ ID NO: 23)	TTCACTGTCCAGCTCTTCC (SEQ ID NO: 24)	115
HSPE1	HSP10	GCAAGCAACGGTGGTAGC (SEQ ID NO: 25)	ACTTTGGTGCCTCCATATTCTG (SEQ ID NO: 26)	117
DNAJB1	HSP40	GAGGAGAAGTTCAAGGAGATCG (SEQ ID NO: 27)	TTAGTACCGCCGCTGCTC (SEQ ID NO: 28)	134
DNAJB12	HSP40	GCAAACCTAGCCCTCAAATTCC (SEQ ID NO: 29)	CCTTGTCATACCGAACTGG (SEQ ID NO: 30)	141
DNAJC14	HSP40	GTGAATGAGTTTCTGTCCAAGC (SEQ ID NO: 31)	GTATCTGGCACTCTTAGGTTCC (SEQ ID NO: 32)	114
HSPA2	HSP70	ACGCTGTGGAGTCCTATACC (SEQ ID NO: 33)	TTCCGCCATCTGGTTCCG (SEQ ID NO: 34)	146
HSPA4	HSP70	TCCTGCCTTAGAAGAGAAACC (SEQ ID NO: 35)	CCCAGTGTGTGTCAAATGC (SEQ ID NO: 36)	132
HSPH1	HSP100	ATGTTGAGTTGCCTATTGAAGC (SEQ ID NO: 37)	CCTCCACCGCATCTTAGC (SEQ ID NO: 38)	141
<u>Transcript</u>				
DNAJC5-667		AGGAGACGGAGTTCTATG (SEQ ID NO: 39)	CACGTTACACCTCAAC (SEQ ID NO: 40)	126
DNAJC5-725		GGCCCTGTTTCATCTTCTG (SEQ ID NO: 41)	GGCACAGACCCTCTCAT (SEQ ID NO: 42)	181
DNAJC5-1049		GGGTTCAACTAAATCCAGGA (SEQ ID NO: 43)	ACGCCATCTCTGTGACTA (SEQ ID NO: 44)	79
DNAJC19-475		GGATCTCCTTATATAGCAGCCA AA (SEQ ID NO: 45)	AGCCTTCCCTCCCAGTGA (SEQ ID NO: 46)	93
DNAJC19-536		ATGCTCATCGGCGAATTATG (SEQ ID NO: 47)	AGCTGGAACGCATAAGAGAA (SEQ ID NO: 48)	161
DNAJC24-628		CTCATTTTAATGGAAGATG (SEQ ID NO: 49)	GTATCACAAGAAATCAGT (SEQ ID NO: 50)	176

TABLE 1-continued

Primers for total expression and alternative splicing analyses of HSP genes			
Gene	Family forward primer 5'-3'	reverse primer 5'-3'	amplicon size (bp)
DNAJC24-2172	CATCCAGATAAACAGAGT (SEQ ID NO: 51)	GGTCCCATATTTCTTAGAT (SEQ ID NO: 52)	161
DNAJC24-1893	CAAAAGAAAGTATCTCATTC (SEQ ID NO: 53)	TAACCTCTTCTGCTTCATC (SEQ ID NO: 54)	176
DNAJB12-1220	ACCGACTGTGAGAGACTATG (SEQ ID NO: 55)	CGGCCTCCAATTCCATTT (SEQ ID NO: 56)	127

TABLE 2

15

TABLE 2-continued

Blastocyst and fertilization rates for genotypic classes of DNAJC15 and DNAJC27 genes				Blastocyst and fertilization rates for genotypic classes of DNAJC15 and DNAJC27 genes			
DNAJC15 SNP	genotype (number of ovaries)	blastocyst rate $\pm$ SE	P value				
SNP 85146(G > A)	AA (202)	0.310 $\pm$ 0.025	0.014	20	AG (201)	0.280 $\pm$ 0.023	
	AG (206)	0.281 $\pm$ 0.025			GG (63)	0.369 $\pm$ 0.037	
	GG (53)	0.401 $\pm$ 0.040					
SNP 85161(G > A)	AA (202)	0.309 $\pm$ 0.023	0.022	25	DNAJC27 SNP	genotype (number of ovaries)	fertilization rate $\pm$ SE
	AG (206)	0.283 $\pm$ 0.024			SNP 35728(G > A)	AA (126)	0.621 $\pm$ 0.029
	GG (61)	0.390 $\pm$ 0.037				AG (257)	0.654 $\pm$ 0.025
SNP 85216(A > G)	GG (203)	0.312 $\pm$ 0.022	0.059	30	SNP 36016(C > G)	GG (103)	0.691 $\pm$ 0.030
	AG (201)	0.282 $\pm$ 0.023				CC (129)	0.622 $\pm$ 0.029
	AA (63)	0.373 $\pm$ 0.036				CG (251)	0.653 $\pm$ 0.026
SNP 85292(C > A)	AA (204)	0.310 $\pm$ 0.023	0.012	30	SNP 38867(A > G)	GG (103)	0.693 $\pm$ 0.030
	AC (201)	0.279 $\pm$ 0.024				AA (133)	0.626 $\pm$ 0.028
	CC (62)	0.395 $\pm$ 0.037				AG (246)	0.650 $\pm$ 0.026
SNP 85300(G > A)	AA (208)	0.312 $\pm$ 0.023	0.066			GG (104)	0.693 $\pm$ 0.030

TABLE 3

Primers used for SNP identification, product sizes, and GenBank accession numbers of amplified genes			
Primer	Primer sequence (5'-3')	Amplicon (bp)	GenBank Accession no.
DNAJC15-1	F: CCGGAGGTCTGCAAAATGGG (SEQ ID NO: 57) R: AACTGCTCGCCTGGTGTGGTC (SEQ ID NO: 58)	589	NC_007310
DNAJC15-2	F: TCACTGAAAATCAGCCAATA (SEQ ID NO: 59) R: CGTACAGAAGAGCCCCAT (SEQ ID NO: 60)	569	NC_007310
DNAJC15-3	F: AATTGCTTTATTACTTTAGCGG (SEQ ID NO: 61) R: AGGGACCATGTCTGTTTGT (SEQ ID NO: 62)	647	NC_007310
DNAJC15-4	F: AAAGTCCCTGTAGAGCTTAG (SEQ ID NO: 63) R: ATAAAGGCACATCACAATA (SEQ ID NO: 64)	571	NC_007310
DNAJC15-5	F: TCCTCTGTCTAGTTCTTG (SEQ ID NO: 65) R: TTCATTATGCCCAATCAGT (SEQ ID NO: 66)	647	NC_007310
DNAJC15-6	F: CCATCCACTTCAGAAAATTC (SEQ ID NO: 67) R: GGGGAAAGATCAGTGCTAGAGT (SEQ ID NO: 68)	620	NC_007310
DNAJC19-1	F: T TTTCCGACCTAGTTTACGG (SEQ ID NO: 69) R: ACTTCTACTTCACCACAGGGA (SEQ ID NO: 70)	670	NC_007299
DNAJC19-2	F: AGCCGCATACCTTTACAATG (SEQ ID NO: 71) R: ATGGGTCACCTCAGATTTCCT (SEQ ID NO: 72)	696	NC_007299
DNAJC19-3	F: TCCCAGAAGAACTGGGTTTG (SEQ ID NO: 73) R: AGATGGAAGTCCCTGGCAGT (SEQ ID NO: 74)	496	NC_007299
DNAJC19-4	F: TTAACTCACTGGAGGTAGG (SEQ ID NO: 75) R: ACCAAAACAGCAAGTAGACC (SEQ ID NO: 76)	597	NC_007299
DNAJC19-5	F: AGGGATTGTTGATAACTGGA (SEQ ID NO: 77) R: TTTATACCAACGCTTTGACT (SEQ ID NO: 78)	611	NC_007299

TABLE 3-continued

Primers used for SNP identification, product sizes, and GenBank accession numbers of amplified genes				
Primer	Primer sequence (5'-3')		Amplicon (bp)	GenBank Accession no.
	(SEQ ID NO: 77)	(SEQ ID NO: 78)		
DNAJC24-1	F: GCTCGGCTGGAACTTGA (SEQ ID NO: 79)	R: CGGTGGATGGCCCTCTAA (SEQ ID NO: 80)	672	NC_007313
DNAJC24-2	F: CCATTCCTTCTCCACTAGT (SEQ ID NO: 81)	R: CTTTGTATTCCTGCTTTGAT (SEQ ID NO: 82)	630	NC_007313
DNAJC24-3	F: CCCTTCTGCTTTGTCCATC (SEQ ID NO: 83)	R: ACCACATTTCTGGGTTGCTC (SEQ ID NO: 84)	538	NC_007313
DNAJC24-4	F: GGAGACTTTTGGCCTAGTGT (SEQ ID NO: 85)	R: ATAAAGTTTTCAGGTGGGAA (SEQ ID NO: 86)	637	NC_007313
DNAJC24-5	F: TAAATAAATTCACCTGAA (SEQ ID NO: 87)	R: ACAATAGCCATGTTTCTGA (SEQ ID NO: 88)	649	NC_007313
DNAJC24-6	F: GGCAACTGTAGAAAGGATAG (SEQ ID NO: 89)	R: TATAAAGAATAAGCACCACA (SEQ ID NO: 90)	539	NC_007313
DNAJC24-7	F: TGGTATTATTATTGGTTTGAC (SEQ ID NO: 91)	R: AGGAGAAAGGATGACAAGG (SEQ ID NO: 92)	531	NC_007313
DNAJC27-1	F: CTCCTCCAGTTCCTACCC (SEQ ID NO: 93)	R: A CAGCCCAGTAAGTTATCAGC (SEQ ID NO: 94)	665	NC_007309
DNAJC27-2	F: TTGAAAACATACCCATATTGG (SEQ ID NO: 95)	R: ATCACTAAAAGGAAGCTCTC (SEQ ID NO: 96)	682	NC_007309
DNAJC27-3	F: CTTCACTCTGCTCCTACTGTCC (SEQ ID NO: 97)	R: CAAAGGGTTGGTTCACCTCTG (SEQ ID NO: 98)	647	NC_007309
DNAJC27-4	F: AGGGACAGGGTAGAAGGC (SEQ ID NO: 99)	R: CAAACATGGCACCAGAAA (SEQ ID NO: 100)	607	NC_007309
DNAJC27-5	F: GGCAACGTGATGAAGCC (SEQ ID NO: 101)	R: GACCTGGAGCCCAGCAAT (SEQ ID NO: 102)	692	NC_007309
DNAJC27-6	F: AGAGTAGGATCATAAGCCATTT (SEQ ID NO: 103)	R: TTCGGTGAAGGAGTAGTGTT (SEQ ID NO: 104)	532	NC_007309
DNAJC27-7	F: TGCAAGAGGTGTTCTGTAT (SEQ ID NO: 105)	R: TTTCAGGGTTCTACTATGT (SEQ ID NO: 106)	666	NC_007309
DNAJC27-8	F: ACATAGTAGAACCCCTGAAAGT (SEQ ID NO: 107)	R: ATGATGCTGCAACAAGGAAA (SEQ ID NO: 108)	500	NC_007309
DNAJC27-9	F: TGCCAAGACAGGTGGGAAAT (SEQ ID NO: 109)	R: CAGGTAGGGTAAGGCGAATG (SEQ ID NO: 110)	613	NC_007309
DNAJC27-10	F: ATTGCCTTACCCTACCTGA (SEQ ID NO: 111)	R: CTGGGAAGTGAAGACCTAA (SEQ ID NO: 112)	644	NC_007309
DNAJC27-11	F: CTTAGGTCTTGCTCAGTTCC (SEQ ID NO: 113)	R: CACAACATCTCCAAGTCCAG (SEQ ID NO: 114)	668	NC_007309
DNAJC27-12	F: ACTTGGAGATGTTGTGCTGC (SEQ ID NO: 115)	R: TATGAACCCCTCTCCCTTT (SEQ ID NO: 116)	565	NC_007309
DNAJC27-13	F: GTGTCAGTATCTGTCCCTAA (SEQ ID NO: 117)	R: TCTATGATCTCAGTCGGTAA (SEQ ID NO: 118)	681	NC_007309

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 118

<210> SEQ ID NO 1
 <211> LENGTH: 4159
 <212> TYPE: DNA

-continued

<213> ORGANISM: Bos taurus

<400> SEQUENCE: 1

```

tagcaattca gatcttgatg tgaaaatgtc agtgcctccc tcaagggtgg aaagctaact      60
atagctcctt gagcgccacc cttccattgc tgattcattt cctgcttccc ttttccttct      120
caaatgggtg gtgtgataac agtaaaaggc ccagataaag tcaggacagt gcatacgcca      180
ctgctgtcta gatgttttgt gtgtgtgtgt ggtttgttgt gagccactca ggtcttaaga      240
tgttttcaac agctgtggat catttgtact caagggggag aataaaaaat aaaaaacaaa      300
cacagagact ggtgagctgg actggttagct tgagtgctaa gcatgaaagt gttcaggact      360
agaagttaga aggtgtatga tgcttaataa tttttagtgt gacagtttgg ggccagtatt      420
tttatatgta agggagctcc tgtttgtcat ttttaacgat tctgaatttg aaattcacct      480
cagttttcct ctggttcttt acgttagaaa taattcattg tgaagaatta tgaaaactga      540
cttcatttta ttttaaggcg agtgacgctg tctgtgctcc ataagtattc tgatggcact      600
gtgtcagaga gcagggggca taccttttaa cgttattttt atttactctg attttttatc      660
atattattta ctaattattg ccaatattag tgaatgctt ttagagctga gtagtatata      720
gttgcttttt taacttattt ttatgaaaat gtatggtttg ataaaacttt gaaggcacc      780
ttggattgct tcacctcctt cagaaagata acttccttca tagaagaatg ttctttcctt      840
aaaaacaaac agaaaaaatt acgtcttggg gaaggactgt tattttattt acctaaccat      900
ccacttcaga aaattctctg gtcagatatt tggcatttga aaggtaatgt ctgaccagtg      960
aaatgtttga gtgtgtttta aattactcct gatcataagt ttatttttct cagagattat     1020
ccttgactgt taggtatctg tagtgcttca ttgaacgagt ttaacttata ctgttgtgat     1080
ttctagagag cttataaaat aatttgtcag ggcattgaatg tgtgggaaaa tatcagagac     1140
atctaactga gtgtttttct ttatttgaga ctttaaaaaa tgtcagtagt gaagtgtcct     1200
ttaatccatt ttcttttctt tgcctcttca ggtgggtctc cttacttagc aacaaaaatt     1260
aatgaagcaa aagacctgct agaagcaacc accaaacact gatggacgct caaggaccac     1320
tctgagggag aagaaggggg gccttataaa agctgtcctg cacattagtc taaaccgtgg     1380
cattcacaat ttcatatgtg caccgaccac agccatttct tccgtgtctg agctatgtga     1440
taataaaaga tgaacagtct agctttgtat tcgctttcag aggagcacat ggaaactcta     1500
gcactgatct ttccccctca ttttcttctt ctctctgata gtcatacatt tgaagacttc     1560
tactactaag ttcccccgac ccagccccct gcagcagtat cattatcagt aatataattg     1620
gaatttggga gcaagagtgt agaatttttt gctgcttata tttatgctcg cattgtccct     1680
tgtttcttca aaggaaattt gcactctgcc tatgtgaagt gctttaaatg ttatgagacg     1740
aaacctgac tgccacccca ggaaagtga ctgaagcttg atggaattta ctaaagggtg     1800
ttttgtaaag cagtgggttag gaaagaacgc tgctctccca gagatcccaa ggctgtagca     1860
tgagaacttg gagagctgct aaaatgatta aaggtttaca gaatgagccc tgtgaggaaa     1920
gcctggaaat tgctgggaga agtctgaggg attcgtactt aattatatag tcttctggtg     1980
ctgtgtactt attcaattat gcattctaca aatatttatt taccacctgt agcgggtgag     2040
gtactgtagg ggtcctcagg agttccaggt gtgaaccagg cacggatcgt gacccaaagg     2100
aacttgaaat tagaggaaag ataaaccaca ggagtactat tagagaggca cagatgatag     2160
gcaaagaagt tcaaaataaa agactatttt aactgggca tcaggggggc gtgtcataga     2220
atcagtagcg tagagctggg ctttaaaaac aatgagatat accatgatag gggtagtca     2280

```

-continued

gaattagcag aaatagggtg tattagttca ttggagtgcg gggttttctc aaagtaacgg	2340
atgaacacac agtctggaag ggctggccct gtaatgtgca gtgcattcaa ttattttttt	2400
tttccccctc aacacagccc ctggttatca tcacttttct aaaagggaga aaagagagag	2460
tcagaatacc agcatatttc ttgtactgct ttacgacgct gggctcatcc caggtaaaca	2520
gaactaccac gtttcatctg tcttaaacgc tcttgtaata gtctagaggc atctttcatt	2580
ttacgacato ctgtagttgg gagttctcag cattggtgat actgactcat tgagctgctg	2640
attctgctt gtggggggct ctctgtgta tggtaggatg attagcaaca tccctgctg	2700
caactacgaa aggccaggag cgtccctctt gttgtgacaa ccgaaaatgt atctagatgt	2760
tgccaaacgt tccttgagg gcaaaatgc ccaggtaga gagccactgc tctaaatgtt	2820
aaacctgttc tccttggtg catgaaataa tggatggcct gacttgaat tgattcttag	2880
gttcaactgaa atacggctct cgttttaaaa tccctgtgt gtccacagtt tttgtgctt	2940
agtatgttt ctggccct gctgcccgg tactttaagt gccctcctga taccagatc	3000
tccactgtt caaagcctgg ggatagttga aggccaaact gaaaagctgt tctctcttt	3060
tctggtgctt ccagcaaaag taatcccatg aattgtcag tctcagcaaa caaggattgt	3120
atctgtcttg gtcactgcta cattgacact caataaatat ttgttaaat aacgaataca	3180
ttcccgtagc actttatttc tcacatttgg taacttgtga acaactctt tgtgtgacat	3240
gacttatttc ttctacaaaa gtacaagaac ctttagagca agaactctgt catttgtctt	3300
tgtaacctcc ttaagagtgg agcacttggg actttttaga catcataaag ttgtaagtac	3360
taaaaaaaat gttttttgaa tactatgctt caaaaaatgt aaccactagt tttgaaagt	3420
atgttatgca atattgggg accctcagaa catacctgtt agttgacatc attttctagt	3480
ccactctgtt tcatgaaaa ttaagtcc cagcactagc ttgaaatttc cttcattcta	3540
ttctcattta tctgttagt gtgtgcccc aaatcagaca cctggtttta agtactatcc	3600
tcctatcaca taatctgggg caattaatct gcaaggcttt aattgcctca tctataaaat	3660
agagaaaata ataacctat ctaaaaagg ccatctagtc aaggctatgg tttttccagt	3720
ggatcatgat ggatgtgaga gttggactgt gaagaaggct gagcgccgaa gaattgatgc	3780
ttttgaactg tgggtgttga gaagactctt gagagtcctt tggactgcaa ggacatccaa	3840
ccagtcctc ctaaggaga tcagtcttgg gtgttcattg gagggactga tgctgaagct	3900
gaaactccaa tactttggcc acctcatgag aagagctgac tcattggaaa agacctgat	3960
gctgggaggg gttgggggca ggaggagaag gggacgacag aggatgagat ggctggatgg	4020
catcaccaac tcgatggatg tgagtttggg taagctccgg gagttggtgc tggacaggga	4080
ggcctggcat gctgcgattc atgggggttc aaagagtcgg acacgactga gtgactgaac	4140
tgaactgaaa aaggttgtt	4159

<210> SEQ ID NO 2

<211> LENGTH: 6741

<212> TYPE: DNA

<213> ORGANISM: Bos taurus

<400> SEQUENCE: 2

ttgctcagtt gtgcccact ctttgcaacc ccatggactg tagcctacta gggtccttgc	60
tccatgggat tctccaggca agagtaccag agtgggttgc catttccttc tccagaggat	120
cttctgacc cagggatcga acccaggtct cccgattcc aggcagacac tttaacctct	180

-continued

gagccactag	ggatgggaag	agggagacag	ttattggtca	cccttgattt	gacaagcgtt	240
actgaggatg	ctttaaacac	ctgacgaaat	tcagatgacc	ttgtagcgta	aagttaattg	300
gtggcttttg	gggttccca	ggtagcacag	tggtaaagaa	tctgcctgcc	catgcaagag	360
atgcaagaga	cttgggttcg	atccctgggt	ggggaagatc	ctctgaagaa	gggaatggcc	420
accactcca	gtattcttgc	ctggagaatt	ccatggacag	aggagcctgg	tggtctacag	480
gccataggtg	tgcaagagt	cgggcacaac	tgagcaacta	atattacttt	tactattgta	540
tatcaaattc	agcttttttg	ttgttttgtc	ctttccctta	tggaccctgc	tccttggtcg	600
actttgtccg	ctgtttgctt	ctggaactac	atgccgtgtc	tctgttcctc	ctctggcctc	660
tgccatttct	ttccctgegg	ctctgtgttt	cccatacccg	gatcaattcc	tgaaacccat	720
cttcttgcat	agtcctcttc	cagggggggc	ttgccctgat	ggcctcttgt	cctcatcgct	780
tcctacagca	attatttgct	gagtgcctaa	tgctgtgag	gcgctaacca	ctaattttca	840
gttgattatg	ttgtataact	cactttgtat	tgtctgtttt	tctacttgaa	tggtacatat	900
tatttgtcca	ttgtttgttt	cccagtcctt	tagtctaacc	tttgagaaag	ggtcacgata	960
atttcagata	tgcttactgc	aagaggtggt	ctgttatcca	ggcagctgaa	atgcagtgga	1020
cagaggggtg	ggctgacctc	cctactgaca	tagagetctg	ctttttcttt	gcagggatga	1080
agtcaataaa	gcataatcga	agcttgctgt	gctgcttcac	cctgacaaat	gtgtagcacc	1140
tggtagttaa	gatgccttca	aagccgttgt	gaatgcccga	acagccctcc	tgaaaaacat	1200
caagtagaaa	ataggaaaaa	aagacaggtg	taggcctcga	gtccaaacag	actttctctg	1260
gaagtaaaat	agccaacatg	ggtttttcc	ccacaaaatt	cttacagccc	ttttactca	1320
cgtgctgtca	tttgtgaatc	agtaatttgg	ctgcctgttt	ccatgtcaca	gacattttgc	1380
agagagacaa	aatgaagaga	accgagtgcc	acttatacac	tgtcgttcat	tttctatttc	1440
tgaatgatga	tagatttttt	ttttttctc	atgagttggt	aagctgtcgg	tatagcatca	1500
cttgggcact	ttgctcagga	aagtccacaa	agtttctgga	agaaggatag	gaggatcctt	1560
ttgctttttc	tttattaaaa	aaacattcat	catcagagga	aaaaaaaaa	aaaatggaag	1620
cagacatagt	agaaccctcg	aaagtgtggt	gactttacaa	gcaagttgcc	cctttacaat	1680
gagtttctaa	ggatgatatt	taagcatcag	tctgcttaat	taatgaatga	actcttgcca	1740
gcccttcgga	agtgaacaaa	ggatgatgct	tcctgagctg	agtgccctt	ttttgacgtt	1800
cggctgtgtc	cccttctcca	tcaccggggg	agagaccagc	cagtcctgag	agtagaggtg	1860
gcggtagccg	caggttctgg	tagactggct	gctgctagac	ctgcccctgg	gtctgaggcc	1920
tttccctggc	cgtgggagtg	ctggtagctg	taggggcata	ttcacctagg	agtctggagt	1980
cagatcccaa	gacacaaatg	cctgcaggtc	aaactggatt	gtgtacacat	tcacagaggg	2040
cctgaccctc	ctggttcggg	cgtgtctgat	gccctaggag	agccccata	ggaggcttat	2100
gcctttcctt	gttgacagat	catttgccaa	gacagtgagg	aaatgcaggg	ttgcgtctca	2160
acagggggcc	tccttctgct	tcctcgtcag	tgtgagctcc	ccttccctct	tctgttcctg	2220
cctgtctctc	caaggcaccg	catctgccac	cacgtctctc	ttctcgaatc	cccaccttcc	2280
ggctattgcc	tcctctgtcc	tgtgcagcct	ttgctggggc	ctggtagacc	agagacaagt	2340
atagatcctt	ggggtggtgg	atggagagat	cttaaggcct	gtgtgggaag	aatttgttac	2400
ttttgtttag	gttccaaatt	tctgatcagt	aatactttta	agctgcagat	tatgaaaatt	2460
aatttcacat	agagaaatcc	attctcttgc	tatcgtctgt	ctctcagtat	ccaagagatg	2520
gatgaagatc	aagtgtcagg	accggccttc	cttcagagac	tcacagaata	gagtcaccac	2580

-continued

gtggcgtggt	acaccatcac	tctcacgtta	ctcagattgt	tagctgtgac	caagctgcac	2640
taattggggc	tttaaagtat	atgaagcttt	gccatcagtg	aattttaagc	agctatccca	2700
tattgattgc	caagtagcat	tcgccttacc	ctacctgaga	agaaaaacca	ctttcattta	2760
attcaaaacc	actgcagctt	actgaaccct	tcctttctcc	agagcagctt	cgctttgtgt	2820
ggctactctt	gggcagcact	gagctggttg	tcaggaaacc	agagcattgt	tttctagtaa	2880
gagttttcat	tttgatatta	aaaatgcttt	tttttcttag	taaaattgag	atatgagata	2940
tgattttttg	tgggattttc	ctcttgccct	ttaattttta	ctagcttctg	ggcctgtaaa	3000
actcaggcca	aagttcttca	gggttttaat	tttgctgttg	gccttcacaa	tggccaaagt	3060
aacaacata	gatctgttct	tgtggcagaa	cttaagagtc	acatgagtga	ggaactgtgc	3120
atggtggtgt	cagcccatat	tgacatgaca	gctcacttgt	gtatgtggcc	ctgtgtaaaa	3180
actggtgcca	gttttcatca	ctgtcattaa	acactagatg	ggtcacctca	ttcagctaca	3240
ttgtgcagat	gagcagttat	agcagataga	gttctctagt	tcttccattc	acatccttat	3300
gtaatgagtg	acttggtgac	atttagcctg	gtcttgatct	ttaggtcttg	ctcagttccc	3360
agcaccttga	ttactaatgt	taaggatggc	catgtactac	ttcatttatg	gaaaagaaag	3420
aaaagaaaga	taaaggttca	gggtcctaga	atttttctga	tttcataaat	cctgagtaga	3480
atgtcaagaa	gtaatgaaat	gaagcctgat	ctgcaatgaa	aagtaacttc	atataaaatt	3540
ttacagacct	ctgtgcatta	atttaaattc	ttggtacaaa	tgtatatttt	ttagttcaac	3600
cttattaaaa	ttcttcagcc	agttaggatt	gcacaataca	tcagtgaagc	taacaactag	3660
cagaataatt	tcaatggcta	aaaaaaattt	tatacagaat	cttggtttttt	caaaaaagaa	3720
caagaagttt	gaacagctgt	tattcaagaa	aagccatttc	ttgggcatct	gagtcaccaa	3780
gaatgaacac	tttaaaaact	tgttagaaat	ccctggaggt	ttttgaaatt	aatcgccaac	3840
ctacctaaga	acttaaaaat	ataccatcct	ttgttctctc	ccttctctct	tcatggtctc	3900
ttttgaatca	atattctcaa	ctgtcaacgt	gtctcttact	tcagcaatgt	cgtttatttc	3960
tttgcatgaa	tgtgtgcttg	gtaaagtctg	gacttggaga	tgttggtgctg	cttcccaaga	4020
taaattgctc	tgattctgta	gattgcactc	tgtgctttta	ggctcacatt	tacctcagga	4080
aactcacttt	agtatgtttt	gttttgtttt	gggttttttt	gcattgaatt	ctcaaagggt	4140
aattatttct	ttatactggt	aaactagtac	ctaatttact	gttcactaag	tgagccatag	4200
ataaggcagc	taaattttctg	aaagaaagaa	aaaataggca	gcccttgatt	gctgtgaatc	4260
agtaataagg	tcctattttc	tcaaagagca	atgacagaat	ctcaagggga	aggaggtatt	4320
tcgggtctcat	gttagaaaag	gggggggttg	tgacacattgc	ctttgacagg	gtgtcagtat	4380
ctgtccccta	agcaatttca	gtgggccccta	gaatgtcact	cccttctcta	ctacactctg	4440
tctagctcac	tctgagtctg	actacatttt	tgtgaaactt	ttgcagagaa	attaagccgc	4500
aaaatggagc	taacctcctt	ccattgcaag	ttttgcaaag	ggaagagggg	ttcatatttc	4560
tcatgatagg	tacctagcac	aattgtaaga	gacagttact	aatgtttata	tgttggtggt	4620
tccattaagc	atttctagag	gagacgaaag	cttaaaacttg	tgggtgtggat	gagttagatc	4680
gtgtctgaca	agcagtgcac	tatgaagctg	tgcaataaag	gctgtgttgt	gaaatgaagc	4740
actggaaaag	tggacagtcc	gggttagctt	cctgcagcaa	ttgcccgcag	gatggctctgc	4800
ccttggccca	ggaatgcaaa	gaaacccttt	gctcttacga	ggtaatttat	gtatttgctt	4860
tctgttgaaa	taattgaaaa	tgttaaagct	aacagctgtc	tatgtcttgt	atagagagtg	4920

-continued

tcatatgtat ttaagttgtg tatttttata aagtaaagcc aaatgccaaa tactggtctg	4980
tgggctgcac tcatttgttg agacactggt ctttgaagca gaagctggcc gttaccgact	5040
gagatcatag ataagacatc ttgctttagt tgccagtggg taaaatgaag gaaatggtat	5100
tcagtgggtcc ctcaggctgt tctaacctta attaatgttg gatgaaagtc tctgctctct	5160
taaagaaaga tactaatatt atgtttgcct ttttaattca aatgaaacta ttgttatctc	5220
attttaataa tgcaggcteta attcagctaa ttagtccctg gacaacagga atccttaagt	5280
aacagactat ggtgaatgcg aatttgaaag taaaagcctt gtaggccag cccacagaag	5340
gcttcaaacc agagctctaa ttttcattgt gagtgacgtt ccctatcacc tgagaatgcg	5400
ctcactcctg gggaaccgct ctaccctctc catactgctg acatgttctg aagcagagct	5460
gccaaatcct cctatgacct ccccttgct ccttggtcac agagcttggc ccgagagatg	5520
ggaatgtcac ccaagctctg gagaaggaaa tggcagccca ctccatggaa tcccatggac	5580
aaaggagcct ggcaagtcca tggggtcgaa agagttggac acgacttagt gactaaacca	5640
gcaccaccac cacctgagat ctttctagtt ggaatctagg aagacagact ttctctcagg	5700
agagacagat aggaggagtc ctctctgta tgctgacaac cgtgatttca acctcatgga	5760
aaaaatcaga gcgaatgaaa cagacattca gagactgtat tggtaggca aagatgttcc	5820
ttgcagtccc cagggtcccc tgaactggct gctctacaag cccatcagct accccagaat	5880
cctcccaatc aatctccact ccttggtttt ctgcttaagt ttgttggagt ttgctttctg	5940
ttacttgcaa ccaaaaggat attagaaaat aatagacatt aagggtcata agcaggaatg	6000
agaaagctat ggtctgtaga ctggtacagt acttgataag taaagtgtga cattttagat	6060
actaagtgtc ggggtccagc cctggttgat ccagggaatt tgaagtgggg acgacgttgg	6120
cgaggatcag gaaacaactg cttaattaaa catatttaa ggatataaag agtaatagaa	6180
tgaggatagc tcagttagga aactcagtgg agaaaggcgg ctgaaataag gatagctcag	6240
tgaggaaatt cagtggagat aagaggctga aataattcag ccagaaggtg agagaaagaa	6300
cgacatgggg agaccaagtt tcagtgaaca aggccgcac tttattttcc aaagtagttt	6360
ttatacctta agttatgcat agaggataat gggggaaggg gtagagtcac gcaaggacag	6420
cagttcctga tcctaatega agccaggctt tcaaacatat catatgcaa agtttagatg	6480
atttacctca tcacttggtc aggaggcctg ttaacatttt aagaaactta tttttctcta	6540
aaggtgatta ttctaaagtc aggcgccagc ctccaaaaa gcattggata aagttgcatt	6600
cctatagggc aaaggtgtgg tgggtacaa caaaaaagaa ttaactcaag ggtccaaggt	6660
tacaaacatt aaagctacta cttacatcaa ttatattaat caatacactg ccaaggacac	6720
agtaggtaag gggtagggag a	6741

<210> SEQ ID NO 3
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer GAPDH-F

 <400> SEQUENCE: 3

tgcccagaat atcatccc

18

<210> SEQ ID NO 4
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: primer GAPDH-R

<400> SEQUENCE: 4

aggtcagatc cacaacag 18

<210> SEQ ID NO 5
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-F

<400> SEQUENCE: 5

ctacgacaag tacggctcac 20

<210> SEQ ID NO 6
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-R

<400> SEQUENCE: 6

ggcagcagca acagtagc 18

<210> SEQ ID NO 7
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer DNAJC15-F

<400> SEQUENCE: 7

aggtcgctac gcatttcag 19

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-R

<400> SEQUENCE: 8

gacttgcttc tcgcctactc 20

<210> SEQ ID NO 9
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-F

<400> SEQUENCE: 9

ggactgacca ttgctgctg 19

<210> SEQ ID NO 10
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-R

<400> SEQUENCE: 10

caaaccacc tctgtaatag c 21

-continued

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-F

<400> SEQUENCE: 11

gaaatatggg accagtagat gc 22

<210> SEQ ID NO 12
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-R

<400> SEQUENCE: 12

tgtaacttct tctgttcat cc 22

<210> SEQ ID NO 13
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-F

<400> SEQUENCE: 13

aacaagcggg caccattcg 19

<210> SEQ ID NO 14
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-R

<400> SEQUENCE: 14

tgaagcagca cagcaagc 18

<210> SEQ ID NO 15
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA5-F

<400> SEQUENCE: 15

caaccaactg ttaccatcaa gg 22

<210> SEQ ID NO 16
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA5-R

<400> SEQUENCE: 16

aaagtgact tcaatctgtg g 21

<210> SEQ ID NO 17
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA9-F

-continued

<400> SEQUENCE: 17

gaccaactgc ctgctgatg

19

<210> SEQ ID NO 18

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSPA9-R

<400> SEQUENCE: 18

gatgccgcct gccttatg

18

<210> SEQ ID NO 19

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSPA8-F

<400> SEQUENCE: 19

cgcagaagcc taccttgg

18

<210> SEQ ID NO 20

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSPA8-R

<400> SEQUENCE: 20

gttgagacca gcaatagtcc c

21

<210> SEQ ID NO 21

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSPA14-F

<400> SEQUENCE: 21

aaccttagca cagtacctag c

21

<210> SEQ ID NO 22

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSPA14-R

<400> SEQUENCE: 22

tgtcagcacc gttcatcag

19

<210> SEQ ID NO 23

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer HSBP1-F

<400> SEQUENCE: 23

catgtccgac cagatcattg

20

<210> SEQ ID NO 24

-continued

<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSBP1-R

<400> SEQUENCE: 24

ttcactgtcc agctcttcc 19

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSBP1-F

<400> SEQUENCE: 25

catgtccgac cagatcattg 20

<210> SEQ ID NO 26
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSBP1-R

<400> SEQUENCE: 26

ttcactgtcc agctcttcc 19

<210> SEQ ID NO 27
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB1-F

<400> SEQUENCE: 27

gaggagaagt tcaaggagat cg 22

<210> SEQ ID NO 28
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB1-R

<400> SEQUENCE: 28

ttagtaccgc cgctgctc 18

<210> SEQ ID NO 29
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB12-F

<400> SEQUENCE: 29

gcaaactagc cctcaaattc c 21

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB12-R

<400> SEQUENCE: 30

-continued

ccttgtcatc accgaactgg 20

<210> SEQ ID NO 31
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC14-F

<400> SEQUENCE: 31

gtgaatgagt ttctgtccaa gc 22

<210> SEQ ID NO 32
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC14-R

<400> SEQUENCE: 32

gtatctggca ctcttaggtt cc 22

<210> SEQ ID NO 33
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA2-F

<400> SEQUENCE: 33

acgctgtgga gtcctatacc 20

<210> SEQ ID NO 34
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA2-R

<400> SEQUENCE: 34

ttccgccatc tggttccg 18

<210> SEQ ID NO 35
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA4-F

<400> SEQUENCE: 35

tcctgcctta gaagagaaac c 21

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPA4-R

<400> SEQUENCE: 36

cccagtggtg tgtcaaatgc 20

<210> SEQ ID NO 37
<211> LENGTH: 22
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPH1-F

<400> SEQUENCE: 37

atgttgagtt gcctattgaa gc 22

<210> SEQ ID NO 38
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer HSPH1-R

<400> SEQUENCE: 38

cctccaccgc attcttagc 19

<210> SEQ ID NO 39
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-667-F

<400> SEQUENCE: 39

aggagacgga gttctatg 18

<210> SEQ ID NO 40
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-667-R

<400> SEQUENCE: 40

cacgttcaca cctcaac 17

<210> SEQ ID NO 41
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-725-F

<400> SEQUENCE: 41

ggccctgttc atcttctg 18

<210> SEQ ID NO 42
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-725-R

<400> SEQUENCE: 42

ggcacagacc ctctcat 17

<210> SEQ ID NO 43
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-1049-F

<400> SEQUENCE: 43

gggttcaact aaatccagga 20

-continued

<210> SEQ ID NO 44
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC5-1049-R

<400> SEQUENCE: 44

acgccatctc tgtgacta

18

<210> SEQ ID NO 45
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-475-F

<400> SEQUENCE: 45

ggatctcctt atatagcagc caaa

24

<210> SEQ ID NO 46
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-475-R

<400> SEQUENCE: 46

agccttcctt cccagtga

18

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-536-F

<400> SEQUENCE: 47

atgctcatcg gcgaattatg

20

<210> SEQ ID NO 48
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-536-R

<400> SEQUENCE: 48

agctggaacg cataagagaa

20

<210> SEQ ID NO 49
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-628-F

<400> SEQUENCE: 49

ctcattttaa tggaagatg

19

<210> SEQ ID NO 50
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: primer DNAJC24-628-R

<400> SEQUENCE: 50

gtatcacaag aaatcagt 18

<210> SEQ ID NO 51
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-2172-F

<400> SEQUENCE: 51

catccagata aacagagt 18

<210> SEQ ID NO 52
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-2172-R

<400> SEQUENCE: 52

ggtcccatat ttcttagat 19

<210> SEQ ID NO 53
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-1893-F

<400> SEQUENCE: 53

caaaagaaaag tatctcatc 20

<210> SEQ ID NO 54
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-1893-R

<400> SEQUENCE: 54

taacttcttc tgcttcac 19

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB12-1220-F

<400> SEQUENCE: 55

accgactgtc agagactatg 20

<210> SEQ ID NO 56
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJB12-1220-R

<400> SEQUENCE: 56

cggcctccaa ttccattt 18

-continued

<210> SEQ ID NO 57
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-1-F

<400> SEQUENCE: 57

ccggaggtct gcaaatggg 19

<210> SEQ ID NO 58
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-1-R

<400> SEQUENCE: 58

aactgctcgc ctggtgctgg tc 22

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-2-F

<400> SEQUENCE: 59

tcactgaaaa tcagccaata 20

<210> SEQ ID NO 60
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-2-R

<400> SEQUENCE: 60

cgtacagaag agccccat 18

<210> SEQ ID NO 61
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-3-F

<400> SEQUENCE: 61

aattgcttta ttacttttagc gg 22

<210> SEQ ID NO 62
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-3-R

<400> SEQUENCE: 62

agggaccatg tctgttttgt 20

<210> SEQ ID NO 63
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC15-4-F

-continued

<400> SEQUENCE: 63

aaagtccttg tagagcttag

20

<210> SEQ ID NO 64

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC15-4-R

<400> SEQUENCE: 64

ataaaggcac atcacaacta

20

<210> SEQ ID NO 65

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC15-5-F

<400> SEQUENCE: 65

tctctctgtc ctagttcttg

20

<210> SEQ ID NO 66

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC15-5-R

<400> SEQUENCE: 66

ttcattatgc ccaaatcagt

20

<210> SEQ ID NO 67

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC15-6-F

<400> SEQUENCE: 67

ccatccactt cagaaaattc

20

<210> SEQ ID NO 68

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC15-6-R

<400> SEQUENCE: 68

ggggaaagat cagtgcctaga gt

22

<210> SEQ ID NO 69

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer DNAJC19-1-F

<400> SEQUENCE: 69

ttttccgacc tagtttacgg

20

<210> SEQ ID NO 70

<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-1-R

<400> SEQUENCE: 70

acttctactt caccacaggg a 21

<210> SEQ ID NO 71
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-2-F

<400> SEQUENCE: 71

agccgcatac ctttacaatg 20

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-2-R

<400> SEQUENCE: 72

atgggtcact tcagattcct 20

<210> SEQ ID NO 73
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-3-F

<400> SEQUENCE: 73

tcccagaaga actgggtttg 20

<210> SEQ ID NO 74
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-3-R

<400> SEQUENCE: 74

agatggaagt ccctggcagt 20

<210> SEQ ID NO 75
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-4-F

<400> SEQUENCE: 75

tttaactcac tggaggtagg 20

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-4-R

<400> SEQUENCE: 76

-continued

accaaaacag caagtagacc 20

<210> SEQ ID NO 77
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-5-F

<400> SEQUENCE: 77

agggattgtt gataactgga 20

<210> SEQ ID NO 78
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC19-5-R

<400> SEQUENCE: 78

tttatacca cgctttgact 20

<210> SEQ ID NO 79
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-1-F

<400> SEQUENCE: 79

gctcggctgg aaacttga 18

<210> SEQ ID NO 80
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-1-R

<400> SEQUENCE: 80

cggtgatgg ccctctaa 18

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-2-F

<400> SEQUENCE: 81

ccatttcctt ctccactagt 20

<210> SEQ ID NO 82
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-2-R

<400> SEQUENCE: 82

cttttgattc ctgctttgat 20

<210> SEQ ID NO 83
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-3-F

<400> SEQUENCE: 83

cccttctgct ttgtccatc 19

<210> SEQ ID NO 84
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-3-R

<400> SEQUENCE: 84

accacatttc tgggttgctc 20

<210> SEQ ID NO 85
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-4-F

<400> SEQUENCE: 85

ggagactttt ggcctagtgt 20

<210> SEQ ID NO 86
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-4-R

<400> SEQUENCE: 86

ataaagtttt caggtgggaa 20

<210> SEQ ID NO 87
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-5-F

<400> SEQUENCE: 87

taaataaatt cccacctgaa 20

<210> SEQ ID NO 88
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-5-R

<400> SEQUENCE: 88

acaatagcca tgttttctga 20

<210> SEQ ID NO 89
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-6-F

<400> SEQUENCE: 89

ggcaactgta gaaaggatag 20

-continued

<210> SEQ ID NO 90
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-6-R

<400> SEQUENCE: 90

tataaagaat aagcaccaca 20

<210> SEQ ID NO 91
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-7-F

<400> SEQUENCE: 91

tggtatttat tattggtttg gac 23

<210> SEQ ID NO 92
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC24-7-R

<400> SEQUENCE: 92

aggagaaagg gatgacaagg 20

<210> SEQ ID NO 93
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-1-F

<400> SEQUENCE: 93

ctcctccagt tccctaccc 19

<210> SEQ ID NO 94
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-1-R

<400> SEQUENCE: 94

acagcccagt aagttatcag c 21

<210> SEQ ID NO 95
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-2-F

<400> SEQUENCE: 95

ttgaaaacat acccatat tt gg 22

<210> SEQ ID NO 96
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-2-R

-continued

<400> SEQUENCE: 96
atcactaaaa ggaagctctc 20

<210> SEQ ID NO 97
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-3-F

<400> SEQUENCE: 97
cttcaccttg ctctactgt cc 22

<210> SEQ ID NO 98
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-3-R

<400> SEQUENCE: 98
caaagggttg gttcacttct g 21

<210> SEQ ID NO 99
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-4-F

<400> SEQUENCE: 99
agggacaggg tagaaggc 18

<210> SEQ ID NO 100
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-4-R

<400> SEQUENCE: 100
caaacatggc accagaaa 18

<210> SEQ ID NO 101
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-5-F

<400> SEQUENCE: 101
ggcaaactgt atgaagcc 18

<210> SEQ ID NO 102
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-5-R

<400> SEQUENCE: 102
gacctggagc ccagcaat 18

<210> SEQ ID NO 103

-continued

<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-6-F

<400> SEQUENCE: 103

agagtaggat cataagccat tt 22

<210> SEQ ID NO 104
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-6-R

<400> SEQUENCE: 104

ttcgggtgaag gagtagtggt 20

<210> SEQ ID NO 105
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-7-F

<400> SEQUENCE: 105

tgcaagaggt gttctgttat 20

<210> SEQ ID NO 106
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-7-R

<400> SEQUENCE: 106

tttcaggggt tctactatgt 20

<210> SEQ ID NO 107
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-8-F

<400> SEQUENCE: 107

acatagtaga acccctgaaa gt 22

<210> SEQ ID NO 108
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-8-R

<400> SEQUENCE: 108

atgatgctgc aacaaggaaa 20

<210> SEQ ID NO 109
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-9-F

<400> SEQUENCE: 109

-continued

tgccaagaca ggtgggaaat 20

<210> SEQ ID NO 110
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-9-R

<400> SEQUENCE: 110

caggtagggt aaggcgaatg 20

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-10-F

<400> SEQUENCE: 111

attcgctta ccctacctga 20

<210> SEQ ID NO 112
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-10-R

<400> SEQUENCE: 112

ctgggaactg agcaagacct aa 22

<210> SEQ ID NO 113
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-11-F

<400> SEQUENCE: 113

cttaggtctt gctcagttcc 20

<210> SEQ ID NO 114
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-11-R

<400> SEQUENCE: 114

cacaacatct ccaagtccag 20

<210> SEQ ID NO 115
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer DNAJC27-12-F

<400> SEQUENCE: 115

acttgagat gttgtgctgc 20

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer DNAJC27-12-R

<400> SEQUENCE: 116

tatgaacccc tcttccttt

20

<210> SEQ ID NO 117
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer DNAJC27-13-F

<400> SEQUENCE: 117

gtgtcagtat ctgtccccta a

21

<210> SEQ ID NO 118
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer DNAJC27-13-R

<400> SEQUENCE: 118

tctatgatct cagtcggtaa

20

What is claimed is:

1. A method for selectively breeding cattle, the method comprising superovulating a female animal, collecting eggs from said superovulated female, in vitro fertilizing said eggs from a suitable male animal, culturing said fertilized eggs into developing embryos, determining the identity of a nucleotide of the HSP gene of the developing embryos corresponding to at least one position selected from the group consisting of positions 1326, 1341, 1396, 1472 and 1480 of SEQ ID NO: 1, wherein the HSP gene comprises a nucleotide sequence of SEQ ID NO: 1, to identify a developing embryo that has guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1, and planting into a suitable female the developing embryo the HSP gene of which has guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1.

2. The method according to claim 1, wherein a developing embryo which is homozygously guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1 is identified and planted into a suitable female.

3. A method for selecting a cattle as a breeder, wherein the HSP gene of the cattle is genotyped, wherein the HSP gene comprises the nucleotide sequence of SEQ ID NO: 1, the method comprising:

obtaining a nucleic acid sample from a cell of the cattle, determining the identity of at least one nucleotide corresponding to a position selected from the group consisting of positions 1326, 1341, 1396, 1472 and 1480 of SEQ ID NO: 1, to identify a cattle the HSP gene of which comprises at least guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1 and

using in a breeding process a cell from the cattle whose HSP gene comprises at least guanine at position 1326,

guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1.

4. The method according to claim 3, wherein a cattle animal is identified and selected whose HSP gene comprises homozygously guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1 and gametes from the cattle whose HSP gene homozygously comprises at least guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1 are used in a breeding process.

5. The method according to claim 3, wherein a bull is selected and its semen used for fertilizing a female animal.

6. The method according to claim 5, wherein the female animal is in vitro fertilized.

7. A method according to claim 5, wherein a multiple ovulation and embryo transfer (MOET) procedure is used.

8. A method according to claim 6, wherein said female animal is also selected and is also homozygous with regard to at least one position selected from the group consisting of positions 1326, 1341, 1396, 1472 and 1480 of SEQ ID NO: 1.

9. A method according to claim 3, wherein a female animal is selected and is fertilized with semen from a suitable bull.

10. A method according to claim 9, wherein the female animal is in vitro fertilized.

11. The method according to claim 10, wherein MOET procedure is used.

12. A dairy cattle breeding method for improved fertilization or embryo survival rate, the method, comprising determining the identity of a nucleotide of the HSP gene, wherein the HSP gene comprises a nucleotide sequence of SEQ ID NO: 1, of a dairy cattle cell or tissue corresponding to at least one position selected from the group consisting of positions 1326, 1341, 1396, 1472 and 1480 of SEQ ID NO: 1, to identify a cattle the HSP gene of which comprises at least guanine at position 1326, guanine at position 1341,

71

adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1, and using in a breeding process a cell or tissue from the cattle whose HSP gene comprises has at least guanine at position 1326, guanine at position 1341, adenosine at position 1396, cytosine at position 1472, or guanine at position 1480 of SEQ ID NO: 1. 5

13. A method according to claim **12**, wherein the dairy cattle cell is an adult cell, an embryo cell, a sperm, an egg, a fertilized egg, or a zygote. 10

14. A method according to claim **12**, wherein the identity of the nucleotide is determined by sequencing the HSP gene, or a fragment thereof comprising at least one position selected from the group consisting of positions 1326, 1341, 1396, 1472 and 1480 of SEQ ID NO. 15

15. A method according to claim **14**, wherein the HSP gene or fragment thereof is isolated from the cell or tissue via amplification by the polymerase chain reaction (PCR) of genomic DNA of the cell or tissue.

16. A method according to claim **14**, wherein the identity of both copies of the HSP gene is determined. 20

* * * * *

72