



US 20200048645A1

(19) **United States**(12) **Patent Application Publication**  
**Hittinger et al.**(10) **Pub. No.: US 2020/0048645 A1**(43) **Pub. Date: Feb. 13, 2020**(54) **YEAST STRAINS WITH SELECTED OR  
ALTERED MITOTYPES AND METHODS OF  
MAKING AND USING THE SAME**(71) Applicant: **Wisconsin Alumni Research  
Foundation, Madison, WI (US)**(72) Inventors: **Chris Todd Hittinger, Madison, WI  
(US); David Peris Navarro, Valencia  
(ES); EmilyClare Patricia Baker,  
Madison, WI (US)**(21) Appl. No.: **16/533,928**(22) Filed: **Aug. 7, 2019****Related U.S. Application Data**(60) Provisional application No. 62/715,498, filed on Aug.  
7, 2018.**Publication Classification**(51) **Int. Cl.**  
*C12N 15/81* (2006.01)  
*C12N 1/18* (2006.01)  
*C12R 1/645* (2006.01)  
(52) **U.S. Cl.**  
CPC ..... *C12N 15/81* (2013.01); *C12R 1/645*  
(2013.01); *C12N 1/18* (2013.01)(57) **ABSTRACT**

Herein we demonstrate that the mitochondrial genome influences temperature tolerance in *Saccharomyces* yeasts. The present invention provides methods for manipulating the mitotype of yeast, including methods to produce synthetic yeast hybrids with a selected mitotype and methods to exchange the native mitochondrial DNA (mtDNA) present in polyploid yeast with mtDNA from a desired source. *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids with selected mitotypes are also provided. The yeast and methods of the present invention may be utilized in a variety of applications, including in fermentation to produce beer and wine.


**Specification includes a Sequence Listing.**

FIG. 1B




FIG. 1D




FIG. 1A




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3




FIG. 4A





FIG. 4C

FIG. 4B

FIG. 5A





FIG. 5B

FIG. 5C

FIG. 6A





FIG. 6B

FIG. 6C



FIG. 7A








FIG. 7B

FIG. 7C

FIG. 8A



8  
G.  
H.



# YEAST STRAINS WITH SELECTED OR ALTERED MITOTYPES AND METHODS OF MAKING AND USING THE SAME

## CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This patent application claims the benefit of priority of U.S. Provisional Patent Application No. 62/715,498, filed Aug. 7, 2018, which is incorporated herein by reference in its entirety.

## STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] This invention was made with government support under 16-CRHF-0-6055 and 17-CRHF-0-6055 awarded by the USDA/NIFA, under 1253634 awarded by the National Science Foundation, and under DE-FC02-07ER64494 and DE-SC0018409 awarded by the US Department of Energy. The government has certain rights in the invention.

## SEQUENCE LISTING

[0003] This application is being filed electronically and includes an electronically submitted Sequence Listing in .txt format. The .txt file contains a sequence listing entitled "2019-08-01\_960296-02452\_SEQ\_Listing.txt" created on Aug. 1, 2019 and is 10,439 bytes in size. The Sequence Listing contained in this .txt file is part of the specification and is hereby incorporated by reference herein in its entirety.

## INTRODUCTION

[0004] Thermal tolerance is an important property in industrial yeast strains that influences the ability of these strains to produce important fermentation products such as beverages, biochemicals, and biofuels. For example, in the beer industry, more than 90% of the beer market consists of lagers, which are brewed with industrial hybrids of *Saccharomyces cerevisiae* x *Saccharomyces eubayanus*. *S. cerevisiae* has been extensively used in fermentation for thousands of years, including in ale production, but pure strains of *S. eubayanus* were not discovered until 2011. Since then, it has become clear that aggressive fermentation of the sugars present in wort are mainly conferred by *S. cerevisiae* genes, whereas the hallmark cold tolerance of lager strains are mainly conferred by *S. eubayanus* genes, with both traits being co-dominant in hybrids. Although brewing strains with colder thermal profiles may be desirable to minimize the production of off-flavors, it often is desirable to conduct fermentations at higher temperatures so as to reduce fermentation times and infrastructure investments. Other than conventional breeding and selection, the inventors are not aware of any other ways to directly manipulate the thermal tolerance of industrial or synthetic yeast strains, such as lager-brewing hybrids. There is thus a need in the art for new methods of altering the thermal tolerance in yeast strains.

## SUMMARY

[0005] In one aspect of the present invention, methods of making a hybrid yeast strain having a selected mitotype are provided. The methods may include treating a first yeast strain with a mitochondrial genome elimination agent to produce a first mitochondrial genome-null yeast strain, and mating the first mitochondrial genome-null yeast strain with

a second yeast strain comprising mitochondrial DNA (mtDNA) to produce the yeast strain having the selected mitotype.

[0006] In another aspect, the present invention relates to methods of making a yeast strain with an altered mitotype. The methods may include treating a first polyploid yeast strain with a mitochondrial genome elimination agent to produce a first mitochondrial genome-null yeast strain, and mating the first mitochondrial genome-null yeast strain with a second karyogamy-deficient yeast strain comprising mitochondrial DNA (mtDNA) to produce the yeast strain having the altered mitotype. In some embodiments of these methods, the mtDNA is supplied by a third, donor yeast strain.

[0007] In a further aspect of the present invention, yeast strains are provided. The yeast strains may include any yeast strain made by one of the methods disclosed herein.

[0008] In another aspect, the present invention relates to *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids comprising mitochondrial DNA (mtDNA) from *Saccharomyces cerevisiae*. Optionally, the *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids may be derived from a lager-brewing strain and include at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of their mitochondria or mtDNA from *Saccharomyces cerevisiae*.

[0009] In another aspect, the present invention relates to synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids comprising mitochondrial DNA (mtDNA) from *Saccharomyces eubayanus*. Optionally, the *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids may be derived from a lager-brewing strain. The hybrids may include at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of their mitochondria or mtDNA from *Saccharomyces eubayanus*. Alternatively, the hybrids may include at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of its mitochondria or mtDNA from *Saccharomyces cerevisiae*.

[0010] In another aspect, the present invention relates to new *Saccharomyces cerevisiae* strains having *Saccharomyces eubayanus* mitotypes. The *Saccharomyces cerevisiae* strains may include genomic DNA and mitochondrial DNA (mtDNA), wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the genomic DNA is from *Saccharomyces cerevisiae* and wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the mtDNA is from *Saccharomyces eubayanus*.

[0011] In another aspect, the present invention relates to new *Saccharomyces eubayanus* strains having *Saccharomyces cerevisiae* mitotypes. The *Saccharomyces eubayanus* strains may include genomic DNA and mitochondrial DNA (mtDNA), wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the genomic DNA is from *Saccharomyces eubayanus* and wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the mtDNA is from *Saccharomyces cerevisiae*.

[0012] In a still further aspect, the present invention relates to methods for making a fermentation product. The methods may include culturing any one of the yeast strains described herein, any one of the *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids disclosed herein, any one of the synthetic *Saccharomyces cerevisiae* x *Saccharomyces*

*eubayanus* hybrids disclosed herein, any one the *Saccharomyces cerevisiae* strains disclosed herein, or any one of the *Saccharomyces eubayanus* strains disclosed herein with a fermentable substrate and at a temperature to produce the fermentation product.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0013]** FIGS. 1A-1D. Relative growth of *S. cerevisiae*, *S. eubayanus*, and their hybrids. Graphs of relative growth scores of strains, combined from all tests. FIGS. 1A and 1B) Relative growth of parent strains carrying their native mtDNA on glucose and glycerol, respectively. FIGS. 1C and 1D) Relative growth of *S. cerevisiae* x *S. eubayanus* hybrids carrying different parental mtDNA on glucose and glycerol, respectively. Error bars represent standard error. Differences in relative growth between hybrids carrying different parental mtDNA with p-values of <0.05 were considered statistically significant and are represented by an asterisk. Parents were not tested for significant differences.

**[0014]** FIGS. 2A-2C. Construction and relative growth of lager cybrids. FIG. 2A) Outline of crosses and strain engineering to produce lager cybrids. Yeast cells represent the nuclear genome, large inner circles represent mtDNA, and small green inner circles represent the HyPr plasmid. Lower case “a” and “α” indicates mating type. Black indicates genetic material from the *S. cerevisiae* karyogamy-deficient strain, red genetic material of *S. cerevisiae* origin, blue of *S. eubayanus* origin, and purple hybrid (i.e. lager) nuclear material. FIGS. 2B and 2C) Growth of a lager strain with native (*S. eubayanus*) mtDNA and lager cybrids with *S. cerevisiae* mtDNA. Error bars represent standard error and asterisks indicate statistically significant differences in growth between the cybrid and lager with native mtDNA (p-value <0.05). FIG. 2B) Growth on glucose FIG. 2C) Growth on glycerol.

**[0015]** FIG. 3. Hybrid mating scheme. Outline of procedure to control the inheritance of mtDNA in crosses of *S. cerevisiae* and *S. eubayanus*. Yeast cells represent the nuclear genome, and inner circles represent mtDNA. Red indicates genetic material of *S. cerevisiae* origin, blue of *S. eubayanus* origin, and purple hybrid nuclear material.

**[0016]** FIGS. 4A-4C. *Sc* x *Se* growth assay. Growth assay for *S. cerevisiae* (laboratory strain) x *S. eubayanus* (type strain) hybrids and parental strains. FIG. 4A) Representative spot assay plates grown at various temperatures on plates containing glucose or glycerol as the sole carbon source. The following strains were tested: *S. cerevisiae* (*Sc*)  $\rho^+$  parent, *Sc*  $\rho^0$  parent, *S. eubayanus* (*Se*)  $\rho^+$  parent, *Se*  $\rho^0$  parent, *Sc*  $\rho^+$  x *Se*  $\rho^0$  hybrid, and *Sc*  $\rho^0$  x *Se*  $\rho^+$  hybrid. FIGS. 4B and 4C) Relative growth of tested strains across all temperatures, combining all replicates. Outer circles and lines represent nuclear genotype, while inner circles represent mtDNA. Highlighted regions represent temperatures where a hybrid of one mitotype had significantly greater relative growth than the hybrid with the alternative mitotype. FIG. 4B) Relative growth of tested strains on glucose. FIG. 4C) Relative growth of tested strains on glycerol.

**[0017]** FIGS. 5A-5C. *ScAle* X *Se* growth assay. Growth assay for *S. cerevisiae*-ale x *S. eubayanus* (type strain) hybrids and parental strains. FIG. 5A) Representative spot assay plates grown at various temperatures on plates containing glucose or glycerol as the sole carbon source. The following strains were tested: *S. cerevisiae*-ale (*ScAle*)  $\rho^+$  parent, *ScAle*  $\rho^0$  parent, *S. eubayanus* (*Se*)  $\rho^+$  parent, *Se*  $\rho^0$

parent, *ScAle* x *Se*  $\rho^0$  hybrid, and *ScAle*  $\rho^0$  x *Se*  $\rho^+$  hybrid. FIGS. 5B and 5C) Relative growth of tested strains across all temperatures, combining all replicates. Outer circles and lines represent nuclear genotype, while inner circles represent mtDNA. Highlighted regions represent temperatures where a hybrid of one mitotype had significantly greater relative growth than the hybrid with the alternative mitotype. FIG. 5B) Relative growth of tested strains on glucose. FIG. 5C) Relative growth of tested strains on glycerol.

**[0018]** FIGS. 6A-6C. *Sc* X *SeNC* growth assay. Growth assay for *S. cerevisiae* (laboratory strain) x *S. eubayanus*-North Carolina hybrids and parental strains. FIG. 6A) Representative spot assay plates grown at various temperatures on plates containing glucose or glycerol as the sole carbon source. The following strains were tested: *S. cerevisiae* (*Sc*)  $\rho^+$  parent, *Sc*  $\rho^0$  parent, *S. eubayanus*-North Carolina (*SeNC*)  $\rho^+$  parent, *SeNC*  $\rho^0$  parent, *Sc*  $\rho^+$  x *SeNC*  $\rho^0$  hybrid, and *Sc*  $\rho^0$  x *SeNC*  $\rho^+$  hybrid. FIGS. 6B and 6C) Relative growth of tested strains across all temperatures, combining all replicates. Outer circles and lines represent nuclear genotype, while inner circles represent mtDNA. Highlighted regions represent temperatures where a hybrid of one mitotype had significantly greater relative growth than the hybrid with the alternative mitotype. FIG. 6B) Relative growth of tested strains on glucose. FIG. 6C) Relative growth of tested strains on glycerol.

**[0019]** FIGS. 7A-7C. *ScAle* X *SeNC*. growth assay. Growth assay for *S. cerevisiae*-ale x *S. eubayanus*-North Carolina hybrids and parental strains. FIG. 7A) Representative spot assay plates grown at various temperatures on plates containing glucose or glycerol as the sole carbon source. The following strains were tested: *S. cerevisiae*-ale (*ScAle*)  $\rho^+$  parent, *ScAle*  $\rho^0$  parent, *S. eubayanus*-North Carolina (*SeNC*)  $\rho^+$  parent, *SeNC*  $\rho^0$  parent, *ScAle*  $\rho^+$  x *SeNC*  $\rho^0$  hybrid, and *ScAle*  $\rho^0$  x *SeNC*  $\rho^+$  hybrid. FIGS. 7B and 7C) Relative growth of tested strains across all temperatures, combining all replicates. Outer circles and lines represent nuclear genotype, while inner circles represent mtDNA. Highlighted regions represent temperatures where a hybrid of one mitotype had significantly greater relative growth than the hybrid with the alternative mitotype. FIG. 7B) Relative growth of tested strains on glucose. FIG. 7C) Relative growth of tested strains on glycerol.

**[0020]** FIGS. 8A-8C. Lager cybrid growth assay. Growth assay for lager cybrids and parental strains. FIG. 8A) Representative spot assay plates grown at various temperatures on plates containing glucose or glycerol as the sole carbon source. The following strains were tested: lager  $\rho^+$  parent, lager  $\rho^0$  parent, *S. cerevisiae* (laboratory strain) (*Sc*)  $\rho^+$  parent, *S. cerevisiae*-ale (*ScAle*)  $\rho^+$  parent, lager  $\rho^{Sc}$  cybrid, and lager  $\rho^{ScAle}$  cybrid. FIGS. 8B and 8C) Relative growth of tested strains across all temperatures combining all replicates. Error bars represent standard error and asterisks indicate statistically significant differences in growth between the cybrid and lager with native mtDNA (p<0.05). FIG. 8B) Relative growth of tested strains, excluding lager  $\rho^0$ , on glucose. FIG. 8C) Relative growth of tested strains, excluding lager  $\rho^0$ , on glycerol.

#### DETAILED DESCRIPTION

**[0021]** When two yeast cells mate, the mitochondrial genome (mtDNA) of both parents is initially present in the zygote. However, within a few cell divisions, a single mtDNA haplotype, or “mitotype” is fixed (Berger and Yaffe

2000), often in a non-random manner (Zweifel and Fangman 1991; Marinoni et al. 1999; Hsu and Chou 2017). In the present application, the inventors demonstrate that the thermal tolerance of yeast is dictated by mitotype and provide methods for controlling or selecting the mitotype. Thus, the ability to control the inheritance of mtDNA allows one to adapt yeast for growth at various temperatures. This ability is a powerful tool for the generation of novel yeast with utility in numerous applications.

**[0022]** The present invention provides new yeast strains with selected mitotypes and new methods for altering or selecting the mitotype of yeast. The inventors have shown that the thermal tolerance of *S. cerevisiae* strains, *S. eubayanus* strains, and hybrids of these species can be controlled by manipulating the parental source of the mtDNA. Specifically, in the non-limiting Examples, the inventors demonstrate that *S. cerevisiae* x *S. eubayanus* strains with *S. cerevisiae* mtDNA perform better at warmer temperatures, while *S. cerevisiae* x *S. eubayanus* strains with *S. eubayanus* mtDNA perform better at colder temperatures. The inventors further show that the thermal profiles of existing industrial lager strains can be manipulated analogously by depleting and replacing the mitochondria of the strains through, for example, cytoduction using a strain of *S. cerevisiae* that can transfer mtDNA to another strain without completing karyogamy. Lager yeast strains with elevated thermal tolerance may be attractive to the lager-brewing industry because they are expected to reduce fermentation times and infrastructure investments. Reduced fermentation times are attractive across all production scales, whereas infrastructure investments are a major obstacle to lager brewing among microbreweries that generally favor ale brewing partly for this reason.

#### Methods of Making Hybrid Yeast Strains with Selected Mitotypes

**[0023]** In one aspect of the present invention, methods of making a yeast strain having a selected mitotype are provided. The methods include treating a first yeast strain with a mitochondrial genome elimination agent to produce a first mitochondrial genome-null yeast strain, and mating the first mitochondrial genome-null yeast strain with a second yeast strain comprising mitochondrial DNA to produce the yeast strain having the selected mitotype.

**[0024]** As used herein, the description “comprising mitochondrial DNA” implies that the yeast contain untreated, replication competent mitochondria comprising intact mitochondrial genomes.

**[0025]** As used herein, “mating” refers to the process of fusing together at least one yeast cell from a first yeast strain with at least one yeast cell from a second yeast strain. Mating may or may not entail karyogamy, or the fusing together of the nuclei in the two parental yeast cells. However, in this particular set of methods, mating produces a “hybrid” yeast strain, which comprises nuclear DNA contributed from both parental strains, as opposed to a cybrid. A “cybrid” is a cell in which the original mitochondrial are substituted with their counterparts from another strain or species using a method such as cytoduction (described below). Thus, a cybrid comprises the nuclear DNA of a single parent yeast cell and the mitochondria of a second donor yeast cell.

**[0026]** Haploid yeast strains with the appropriate mating types may mate naturally. Thus, in preferred embodiments

of these methods, two haploid yeast strains are mated to produce a new hybrid yeast strain with the selected mitotype.

**[0027]** As used herein, “mitotype” refers to the parental yeast strain from which the mitochondrial DNA in a yeast strain originated. For example, if a *Saccharomyces cerevisiae* yeast strain having mitochondria (and mtDNA) is mated with *Saccharomyces eubayanus* that lacks mtDNA, the resulting hybrid yeast strain will have a *Saccharomyces cerevisiae* “mitotype” (i.e., mtDNA originated from the *Saccharomyces cerevisiae* parent).

**[0028]** As used herein, “treating” refers to contacting, for example, a yeast strain, with a particular substance or substances. Contacting encompasses administration to a cell directly or adding an agent to a culture comprising the yeast. As used herein, a “mitochondrial genome elimination agent” refers to an agent that disrupts yeast mitochondrial DNA, allowing the mitochondrial genome of a yeast strain to be removed. To control the inheritance of mtDNA during yeast mating, the present inventors generated “mitochondrial genome-null yeast strains” ( $\rho^0$ ) (mtDNA completely absent) to mate with strains having mitochondria ( $\rho^+$ ) so that mtDNA from only the  $\rho^+$  parent would be present in offspring. See, e.g., FIG. 3. In the non-limiting Examples, mitochondrial genome-null yeast strains ( $\rho^0$ ) were generated by treating  $\rho^+$  parent strains with ethidium bromide to generate respiration-deficient strains. Respiration-deficient strains were screened for by the absence of growth on glycerol, and the complete removal of mtDNA was confirmed by DAPI staining. Suitable “mitochondrial genome elimination agents” in accordance with the present invention may include, without limitation, ethidium bromide or antifolate cocktails that include methotrexate and sulfanilamide RRR.

**[0029]** The first yeast strain and the second yeast strain of the present method may be the same or different yeast species (defined below). In some embodiments, the first yeast strain may be the same yeast species as the second yeast strain but be a different strain of that yeast species. In certain embodiments, the methods are used to create synthetic “lager hybrids” of *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* with selected mitotypes. In one preferred embodiment, the first yeast strain is *Saccharomyces cerevisiae* and the second yeast strain is *Saccharomyces eubayanus*. In a second preferred embodiment, the first yeast strain is *Saccharomyces eubayanus* and the second yeast strain is *Saccharomyces cerevisiae*.

#### Methods of Making Polyploid Cybrids

**[0030]** In another aspect, the present invention provides methods of making a polyploid cybrid yeast strain with an altered mitotype. These methods involve treating a first polyploid yeast strain with a mitochondrial genome elimination agent to produce a first mitochondrial genome-null “acceptor” yeast strain, and mating the acceptor yeast strain with a second “shuttle” yeast strain comprising mitochondrial DNA (mtDNA). The shuttle yeast strain must be a karyogamy-deficient yeast strain, such that the resulting yeast strain with an altered mitotype is a “cybrid” comprising nuclear DNA of the acceptor yeast strain and the mtDNA delivered in the cytoplasm of the shuttle yeast strain.

**[0031]** A “karyogamy-deficient yeast strain” refers to a yeast strain that may not perform karyogamy or the fusing together of the nuclei in the two parental yeast cells. The

“karyogamy-deficient yeast strain” may include mutations in genes required for karyogamy including, without limitation, a *kar1-1* mutation. In the present invention, mating the first acceptor yeast strain to a second karyogamy-deficient yeast strain allows for mixing of cytoplasm between the mated cells, while preventing fusion between the nuclear genomes, ultimately leading to progeny with mixed cytoplasm, but only one nuclear background. In this way, the mitochondria from the shuttle strain may be transferred into the mitochondrial genome-null acceptor yeast strain by mating the yeast and selecting for functional mtDNA (by growth on glycerol, a non-fermentable carbon source) and nuclear background of the acceptor yeast strain. This transfer of mitochondria from one cell to another is referred to as “cytoduction”.

**[0032]** In certain embodiments, the karyogamy-deficient shuttle strain may deliver its native mitochondria (or mitochondrial DNA) to the acceptor strain. In other embodiments, the shuttle strain may deliver mitochondria from a “donor” yeast strain. The donor mitochondria may be introduced into the shuttle yeast strain by mating the donor yeast strain with the shuttle strain (see, e.g., FIG. 2A). In this way, the mitochondria from the donor strain may be transferred into the shuttle strain, forming a cybrid with the desired mitotype by cytoduction. Including this optional, additional step before the shuttle strain is mated with the acceptor yeast strain allows the mitochondria of any suitable yeast strain to ultimately be transferred to the acceptor yeast strain. Thus, using this two-step cytoduction method, any yeast cell with a desirable genome (i.e., the acceptor yeast strain) may be altered to have any desirable mitotype (either that of the shuttle or the donor strain). For instance, in the Examples, the applicants demonstrate that by replacing the mitochondria of an acceptor yeast strain of *Saccharomyces eubayanus* origin (a lager strain) with the mitochondria of a *Saccharomyces cerevisiae* donor strain via a karyogamy-deficient *Saccharomyces cerevisiae* shuttle strain, a cybrid lager yeast strain with elevated thermal tolerance may be produced.

**[0033]** Importantly, the acceptor yeast strains of the present invention may be any polyploid yeast strain. As used herein, “polyploid” refers to cells with multiple sets of chromosomes. Thus, a polyploid yeast strain has a genome that is of diploid or higher ploidy (i.e., diploid (2N), triploid (3N), tetraploid (4N), pentaploid (5N), hexaploid (6N), heptaploid (7N), octaploid (8N), nonaploid (9N), decaploid (10N), 11N, 12N, 13N, 14N, 15N, 16N, or more). Preferably, the acceptor yeast strain are allopolyploid. As used herein, “allopolyploid” refers to hybrid cells with two or more sets of chromosomes derived from two different species. This definition includes allodiploids, allotriploids, allotetraploids, allopentaploids, allohexaploids, and allopolyploids with a higher number of haploid sets.

**[0034]** The inventors envision that these methods can be used to alter industrial lager strains, such as *Saccharomyces pastorianus* and *Saccharomyces carlsbergensis*, to adapt them for fermentation at various temperatures. Because these strains have complex allopolyploid genomes, it is challenging to work with them using traditional genetic methods. Mating does not usually occur in polyploid yeasts because they contain both MATa and MATα at their mating type locus. Thus, to mate polyploid strains for cytoduction, the MAT locus must first be homozygosed. In certain embodiments, this is accomplished using the HyPr method, which is detailed in U.S. Patent Publication 2018/

0127784, the contents of which is incorporated herein in its entirety. For instance, in the Examples, the MAT locus of lager genome-null acceptor strains was homozygosed using a HyPr (Hybrid Production) plasmid (pHCT2). Briefly, a “HyPr plasmid” may include a polynucleotide comprising a promoter operably connected to an HO polynucleotide encoding a yeast Ho protein, a selectable marker cassette, and a yeast origin of replication.

**[0035]** The terms “promoter,” “promoter region,” or “promoter sequence” refer generally to transcriptional regulatory regions of a gene, which may be found at the 5' or 3' side of a gene such as the HO polynucleotide, or within the coding region of the gene (i.e., HO polynucleotide). Typically, a promoter is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. The typical 5' promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence is a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

**[0036]** In some embodiments, the promoters within the polynucleotides of the present invention may be operably connected to the HO polynucleotide. As used herein, a promoter is “operably connected to” or “operably linked to” when it is placed into a functional relationship with a second polynucleotide sequence. For instance, a promoter is operably connected to an HO polynucleotide if the promoter is connected to the HO polynucleotide such that it may effect transcription of the HO polynucleotide coding sequence. In various embodiments, the HO polynucleotides may be operably linked to at least 1, at least 2, at least 3, at least 4, at least 5, or at least 10 promoters.

**[0037]** Promoters useful in the practice of the present invention include, but are not limited to, constitutive, inducible, temporally-regulated, and chemically regulated promoters. Preferably, the promoters are inducible. Suitable inducible promoters for expression in yeast include, without limitation, galactose inducible promoters (i.e., GAL1) and doxycycline-inducible promoters. Those of skill in the art are familiar with a wide variety of additional promoters for use in various yeast species.

**[0038]** The HyPr plasmid may include an HO polynucleotide encoding a yeast Ho protein. Yeast Ho proteins are site-specific endonucleases that produce a double-strand break in the MAT locus. The double-strand break is followed by a unidirectional gene conversion event that replaces the information at the MAT locus by information copied from either of the two homologous loci (HMR and HML) or from another copy of the MAT locus. Alternatively, the cut chromosome may be lost, leading to a mating type change. Yeast Ho proteins may be any of the Ho proteins found in any yeast species including, without limitation, those yeast species closely related to *Candida glabrata* and *Naumovozyma castellii*. Suitably, the protein sequence of an exemplary yeast Ho protein from *Saccharomyces cerevisiae* is indicated in SEQ ID NO: 14. In some embodiments, the yeast Ho protein comprises SEQ ID NO: 11 or a mutant, variant, derivative, or fragment thereof.

**[0039]** As used herein, a “protein,” “polypeptide,” or “peptide” may be used interchangeably to refer to a polymer

of amino acids. A “protein” as contemplated herein typically comprises a polymer of naturally occurring amino acids (e.g., alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine).

**[0040]** A yeast Ho protein may include “mutant” proteins, variants, and derivatives thereof. As used herein the term “wild-type” is a term of the art understood by skilled persons and means the typical form of an organism, strain, gene, or characteristic as it occurs in nature as distinguished from mutant or variant forms. As used herein, a “variant,” “mutant,” or “derivative” refers to a polypeptide molecule having an amino acid sequence that differs from a reference protein or polypeptide molecule. A variant or mutant may have one or more insertions, deletions, or substitutions of an amino acid residue relative to a reference molecule. A variant or mutant may include a fragment of a reference molecule. For example, a yeast Ho protein mutant or variant protein may have one or more insertions, deletions, or substitution of at least one amino acid residue relative to the yeast Ho “wild-type” protein. The protein sequences of a “wild-type” yeast Ho protein from *Saccharomyces cerevisiae* is presented as SEQ ID NO: 14. This sequence may be used as a reference sequence.

**[0041]** A yeast Ho protein may be a full-length protein or may be fragments of the full-length protein. As used herein, a “fragment” is a portion of an amino acid sequence, which is identical in sequence to but shorter in length than a reference sequence. A fragment may comprise up to the entire length of the reference sequence, minus at least one amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous amino acid residues of a reference protein, respectively. In some embodiments, a fragment may comprise at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 150, 250, or 500 contiguous amino acid residues of a reference protein. Fragments may be preferentially selected from certain regions of a molecule. The term “at least a fragment” encompasses the full-length polypeptide. A fragment may include an N-terminal truncation, a C-terminal truncation, or both truncations relative to the full-length yeast Ho protein. Preferably, a fragment of a yeast Ho protein includes amino acid residues required for recognition and cleavage of the MAT locus site.

**[0042]** A “deletion” in a yeast Ho protein refers to a change in the amino acid sequence resulting in the absence of one or more amino acid residues. A deletion may remove at least 1, 2, 3, 4, 5, 10, 20, 50, 100, 200, or more amino acids residues. A deletion may include an internal deletion and/or a terminal deletion (e.g., an N-terminal truncation, a C-terminal truncation or both of a reference polypeptide).

**[0043]** “Insertions” and “additions” in a yeast Ho protein refer to changes in an amino acid sequence resulting in the addition of one or more amino acid residues. An insertion or addition may refer to 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, or more amino acid residues. A variant of a yeast Ho protein may have N-terminal insertions, C-terminal insertions, internal insertions, or any combination of N-terminal insertions, C-terminal insertions, and internal insertions.

**[0044]** Regarding proteins, the phrases “percent identity,” “% identity,” and “% sequence identity” refer to the percentage of residue matches between at least two amino acid sequences aligned using a standardized algorithm. Methods

of amino acid sequence alignment are well known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail below, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide. Percent identity for amino acid sequences may be determined as understood in the art. (See, e.g., U.S. Pat. No. 7,396,664). A suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST), which is available from several sources, including the NCBI, Bethesda, Md., at its website. The BLAST software suite includes various sequence analysis programs including “blastp,” that is used to align a known amino acid sequence with other amino acids sequences from a variety of databases. As described herein, variants, mutants, or fragments (e.g., a yeast Ho protein variant, mutant, or fragment thereof) may have 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 80%, 70%, 60%, or 50% amino acid sequence identity relative to a reference molecule (e.g., relative to a full-length yeast Ho protein (SEQ ID NO: 14)).

**[0045]** Protein sequence identity may be measured over the length of an entire defined protein sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined protein sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70, or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

**[0046]** The amino acid sequences of the yeast Ho protein variants, mutants, or derivatives as contemplated herein may include conservative amino acid substitutions relative to a reference amino acid sequence. For example, a variant, mutant, or derivative polypeptide may include conservative amino acid substitutions relative to a reference molecule. “Conservative amino acid substitutions” are those substitutions that are a substitution of an amino acid for a different amino acid where the substitution is predicted to interfere least with the properties of the reference polypeptide. In other words, conservative amino acid substitutions substantially conserve the structure and the function of the reference polypeptide. Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

**[0047]** The HO polynucleotides encoding the yeast Ho proteins, fragments, variants, mutants, or derivatives thereof may be any polynucleotide encoding the appropriate yeast Ho protein amino acid sequence. Those of skill in the art also understand the degeneracy of the genetic code and that a variety of polynucleotides can encode the same polypeptide. In some embodiments, the polynucleotides may be codon-optimized for expression in a particular yeast cell. While particular nucleotide sequences, which are found in *Saccharomyces cerevisiae* (i.e., SEQ ID NO: 15), are disclosed, herein any nucleotide sequences may be used which encode



a desired form of the yeast Ho proteins described herein. Thus, non-naturally occurring sequences may be used. These may be desirable, for example, to enhance expression in a particular yeast species. Computer programs for generating degenerate coding sequences are available and can be used for this purpose as well as other means.

**[0048]** The HyPr plasmid may also include a selectable marker cassette. The selectable marker cassette may confer resistance to nourseothricin (natMX4) or hygromycin (hphMX). Other selectable markers conferring resistance to other antibiotics, such as kanMX, amdS, TK, Sh ble or ble, which confer resistance to G418, fluoroacetamide, antifolates, Zeocin or phleomycin, respectively, could also be used. Those of skill in the art will appreciate that additional combinations of selectable markers can be used as well. Other forms of selectable markers may be used such as markers that provide a growth advantage or colorimetric selection other than antibiotic resistance. The selectable marker cassettes include a polynucleotide encoding the selectable marker operably connected to a promoter capable of inducing transcription of the selectable marker.

**[0049]** The HyPr plasmid may also include a yeast origin of replication to allow replication of the polynucleotides in a particular yeast species. Suitably, the yeast origin of replication is functional across many yeast species including, without limitation, all *Saccharomyces* species. Exemplary yeast origin of replications may include, without limitation, KARS101 from *Kluyveromyces lactis* and *S. cerevisiae* CEN and ARS sequences to improve stability in multiple yeast species.

**[0050]** The “yeast strains” used in any method of the present invention (i.e., first yeast strain, second yeast strain, acceptor yeast strain, karyogamy-deficient shuttle yeast strain, donor yeast strain) may be from the family Saccharomycetaceae. In some embodiments, the yeast strains are from the genus *Saccharomyces*. Suitable *Saccharomyces* species may include, without limitation, *Saccharomyces cerevisiae*, *Saccharomyces eubayanus*, *Saccharomyces paradoxus*, *Saccharomyces mikatae*, *Saccharomyces jurei*, *Saccharomyces arboricola*, *Saccharomyces bayanus*, *Saccharomyces pastorianus*, *Saccharomyces carlsbergensis*, *Saccharomyces kudriavzevii*, and *Saccharomyces uvarum*. Suitable *Saccharomyces* species may also include any taxonomic synonyms of these species or any newly discovered species to be members of the genus *Saccharomyces*. Suitable yeast strains may include those that are cold adapted, such as *Saccharomyces bayanus*, *Saccharomyces kudriavzevii*, *Saccharomyces pastorianus*, *Saccharomyces carlsbergensis*, *Saccharomyces uvarum*, and *Saccharomyces eubayanus*. Suitable yeast strains may also include yeast hybrids including, without limitation, *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids. Throughout the present application, the nomenclature Species 1 x Species 2 will be used to refer to interspecies hybrids (as suggested by Nguyen and Boekhout, 2017). In certain preferred embodiments, the yeast strain is a *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid, including those commonly used in the beer industry, including *Saccharomyces bayanus*, *Saccharomyces pastorianus*, and *Saccharomyces carlsbergensis*.

**[0051]** A “yeast species,” as used herein, may refer to a substantially pure yeast species or to a yeast strain that is a mixed yeast species. A substantially pure yeast species may contain greater than 95%, 96%, 97%, 98%, 99%, or 99.9% chromosomal content from a single yeast species. For

example, strains of *Saccharomyces cerevisiae* that contain greater than 95%, 96%, 97%, 98%, 99%, or 99.9% *Saccharomyces cerevisiae* chromosomal content may be considered a substantially pure yeast species and would be considered to be a *Saccharomyces cerevisiae* species. A mixed yeast species may contain substantial chromosomal content from 2 or more different yeast species. For example, a particular yeast strain may have 90% chromosomal content from *Saccharomyces uvarum* and 5% chromosomal content from *Saccharomyces cerevisiae* and 5% chromosomal content from *Saccharomyces eubayanus*. In the case of mixed species, and in accordance with the present invention, the yeast species of a yeast cell from a mixed yeast species is determined by which species contributes the majority of the chromosomal content in the cell. Thus, in the preceding mixed species example, the exemplary yeast strain would be considered a *Saccharomyces uvarum* species.

**[0052]** As used herein, a “different yeast species” refers to yeast species that are not the same as determined in view of the “yeast species” definitions provided herein. For example, if the first yeast species was a yeast strain having 90% chromosomal content from *Saccharomyces uvarum* and 5% chromosomal content from *Saccharomyces cerevisiae* and 5% chromosomal content from *Saccharomyces eubayanus* and the second yeast species was a substantially pure *Saccharomyces uvarum* strain, the first yeast species and the second yeast species would be considered the same—*Saccharomyces uvarum*. Examples of embodiments where the first yeast species and the second yeast species would be considered different might include the first yeast species being *Saccharomyces cerevisiae* and the second yeast species being *Saccharomyces eubayanus*. In some embodiments, the first yeast strain is *Saccharomyces cerevisiae* and the second yeast strain is *Saccharomyces eubayanus*. In some embodiments, the first yeast strain is *Saccharomyces eubayanus* and the second yeast strain is *Saccharomyces cerevisiae*.

#### Compositions (“Yeast Strains”)

**[0053]** In a further aspect of the present invention, yeast strains are provided. The yeast strains may include any yeast strains made by one of the methods disclosed herein.

**[0054]** In another aspect, the present invention relates to *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids comprising mitochondrial DNA (mtDNA) from *Saccharomyces cerevisiae*. Lager style beers are fermented at low temperatures, and it has been known for some time that lager yeasts inherited the ability to ferment at these low temperatures from their *S. eubayanus* parent. Lager hybrids also inherited their mtDNA from the *S. eubayanus* parent. The present inventors, however, have been able to alter the mitotype of lager hybrid strains from a *Saccharomyces eubayanus* mitotype to a *Saccharomyces cerevisiae* mitotype. Such lager hybrid strains have superior growth over lager hybrid strains with *S. eubayanus* mtDNA when grown at high temperatures. See, e.g. Examples.

**[0055]** The *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids including mitochondria or mitochondrial DNA (mtDNA) from *Saccharomyces cerevisiae* may include at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of its mitochondria or mtDNA from *Saccharomyces cerevisiae*. Suitably, the *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid



may include all of its mitochondria or mitochondria DNA (mtDNA) from *Saccharomyces cerevisiae*.

**[0056]** The present invention also encompasses synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids with mitochondrial DNA (mtDNA) from *Saccharomyces eubayanus*. As used herein, “synthetic” refers to hybrids that may be produced using laboratory methods as opposed to those hybrids found in nature. The present inventors envision that the methods disclosed herein may also be used to generate new synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids having new *Saccharomyces eubayanus* mitotypes that may be expected to produce lager beers at new optimal temperatures or lower temperatures.

**[0057]** The synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids including mitochondria or mitochondrial DNA (mtDNA) from *Saccharomyces eubayanus* may include at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of its mitochondria or mtDNA from *Saccharomyces eubayanus*. Suitably, the synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid may include all of its mitochondria or mitochondrial DNA (mtDNA) from *Saccharomyces eubayanus*.

**[0058]** In some embodiments, the “*Saccharomyces cerevisiae* x *Saccharomyces eubayanus*” or “synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus*” hybrid is or may be derived from a lager-brewing strain. As used herein, a “lager-brewing strain” refers to a *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid strains historically used in the production of lager style beers, such as *Saccharomyces pastorianus* and *Saccharomyces carlsbergensis*. The lager-brewing strains may have a *Saccharomyces eubayanus* mitotype. In another aspect, the present invention relates to new *Saccharomyces cerevisiae* strains having *Saccharomyces eubayanus* mitotypes. The *Saccharomyces cerevisiae* strains may include genomic DNA and mitochondrial DNA (mtDNA), wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the genomic DNA is from *Saccharomyces cerevisiae* and wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the mtDNA is from *Saccharomyces eubayanus*.

**[0059]** In another aspect, the present invention relates to new *Saccharomyces eubayanus* strains having *Saccharomyces cerevisiae* mitotypes. The *Saccharomyces eubayanus* strains may include genomic DNA and mitochondrial DNA (mtDNA), wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the genomic DNA is from *Saccharomyces eubayanus* and wherein at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% of the mtDNA is from *Saccharomyces cerevisiae*.

#### Methods of Making a Fermentation Product

**[0060]** In a still further aspect, the present invention relates to methods for making a fermentation product. The methods may include the culturing any one of the yeast strains described herein (see “Compositions” section above), any one of the *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids disclosed herein, any one of the synthetic *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids disclosed herein, any one the *Saccharomyces cerevisiae* strains disclosed herein, or any one of the *Saccha-*

*romyces eubayanus* strains disclosed herein with a fermentable substrate and at a temperature to produce the fermentation product.

**[0061]** As used herein, “culturing” refers to mixing the yeast strains or *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids into any medium including a fermentable substrate. The fermentable substrate may include a carbohydrate, wort, and/or malt extract.

**[0062]** The fermentation product may be a beer product, a wine product, an alcoholic beverage, a biochemical, or a biofuel. In some embodiments, the fermentation product is a lager beer.

**[0063]** The yeast strains or *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid may be cultured at a temperature at, below, or above 36° F., 37° F., 38° F., 39° F., 40° F., 41° F., 42° F., 43° F., 44° F., 45° F., 46° F., 47° F., 48° F., 49° F., 50° F., 51° F., 52° F., 53° F., 54° F., 55° F., 56° F., 57° F., 58° F., 59° F., 60° F., 61° F., 62° F., 63° F., 64° F., 65° F., 66° F., 67° F., 68° F., 69° F., 70° F., 75° F., 80° F., 85° F., 90° F., 95° F., or 100° F. Suitably, the temperature may be between about 40° F. and about 50° F., about 50° F. and about 58° F., about 55° F. and about 60° F., about 56° F. and about 61° F., about 57° F. and about 62° F., about 58° F. and about 63° F., about 59° F. and about 64° F., about 60° F. and about 65° F., about 61° F. and about 66° F., about 62° F. and about 67° F., about 63° F. and about 68° F., about 64° F. and about 69° F., about 65° F. and about 70° F., about 66° F. and about 71° F., about 67° F. and about 72° F., about 68° F. and about 73° F., about 69° F. and about 74° F., about 70° F. and about 75° F., about 75° F. and about 80° F., about 80° F. and about 85° F., about 85° F. and about 90° F., about 90° F. and about 95° F., about 95° F. and about 100° F. or any range therein. In some embodiments, the temperature is between about 40° F. and about 100° F. or any range therein.

**[0064]** Suitably, in some embodiments, when the yeast strain or hybrid includes *Saccharomyces eubayanus* mitochondrial DNA (mtDNA), the yeast strain or hybrid is grown at temperatures below 60° F. or between about 40° F. and about 50° F., about 50° F. and about 58° F., about 55° F. and about 60° F., or any range therein.

**[0065]** Suitably, in some embodiments, when the yeast strain or hybrid includes *Saccharomyces cerevisiae* mitochondrial DNA (mtDNA), the yeast strain or hybrid is grown at temperatures above 60° F. or between about 60° F. and about 65° F., about 61° F. and about 66° F., about 62° F. and about 67° F., about 63° F. and about 68° F., about 64° F. and about 69° F., about 65° F. and about 70° F., about 66° F. and about 71° F., about 67° F. and about 72° F., about 68° F. and about 73° F., about 69° F. and about 74° F., about 70° F. and about 75° F., about 75° F. and about 80° F., about 80° F. and about 85° F., about 85° F. and about 90° F., about 90° F. and about 95° F., about 95° F. and about 100° F. or any range therein.

**[0066]** The present disclosure is not limited to the specific details of construction, arrangement of components, or method steps set forth herein. The compositions and methods disclosed herein are capable of being made, practiced, used, carried out and/or formed in various ways that will be apparent to one of skill in the art in light of the disclosure that follows. The phraseology and terminology used herein is for the purpose of description only and should not be regarded as limiting to the scope of the claims. Ordinal indicators, such as first, second, and third, as used in the description and the claims to refer to various structures or

method steps, are not meant to be construed to indicate any specific structures or steps, or any particular order or configuration to such structures or steps. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to facilitate the disclosure and does not imply any limitation on the scope of the disclosure unless otherwise claimed. No language in the specification, and no structures shown in the drawings, should be construed as indicating that any non-claimed element is essential to the practice of the disclosed subject matter. The use herein of the terms “including,” “comprising,” or “having,” and variations thereof, is meant to encompass the elements listed thereafter and equivalents thereof, as well as additional elements. Embodiments recited as “including,” “comprising,” or “having” certain elements are also contemplated as “consisting essentially of” and “consisting of” those certain elements.

**[0067]** Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. For example, if a concentration range is stated as 1% to 50%, it is intended that values such as 2% to 40%, 10% to 30%, or 1% to 3%, etc., are expressly enumerated in this specification. These are only examples of what is specifically intended, and all possible combinations of numerical values between and including the lowest value and the highest value enumerated are to be considered to be expressly stated in this disclosure. Use of the word “about” to describe a particular recited amount or range of amounts is meant to indicate that values very near to the recited amount are included in that amount, such as values that could or naturally would be accounted for due to manufacturing tolerances, instrument and human error in forming measurements, and the like. All percentages referring to amounts are by weight unless indicated otherwise.

**[0068]** No admission is made that any reference, including any non-patent or patent document cited in this specification, constitutes prior art. In particular, it will be understood that, unless otherwise stated, reference to any document herein does not constitute an admission that any of these documents forms part of the common general knowledge in the art in the United States or in any other country. Any discussion of the references states what their authors assert, and the applicant reserves the right to challenge the accuracy and pertinence of any of the documents cited herein. All references cited herein are fully incorporated by reference in their entirety, unless explicitly indicated otherwise. The present disclosure shall control in the event there are any disparities between any definitions and/or description found in the cited references.

**[0069]** Unless otherwise specified or indicated by context, the terms “a,” “an,” and “the” mean “one or more.” For example, “a protein” or “an RNA” should be interpreted to mean “one or more proteins” or “one or more RNAs,” respectively.

**[0070]** The following examples are meant only to be illustrative and are not meant as limitations on the scope of the invention or of the appended claims.

## EXAMPLES

### Example 1

#### Impact of Mitochondria on Temperature Tolerance in Industrial and Synthetic Lager-Brewing Yeast Hybrids

**[0071]** A growing body of observational and experimental data supports the role of mitochondria in thermal adaptation. In *Saccharomyces* yeasts, relatively little work has examined the role of mitochondrial DNA (mtDNA) in temperature tolerance, and most of this work has focused on tolerance to high temperatures. The yeasts of the genus *Saccharomyces* can be broadly divided between cryotolerant and thermotolerant species. The genetic and molecular basis of the ability to grow at low temperatures in cryotolerant *Saccharomyces* yeasts is still poorly understood and what role mtDNA may play has not been tested. The industrial yeast strains used in brewing lager-style beers were formed by hybridization between the thermotolerant species *Saccharomyces cerevisiae* and the cryotolerant species *Saccharomyces eubayanus* and inadvertently selected by brewers over numerous generations. Lager-style beers are fermented at low temperatures, and it has been known for some time that lager yeasts inherited the ability to ferment at these low temperatures from their *S. eubayanus* parent. Intriguingly, lager hybrids also inherited their mtDNA from the *S. eubayanus* parent. In the following study, we assayed the influence of parental mitotype on relative growth in synthetic *S. cerevisiae* x *S. eubayanus* hybrids across a wide range of temperatures. We also examined the impact of exchanging the native mtDNA present in an industrial lager-brewing hybrid with mtDNA from *S. cerevisiae*. In this way we demonstrated that mtDNA influences the growth of *Saccharomyces* hybrids, at both high and low temperatures, and showed a strong influence of mitotype in an industrial lager hybrid.

**[0072]** Suitable thermal tolerance is a critical component of how organisms adapt to their environment. Studies have begun to establish the link between variation in mitochondrial DNA (mtDNA) sequence (mitotypes), mitochondrial function, and temperature adaptation between populations, particularly in metazoans. The mitochondrial climatic adaptation hypothesis (Camus et al. 2017) posits that functional variation between mitotypes plays an important role in shaping adaptation of organisms to their thermal environment. Support for this hypothesis comes from both indirect and, increasingly, direct lines of evidence. Clines of mitotypes along temperature gradients or associations between mitotype and distinct thermal environments have been observed for numerous metazoan species (Cheviron and Brumfield 2009; DuBay and Witt 2014; Quintela et al. 2014; Silva et al. 2014; Bans et al. 2016), including *Drosophila* (Camus et al. 2017), salmon (Consuegra et al. 2015), shrews (Fontanillas et al. 2005), whales (Foote et al.

**[0073]** 2011), and humans (Mishmar et al. 2003) among others. Experiments in invertebrates have demonstrated directly that different mitotypes can alter temperature tolerance (Willett 2011; Pichaud et al. 2013), and more recently, direct experimental evidence has emerged for the role of mitotype in thermal adaption in natural environments (Dingley et al. 2014; Camus et al. 2017).

**[0074]** Genetically dissecting mitochondrially-encoded traits in metazoans is difficult, due to their obligately sexual

lifestyle and uniparental mitochondrial inheritance. In addition, mitochondrial candidate loci are difficult to identify, unless there are only a small number of nucleotide differences between mitotypes, and differences in non-coding regions are even more difficult to detect (Camus et al. 2017). Furthermore, the results of experiments with mitochondria in animals can be confounded by sex- and tissue-specific differences in mitochondrial function (Fontanillas et al. 2005; Wolff et al. 2016; Camus et al. 2017). For fine scale genetic dissection of traits linked to mtDNA, a more genetically tractable system is desirable.

**[0075]** Recent work has shown that mitotype can also play a role in thermotolerance in the model fungal yeast genus *Saccharomyces* (Paliwal et al. 2014; Špirek et al. 2014; Wolters et al. 2018). The *Saccharomyces* genus consists of eight known species (Liti et al. 2006; Hittinger 2013; Naseeb et al. 2017), which can be broadly divided between cryotolerant and thermotolerant species. Thermotolerant strains (maximum growth temperature  $\geq 36^\circ\text{C}$ .) form a clade that includes the standard model organism *Saccharomyces cerevisiae* (Gonçalves et al. 2011; Salvadó et al. 2011). To date, the genetics of temperature preference, particularly preference for cold temperatures, in *Saccharomyces* yeasts has been difficult to ascertain. Only three candidate loci have been identified as supporting cryotolerance in *Saccharomyces*, two in *S. cerevisiae* and one in a hybrid between the thermotolerant species *S. cerevisiae* and the distantly related cryotolerant species *S. eubayanus* (Yamagishi et al. 2010; Libkind et al. 2011; Gibson et al. 2013; Paget et al. 2014). Recent studies have found that both within and between species variation in mitotype can impact thermotolerance in *Saccharomyces*. Most work in this area has focused on the impact of intraspecies variation in mitotype within *S. cerevisiae* (Paliwal et al. 2014; Wolters et al. 2018) or on interspecies differences between *S. cerevisiae* and its thermotolerant sister species *S. paradoxus* (Leducq et al. 2017), though some work has also investigated more distant genetic relationships within *Saccharomyces* (Špirek et al. 2014). These studies have been largely concerned with mitochondrial function under heat-related stress ( $\sim 37^\circ\text{C}$ .). However, mitotype could influence temperature tolerance in *Saccharomyces* across a broad range of temperatures, not just at thermal extremes. Indeed, evidence from arctic species suggests that mitochondrial adaption specifically to cold conditions is common (Foote et al. 2011; Garvin et al. 2011; Melo-Ferreira et al. 2014). Furthermore, in a recent study of hybrids of *S. cerevisiae* and the cryotolerant species *Saccharomyces uvarum* when allele specific expression was measured at both  $22^\circ\text{C}$ . and  $37^\circ\text{C}$ ., an unexpected abundance of mitochondrial genes were identified as having allele-specific differences in expression, not only at  $37^\circ\text{C}$ ., but also at  $22^\circ\text{C}$ ., where both *S. cerevisiae*, *S. uvarum*, and their hybrid grow robustly (Li and Fay 2017). These results suggest the importance of mitochondrial DNA, even at moderate temperatures.

**[0076]** Among the other cryotolerant species of the *Saccharomyces* genus are *Saccharomyces eubayanus*, *Saccharomyces arboricola*, and *Saccharomyces kudriavzevii*. Together, *S. uvarum* and its sister species *S. eubayanus* form the small “*Saccharomyces bayanus*” clade of *Saccharomyces* (Hittinger 2013), which diverged from *S. cerevisiae* roughly 20 million years ago (Kellis et al. 2003). This amount of divergence represents approximately the equivalent genetic divergence between humans and chickens (Du-

jon 2006). While the thermotolerant species *S. cerevisiae* is best known for its role in human-associated fermentations, most commercial brewing occurs using cryotolerant *S. cerevisiae* x *S. eubayanus* hybrids in the production of lager-style beers (Libkind et al. 2011). These lager-brewing hybrids are distinguished by the tendency of the yeast to drop to the bottom of fermentations (bottom fermenting), a distinct flavor profile, and robust fermentation at low temperatures ( $\sim 7\text{--}15^\circ\text{C}$ .) (Tenge 2009). In comparison, ale-brewing yeasts tend to float at the top of fermentations and are used to brew at relatively high temperatures ( $15\text{--}24^\circ\text{C}$ .). Most ale strains have been found to consist of pure *S. cerevisiae* genetic material, though some brewing strains classified as ales and isolated from low-temperature regions of Europe, have been determined to be *S. cerevisiae* x *S. kudriavzevii* hybrids (Peris, Belloch, et al. 2012; Peris et al. 2018). Perhaps not surprisingly, the *S. cerevisiae* component of lager yeasts has been found to be most similar to other strains of *S. cerevisiae* used in beer brewing (Dunn and Sherlock 2008; Gonçalves et al. 2016). Among strains of *S. eubayanus* that have been characterized, strains belonging to the Holarctic lineage have been identified as being closely related to the population of *S. eubayanus* that gave rise to lager yeasts (Bing et al. 2014; Peris & Langdon et al. 2016). **[0077]** With the discovery of the wild-stock of *S. eubayanus* (Libkind et al. 2011), there is substantial interest in developing novel lager-brewing hybrids (Hebly et al. 2015; Krogerus et al. 2015; Krogerus, Magalhães, et al. 2017; Hittinger et al. 2018) and, therefore, in understanding the genetics of brewing-relevant traits, such as temperature tolerance. Intriguingly, the two lineages of lager-brewing yeast and other industrial hybrids inherited their mtDNA from their cryotolerant parent, *S. eubayanus* (Nakao et al. 2009; Baker et al. 2015; Okuno et al. 2016; Peris et al. 2017), but the influence of mtDNA on cryotolerance in lager-brewing yeast is unknown. Here, to determine whether mtDNA plays a role in temperature tolerance in hybrids of *S. cerevisiae* and *S. eubayanus*, we tested relative growth of newly created synthetic hybrids inheriting different parental mitotypes. We also directly tested the influence of mtDNA in an industrial lager strain by replacing the *S. eubayanus* mitotype with mtDNA from *S. cerevisiae*.

**[0078]** The genetic tools available in *Saccharomyces* yeasts permitted us to manipulate the inheritance of mtDNA in identical nuclear backgrounds. We find that, when mtDNA from the thermotolerant parent, *S. cerevisiae*, is inherited, hybrids have superior growth over hybrids with *S. eubayanus* mtDNA when grown at high temperatures. Likewise, hybrids with mtDNA from the cryotolerant parent, *S. eubayanus*, have growth superior to hybrids with *S. cerevisiae* mtDNA at low temperatures. In this way, we show that mitotype directly influences the relative ability of otherwise identical strains to grow at different temperatures.

## Results

**[0079]** *S. cerevisiae* and *S. eubayanus* Parent Strains are Thermotolerant and Cryotolerant, Respectively.

**[0080]** In order to establish relative differences in growth between *S. cerevisiae*, *S. eubayanus*, and their hybrids carrying different mitotypes, dilution series of each set of hybrids and parents were spotted onto plates containing either glucose or glycerol as the sole carbon source. As a non-fermentable carbon source, glycerol forces yeasts to utilize their mitochondria via respiration, rather than relying

on alcoholic fermentation, their preferred metabolic process of *Saccharomyces* (Crabtree/Warburg Effect) (Dashko et al. 2014). On glucose,  $\rho^0$  strains followed the same patterns of growth as their  $\rho^+$  parent but grew less at all temperatures (FIGS. 4A-4C), reflecting the well-known “petite” phenotype of respiratorily deficient cells (Merico et al. 2007).  $\rho^0$  strains were completely unable to grow on glycerol.

**[0081]** On both glucose and glycerol, the *S. eubayanus* and *S. cerevisiae* parents had opposite temperature responses (FIGS. 1A-1B). *S. eubayanus* strains grew at all temperatures, except 37° C., while *S. cerevisiae* strains began to decline in relative growth at 15° C. and were completely unable to grow at 4° C., a temperature at which the *S. eubayanus* strains still grew well (FIGS. 1A-1B and FIGS. 4A-4C, 5A-5C, 6A-6C, and 7A-7C). These results are consistent with the description of *S. eubayanus* as a cryotolerant species (Libkind et al. 2011), with a similar temperature range to its sister species, *S. uvarum*, and are also consistent with the well-known thermotolerant nature of *S. cerevisiae* strains (Gonçalves et al. 2011; Salvadó et al. 2011). Strain-specific differences were also apparent. *Sc* and *SeNC* both grew relatively weakly compared to the other parental strains (FIGS. 1A-1B). For *Sc*, poor growth was likely driven by the presence of multiple auxotrophies, but the reason for *SeNC*’s poor performance is unknown.

#### Mitotype Influences Temperature Preference in Hybrids According to the Parental Temperature Profile.

**[0082]** In general, heterosis was clear for hybrids grown on glucose across all temperatures tested (FIG. 1C and 1D, FIGS. 4A-4C), as has been previously observed for synthetic *S. cerevisiae* x *S. eubayanus* hybrids (Hebly et al. 2015). While relative growth was typically high for hybrids of both mitotypes on glucose, subtle differences were apparent (FIG. 1C). Hybrids carrying *S. cerevisiae* mtDNA had significantly greater growth than hybrids carrying *S. eubayanus* mtDNA between 22 and 37° C., while at 4 and 10° C., it was hybrids with *S. eubayanus* mtDNA that had significantly greater growth. There was no significant difference in growth between hybrids grown at 15° C. on glucose. These same trends were also seen on glycerol but were exaggerated (FIG. 1D), with significant differences between mitotypes at all temperatures. On glycerol, relative growth was greater for *S. eubayanus* mitotype hybrids between 4 and 15° C., while relative growth was greater for hybrids with *S. cerevisiae* mtDNA between 22 and 37° C.

**[0083]** Relative growth patterns for hybrids of different mitotypes were consistent across individual crosses (FIGS. 1A-1B and FIGS. 4A-4C, 5A-5C, 6A-6C, and 7A-7C) and when the data were analyzed statistically in aggregate (FIGS. 1C and 1D). Hybrids carrying mtDNA inherited from the *S. eubayanus* parent, whether from the type strain or the North Carolinian strain, had relatively greater growth at low temperatures compared to hybrids carrying mtDNA inherited from either *S. cerevisiae* parent. Conversely, hybrids with *S. cerevisiae* mtDNA, regardless of whether the mtDNA was from the laboratory strain or the ale strain, had relatively more growth at high temperatures compared to hybrids with *S. eubayanus* mtDNA, with the exception of the *ScAle* X *SeNC*  $\rho^{ScAle}$  hybrid, which had a substantial growth defect at 37° C. (FIGS. 7A-7C). The growth of the *ScAle* X *SeNC*  $\rho^{ScAle}$  at other temperatures was otherwise consistent with *S. cerevisiae* mtDNA supporting greater growth at high temperatures (FIG. 7B). On glycerol at 37°

C., despite still displaying a clear growth defect, the *ScAle* X *SeNC*  $\rho^{ScAle}$  hybrid still had greater growth than the *ScAle* X *SeNC*  $\rho^{SeNC}$  hybrid (FIG. 7C). These results strongly support the contribution of mtDNA to temperature tolerance in *S. cerevisiae* x *S. eubayanus* hybrids, despite strain-specific differences.

#### *S. cerevisiae* Mitochondrial DNA Improves Thermotolerance of an Industrial Lager Strain

**[0084]** Unlike the synthetic hybrids assayed above, the *S. cerevisiae* and *S. eubayanus* nuclear genomes of industrial lager-brewing yeast hybrids have been evolving in to lagering conditions for numerous generations (Meussdoerffer 2009; Gibson and Liti 2015). As a result, the effect of mtDNA on temperature tolerance may not be the same in these industrial hybrids as it was for newly generated hybrids. To test if mtDNA still play a role in temperature tolerance in lager-brewing yeast, the native lager mtDNA of *S. eubayanus* origin (Nakao et al. 2009; Peris et al. 2014; Baker et al. 2015; Okuno et al. 2016; Peris et al. 2017) was replaced with *S. cerevisiae* mtDNA from *Sc* and *ScAle* yeasts (FIG. 2A). Consistent with the results for our synthetic hybrids, lager yeasts carrying *S. cerevisiae* mtDNA had greater growth at higher temperatures and increased sensitivity to colder temperatures, with the effect greatly exaggerated on glycerol (FIGS. 2B and 2C). Differences between  $\rho^{Sc}$  and  $\rho^{ScAle}$  lager cybrids were also clear, especially on glucose. On glucose, there was no difference in growth between lager yeast carrying its native (*S. eubayanus*) mtDNA and those carrying *Sc* mtDNA, except at temperature extremes (FIG. 2B). In contrast, lager  $\rho^{ScAle}$  strains grew significantly less than the lager strain with its native mtDNA at most temperatures (FIG. 2B), despite the fact that the *ScAle*  $\rho^+$  strain displayed relatively robust growth across most temperatures in comparison to the industrial lager yeast with its native mtDNA (FIG. 8B).

#### Discussion

##### Mitotype Influences Temperature Tolerance in Synthetic Hybrids

**[0085]** Overall, hybrids had an increased range of temperatures they could tolerate compared to their parent strains, regardless of which mtDNA they carried. This heterosis was most evident at temperature extremes. On glucose at 37° C., hybrids grew most like their *S. cerevisiae* parent, while under 15° C., they grew like their *S. eubayanus* parent. These results support a strong role for the nuclear genome in temperature tolerance and indicate a certain amount of codominance between alleles supporting thermotolerance and cryotolerance. While this overall robustness to temperature was observed regardless of which mtDNA a hybrid carried, there were clear and consistent differences in relative growth between hybrids of different mitotypes. At higher temperatures, the *S. cerevisiae* mitotypes permitted increased growth relative to the *S. eubayanus* mitotypes, while the same was true for *S. eubayanus* mitotypes at lower temperatures, correlating with the relative cryotolerance and thermotolerance of their respective species of origin. Since the nuclear component is identical between hybrids of the same cross, these differences must be due to differences encoded in their mtDNA. While trends in temperature preference were apparent on both respiratory and fermentable carbon sources, the effect was exaggerated on respiratory media where mitochondrial respiration is required for

growth. These results were consistent across multiple strain backgrounds, indicating the generality of mtDNA effects on temperature preferences between these species.

Putative Strain-Specific Cytonuclear Incompatibilities Between *S. cerevisiae* and *S. eubayanus*

**[0086]** Out of six *SeNC*  $\rho^0$  strains tested, we were only able to generate a small number of hybrids with *S. cerevisiae* strains that carried *S. cerevisiae* mtDNA, and then only with a single *SeNC*  $\rho^0$  strain (yHEB1638). Compared to *Se*  $\rho^-$  and  $\rho^0$  strains and the *SeNC*  $\rho^+$  parent, where every mating attempt with *S. cerevisiae* strains produced hybrids, successful mating with yHEB1638 was sporadic, with only one out of ten mating attempts resulting in respiratorily competent hybrids. Difficulty forming hybrids was not the only unusual characteristic of the *S. cerevisiae* x *SeNC* hybrids with *S. cerevisiae* mtDNA. While the *Sc* X *SeNC*  $\rho^{Sc}$  hybrid had high relative growth at 37° C., like other hybrids carrying *S. cerevisiae* mitochondria, relative growth for the *ScAle* X *SeNC*  $\rho^{ScAle}$  hybrid plummeted at 37° C. Interestingly, even with this severe temperature-related growth defect, the *ScAle* mitotype still supported greater growth at 37° C. on glycerol than the *SeNC* mitotype. Because we were only able to form *S. cerevisiae* mtDNA carrying hybrids with one *SeNC*  $\rho^0$  strain, it is unclear if this temperature-dependent growth defect is specific to the yHEB1638 background or general to all *ScAle* X *SeNC*  $\rho^{ScAle}$  crosses. Even if the defect is specific to yHEB1638 and not *SeNC* in general, it is interesting that it was only detrimental in the *ScAle* background, as the *Sc* X *SeNC*  $\rho^{Sc}$  hybrid did not have the same sensitivity to 37° C., despite sharing the same *SeNC*  $\rho^0$  parent. Other studies have also found mitotype-related defects in temperature both within and between species in *Saccharomyces* in the study of cybrids (Paliwal et al. 2014; Špirek et al. 2014), though not interspecies hybrids as we have here. The potential for dominant cytonuclear incompatibilities could explain why, in hybrids of *Saccharomyces*, it has been observed that there is a tendency for there to be greater loss of nuclear genetic material from the parental genome that did not contribute the mtDNA (Marinoni et al. 1999; Peris, Lopes, et al. 2012; Peris et al. 2018). Intriguingly, another study also recently uncovered a strain-specific incompatibility between *S. cerevisiae* and *S. eubayanus* (Mertens et al. 2015). Taken together, these results imply that strain-specific incompatibilities exist between *S. eubayanus* and *S. cerevisiae* that prevent them from mating and/or forming viable offspring, which might be condition-specific in some cases.

#### Influence and Origin of Mitotype in Industrial Lager Yeasts

**[0087]** The impact of mtDNA on lager strain temperature tolerance was broadly similar to what was observed for synthetic hybrids, with some differences. In synthetic crosses of *S. eubayanus* and *S. cerevisiae*, hybrids generally experienced robust growth across all temperatures, regardless of which mtDNA they carried, though mtDNA was more important at temperature extremes, especially on glycerol (FIGS. 1C-1D). This observation supports a strong role of the nuclear genome in supporting general temperature tolerance. In contrast, the industrial lager-brewing hybrid W34/70 was unable to grow at 37° C. and steadily declined in relative growth as temperature increased. Swapping the native *S. eubayanus* mtDNA for *S. cerevisiae* mtDNA, increased tolerance to high temperatures on glycerol and, to a lesser extent, on glucose, but the relative growths of the

cybrids still declined precipitously as the temperature increased, and they were unable to grow at 37° C. Based on these results, it is likely that, after many generation of selection for cold fermentation, the nuclear genes necessary to support growth at higher temperatures are no longer functional, present, and/or adequately expressed in the industrial hybrid. As a result, any contribution to thermotolerance from the mtDNA is relatively minor on fermentable carbon and insufficient to rescue growth at temperature extremes on either carbon source.

**[0088]** These results are particularly interesting because W34/70 is part of the Froberg or Group II (Dunn and Sherlock 2008; Nakao et al. 2009) lineage of industrial lager yeasts. Compared to the Saaz or Group I lineage of industrial lager yeasts, the Froberg lineage has a relatively larger *S. cerevisiae* contribution to its genome (Dunn and Sherlock 2008). The higher *S. cerevisiae* contribution has been associated with relatively greater thermotolerance among Froberg lineage strains (Dunn and Sherlock 2008; Gibson et al. 2013; Walther et al. 2014). This correlation suggests that, even in a relatively thermotolerant industrial lager strain, the capacity of the *S. cerevisiae* nuclear genome to provide thermotolerance has been substantially reduced compared to the ancestral hybrid. It is nonetheless noteworthy that, even in a genetic background where the nuclear component of thermotolerance has been greatly diminished, mtDNA still plays a clear role in temperature tolerance.

**[0089]** It is tempting to speculate about what factors might have favored the retention of *S. eubayanus* mtDNA over *S. cerevisiae* mtDNA in present-day industrial lager-brewing hybrids. Given the difference in growth between our synthetic hybrids and the industrial lager hybrid and cybrids, it is likely that substantial changes occurred with regard to temperature tolerance over the course of adaption to lagering conditions. It is also evident that much of this change is attributable to changes within the nuclear genome. Even so, the mtDNA inherited still has a significant impact on temperature tolerance in all strains tested, with the *S. eubayanus* mtDNA favoring growth at lower temperatures. Increased cold tolerance could have given hybrids carrying the *S. eubayanus* mtDNA a selective advantage at the lower temperatures and high population densities at which lagers are brewed.

**[0090]** It is interesting to consider that interspecies incompatibilities, along with the ability to grow at low temperatures, might also have been a factor driving the retention of *S. eubayanus* mtDNA in industrial lagers. Of our synthetic hybrids, the *ScAle* X *SeNC* hybrids are the most genetically similar to the strains that gave rise to industrial lager hybrids. Like other hybrids tested, at low temperatures those that carried *S. eubayanus* type mtDNA had a growth advantage over hybrids that carried *S. cerevisiae* type mtDNA. Unlike other hybrids, the *ScAle* X *SeNC* hybrids with *ScAle* mtDNA had a severe growth defect at 37° C., the highest temperature assayed. As discussed above, we cannot be certain if this is a strain-specific defect or one general to all *ScAle* X *SeNC*  $\rho^{ScAle}$  strains. However, if a high-temperature growth defect is common to hybrids between *S. cerevisiae* ale strains and Holarctic lineage *S. eubayanus* strains that inherit the *ScAle* mitotype, those hybrids that inherited the *S. eubayanus* mitotype could have had another significant advantage above and beyond superior growth at lower temperatures.

**[0091]** One of the initial stages of beer manufacturing is the production of wort by boiling malted grain to extract the sugar component (Krottenthaler et al. 2009). Today, industrial brewers use modern cooling systems to cool the wort after boiling (Schu 2009), but historically, wort was cooled in open troughs, allowing air to pass freely over the hot liquid (Unger 2004: 167). This process exposed the wort to microbes that could colonize and ferment the wort into beer; a similar process is still used in the manufacture of lambics (Burberg and Zarnkow 2009) and American coolship ales (Bokulich and Bamforth 2013). In this scenario, the hybrids with the *S. eubayanus* mtDNA would, not only have had an advantage at the lower temperatures, but would have had an immediate advantage in being able to colonize the wort while it was still too hot for hybrids with *ScAle* mtDNA. The ability to colonize the wort early and continue rapid growth as the temperature cooled could have given hybrids with *S. eubayanus* mtDNA an insurmountable advantage, not only over their pure *S. cerevisiae* and *S. eubayanus* parents, but also over other hybrids carrying *S. cerevisiae* mtDNA.

#### Conclusions

**[0092]** We have shown that mtDNA can have a significant impact on the thermotolerance of hybrids between *S. cerevisiae* and *S. eubayanus*. The identification of a role for mtDNA in temperature adaptation in *Saccharomyces* yeasts offers a new genetically and experimentally tractable tool outside of metazoan systems with which to investigate the mitochondrial climatic adaptation hypothesis (Camus et al. 2017). A particularly exciting possibility from recent work in *S. cerevisiae* is the potential to map differences in thermotolerance, not only to nuclear loci, but to mitochondrially-encoded sequences as well by taking advantage of natural heteroplasmy and mtDNA recombination (Wolters et al. 2018).

**[0093]** While the finding that mtDNA influenced temperature preference in *S. cerevisiae* x *S. eubayanus* hybrids was general across the different strains tested, clear background-dependent difference were also observed. Given the interest in creating new lager hybrids for industrial use (Hebly et al. 2015; Krogerus et al. 2015; Mertens et al. 2015; Krogerus et al. 2016; Krogerus, Seppänen-Laakso, et al. 2017; Nikulin et al. 2018), it is clear that strain background, not only of the *S. cerevisiae* parent, but also the *S. eubayanus* parent, and the inheritance of mtDNA should all be important consid-

erations in strain construction. In addition to establishing a role for mtDNA in cryotolerance in lager yeast, we found that potential strain-specific incompatibilities suggest that *S. cerevisiae* x *S. eubayanus* hybrids could be a productive system for the study of genetic incompatibilities between species, particularly with regard to within species variation in such incompatibilities.

#### Methods

##### Yeast Strains and Strain Construction

**[0094]** Not all strains within a species are equally thermotolerant or cryotolerant, and different strains of *S. cerevisiae* can have 4° C. or more difference between their optimum growth temperatures (Salvadó et al. 2011). Since mitotype has been found to be important, at least at temperature extremes (Paliwal et al. 2014; Špirek et al. 2014; Leducq et al. 2017; Wolters et al. 2018), when determining thermotolerance in different strains of *S. cerevisiae*, we decided to include strains from different populations in our study. In addition to a laboratory strain of *S. cerevisiae* and a monosporic derivative of the type strain of *S. eubayanus*, an ale strain of *S. cerevisiae* and a strain of *S. eubayanus* isolated from North Carolina were also included (Hittinger and Carroll 2007; Libkind et al. 2011; Peris & Langdon et al. 2016). These two additional strains were chosen for their relative similarity to the parents of lager-brewing yeast hybrids.

**[0095]** Specifically, FM1283 (Sc) is descended from BY4724, which is itself a derivative of S288C (Brachmann et al. 1998; Hittinger and Carroll 2007). WLP530B (SeAle), is a commercial ale strain; its' pure *S. cerevisiae* background was confirmed by whole genome sequencing and assembly of reads to a concatenated pan-*Saccharomyces* reference genome, by use of the program sppIDer (<https://www.biorxiv.org/content/early/2018/05/30/333815>). FM1318 (Se) is a monosporic derivative of the taxonomic type strain of *S. eubayanus*, CBS 12357<sup>T</sup> (Libkind et al. 2011). The strain yHRVM108 (SeNC) was isolated from North Carolina and identified as being a close relative of the *S. eubayanus* parent of lager-brewing yeast hybrids (Peris and Langdon et al. 2016). W34/70 (Weihenstephan 34/70 or yHAB47) is an industrial strain belonging to the Froberg lineage of lager-brewing yeast hybrids (Peris and Langdon et al. 2016). All strains used in this study are listed in (Table 1).

TABLE 1

Strains and plasmids used in this Example								
Strain	Synonym	Species	Background	ρ status	MAT	Markers	Description	Source
FM1283	yHWA117	<i>S. cerevisiae</i>	BY4724 (S288C)	native mtDNA	a	ura3-Δ lys2-Δ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub>	Laboratory strain	Hittinger & Carroll ( <i>Nature</i> , 2007)
yHRVM481	—	<i>S. cerevisiae</i>	FM1283	null	a	ura3-Δ lys2-Δ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub>	ρ <sup>0</sup> laboratory strain	This study
yHRVM483	—	<i>S. cerevisiae</i>	FM1283	null	a	ura3-Δ lys2-Δ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub>	ρ <sup>0</sup> laboratory strain	This study

TABLE 1-continued

Strains and plasmids used in this Example								
Strain	Synonym	Species	Background	$\rho$ status	MAT	Markers	Description	Source
yHRVM485	—	<i>S. cerevisiae</i>	FM1283	null	a	ura3- $\Delta$ lys2- $\Delta$ $P_{TDH3}$ - yEBFP- T <sub>CYC1</sub>	$\rho^0$ laboratory strain	This study
WLP530B	yHAB48	<i>S. cerevisiae</i>	—	native mtDNA	a/ $\alpha$	—	Brewing isolate (Belgian Ale)	White Labs
yHWA245	—	<i>S. cerevisiae</i>	RM11-1a	native mtDNA	a	ade2- $\Delta$ ::HERP1 leu2- $\Delta$ ura3- $\Delta$ ho $\Delta$ ::KanMX	Laboratory strain used to amplify the KanMX cassette with overhangs to the HO locus for allele replacement in WLP530B	Alexander et al. ( <i>Genetics</i> , 2014)
yHEB1528	—	<i>S. cerevisiae</i>	WLP530B	native mtDNA	a	ho $\Delta$ ::KanMX	Stable haploid of brewing isolate	This study
yHEB1632	—	<i>S. cerevisiae</i> - ale	WLP530B	null	a	ho $\Delta$ ::KanMX	$\rho^0$ stable haploid of brewing isolate	This study
yHEB1621	—	<i>S. cerevisiae</i> - ale	WLP530B	null	a	ho $\Delta$ ::KanMX	$\rho^0$ stable haploid of brewing isolate	This study
yHEB1623	—	<i>S. cerevisiae</i> - ale	WLP530B	null	a	ho $\Delta$ ::KanMX	$\rho^0$ stable haploid of brewing isolate	This study
FM1318	yHEB10	<i>S. eubayanus</i>	—	native mtDNA	a/ $\alpha$	—	Monosporic derivative of the type strain	Libkind & Hittinger et al. ( <i>PNAS</i> , 2011)
yHEB162	—	<i>S. eubayanus</i>	FM1318	native mtDNA	$\alpha$	ho $\Delta$ ::NatMX	Stable haploid of FM1318	This study
yHEB1611	—	<i>S. eubayanus</i>	yHEB162	null	$\alpha$	ho $\Delta$ ::NatMX	$\rho^0$ stable haploid of FM1318	This study
yHEB1613	—	<i>S. eubayanus</i>	yHEB162	null	$\alpha$	ho $\Delta$ ::NatMX	$\rho^0$ stable haploid of FM1318	This study
yHEB1614	—	<i>S. eubayanus</i>	yHEB162	null	$\alpha$	ho $\Delta$ ::NatMX	$\rho^0$ stable haploid of FM1318	This study
yHRVM108	yHEB449	<i>S. eubayanus</i>	—	native mtDNA	a/ $\alpha$	—	Holarctic population isolate	Peris & Langdon et al. ( <i>PLOS</i> <i>Genetics</i> , 2016)
yHEB1606	—	<i>S. eubayanus</i>	yHRVM108	native mtDNA	$\alpha$	ho $\Delta$ ::NatMX	Stable haploid of yHRVM108	This study
yHEB1633	—	<i>S. eubayanus</i>	yHEB1606	null		ho $\Delta$ ::NatMX	$\rho^0$ stable haploid of yHRVM108; potentially <i>S. cerevisiae</i> $\rho$ incompatible	This study
yHEB1634	—	<i>S. eubayanus</i>	yHEB1606	null		ho $\Delta$ ::NatMX	$\rho^0$ stable haploid of yHRVM108; potentially	This study

TABLE 1-continued

Strains and plasmids used in this Example								
Strain	Synonym	Species	Background	$\rho$ status	MAT	Markers	Description	Source
yHEB1635	—	<i>S. eubayanus</i>	yHEB1606	null		ho $\Delta$ ::NatMX	<i>S. cerevisiae</i> $\rho$ incompatible $\rho^0$ stable haploid of yHRVM108; potentially <i>S. cerevisiae</i> $\rho$ incompatible	This study
yHEB1636	—	<i>S. eubayanus</i>	yHEB1606	null		ho $\Delta$ ::NatMX	<i>S. cerevisiae</i> $\rho$ incompatible $\rho^0$ stable haploid of yHRVM108; potentially <i>S. cerevisiae</i> $\rho$ incompatible	This study
yHEB1637	—	<i>S. eubayanus</i>	yHEB1606	null		ho $\Delta$ ::NatMX	<i>S. cerevisiae</i> $\rho$ incompatible $\rho^0$ stable haploid of yHRVM108; potentially <i>S. cerevisiae</i> $\rho$ incompatible	This study
yHEB1638	—	<i>S. eubayanus</i>	yHEB1606	null	$\alpha$	ho $\Delta$ ::NatMX	<i>S. cerevisiae</i> $\rho$ incompatible $\rho^0$ stable haploid of yHRVM108; <i>S. cerevisiae</i> $\rho$ compatible	This study
MCC123	yHEB879	<i>S. cerevisiae</i>	—	null	a	ade2-1, ura3-52, kar1-1	$\rho^0$ karyogamy deficient	Thorsness & Fox ( <i>Genetics</i> , 1993)
MCC109	yHEB880	<i>S. cerevisiae</i>	—	null	$\alpha$	ade2-1, ura3-52, kar1-1	$\rho^0$ karyogamy deficient	Costanzo & Fox ( <i>Mol. Cell Biol.</i> , 1993)
yHEB1752	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	FM1283 $\times$ yHEB1611	<i>S. cerevisiae</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study
yHEB1749	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	FM1283 $\times$ yHEB1613	<i>S. cerevisiae</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study
yHEB1746	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	FM1283 $\times$ yHEB1614	<i>S. cerevisiae</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study
yHEB1127	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	yHRVM481 $\times$ yHEB162	<i>S. eubayanus</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study
yHEB1737	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	yHRVM483 $\times$ yHEB162	<i>S. eubayanus</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study
yHEB1740	—	<i>S. cerevisiae</i> $\times$ <i>S. eubayanus</i>	yHRVM485 $\times$ yHEB162	<i>S. eubayanus</i> mtDNA	a/ $\alpha$	ura3- $\Delta$ lys2- $\Delta$ P <sub>TDH3</sub> -yEBFP-T <sub>CYC1</sub> ho $\Delta$ ::NatMX	synthetic hybrid	This study



TABLE 1-continued

Strains and plasmids used in this Example								
Strain	Synonym	Species	Background	$\rho$ status	MAT	Markers	Description	Source
yHEB1768	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1528 × yHEB1611	<i>S. cerevisiae</i> -ale mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
yHEB1766	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1528 × yHEB1613	<i>S. cerevisiae</i> -ale mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
yHEB1764	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1528 × yHEB1614	<i>S. cerevisiae</i> -ale mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
yHEB1761	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1632 × yHEB162	<i>S. eubayanus</i> mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
yHEB1758	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1621 × yHEB162	<i>S. eubayanus</i> mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
yHEB1755	—	<i>S. cerevisiae</i> × <i>S. eubayanus</i>	yHEB1623 × yHEB162	<i>S. eubayanus</i> mtDNA	a/α	hoΔ::KanMX/ hoΔ::NatMX	synthetic hybrid	This study
W34/70	yHAB47	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	Weihenstephan 34/70	native mtDNA	a/α	—	industrial lager-brewing strain; Froberg (Group II) lineage	Peris & Langdon et al. ( <i>PLOS Genetics</i> , 2016)
yHEB1626	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHAB47	null	a/α	—	$\rho^0$ industrial lager-brewing strain	This study
yHEB1627	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHAB47	null	a/α	—	$\rho^0$ industrial lager-brewing strain	This study
yHEB1628	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHAB47	null	a/α	—	$\rho^0$ industrial lager-brewing strain	This study
pHCT2	pHEB12	plasmid	—	—	—	NatMX	HyPr plasmid; for doxycycline-inducible mating type switching in <i>Saccharomyces</i> ; GenBank KT725395	Alexander et al. 2016 ( <i>Fungal Genet. Biol</i> )
yHEB1793	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1626	null	a/α	[pHCT2 (HyPr)]	mating competent $\rho^0$ lager strain	This study
yHEB1797	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1627	null	a/α	[pHCT2 (HyPr)]	mating competent $\rho^0$ lager strain	This study
yHEB1798	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1628	null	a/α	[pHCT2 (HyPr)]	mating competent $\rho^0$ lager strain	This study
yHEB1800	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1628	null	a/α	[pHCT2 (HyPr)]	mating competent $\rho^0$ lager strain	This study
yHEB1835	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1793	<i>S. cerevisiae</i> -ale mtDNA	a/α	[pHCT2 (HyPr)]	lager cybrid	This study
yHEB1827	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1797	<i>S. cerevisiae</i> -ale mtDNA	a/α	[pHCT2 (HyPr)]	lager cybrid	This study
yHEB1828	—	<i>S. cerevisiae</i> (2n) × <i>S. eubayanus</i> (2n)	yHEB1798	<i>S. cerevisiae</i> -ale mtDNA	a/α	[pHCT2 (HyPr)]	lager cybrid	This study

TABLE 1-continued

Strains and plasmids used in this Example								
Strain	Synonym	Species	Background	$\rho$ status	MAT	Markers	Description	Source
yHEB1839	—	<i>S. cerevisiae</i> (2n) $\times$ <i>S. eubayanus</i> (2n)	yHEB1793	<i>S. cerevisiae</i> -mtDNA	a/ $\alpha$	[pHCT2 (HyPr)]	lager cybrid	This study
yHEB1841	—	<i>S. cerevisiae</i> (2n) $\times$ <i>S. eubayanus</i> (2n)	yHEB1797	<i>S. cerevisiae</i> -mtDNA	a/ $\alpha$	[pHCT2 (HyPr)]	lager cybrid	This study
yHEB1843	—	<i>S. cerevisiae</i> (2n) $\times$ <i>S. eubayanus</i> (2n)	yHEB1800	<i>S. cerevisiae</i> -mtDNA	a/ $\alpha$	[pHCT2 (HyPr)]	lager cybrid	This study

**[0096]** To facilitate strain crossing, stable haploid *ScAle*, *Se*, and *SeNC* strains were generated by replacing one allele of the HO locus with a selectable marker by standard lithium acetate transformation (Gietz and Woods 2002; Alexander et al. 2014), with modifications made for transforming *S. eubayanus* (see below). Successful replacement of the HO locus was confirmed by PCR with primers specific to the HO locus (Table 2). The resultant strains were sporulated and individual tetrads dissected using a Singer Sporeplay. *ScAle*

was sporulated in liquid sporulation medium (1% potassium acetate, 0.005% zinc acetate) and grown at room temperature (~22° C.) before dissecting after 4-5 days. To sporulate *Se* and *SeNC*, 200  $\mu$ L of saturated culture were plated onto a YPD (1% yeast extract, 2% peptone, 2% glucose) plate and grown at room temperature for 3-5 days before dissecting tetrads. Strains lacking the HO coding sequence were selected for by growth on YPD+antibiotic, and the mating type was determined by mating with tester strains.

TABLE 2

Oligonucleotides used in this Example		
Name	Sequence (SEQ ID NO:)	Description
oHECP B15	AATACATACAACCTTAC TTTTTCAAATTAATT TACATACTAGATCTGT TTAGCTTGCCCTT (SEQ ID NO: 1)	amplifies MX-driven drug markers with overhangs to the <i>S. eubayanus</i> HO locus for allele replacement by homologous recombination.
oHECP B16	TCTATATAGACAACAA CCACTTCCACTAGCCT TTAAGCGAGCTCGTTT TCGACACTGGAT (SEQ ID NO: 2)	amplifies MX-driven drug markers with overhangs to the <i>S. eubayanus</i> HO locus for allele replacement by homologous recombination.
oHWA 565	CTATGGTTTACGAAAT GATCCACG (SEQ ID NO: 3)	primer specific to 450 bp upstream of <i>S. cerevisiae</i> HO locus; used to amplify region around HO locus with selection marker for allele replacement by homologous recombination, to confirm allele replacement, and to confirm the absence of $\rho$ carrier DNA in lager cybrids.
oHWA 566	CACTGACCCAGTCTTG TCTTC (SEQ ID NO: 4)	primer specific to 540 bp downstream of <i>S. cerevisiae</i> HO locus; used to amplify region around HO locus with selection marker for allele replacement by homologous recombination, to confirm allele replacement, and to confirm the absence of $\rho$ carrier DNA in lager cybrids.
oHWA 568	TTTGCAATCGAAGAC CCAT (SEQ ID NO: 5)	primer internal to oHWA565 and oHWA566; used for sequencing to confirm the absence of $\rho$ carrier DNA in lager cybrids.
oHMB3	GTTTCTGGCCGAGCTA CAAG (SEQ ID NO: 6)	primer specific to 260 bp upstream of <i>S. eubayanus</i> HO locus; used to confirm allele replacement.
oHMB4	CAAGCCATGTCTTCT CGTT (SEQ ID NO: 7)	primer specific to 410 bp upstream of <i>S. eubayanus</i> HO locus; used to confirm allele replacement.
oHECP B148	TTTGAATATCAATGAA AATGCC (SEQ ID NO: 8)	primer specific to the KAR1 locus of <i>S. cerevisiae</i> ; used to amplify the KAR1 locus to confirm absence of $\rho$ carrier DNA in lager cybrids.

TABLE 2-continued

Oligonucleotides used in this Example		
Name	Sequence (SEQ ID NO:)	Description
oHECP B149	TTAAACCTATAATAC ACATATATATTGC (SEQ ID NO: 9)	primer specific to the KAR1 locus of <i>S. cerevisiae</i> ; used to amplify and sequence the KAR1 locus to confirm absence of $\rho$ carrier DNA in lager cybrids.
oHDP2 5	TGCGCCAAGTGTCTGA AGAACAACCTGGGA (SEQ ID NO: 10)	primer general to the GAL4 locus of <i>Saccharomyces</i> ; used to amplify and sequence the GAL4 locus to confirm absence of $\rho$ carrier DNA in lager cybrids (Peris et al. 2012).
oHDP2 6	GCGATTTCAATCTGGT TATTATACAACATCAT (SEQ ID NO: 11)	primer general to the GAL4 locus of <i>Saccharomyces</i> ; used to amplify the GAL4 locus to confirm absence of $\rho$ carrier DNA in lager cybrids (Peris et al. 2012).
ITS1	TCCGTAGTGGAACCTG CGG (SEQ ID NO: 12)	standard primer to amplify out the 5.8S rDNA (ITS) sequence from fungi; used to amplify ITS region to confirm successful creation of synthetic hybrids (McCullough et al. 1998).
ITS4	TCCTCCGCTTATTGAT ATGC (SEQ ID NO: 13)	standard primer to amplify out the 5.8S rDNA (ITS) sequence from fungi; used to amplify and sequence ITS region to confirm successful creation of synthetic hybrids (McCullough et al. 1998).

*Saccharomyces* Lithium Acetate Transformation Protocol  
(Gietz and Woods 2002) Adapted for *S. eubayanus*

Materials:

- [0097] 1 M lithium acetate
- [0098] 100 mM lithium acetate
- [0099] 50% (w/v) PEG-4000
- [0100] Boiled single-stranded salmon sperm DNA (ssssDNA) (10 mg/mL)
- [0101] 100% Ethanol

Procedure:

- [0102] 1. Grow yeast overnight in 3 mL of YPD or other culturing medium.
- [0103] 2. The next day, inoculate 50 mL of fresh YPD with enough overnight culture to bring the OD to 0.25 and shake at 250 rpm at room temperature.
- [0104] Some strains of *S. eubayanus* can tolerate growth at 30° C. and will grow faster at this temperature, but other strains are sensitive and will grow much more slowly.
- [0105] 3. After four hours, take an OD<sub>600</sub> reading. If between 0.75 and 1.0 OD, continue to step 4; otherwise, allow to continue shaking.
- [0106] For slow growing strains, it may take several more hours to reach the appropriate OD.
- [0107] 4. Wash yeast cells in H<sub>2</sub>O.
- [0108] 5. Resuspend yeast cells in 950  $\mu$ L 100 mM lithium acetate.
- [0109] 6. Aliquot 100  $\mu$ L of cell suspension into Eppendorf tubes.
- [0110] 7. Spin at maximum speed in microcentrifuge for 1 min, then remove supernatant.
- [0111] 8. Add in this order:
  - [0112] 240  $\mu$ L 50% PEG-4000,
  - [0113] 36  $\mu$ L 1 M lithium acetate,
  - [0114] 43  $\mu$ L of DNA solution to be transformed,
  - [0115] 5  $\mu$ L boiled sssDNA.

- [0116] 9. Gently resuspend pellet in transformation mixture.
- [0117] 10. Heat shock for 55 minutes at 34° C.
- [0118] 37° C. works almost as well.
- [0119] 11. Add 36  $\mu$ L 100% Ethanol.
- [0120] 12. Heat shock for another 5 minutes.
- [0121] 13. After heat shock, briefly spin down and remove supernatant.
- [0122] 14. Immediately resuspend in 600  $\mu$ L of YPD.

Recovery Method 1

- [0123] 1) Incubate at room temperature for 3 hours on wheel or shaker.
- [0124] 2) Plate 200  $\mu$ L of transformation suspension to each of three selective media plates.
- [0125] 3) Incubate at room temperature.
- [0126] The strain can also be grown at 30° C. if the strain is not heat sensitive.

Recovery Method 2

- [0127] You MUST use this method for counter-selection (e.g. selecting for the loss of URA3 by 5-FOA resistance) to allow for protein turnover.
- [0128] 1) Plate immediately to YPD and allow to grow overnight.
- [0129] 2) The next day, replica-plate to selective media.
- [0130] Note: The spatial separation provided by this method guarantees that transformants are independent.
- [0131] If successful, colonies will generally appear after two or three days.
- [0132] Depending on the sensitivity of a strain to a given selection regime, it could take a week or more for colonies to appear, with new colonies appearing up to two weeks after transformation.

[0133] Gietz D. R., Woods R. A., 2002 Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* 350: 87-96.

#### Synthetic Hybrids

[0134] To test the effect of mitotype on temperature tolerance in *S. cerevisiae* x *S. eubayanus* hybrids, we made sets of hybrids containing mtDNA from one parent or the other. When two  $\rho^+$  yeast cells mate, the mtDNA of both parents is present in the zygote, but a single mtDNA haplotype is rapidly fixed after only a few cell divisions (Berger and Yaffe 2000). Which mtDNA haplotype is fixed often happens in a non-random manner (Zweifel and Fangman 1991; Marinoni et al. 1999; Hsu and Chou 2017), and recombinant mtDNAs are also possible, even common (Berger and Yaffe 2000; Wolters et al. 2018). To control the inheritance of mtDNA in synthetic hybrids, we generated  $\rho^0$  (mtDNA completely absent) strains to mate with  $\rho^+$  strains, so that mtDNA from only the  $\rho^+$  parent would be present in hybrids (FIG. 3).  $\rho^0$  strains were generated by treating  $\rho^+$  parent strains with ethidium bromide (Fox et al. 1991). Respiration-deficient strains were screened for by the absence of growth on glycerol, and the complete removal of mtDNA was confirmed by DAPI staining (Eckert-Boulet et al. 2011). Because of the mutagenic nature of ethidium bromide, to control for the effect of any spurious mutations, we generated  $\rho^0$  strains of each parent strain in triplicate.

[0135] Hybrids were made by mating a  $\rho^0$  strain of one species with a  $\rho^+$  strain of the opposite mating type of the other species. Mating was performed by mixing the parent strains together on a YPD plate and letting them mate overnight. Allowing mating to occur for one or two more days and/or at 30° C. sometimes improved mating efficiency. Hybrids were selected by growth on glycerol and resistance to the appropriate antibiotics. When appropriate drug selection markers were not present in the parental genomes, zygotes were picked manually and tested for growth on glycerol to confirm retention of functional mitochondria. The hybrid nature of all strains was confirmed by ITS sequencing (Table 2) (McCullough et al. 1998; Sylvester et al. 2015). To ensure the maintenance of functional mitochondria, hybrid strains were grown only on media with glycerol as the sole carbon source, except for during experiments.

[0136] In general, the different *S. cerevisiae* and *S. eubayanus* backgrounds and mitotypes readily formed hybrids, although mtDNA could be lost if hybrids were not grown on non-fermentable media. The exception was for crosses attempted between *S. eubayanus*-North Carolina (SeNC)  $\rho^0$  strains and *S. cerevisiae*  $\rho^+$  strains (both the lab and ale strains). Hybrids between SeNC  $\rho^0$  and *S. cerevisiae*  $\rho^+$  strains were attempted multiple times (>50 attempts total) with six independently generated SeNC  $\rho^0$  strains. Out of these attempts, only 4 successful hybrids were formed, one between yHEB1528 (ScAle  $\rho^+$ ) and yHEB1638 (SeNC  $\rho^0$ ) and three between yHWA117 (Sc  $\rho^+$ ) and yHEB1638 (SeNC  $\rho^0$ ). There was no similar difficulty producing the same hybrids with *S. eubayanus* mitochondrial genomes.

[0137] It is not clear if the ability to form respiratorily competent hybrids is unique to yHEB1638, as even hybrids with this strain took multiple attempts to achieve. Because the ethidium bromide used to generate  $\rho^0$  strains is broadly mutagenic, it is likely that yHEB1638 has a number of

mutations differentiating it from the other SeNC  $\rho^0$  strains we generated. It is possible that one of these changes allowed yHEB1638 to maintain functional mtDNA in hybrids with *S. cerevisiae* carrying *S. cerevisiae* mtDNA, whereas other SeNC  $\rho^0$  strains could not. We include the results of growth assays with the hybrids made using yHEB1638 to determine if they follow the same general trends as other hybrid strains, with the caveat that the results from these experiments cannot be verified by hybrids made from independently generated SeNC  $\rho^0$  strains.

#### Mitochondrial Transfers

[0138] To produce strains with a lager yeast nuclear background and *S. cerevisiae* mtDNA (cybrids), karyogamy-deficient (kar1-1)  $\rho^0$  strains (Conde and Fink 1976; Costanzo and Fox 1993; Thorsness and Fox 1993) were used to transfer mitochondria from a donor *S. cerevisiae* strain to a  $\rho^0$  lager strain (FIG. 2A), which were constructed as described above. Briefly, the lack of karyogamy in crosses with kar1-1 mutants allows the mixing of cytoplasm between mated cells, while preventing fusion between the nuclear genomes, ultimately leading to progeny with mixed cytoplasm, but only one nuclear background. In this way, donor mitochondria from *S. cerevisiae* strains were transferred into the kar1-1  $\rho^0$  strains by mating yeast as above and selecting for functional mtDNA (by growth on glycerol, a non-fermentable carbon source) and the kar1-1 background, while selecting against the donor strain background (FIG. 2A). Since the *S. cerevisiae*-ale strain, WLP530B (ScAle)), and its derivatives are prototrophic, and the kar1-1 strains (MCC109 and MCC123) are auxotrophic for ura3, we were able to select for the kar1-1 background and simultaneously select against the ScAle background by selecting for resistance to 5-fluoro-orotic acid (FOA). To select for kar1-1 background strains carrying mtDNA from the *S. cerevisiae* laboratory strain, FM1283 (Sc), strains were grown on minimal media supplemented with adenine. Because the kar1-1 strains are ade2-1 auxotrophs while Sc is auxotrophic for lys2, this selected for the kar1-1 genetic background and against the Sc genetic background. The medium was also supplemented with uracil for which both the kar1-1 strains and Sc are auxotrophic.

[0139] Because lager yeasts contain both MATa and MAT $\alpha$  at their mating type locus, mating does not usually occur. To mate polyploid lagers to the kar1-1  $\rho^+$  strains for mitochondria transfer, the MAT locus had to first be homozygosed. The MAT locus of lager  $\rho^0$  strains was homozygosed using a HyPr (Hybrid Production) plasmid (pHCT2) to induce mating type switching (Alexander et al. 2016). Cybrids, strains with a single nuclear background and mitochondria from a donor strain, were selected for by selecting against the kar1-1 background. To confirm that only lager genetic material was present in the resulting cybrids, three loci throughout the lager genome were sequenced to confirm that they contained only lager alleles (Table 2). As with hybrids, cybrids were also cultured on glycerol, except for during experiments, to ensure maintenance of mtDNA.

#### Growth Assays

[0140] Each hybrid and cybrid was constructed three times with an independently generated  $\rho^0$  parent. Each of these independent hybrids was tested three times at each tempera-

ture. In total, combining biological and technical replicates, each hybrid cross was tested a total of nine times at each temperature, with some exceptions. Since there was only one SeNC  $\rho^0$  strain with which we were able to successfully form hybrids containing *S. cerevisiae* mtDNA, only one biological replicate for each *S. cerevisiae* strain was formed with SeNC, which each had three technical replicates at each temperature. Consequently, these hybrids (*Sc* X SeNC  $\rho^{ScAle}$  and *ScAle* X SeNC  $\rho^{ScAle}$ ) only had three replicates total at each temperature. In addition, because of contamination or poor photo quality a small number of replicates ( $n=5$ ) had to be discarded. These were: two for *Sc* X SeNC experiments on glycerol, one at 22° C. and one at 37° C.; and three for lager cybrid experiments at 4° C., one replicate growing on glucose and two growing on glycerol.

[0141] Yeast strains were grown in liquid synthetic complete (SC) medium (0.17% yeast nitrogen base, 0.5% ammonium sulfate, 0.2% complete drop out mix). Strains containing their native mtDNA and  $\rho^0$  strains were grown with 2% glucose, while hybrids and cybrids were grown with 2% glycerol and 2% ethanol to force the maintenance of mtDNA. After reaching saturation, cells were washed in either water or deflocculation buffer (20 mM citrate, 5 mM EDTA) and resuspended in either SC (without carbon) or deflocculation buffer to an OD<sub>600</sub> of 1+/-0.05. Due to the extremely flocculent nature of *ScAle*, cultures had to be washed and resuspended in deflocculation buffer. For consistency, all strains used in experiments with *ScAle* were treated identically with buffer. Yeast strains were plated in a dilution series of OD<sub>600</sub>=1.0, 10<sup>-1</sup>, 10<sup>-2</sup>, 10<sup>-3</sup>, and 10<sup>-4</sup>. Dilutions were plated onto SC plates containing either 2% glucose or 2% glycerol as the sole carbon source. Plates were grown at 4, 10, 15, 22, 30, and 37° C. Lager cybrids were also grown at 33.5° C. Plates were grown until at least one strain on a plate showed growth at all five dilutions or after they had been allowed to grow for more than two months, whichever came first.

#### Analysis of Growth Assays.

[0142] To determine how well different strains grew relative to each other, the combined intensity (a proxy for growth) of the first and second dilutions (OD<sub>600</sub>=1 and 10<sup>-1</sup>) were measured using custom CellProfiler pipelines (Lamprecht, Sabatini, & Carpenter, 2007; www.cellprofiler.org), and the values were combined. To be able to compare growth between plates, which may have differences in absolute intensity, growth on each plate was normalized by dividing by the strain with the highest measured combined intensity on each plate. This procedure created a relative growth score for each strain that was used to compare growth across different replicates. Statistically significant differences in growth were tested for using the Wilcoxon rank-sum test, as implemented in R version 3.4.3 (R Development Core Team 2017), and corrected for multiple tests using the Benjamini-Hochberg procedure (Benjamini and Hochberg 1995), as implemented in R version 3.4.3. P-values  $\leq 0.05$  were considered significant.

#### REFERENCES

[0143] Alexander W G, Doering D T, Hittinger C T. 2014. High-efficiency genome editing and allele replacement in prototrophic and wild strains of *Saccharomyces*. *Genetics* 198:859-866.

- [0144] Alexander W G, Peris D, Pfannenstiel B T, Oplente D A, Kuang M, Hittinger C T. 2016. Efficient engineering of marker-free synthetic allotetraploids of *Saccharomyces*. *Fungal Genet. Biol.* 89:10-17.
- [0145] Baker E, Wang B, Bellora N, Peris D, Hulfachor A B, Koshalek J A, Adams M, Libkind D, Hittinger C T. 2015. The Genome Sequence of *Saccharomyces eubayanus* and the Domestication of Lager-Brewing Yeasts. *Mol. Biol. Evol.* 32:2818-2831.
- [0146] Baris T Z, Blier P U, Pichaud N, Crawford D L, Oleksiak M F. 2016. Gene by environmental interactions affecting oxidative phosphorylation and thermal sensitivity. *Am. J. Physiol. Integr. Comp. Physiol.* 311:R157-R165.
- [0147] Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B* 57:289-300.
- [0148] Berger K H, Yaffe M P. 2000. Mitochondrial DNA inheritance in *Saccharomyces cerevisiae*. *Trends Microbiol.* 8:508-513.
- [0149] Bing J, Han P-J, Liu W-Q, Wang Q-M, Bai F-Y. 2014. Evidence for a Far East Asian origin of lager beer yeast. *Curr. Biol.* 24:R380-1.
- [0150] Bokulich N A, Bamforth C W. 2013. The microbiology of malting and brewing. *Microbiol. Mol. Biol. Rev.* 77:157-172.
- [0151] Brachmann C B, Davies A, Cost G J, Caputo E, Li J, Hieter P, Boeke J D. 1998. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* 14:115-132.
- [0152] Burberg F, Zarnkow M. 2009. Special Production Methods. In: Esslinger H M, editor. *Handbook of brewing processes, technology, markets*. Weinheim (Germany). p. 250.
- [0153] Camus M F, Wolff J N, Sgrò C M, Dowling D K. 2017. Experimental Support That Natural Selection Has Shaped the Latitudinal Distribution of Mitochondrial Haplotypes in Australian *Drosophila melanogaster*. *Mol. Biol. Evol.* 34:2600-2612.
- [0154] Cheviron Z A, Brumfield R T. 2009. MIGRATION-SELECTION BALANCE AND LOCAL ADAPTATION OF MITOCHONDRIAL HAPLOTYPES IN RUFOUS-COLLARED SPARROWS (*Zonotrichia capensis*) ALONG AN ELEVATIONAL GRADIENT. *Evolution* (N.Y.). 63:1593-1605.
- [0155] Conde J, Fink G R. 1976. A mutant of *Saccharomyces cerevisiae* defective for nuclear fusion. *Proc. Natl. Acad. Sci. U. S. A.* 73:3651-3655.
- [0156] Consuegra S, John E, Verspoor E, de Leaniz C G. 2015. Patterns of natural selection acting on the mitochondrial genome of a locally adapted fish species. *Genet. Sel. Evol.* 47:58.
- [0157] Costanzo M C, Fox T D. 1993. Suppression of a defect in the 5' untranslated leader of mitochondrial COX3 mRNA by a mutation affecting an mRNA-specific translational activator protein. *Mol. Cell. Biol.* 13:4806-4813.
- [0158] Dashko S, Zhou N, Compagno C, Piškur J. 2014. Why, when, and how did yeast evolve alcoholic fermentation? *FEMS Yeast Res.* 14:826-832.
- [0159] Dingley S D, Polyak E, Ostrovsky J, Srinivasan S, Lee I, Rosenfeld A B, Tsukikawa M, Xiao R, Selak M A, Coon J J, et al. 2014. Mitochondrial DNA variant in

- COX1 subunit significantly alters energy metabolism of geographically divergent wild isolates in *Caenorhabditis elegans*. *J. Mol. Biol.* 426:2199-2216.
- [0160] DuBay S G, Witt C C. 2014. Differential high-altitude adaptation and restricted gene flow across a mid-elevation hybrid zone in Andean tit-tyrant flycatchers. *Mol. Ecol.* 23:3551-3565.
- [0161] Dujon B. 2006. Yeasts illustrate the molecular mechanisms of eukaryotic genome evolution. *Trends Genet.* 22:375-387.
- [0162] Dunn B, Sherlock G. 2008. Reconstruction of the genome origins and evolution of the hybrid lager yeast *Saccharomyces pastorianus*. *Genome Res.* 18:1610-1623.
- [0163] Eckert-Boulet N, Rothstein R, Lisby M. 2011. Cell biology of homologous recombination in yeast. *Methods Mol. Biol.* 745:523-536.
- [0164] Fontanillas P, Dépraz A, Giorgi M S, Perrin N. 2005. Nonshivering thermogenesis capacity associated to mitochondrial DNA haplotypes and gender in the greater white-toothed shrew, *Crocidura russula*. *Mol. Ecol.* 14:661-670.
- [0165] Foote A D, Morin P A, Durban J W, Pitman R L, Wade P, Willerslev E, Gilbert M T P, da Fonseca R R. 2011. Positive selection on the killer whale mitogenome. *Biol. Lett.* 7:116-118.
- [0166] Fox T D, Folley L S, Mulero J J, McMullin T W, Thorsness P E, Hedin L O, Costanzo M C. 1991. Analysis and manipulation of yeast mitochondrial genes. *Methods Enzymol.* 194:149-165.
- [0167] Garvin M R, Bielawski J P, Gharrett A J. 2011. Positive Darwinian Selection in the Piston That Powers Proton Pumps in Complex I of the Mitochondria of Pacific Salmon. *PLoS One* 6:e24127.
- [0168] Gibson B, Liti G. 2015. *Saccharomyces pastorianus*: genomic insights inspiring innovation for industry. *Yeast* 32:17-27.
- [0169] Gibson B R, Storgårds E, Krogerus K, Vidgren V. 2013. Comparative physiology and fermentation performance of Saaz and Froberg lager yeast strains and the parental species *Saccharomyces eubayanus*. *Yeast* 30:255-266.
- [0170] Gietz D R, Woods R A. 2002. Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* 350:87-96.
- [0171] Gonçalves M, Pontes A, Almeida P, Barbosa R, Serra M, Libkind D, Hutzler M, Gonçalves P, Sampaio J P. 2016. Distinct Domestication Trajectories in Top-Fermenting Beer Yeasts and Wine Yeasts. *Curr. Biol.* 26:2750-2761.
- [0172] Gonçalves P, Valério E, Correia C, de Almeida J M G C F, Sampaio J P. 2011. Evidence for divergent evolution of growth temperature preference in sympatric *Saccharomyces* species. *PLoS One* 6:e20739.
- [0173] Heblly M, Brickwedde A, Bolat I, Driessen M R M, de Hulster E A F, van den Broek M, Pronk J T, Geertman J-M, Daran J-M, Daran-Lapujade P. 2015. *S. cerevisiae* x *S. eubayanus* interspecific hybrid, the best of both worlds and beyond. *FEMS Yeast Res.* 15:fov005.
- [0174] Hittinger C T. 2013. *Saccharomyces* diversity and evolution: a budding model genus. *Trends Genet.* 29:309-317.
- [0175] Hittinger C T, Carroll S B. 2007. Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* 449:677-681.
- [0176] Hittinger C T, Steele J L, Ryder D S. 2018. Diverse yeasts for diverse fermented beverages and foods. *Curr. Opin. Biotechnol.* 49:199-206.
- [0177] Hsu Y-Y, Chou J-Y. 2017. Environmental Factors Can Influence Mitochondrial Inheritance in the *Saccharomyces* Yeast Hybrids. *PLoS One* 12:e0169953.
- [0178] Kellis M, Patterson N, Endrizzi M, Birren B, Lander E S. 2003. Sequencing and comparison of yeast species to identify genes and regulatory elements. *Nature* 423:241-254.
- [0179] Krogerus K, Arvas M, De Chiara M, Magalhães F, Mattinen L, Oja M, Vidgren V, Yue J-X, Liti G, Gibson B. 2016. Ploidy influences the functional attributes of de novo lager yeast hybrids. *Appl. Microbiol. Biotechnol.* 100:7203-7222.
- [0180] Krogerus K, Magalhães F, Vidgren V, Gibson B. 2015. New lager yeast strains generated by interspecific hybridization. *J. Ind. Microbiol. Biotechnol.* 42:769-778.
- [0181] Krogerus K, Magalhães F, Vidgren V, Gibson B. 2017. Novel brewing yeast hybrids: creation and application. *Appl. Microbiol. Biotechnol.* 101:65-78.
- [0182] Krogerus K, Seppänen-Laakso T, Castillo S, Gibson B. 2017. Inheritance of brewing-relevant phenotypes in constructed *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrids. *Microb. Cell Fact.* 16:66.
- [0183] Krottenthaler M, Back W, Zarnkow M. 2009. Wort Production. In: Esslinger H M, editor. *Handbook of brewing processes, technology, markets*. Weinheim (Germany). p. 165-205.
- [0184] Lamprecht M R, Sabatini D M, Carpenter A E. 2007. CellProfiler: free, versatile software for automated biological image analysis. *Biotechniques* 42:71-75.
- [0185] Leducq J-B, Henault M, Charron G, Nielly-Thibault L, Terrat Y, Fiumera H L, Shapiro B J, Landry C R. 2017. Mitochondrial Recombination and Introgression during Speciation by Hybridization. *Mol. Biol. Evol.* 34:1947-1959.
- [0186] Li X C, Fay J C. 2017. Cis-Regulatory Divergence in Gene Expression between Two Thermally Divergent Yeast Species. *Genome Biol. Evol.* 9:1120-1129.
- [0187] Libkind D, Hittinger C T, Valério E, Gonçalves C, Dover J, Johnston M, Gonçalves P, Sampaio J P. 2011. Microbe domestication and the identification of the wild genetic stock of lager-brewing yeast. *Proc. Natl. Acad. Sci. U. S. A.* 108:14539-14544.
- [0188] Liti G, Barton D B H, Louis E J. 2006. Sequence diversity, reproductive isolation and species concepts in *Saccharomyces*. *Genetics* 174:839-850.
- [0189] Marinoni G, Manuel M, Petersen R F, Hvidtfeldt J, Sulo P, Piskur J. 1999. Horizontal Transfer of Genetic Material among *Saccharomyces* Yeasts Horizontal Transfer of Genetic Material among *Saccharomyces* Yeasts. 181:6488-6496.
- [0190] McCullough M J, Clemons K V, McCusker J H, Stevens D A. 1998. Intergenic transcribed spacer PCR ribotyping for differentiation of *Saccharomyces* species and interspecific hybrids. *J. Clin. Microbiol.* 36:1035-1038.
- [0191] Melo-Ferreira J, Vilela J, Fonseca M M, da Fonseca R R, Boursot P, Alves P C. 2014. The Elusive Nature

- of Adaptive Mitochondrial DNA Evolution of an Arctic Lineage Prone to Frequent Introgression. *Genome Biol. Evol.* 6:886-896.
- [0192] Merico A, Sulo P, Piškur J, Compagno C. 2007. Fermentative lifestyle in yeasts belonging to the *Saccharomyces* complex. *FEBS J.* 274:976-989.
- [0193] Mertens S, Steensels J, Saels V, De Rouck G, Aerts G, Verstrepen K J. 2015. A large set of newly created interspecific *Saccharomyces* hybrids increases aromatic diversity in lager beers. *Appl. Environ. Microbiol.* 81:8202-8214.
- [0194] Meussdoerffer F G. 2009. A Comprehensive History of Beer Brewing. In: Esslinger H M, editor. *Handbook of brewing processes, technology, markets.* Weinheim (Germany): Wiley-VCH. p. 28,31-32.
- [0195] Mishmar D, Ruiz-Pesini E, Golik P, Macaulay V, Clark A G, Hosseini S, Brandon M, Easley K, Chen E, Brown M D, et al. 2003. Natural selection shaped regional mtDNA variation in humans. *Proc. Natl. Acad. Sci. U. S. A.* 100:171-176.
- [0196] Nakao Y, Kanamori T, Itoh T, Kodama Y, Rainieri S, Nakamura N, Shimonaga T, Hattori M, Ashikari T. 2009. Genome sequence of the lager brewing yeast, an interspecies hybrid. *DNA Res.* 16:115-129.
- [0197] Naseeb S, James S A, Alsammar H, Michaels C J, Gini B, Nueno-Palop C, Bond C J, McGhie H, Roberts I N, Delneri D. 2017. *Saccharomyces jurei* sp. nov., isolation and genetic identification of a novel yeast species from *Quercus robur*. *Int. J. Syst. Evol. Microbiol.* 67:2046-2052.
- [0198] Nguyen H V and Boekhout T. 2017. Characterization of *Saccharomyces uvarum* (Beijerinck, 1898) and related hybrids: assessment of molecular markers that predict the parent and hybrid genomes and a proposal to name yeast hybrids. *FEMS Yeast Res.* 17(2).
- [0199] Nikulin J, Krogerus K, Gibson B. 2018. Alternative *Saccharomyces* interspecies hybrid combinations and their potential for low-temperature wort fermentation. *Yeast* 35:113-127.
- [0200] Okuno M, Kajitani R, Ryusui R, Morimoto H, Kodama Y, Itoh T. 2016. Next-generation sequencing analysis of lager brewing yeast strains reveals the evolutionary history of interspecies hybridization. *DNA Res.* 23:67-80.
- [0201] Paget C M, Schwartz J-M, Delneri D. 2014. Environmental systems biology of cold-tolerant phenotype in *Saccharomyces* species adapted to grow at different temperatures. *Mol. Ecol.* 23:5241-5257.
- [0202] Paliwal S, Fiumera A C, Fiumera H L. 2014. Mitochondrial-nuclear epistasis contributes to phenotypic variation and coadaptation in natural isolates of *Saccharomyces cerevisiae*. *Genetics* 198:1251-1265.
- [0203] Peris D, Arias A, Orlic S, Belloch C, Pérez-Través L, Querol A, Barrio E. 2017. Mitochondrial introgression suggests extensive ancestral hybridization events among *Saccharomyces* species. *Mol. Phylogenet. Evol.* 108:49-60.
- [0204] Peris D, Belloch C, Lopandić K, Alvarez-Pérez J M, Querol A, Barrio E. 2012. The molecular characterization of new types of *Saccharomyces cerevisiae* x *S. kudriavzevii* hybrid yeasts unveils a high genetic diversity. *Yeast* 29:81-91.
- [0205] Peris D, Langdon Q K, Moriarty R V., Sylvester K, Bontrager M, Charron G, Leducq J-B, Landry C R, Libkind D, Hittinger C T. 2016. Complex Ancestries of Lager-Brewing Hybrids Were Shaped by Standing Variation in the Wild Yeast *Saccharomyces eubayanus*. *PLOS Genet.* 12:e1006155.
- [0206] Peris D, Lopes C A, Belloch C, Querol A, Barrio E. 2012. Comparative genomics among *Saccharomyces cerevisiae* x *Saccharomyces kudriavzevii* natural hybrid strains isolated from wine and beer reveals different origins. *BMC Genomics* 13:407.
- [0207] Peris D, Pérez-Torrado R, Hittinger C T, Barrio E, Querol A. 2018. On the origins and industrial applications of *Saccharomyces cerevisiae* x *Saccharomyces kudriavzevii* hybrids. *Yeast* 35:51-69.
- [0208] Peris D, Sylvester K, Libkind D, Gonçalves P, Sampaio J P, Alexander W G, Hittinger C T. 2014. Population structure and reticulate evolution of *Saccharomyces eubayanus* and its lager-brewing hybrids. *Mol. Ecol.* 23:2031-2045.
- [0209] Pichaud N, Ballard J W O, Tanguay R M, Blier P U. 2013. Mitochondrial haplotype divergences affect specific temperature sensitivity of mitochondrial respiration. *J. Bioenerg. Biomembr.* 45:25-35.
- [0210] Quintela M, Johansson M P, Kristjánsson B K, Barreiro R, Laurila A. 2014. AFLPs and Mitochondrial Haplotypes Reveal Local Adaptation to Extreme Thermal Environments in a Freshwater Gastropod. *PLoS One* 9:e101821.
- [0211] R Development Core Team R. 2017. R: A Language and Environment for Statistical Computing.
- [0212] Salvadó Z, Arroyo-López F N, Guillamón J M, Salazar G, Querol A, Barrio E. 2011. Temperature adaptation markedly determines evolution within the genus *Saccharomyces*. *Appl. Environ. Microbiol.* 77:2292-2302.
- [0213] Schu G. 2009. Energy. In: Esslinger H M, editor. *Handbook of brewing processes, technology, markets.* Weinheim (Germany). p. 648-650.
- [0214] Silva G, Lima F P, Martel P, Castilho R. 2014. Thermal adaptation and clinal mitochondrial DNA variation of European anchovy. *Proceedings. Biol. Sci.* 281: 20141093-20141093.
- [0215] Špírek M, Poláková S, Jatzová K, Sulo P. 2014. Post-zygotic sterility and cytonuclear compatibility limits in *S. cerevisiae* xenomitochondrial cybrids. *Front. Genet.* 5:454.
- [0216] Sylvester K, Wang Q-M, James B, Mendez R, Hulfachor A B, Hittinger C T. 2015. Temperature and host preferences drive the diversification of *Saccharomyces* and other yeasts: a survey and the discovery of eight new yeast species. *FEMS Yeast Res.* 15:fov002.
- [0217] Tenge C. 2009. Yeast. In: Esslinger H M, editor. *Handbook of brewing processes, technology, markets.* Weinheim (Germany): Wiley-VCH. p. 120, 123.
- [0218] Thorsness P E, Fox T D. 1993. Nuclear mutations in *Saccharomyces cerevisiae* that affect the escape of DNA from mitochondria to the nucleus. *Genetics* 134:21-28.
- [0219] Unger R W. 2004. *Beer in the Middle Ages and the Renaissance.* Philadelphia, Pa.: University of Pennsylvania Press

- [0220] Walther A, Hesselbart A, Wendland J. 2014. Genome Sequence of *Saccharomyces carlsbergensis*, the World's First Pure Culture Lager Yeast. *G3* 4:783-793.
- [0221] Willett C S. 2011. The nature of interactions that contribute to postzygotic reproductive isolation in hybrid copepods. *Genetica* 139:575-588.
- [0222] Wolff J N, Pichaud N, Camus M F, Côté G, Blier P U, Dowling D K. 2016. Evolutionary implications of mitochondrial genetic variation: mitochondrial genetic effects on OXPHOS respiration and mitochondrial quantity change with age and sex in fruit flies. *J. Evol. Biol.* 29:736-747.
- [0223] Wolters J F, Charron G, Gaspary A, Landry C R, Fiumera A C, Fiumera H L. 2018. Mitochondrial Recombination Reveals Mito-Mito Epistasis in Yeast. *Genetics* 209:307-319.
- [0224] Yamagishi H, Ohnuki S, Nogami S, Ogata T, Ohya Y. 2010. Role of bottom-fermenting brewer's yeast KEX2 in high temperature resistance and poor proliferation at low temperatures. *J. Gen. Appl. Microbiol.* 56:297-312.
- [0225] Zweifel S G, Fangman W L. 1991. A nuclear mutation reversing a biased transmission of yeast mitochondrial DNA. *Genetics* 128:241-249.

---

 SEQUENCE LISTING
 

---

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oHECPB15 primer

<400> SEQUENCE: 1

aatacatata acttactttt tcaaaattaa ttacatact agatctgttt agcttgctt 60

<210> SEQ ID NO 2

<211> LENGTH: 60

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oHECPB16 primer

<400> SEQUENCE: 2

tctatataga caacaaccac ttccactagc cttaagcga gctcgtttc gacactggat 60

<210> SEQ ID NO 3

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oHWA565 primer

<400> SEQUENCE: 3

ctatggttta cgaaatgatc cacg 24

<210> SEQ ID NO 4

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oHWA566 primer

<400> SEQUENCE: 4

cactgaccca gtcttgctt c 21

<210> SEQ ID NO 5

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: oHWA568 primer

<400> SEQUENCE: 5



-continued

---

tttgcaaadc gaagacccat 20

<210> SEQ ID NO 6  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHMB3 primer

<400> SEQUENCE: 6

gtttctggcc gagctacaag 20

<210> SEQ ID NO 7  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHMB4 primer

<400> SEQUENCE: 7

caaggccatg tcttctcggt 20

<210> SEQ ID NO 8  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHECPB148 primer

<400> SEQUENCE: 8

tttgaatadc aatgaaaatg cc 22

<210> SEQ ID NO 9  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHECPB149 primer

<400> SEQUENCE: 9

ttaaaaccta taatacacat atatattgc 29

<210> SEQ ID NO 10  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHDP25 primer

<400> SEQUENCE: 10

tgcgccaagt gctgaagaa caactggga 29

<210> SEQ ID NO 11  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: oHDP26 primer

<400> SEQUENCE: 11

gcgatttcaa tctggttatt atacaacatc at 32

<210> SEQ ID NO 12

-continued

---

<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: ITS1 primer

<400> SEQUENCE: 12

tccgtaggtg aacctgcgg

19

<210> SEQ ID NO 13  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: ITS4 primer

<400> SEQUENCE: 13

tcctccgctt attgatatgc

20

<210> SEQ ID NO 14  
<211> LENGTH: 586  
<212> TYPE: PRT  
<213> ORGANISM: *Saccharomyces cerevisiae*  
<220> FEATURE:  
<221> NAME/KEY: MISC\_FEATURE  
<222> LOCATION: (1)..(586)  
<223> OTHER INFORMATION: HO protein

<400> SEQUENCE: 14

Met Leu Ser Glu Asn Thr Thr Ile Leu Met Ala Asn Gly Glu Ile Lys  
1 5 10 15  
Asp Ile Ala Asn Val Thr Ala Asn Ser Tyr Val Met Cys Ala Asp Gly  
20 25 30  
Ser Ala Ala Arg Val Ile Asn Val Thr Gln Gly Tyr Gln Lys Ile Tyr  
35 40 45  
Asn Ile Gln Gln Lys Thr Lys His Arg Ala Phe Glu Gly Glu Pro Gly  
50 55 60  
Arg Leu Asp Pro Arg Arg Arg Thr Val Tyr Gln Arg Leu Ala Leu Gln  
65 70 75 80  
Cys Thr Ala Gly His Lys Leu Ser Val Arg Val Pro Thr Lys Pro Leu  
85 90 95  
Leu Glu Lys Ser Gly Arg Asn Ala Thr Lys Tyr Lys Val Arg Trp Arg  
100 105 110  
Asn Leu Gln Gln Cys Gln Thr Leu Asp Gly Arg Ile Ile Ile Ile Pro  
115 120 125  
Lys Asn His His Lys Thr Phe Pro Met Thr Val Glu Gly Glu Phe Ala  
130 135 140  
Ala Lys Arg Phe Ile Glu Glu Met Glu Arg Ser Lys Gly Glu Tyr Phe  
145 150 155 160  
Asn Phe Asp Ile Glu Val Arg Asp Leu Asp Tyr Leu Asp Ala Gln Leu  
165 170 175  
Arg Ile Ser Ser Cys Ile Arg Phe Gly Pro Val Leu Ala Gly Asn Gly  
180 185 190  
Val Leu Ser Lys Phe Leu Thr Gly Arg Ser Asp Leu Val Thr Pro Ala  
195 200 205  
Val Lys Ser Met Ala Trp Met Leu Gly Leu Trp Leu Gly Asp Ser Thr  
210 215 220

-continued

Thr	Lys	Glu	Pro	Glu	Ile	Ser	Val	Asp	Ser	Leu	Asp	Pro	Lys	Leu	Met
225					230					235					240
Glu	Ser	Leu	Arg	Glu	Asn	Ala	Lys	Ile	Trp	Gly	Leu	Tyr	Leu	Thr	Val
			245						250					255	
Cys	Asp	Asp	His	Val	Pro	Leu	Arg	Ala	Lys	His	Val	Arg	Leu	His	Tyr
			260					265					270		
Gly	Asp	Gly	Pro	Asp	Glu	Asn	Arg	Lys	Thr	Arg	Asn	Leu	Arg	Lys	Asn
		275					280					285			
Asn	Pro	Phe	Trp	Lys	Ala	Val	Thr	Ile	Leu	Lys	Phe	Lys	Arg	Asp	Leu
	290					295					300				
Asp	Gly	Glu	Lys	Gln	Ile	Pro	Glu	Phe	Met	Tyr	Gly	Glu	His	Ile	Glu
305					310					315					320
Val	Arg	Glu	Ala	Phe	Leu	Ala	Gly	Leu	Ile	Asp	Ser	Asp	Gly	Tyr	Val
			325						330					335	
Val	Lys	Lys	Gly	Glu	Gly	Pro	Glu	Ser	Tyr	Lys	Ile	Ala	Ile	Gln	Thr
			340					345					350		
Val	Tyr	Ser	Ser	Ile	Met	Asp	Gly	Ile	Val	His	Ile	Ser	Arg	Ser	Leu
		355					360					365			
Gly	Met	Ser	Ala	Thr	Val	Thr	Thr	Arg	Ser	Ala	Arg	Glu	Glu	Ile	Ile
	370					375					380				
Glu	Gly	Arg	Lys	Val	Gln	Cys	Gln	Phe	Thr	Tyr	Asp	Cys	Asn	Val	Ala
385					390					395					400
Gly	Gly	Thr	Thr	Ser	Gln	Asn	Val	Leu	Ser	Tyr	Cys	Arg	Ser	Gly	His
				405					410					415	
Lys	Thr	Arg	Glu	Val	Pro	Pro	Ile	Ile	Lys	Arg	Glu	Pro	Val	Tyr	Phe
			420					425					430		
Ser	Phe	Thr	Asp	Asp	Phe	Gln	Gly	Glu	Ser	Thr	Val	Tyr	Gly	Leu	Thr
		435					440					445			
Ile	Glu	Gly	His	Lys	Asn	Phe	Leu	Leu	Gly	Asn	Lys	Ile	Glu	Val	Lys
	450					455					460				
Ser	Cys	Arg	Gly	Cys	Cys	Val	Gly	Glu	Gln	Leu	Lys	Ile	Ser	Gln	Lys
465					470					475					480
Lys	Asn	Leu	Lys	His	Cys	Val	Ala	Cys	Pro	Arg	Lys	Gly	Ile	Lys	Tyr
				485					490					495	
Phe	Tyr	Lys	Asp	Trp	Ser	Gly	Lys	Asn	Arg	Val	Cys	Ala	Arg	Cys	Tyr
			500					505					510		
Gly	Arg	Tyr	Lys	Phe	Ser	Gly	His	His	Cys	Ile	Asn	Cys	Lys	Tyr	Val
		515					520					525			
Pro	Glu	Ala	Arg	Glu	Val	Lys	Lys	Ala	Lys	Asp	Lys	Gly	Glu	Lys	Leu
	530					535					540				
Gly	Ile	Thr	Pro	Glu	Gly	Leu	Pro	Val	Lys	Gly	Pro	Glu	Cys	Ile	Lys
545					550					555					560
Cys	Gly	Gly	Ile	Leu	Gln	Phe	Asp	Ala	Val	Arg	Gly	Pro	His	Lys	Ser
				565					570					575	
Cys	Gly	Asn	Asn	Ala	Gly	Ala	Arg	Ile	Cys						
				580				585							

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 1758

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Saccharomyces cerevisiae

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: misc\_feature

-continued

&lt;222&gt; LOCATION: (1) .. (1758)

&lt;223&gt; OTHER INFORMATION: HO polynucleotide

&lt;400&gt; SEQUENCE: 15

```

atgcttttctg aaaacacgac tattctgatg gctaacgggtg aaattaaaga catcgcaaac      60
gtcacggcta actettacgt tatgtgcgca gatggctcgg ctgcccgcgt cataaatgtc      120
acacagggct atcagaaaat ctataatata cagcaaaaaa ccaaacacag agcttttgaa      180
ggatgaacctg gtatggttaga tcccaggcgt agaacagttt atcagcgtct tgcattacaa      240
tgtactgcag gtcataaatt gtcagtcagg gtccctacca aacctactgt ggaaaaaagt      300
ggtagaaaatg ccaccaaata taaagtgaga tggagaaatc tgcagcaatg tcagacgctt      360
gatggttaga taataataat tccaaaaaac catcataaga cattoccaat gacagttgaa      420
ggtagagtttg ccgcaaaacg cttcatagaa gaaatggagc gctctaaagg agaatatctc      480
aactttgaca ttgaagttag agatttgatg tatcttgatg ctcaattgag aattttctagc      540
tgcataagat ttggtccagt actcgcagga aatggtggtt tatctaaatt tctcactgga      600
cgtagtgaac ttgtaactcc tgcgtgtaaa agtatggctt ggatgcttgg tctgtgggta      660
ggtagacagta caacaaaaga gccagaaatc tcagtagata gcttgatcc taagctaag      720
gagagtttaa gagaaaatgc gaaaatctgg ggtctctacc ttacggtttg tgacgatcac      780
gttccgctac gtgcaaaaca tgtaaggctt cattatggag atggtccaga tgaaaacagg      840
aagacaagga atttgaggaa aaataatcca ttctggaaag ctgtcacaat tttaaagttt      900
aaaagggatc ttgatggaga gaagcaaatc cctgaattta tgtacggcga gcatatagaa      960
gttctgtaag cattcttagc cggcttgatc gactcagatg ggtacgttgt gaaaaagggc     1020
gaagccctg aatcttataa aatagcaatt caaactgttt attcatccat tatggacgga     1080
attgtccata tttcaagatc tcttggtatg tcagctactg tgacgaccag gtcagctagg     1140
gaggaaatca ttgaaggaag aaaagtccaa tgtcaattta catacgactg taatgttgct     1200
gggggaacaa cttcacagaa tgttttgtca tattgtcgaa gtggtcacaa aacaagagaa     1260
gttccgcaa tttataaaaag ggaaccgta tatttcagct tcacggatga tttccagggt     1320
gagagtactg tatatgggct tacgatagaa ggccataaaa atttcttgct tggcaacaaa     1380
atagaagtga aatcatgtcg aggctgctgt gtgggagaac agcttaaaat atcacaaaaa     1440
aagaatctaa aacactgtgt tgcttgctcc agaaaggga tcaagtatct ttataaagat     1500
tggagtggta aaaatcgagt atgtgctaga tgctatggaa gatacaaat cagcgggtcat     1560
cactgtataa attgcaagta tgtaccagaa gcacgtgaag tgaaaaaggc aaaagacaaa     1620
ggcgaataat tgggcattac gcccgaaagg ttgccagta aaggaccaga gtgtataaaa     1680
tgtggcggaa tcttacagtt tgatgctgtc cgcgggcctc ataagagttg tggtataaac     1740
gcaggtgcgc gcatctgc                                     1758

```

1. A method of making a hybrid yeast strain having a selected mitotype comprising:

treating a first yeast strain with a mitochondria elimination agent to produce a first mitochondria-null yeast strain, and

mating the first mitochondria-null yeast strain with a second yeast strain comprising mitochondria to produce the yeast strain having the selected mitotype.

2. The method of claim 1, wherein the mitochondria elimination agent is ethidium bromide.

3. The method of claim 1, wherein the first yeast strain and the second yeast strain are different yeast species.

4. The method of claim 1, wherein the first yeast strain and the second yeast strain are haploid.

5. (canceled)

6. The method of claim 1, wherein the first yeast strain and the second yeast strain are from the genus *Saccharomyces*.

7. The method of claim 6, wherein the first yeast strain and/or the second yeast strain are selected from the group consisting of *Saccharomyces cerevisiae*, *Saccharomyces paradoxus*, *Saccharomyces mikatae*, *Saccharomyces arboricola*, *Saccharomyces jurei*, *Saccharomyces kudriavzevii*, *Saccharomyces uvarum*, and *Saccharomyces eubayanus*.

8. (canceled)

9. The method of claim 6, wherein the first yeast strain is *Saccharomyces cerevisiae* and the second yeast strain is *Saccharomyces eubayanus*.

10. The method of claim 6, wherein the first yeast strain is *Saccharomyces eubayanus* and the second yeast strain is *Saccharomyces cerevisiae*.

11. A method of making a yeast strain with an altered mitotype comprising:

treating a first yeast strain with a mitochondria elimination agent to produce a first mitochondria-null yeast strain, and

mating the first mitochondria-null yeast strain with a second karyogamy-deficient yeast strain comprising mitochondrial DNA (mtDNA) to produce the yeast strain having the altered mitotype, wherein the first yeast strain is polyploid.

12. The method of claim 11, wherein the first mitochondria-null yeast strain comprises a HyPr plasmid.

13. The method of claim 11, wherein the mitochondria elimination agent is ethidium bromide.

14. The method of claim 11, wherein the second karyogamy-deficient yeast strain comprises a *kar1-1* mutation.

15. The method of claim 11, wherein the mtDNA is from a donor yeast strain.

16. The method of claim 11, wherein the first yeast strain is a *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid.

17. The method of claim 16, wherein the first yeast strain is a lager brewing strain.

18. (canceled)

19. The method of claim 11, wherein the second karyogamy-deficient yeast strain and/or the donor yeast strain is a *Saccharomyces cerevisiae* strain.

20. (canceled)

21. A yeast strain made by the method of claim 1.

22. A *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid comprising mitochondria DNA (mtDNA), wherein at least 90% of the mtDNA is from *Saccharomyces cerevisiae*.

23. (canceled)

24. (canceled)

25. A *Saccharomyces cerevisiae* x *Saccharomyces eubayanus* hybrid comprising mitochondrial DNA (mtDNA), wherein at least 90% of the mtDNA is from *Saccharomyces eubayanus*.

26. (canceled)

27. (canceled)

28. A method for making a fermentation product comprising culturing the yeast strain of claim 21 with a fermentable substrate and at a temperature to produce the fermentation product.

29. (canceled)

30. (canceled)

31. (canceled)

32. (canceled)

33. (canceled)

\* \* \* \* \*