



US 20210071186A1

(19) **United States**(12) **Patent Application Publication** (10) **Pub. No.: US 2021/0071186 A1**
(43) **Pub. Date: Mar. 11, 2021**
Petersen et al.(54) **METHODS OF GENE EDITING AND
TRANSFORMING CANNABIS**(71) Applicant: **Wisconsin Alumni Research
Foundation**, Madison, WI (US)(72) Inventors: **Michael W. Petersen**, Merrimac, WI
(US); **Edward James Williams**,
Madison, WI (US); **Robert Harnish**,
Middleton, WI (US); **Heidi Flewelling**
Kaepler, Oregon, WI (US); **Brian**
Martinell, Mt. Horeb, WI (US); **Ray**
Collier, Middleton, WI (US); **Frank**
McFarland, Madison, WI (US); **Shawn**
Michael Kaepler, Oregon, WI (US)(73) Assignee: **Wisconsin Alumni Research
Foundation**, Madison, WI (US)(21) Appl. No.: **16/930,936**(22) Filed: **Jul. 16, 2020****Related U.S. Application Data**(60) Provisional application No. 62/982,522, filed on Feb.
27, 2020, provisional application No. 62/906,210,
filed on Sep. 26, 2019, provisional application No.
62/875,311, filed on Jul. 17, 2019.**Publication Classification**(51) **Int. Cl.**
C12N 15/82 (2006.01)
C07K 14/415 (2006.01)
(52) **U.S. Cl.**
CPC **C12N 15/8201** (2013.01); **C07K 14/415**
(2013.01)(57) **ABSTRACT**Disclosed herein are methods for the production of *Canna-
bis* meristem explants from dry seeds. Also described are
methods of transforming and gene editing using the *Can-
nabis* meristem explants disclosed herein.**Specification includes a Sequence Listing.**

FIG. 1




FIG. 2



FIG. 3

May 9, 2019 Cannabis meristem transformation




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

Cannabis seed and meristem explants




FIG. 9

Cannabis meristem explants on B5 medium; post 4 day co-culture after inoculation with Ar18r12v / Dicot Binary 19




FIG. 10

Transient GUS expression in *Cannabis* meristem explants transformed with
Ar18r12v / Dicot Binary 19




FIG. 11

Transient GUS expression in *Cannabis* meristem explants transformed with
Ar18r12v / Dicot Binary 19 (explants de-stained in 70% EtOH)




FIG. 12

Non-inoculated *Cannabis* seed sanitized in 20% Clorox on BS media (left)
Non-inoculated meristem explants on BS media (right)
• POC of viability of *Cannabis* seeds, meristem explants for transformation



FIG. 13

Spectinomycin sensitive (bleaching) phenotype visible in inoculated *Cannabis* meristem explants on 150 mg/L spectinomycin B5

- POC of use of *aadA* / spectinomycin resistance as a selectable marker in *Cannabis*
- Imaged 12 April 2019






FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18



FIG. 20



FIG. 21





FIG. 23






FIG. 25






FIG. 26

FIG. 27




FIG. 28




FIG. 29



FIG. 30



FIG. 31



FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37



FIG. 38



FIG. 39






FIG. 40




FIG. 41




FIG. 42



FIG. 43



FIG. 44



FIG. 45




FIG. 46




FIG. 47




FIG. 48




FIG. 49




FIG. 50




FIG. 51






FIG. 53



FIG. 54



Ln 1 = Low molecular weight ladder
 Ln 2 = blank
 Ln 3 = 50:50 reagent negative control
 Ln 4 = Cannabis leaf negative control
 Ln 5 = blank
 Ln 6 = T0 Cannabis WP001181-1 leaf 1
 Ln 7 = T0 Cannabis WP001181-1 leaf 2
 Ln 8 = T0 Cannabis WP001181-1 leaf 3
 Ln 9 = T0 Cannabis WP001181-1 leaf 4
 Ln 10 = T0 Cannabis WP001181-1 leaf 5
 Ln 11 = T0 Cannabis WP001181-1 leaf 6
 Ln 12 = T0 Cannabis WP001181-1 leaf 7
 Ln 13 = T0 Cannabis WP001181-1 leaf 8
 Ln 14 = T0 Cannabis WP001181-1 leaf 9
 Ln 15 = T0 Cannabis WP001181-1 leaf 10
 Ln 16 = blank
 Ln 17 = 2 ng plasmid DNA
 Ln 18 = blank
 Ln 19 = Low molecular weight ladder

FIG. 55




FIG. 56




FIG. 57




FIG. 58



FIG. 59




FIG. 60





FIG. 61




FIG. 62




FIG. 63




FIG. 64




FIG. 64 CONTINUED

[illegible]

FIG. 65

	(1)	1	10	20	30	40	50	60	70	80	90	100	110	120	130	140	150	160	
CBDA synthase (active)	(1)																		
CBDA synthase (Genbank XM_030624886.1)	(1)																		
CBDA Synthase (ccbr Fhola)	(1)																		
CBDA Synthase (Genbank KJ469374.1)	(1)																		
CBDA synthase (inactive)	(1)																		
Consensus	(1)	ARGNAGTCTCAACATTCCTCTTGGPTTGTTCGACGATAATATTTCTTTCTCAATATCAAACTTCATTCTCAATCTCGAGAAACCTTCCTTMAATGCTTCCTGCAATATATTCCTCAATATGCAACAACTCAAACTCTGATACACTC																	
	(159)	159	170	180	190	200	210	220	230	240	250	260	270	280	290	300	310	321	
CBDA synthase (active)	(159)																		
CBDA synthase (Genbank XM_030624886.1)	(159)																		
CBDA Synthase (ccbr Fhola)	(159)																		
CBDA Synthase (Genbank KJ469374.1)	(159)																		
CBDA synthase (inactive)	(155)																		
Consensus	(159)	CACGCAAAACAACCCATTGTATATGCTGCTCTTAAATTCGACAAATGACAGATCTTAGATTAGGCTCGACCAACCCCAAAACCACTCTTATCTGTCACCTCTTCACAGTGTCTCTATATCAAGGCACTATTCTATGCTCCAGAAAGTTGGCTTCAGACTT																	
	(319)	319	330	340	350	360	370	380	390	400	410	420	430	440	450	460	470	481	
CBDA synthase (active)	(319)																		
CBDA synthase (Genbank XM_030624886.1)	(319)																		
CBDA Synthase (ccbr Fhola)	(319)																		
CBDA Synthase (Genbank KJ469374.1)	(319)																		
CBDA synthase (inactive)	(314)																		
Consensus	(319)	ATTGCAACTCGAAGTGTGGTCATGATTCTGAGGCATGCTCATACATCTCAASTGCCATTCTTTATAGTGAATTGAGAAACATCCCTCAATCAAATAGATGTTTCATAGCAAACTCATGGGTGAAGCGAGCTATCTTCGAGAGTTTATATT																	
	(479)	479	490	500	510	520	530	540	550	560	570	580	590	600	610	620	630	641	
CBDA synthase (active)	(479)																		
CBDA synthase (Genbank XM_030624886.1)	(479)																		
CBDA Synthase (ccbr Fhola)	(479)																		
CBDA Synthase (Genbank KJ469374.1)	(479)																		
CBDA synthase (inactive)	(474)																		
Consensus	(479)	ATTGCGTTTAATGAGAAATGAGAACTCTTAGTTTGGTGTGAGTATGGCTACCTGTTTACCGAGTGGACACTTTGGTGGAGAGGCTATGAGCACTTGAATGAGAAAGCTATGGCTTCGCGTGTGATAATATCATTGATGACACTTAGTCAAACTTCATGG																	
	(641)	641	650	660	670	680	690	700	710	720	730	740	750	760	770	780	790	803	
CBDA synthase (active)	(641)																		
CBDA synthase (Genbank XM_030624886.1)	(641)																		
CBDA Synthase (ccbr Fhola)	(641)																		
CBDA Synthase (Genbank KJ469374.1)	(641)																		
CBDA synthase (inactive)	(636)																		
Consensus	(641)	GAAAGAGTCTAGATCGAABATCTATGCGGGAAGATCTCTTTGGGCTTTACGTGGTGGTGGAGCAGAACTCTCGGAATCTGTAGACATGAAABATTAGACTGGTTGCTGTGCCAAAGTCTACTATGTTTGTGTTTAAAGATCATGAGATACATGAT																	
	(800)	800	810	820	830	840	850	860	870	880	890	900	910	920	930	940	950	962	
CBDA synthase (active)	(797)																		
CBDA synthase (Genbank XM_030624886.1)	(797)																		
CBDA Synthase (ccbr Fhola)	(797)																		
CBDA Synthase (Genbank KJ469374.1)	(797)																		
CBDA synthase (inactive)	(795)																		
Consensus	(800)	ATGAGTGTGTCAGCTTAGTTAACAATGCAAAATATTCCTCAACAGTATGAAAGATTTATTACATAGGACTACTTCAACTAGAACATTAAGATATCAAGGGAGAMTANGACAGCAATACCACTATCTCTCTTCAGTTTCTCTTGGTGGAT																	
	(961)	961	970	980	990	1000	1010	1020	1030	1040	1050	1060	1070	1080	1090	1100	1110	1123	
CBDA synthase (active)	(958)																		
CBDA synthase (Genbank XM_030624886.1)	(958)																		
CBDA Synthase (ccbr Fhola)	(958)																		
CBDA Synthase (Genbank KJ469374.1)	(958)																		
CBDA synthase (inactive)	(956)																		
Consensus	(961)	GTTGATAGTCTAGTGCATGATGAGCAAGACTTTTCTGAGTGGGTATTAAGAAACGAGTTGGCAGCAATTTGAGTGGATTTGATCTATCTTCTATAGTGGTGTGTAAATTAAGCAGCTGCAAAATTTTAACAGGAAATTTGCTTGTATGATCT																	

FIG. 65 CONTINUED




FIG. 66

BraLTP2 Original sequence: Type IIS recognition sequences: **BsaI**, **BpiI**, **BsmBI**

ATGGCGACAGGTTCTCGTGTTCGATCGGTCTAGCAATGATCCTCATAATCTCAGGAGAACTGCTAGTTCAGGGCAAGGA
ACGTGCCAAGGAGACATAGAGGGTCTGATGAGAGAATGTGCGGTCTACGTCCAGCGTCCAGGCCCAAAGGTAAACCCATC
CGCAGCGTGTGCAAAGTCGTCAAGAGATCAGACATCCCCTGCGCATGTGGCCGTATCACACCCTCGGTTCAAAAAATGAT
AGACATGAATAAGGTTGTTCTTGTCACCTTCCTTTGTGGGAGGCCTCTCGCTCATGGTACCAAGTGTGGAAGCTACATTGTG
CCATGA

BraLTP2 CDS1 STOP Part for synthesis: Type IIS recognition sequences: **BsaI**, **BpiI**, **BsmBI**; **START**; **STOP**; **XXXX**
= motif liberated by **BsaI** digestion during Golden Gate Level 1 assembly of Transcriptional Unit.

GGTCTCAAATGGCGACAGGTTCTCGTGTTCGATCGGTCTAGCAATGATCCTCATAATCTCAGGAGAACTGCTAGTTCAGG
GCAAGGAACGTGCCAAGGAGACATAGAGGGTCTGATGAGAGAATGTGCGGTCTACGTCCAGCGTCCAGGCCCAAAGGTA
AACCCATCCGCAGCGTGTGCAAAGTCGTCAAGAGATCAGACATCCCCTGCGCATGTGGCCGTATCACACCCTCGGTTCAAA
AAATGATAGACATGAATAAGGTTGTTCTTGTCACCTTCCTTTGTGGGAGGCCTCTCGCTCATGGTACCAAGTGTGGAAGCTA
CATTGTGCCATGAGCTTAGAGACC

BraLTP2 CDS1 ns Part for synthesis: Type IIS recognition sequences: **BsaI**, **BpiI**, **BsmBI**; **START**; **STOP**; **XXXX** =
motif liberated by **BsaI** digestion during Golden Gate Level 1 assembly of Transcriptional Unit; **AG** replaces STOP
(and produces a Serine-Serine linker)

GGTCTCAAATGGCGACAGGTTCTCGTGTTCGATCGGTCTAGCAATGATCCTCATAATCTCAGGAGAACTGCTAGTTCAGG
GCAAGGAACGTGCCAAGGAGACATAGAGGGTCTGATGAGAGAATGTGCGGTCTACGTCCAGCGTCCAGGCCCAAAGGTA
AACCCATCCGCAGCGTGTGCAAAGTCGTCAAGAGATCAGACATCCCCTGCGCATGTGGCCGTATCACACCCTCGGTTCAAA
AAATGATAGACATGAATAAGGTTGTTCTTGTCACCTTCCTTTGTGGGAGGCCTCTCGCTCATGGTACCAAGTGTGGAAGCTA
CATTGTGCCAATTCGAGAGACC

FIG. 67

CsPT1 Original sequence₁₋₁: Type IIS recognition sequences:   

ATGGGACTCTCATCAGTTTGTACCTTTTCATTTCAAACCTAATTACCATACTTTATTAAATCCTCACAATAATAATCCCAAAACCTCATTATTA
TGTTATCGACACCCCAAAACACCAATTAATACTCTTACAATAATTTTCCCTCTAAACATTGCTCCACCAAGAGTTTTCATCTACAAAACAA
ATGCTCAGAATCATTATCAATCGCAAAAAATTCATTAGGGCAGCTACTACAAATCAAACCTGAGCCTCCAGAATCTGATAATCATTGAGTA
GCAACTAAAATTTTAACTTTGGGAAGGCATGTTGGAAACTTCAAAGACCATATACAATCATAGCATTACTTTCATGCGCTTGTGGATTGT
TTGGGAAAGAGTTGTTGCATAACACAAATTTAATAAGTTGGTCTCTGATGTTCAAGGCATTCTTTTTTTGGTGGCTGTATTATGCATTGCT
TCTTTTACAACCTACCATCAATCAGATTACGATCTTCACATTGACAGAATAACAAGCCTGATCTACCACTAGCTTCAGGGGAAATATCAG
TAAACACAGCTTGGATTATGAGCATAATTGTGGCACTGTTTGGATTGATAATAACTATAAAAATGAAGGGTGGACCACTCTATATATTTG
GCTACTGTTTTGGTATTTTGGTGGGATTGTCTATTCTGTTCCACCATTTAGATGGAAGCAAAATCCTTCCACTGCATTCTTCTCAATTTCC
TGGCCCATATTATTACAAATTTACATTTTATTATGCCAGCAGAGCAGCTCTTGGCCTACCATTGAGTTGAGGCCTCTTTTACTTCTCTGCTG
TAGCATTATGAAATCAATGGGTTGAGCTTTGGCTTTAATCAAAGATGCTTCAGACGTTGAAGGCGACACTAAATTTGGCATATCAACCTT
GGCAAGTAAATATGGTTCCAGAACTTGACATTATTGTTCTGGAATTGTTCTCTATCTATGTGGCTGCTATACTTGTCTGGGATTATCT
GGCCCCAGGCTTTCAACAGTAACGTAATGTTACTTTCTCATGCAATCTTAGCATTTTGGTTAATCCTCCAGACTCGAGATTTTGGCTTAACA
AATTACGACCCGGAAGCAGGCAGAGAAGATTTACGAGTTCATGTGGAAGCTTTATTATGCTGAATATTTAGTATATGTTTTCATATAA

CsPT1 CD51 STOP Part for synthesis: Type IIS recognition sequences:      XXXX = motif liberated by BsaI digestion during Golden Gate Level 1 assembly of Transcriptional Unit.

GGTCTCAAATGGGACTCTCATCAGTTTGTACCTTTTCATTTCAAACCTAATTACCATACTTTATTAAATCCTCACAATAATAATCCCAAAACCT
CATTATTATGTTATCGACACCCCAAAACACCAATTAATACTCTTACAATAATTTTCCCTCTAAACATTGCTCCACCAAGAGTTTTCATCTAC
AAAACAAATGCTCAGAATCATTATCAATCGCAAAAAATTCATTAGGGCAGCTACTACAAATCAAACCTGAGCCTCCAGAATCTGATAATC
ATTCAGTAGCAACTAAAATTTTAACTTTGGGAAGGCATGTTGGAAACTTCAAAGACCATATACAATCATAGCATTACTTTCATGCGCTTG
TGGATTGTTTGGGAAAGAGTTGTTGCATAACACAAATTTAATAAGTTGGTCTCTGATGTTCAAGGCATTCTTTTTTTGGTGGCTGTATTA
TGCATTGCTCTTTTACAACCTACCATCAATCAGATTACGATCTTCACATTGACAGAATAACAAGCCTGATCTACCACTAGCTTCAGGGG
AAATATCAGTAAACACAGCTTGGATTATGAGCATAATTGTGGCACTGTTTGGATTGATAATAACTATAAAAATGAAGGGTGGACCACTCT
ATATATTTGGCTACTGTTTTGGTATTTTGGTGGGATTGTCTATTCTGTTCCACCATTTAGATGGAAGCAAAATCCTTCCACTGCATTCTTC
TCAATTTCTGGCCCATATTATTACAAATTTACATTTTATTATGCCAGCAGAGCAGCTCTTGGCCTACCATTGAGTTGAGGCCTCTCTTTA
CTTCTCTGCTAGCATTATGAAATCAATGGGTTGAGCTTTGGCTTTAATCAAAGATGCTTCAGACGTTGAAGGCGACACTAAATTTGGCAT
ATCAACCTTGCCAAGTAAATATGGTTCCAGAACTTGACATTATTTGTTCTGGAATTGTTCTCTATCTATGTGGCTGCTATACTTGTCTG
GGATTATCTGGCCCCAGGCTTTCAACAGTAACGTAATGTTACTTTCTCATGCAATCTTAGCATTTTGGTTAATCCTCCAGACTCGAGATTTT
GGCTTAACAAATTACGACCCGGAAGCAGGCAGAGAAGATTTACGAGTTCATGTGGAAGCTTTATTATGCTGAATATTTAGTATATGTTTTCA
TATAAGCTTAGAGACC

CsPT1 CD51 ns Part for synthesis: Type IIS recognition sequences:      XXXX = motif liberated by BsaI digestion during Golden Gate Level 1 assembly of Transcriptional Unit;  replaces STOP (and produces a Serine-Serine linker)

GGTCTCAAATGGGACTCTCATCAGTTTGTACCTTTTCATTTCAAACCTAATTACCATACTTTATTAAATCCTCACAATAATAATCCCAAAACCT
CATTATTATGTTATCGACACCCCAAAACACCAATTAATACTCTTACAATAATTTTCCCTCTAAACATTGCTCCACCAAGAGTTTTCATCTAC
AAAACAAATGCTCAGAATCATTATCAATCGCAAAAAATTCATTAGGGCAGCTACTACAAATCAAACCTGAGCCTCCAGAATCTGATAATC
ATTCAGTAGCAACTAAAATTTTAACTTTGGGAAGGCATGTTGGAAACTTCAAAGACCATATACAATCATAGCATTACTTTCATGCGCTTG
TGGATTGTTTGGGAAAGAGTTGTTGCATAACACAAATTTAATAAGTTGGTCTCTGATGTTCAAGGCATTCTTTTTTTGGTGGCTGTATTA
TGCATTGCTCTTTTACAACCTACCATCAATCAGATTACGATCTTCACATTGACAGAATAACAAGCCTGATCTACCACTAGCTTCAGGGG
AAATATCAGTAAACACAGCTTGGATTATGAGCATAATTGTGGCACTGTTTGGATTGATAATAACTATAAAAATGAAGGGTGGACCACTCT
ATATATTTGGCTACTGTTTTGGTATTTTGGTGGGATTGTCTATTCTGTTCCACCATTTAGATGGAAGCAAAATCCTTCCACTGCATTCTTC
TCAATTTCTGGCCCATATTATTACAAATTTACATTTTATTATGCCAGCAGAGCAGCTCTTGGCCTACCATTGAGTTGAGGCCTCTCTTTA
CTTCTCTGCTAGCATTATGAAATCAATGGGTTGAGCTTTGGCTTTAATCAAAGATGCTTCAGACGTTGAAGGCGACACTAAATTTGGCAT
ATCAACCTTGCCAAGTAAATATGGTTCCAGAACTTGACATTATTTGTTCTGGAATTGTTCTCTATCTATGTGGCTGCTATACTTGTCTG
GGATTATCTGGCCCCAGGCTTTCAACAGTAACGTAATGTTACTTTCTCATGCAATCTTAGCATTTTGGTTAATCCTCCAGACTCGAGATTTT
GGCTTAACAAATTACGACCCGGAAGCAGGCAGAGAAGATTTACGAGTTCATGTGGAAGCTTTATTATGCTGAATATTTAGTATATGTTTTCA
TATAAGCTTAGAGACC

FIG. 68

CsOMT1 CDS Original sequence: Type IIS recognition sequences: **BsaI**, **BsaI**, **BsmBI**

ATGGGTTCAACAGGAATAGAGACCCAAATGACCCCAACCCAAATATCCGACGAAGAAGCCAACTCTTCGCCATGCAATTAGCCAGTGC
CTCAGTCTTACCCATGGTTCTCAAAGCAGCTTTAGAGCTCGACCTCTTGGAGATCATAGCCAAGGCCGGTCCAGGCGGTTCTCTCACCT
TCCGACATAGCTCAACAGCTTCGACTCAGAACCCAGACGCCCGGTGATGCTGGACCGGATGCTGAGACTGTTGGCTAGCTACAAAGT
GGTGACGTAAGCTCGCTGCGTGAGCGT**GGG**CGGAAGAGGAAGGGAAGGTGGAGAGGCTTTATGGGTTGGCTCCGGTGAGTAAATA
TCTGACGAAGAATGAAGATGGAGTCTCCATTGCTCCTCTTGTCTCATGAACCAGGATAAGGTTCTTATGGAGAGTTGGTATCACTTAAA
AGATGCAGTACTTGATGGAGGAATACCTTTCAACAAGGCATATGGAATGACAGCATTGAATATCATGGAACCGATCAAAGGTTCAATA
AAATCTTTAATAGAGGAATGTCCGACCACTCGACTATTACCATGAAGAAATCTCGAACTTACAAGGGTTTCGAGGGTCTTAAGTGA
TTGTTGATGTTGGTGGTGGTACTGGAGCTGTTGTTAACATGAT**GGG**TAAGTACCTACTATTAAGGGTATTAACTTCGATTGGCTCA
TGTCATCGAAGATGCACCTCCATTGACCGGTGTAGAGCATGTTGGAGGAGACATGTTGTAAGTGTACCAAAAGGAGATGCAATTTTCAT
GAASTGGATTGGCATGATTGGAGCGATGAACACTGCTTGAAATCTTGAAGAACTGCCACGCTGCACCTGCCGAACACGGAAGAGTGA
TCGTGGCGGAGTGCAATCTTCGGGTGGCACCAGGACTCGAGCTTGCCACAAAGAGTACGGTCCACATTGATGTGATCATGTTGGCCATA
ACCTGGTGGCAAAGAGAGAGAACAGAGAAAGAGTTTGAGGCATTGGCTAAGGGAGCTGGCTTTAAAGGCTTCAAAGTCCATTGCAATGC
TTTCAATACCCATATCATGGAATTTCTCAAGACCATTTAA

CsOMT21 CDS1 STOP Part for synthesis: Type IIS recognition sequences: **BsaI**, **BsaI**, **BsmBI**, **START**, **STOP**, XXXX = motif liberated by BsaI digestion during Golden Gate Level 1 assembly of Transcriptional Unit; **GG** = mutations to eliminate internal Type IIS restriction endonuclease recognition motifs.

GGTCTCAA**GG**GGTTCAACAGGAATAGAGACCCAAATGACCCCAACCCAAATATCCGACGAAGAAGCCAACTCTTCGCCATGCAATTAG
CCAGTGCCCTCAGTCTTACCCATGGTTCTCAAAGCAGCTTTAGAGCTCGACCTCTTGGAGATCATAGCCAAGGCCGGTCCAGGCGGTTTC
TCTCACCTTCCGACATAGCTCAACAGCTTCGACTCAGAACCCAGACGCCCGGTGATGCTGGACCGGATGCTGAGACTGTTGGCTAGCT
ACAACTGGTGACGTAAGCTCGCTGCGTGAGCGT**GGG**CGGAAGAGGAAGGGAAGGTGGAGAGGCTTTATGGGTTGGCTCCGGTGGA
GTAAATATCTGACGAAGAATGAAGATGGAGTCTCCATTGCTCCTCTTGTCTCATGAACCAGGATAAGGTTCTTATGGAGAGTTGGTATC
ACTTAAAGATGCAGTACTTGATGGAGGAATACCTTTCAACAAGGCATATGGAATGACAGCATTGAATATCATGGAACCGATCAAAGG
TTCAATAAAATCTTTAATAGAGGAATGTCCGACCACTCGACTATTACCATGAAGAAATCTCGAACTTACAAGGGTTTCGAGGGTCTTA
ACTCGATTGTTGATGTTGGTGGTGGTACTGGAGCTGTTGTTAACATGAT**GGG**TAAGTACCTACTATTAAGGGTATTAACTTCGATT
GCCTCATGTATCGAAGATGCACCTCCATTGACCGGTGTAGAGCATGTTGGAGGAGACATGTTTGAAGTGTACCAAAAGGAGATGCAA
TTTTATGAAGTGGATTGGCATGATTGGAGCGATGAACACTGCTTGAAATCTTGAAGAACTGCCACGCTGCACCTGCCGAACACGGAA
AAGTGATCGTGGCGGAGTGCAATCTTCGGGTGGCACCAGGACTCGAGCTTGCCACAAAGAGTACGGTCCACATTGATGTGATCATGTTG
GCCATAACCTGGTGGCAAAGAGAGAGAACAGAGAAAGAGTTTGAGGCATTGGCTAAGGGAGCTGGCTTTAAAGGCTTCAAAGTCCATT
GCAATGCTTTCAATACCCATATCATGGAATTTCTCAAGACCATTT**TAAGCTTAGAGACC**

CsOMT1 CDS1 ns Part for synthesis: Type IIS recognition sequences: **BsaI**, **BsaI**, **BsmBI**, **START**, **STOP**, XXXX = motif liberated by BsaI digestion during Golden Gate Level 1 assembly of Transcriptional Unit; **GG** = mutations to eliminate internal Type IIS restriction endonuclease recognition motifs; **GG** replaces STOP (and produces a Serine-Serine linker)

GGTCTCAA**GG**GGTTCAACAGGAATAGAGACCCAAATGACCCCAACCCAAATATCCGACGAAGAAGCCAACTCTTCGCCATGCAATTAG
CCAGTGCCCTCAGTCTTACCCATGGTTCTCAAAGCAGCTTTAGAGCTCGACCTCTTGGAGATCATAGCCAAGGCCGGTCCAGGCGGTTTC
TCTCACCTTCCGACATAGCTCAACAGCTTCGACTCAGAACCCAGACGCCCGGTGATGCTGGACCGGATGCTGAGACTGTTGGCTAGCT
ACAACTGGTGACGTAAGCTCGCTGCGTGAGCGT**GGG**CGGAAGAGGAAGGGAAGGTGGAGAGGCTTTATGGGTTGGCTCCGGTGGA
GTAAATATCTGACGAAGAATGAAGATGGAGTCTCCATTGCTCCTCTTGTCTCATGAACCAGGATAAGGTTCTTATGGAGAGTTGGTATC
ACTTAAAGATGCAGTACTTGATGGAGGAATACCTTTCAACAAGGCATATGGAATGACAGCATTGAATATCATGGAACCGATCAAAGG
TTCAATAAAATCTTTAATAGAGGAATGTCCGACCACTCGACTATTACCATGAAGAAATCTCGAACTTACAAGGGTTTCGAGGGTCTTA
ACTCGATTGTTGATGTTGGTGGTGGTACTGGAGCTGTTGTTAACATGAT**GGG**TAAGTACCTACTATTAAGGGTATTAACTTCGATT
GCCTCATGTATCGAAGATGCACCTCCATTGACCGGTGTAGAGCATGTTGGAGGAGACATGTTTGAAGTGTACCAAAAGGAGATGCAA
TTTTATGAAGTGGATTGGCATGATTGGAGCGATGAACACTGCTTGAAATCTTGAAGAACTGCCACGCTGCACCTGCCGAACACGGAA
AAGTGATCGTGGCGGAGTGCAATCTTCGGGTGGCACCAGGACTCGAGCTTGCCACAAAGAGTACGGTCCACATTGATGTGATCATGTTG
GCCATAACCTGGTGGCAAAGAGAGAGAACAGAGAAAGAGTTTGAGGCATTGGCTAAGGGAGCTGGCTTTAAAGGCTTCAAAGTCCATT
GCAATGCTTTCAATACCCATATCATGGAATTTCTCAAGACCATTT**TTTGAGAGACC**

FIG. 69

CsPT3 CDS Original sequence: Type IIS recognition sequences:   

ATGGTGTCTCATCAGTTTGTAGTTTTCATCCTCCCTTGGAACTAATTTTAAATTAGTTCCTCGTAGTAATTTAAGGCATCATCTTCTCAT
TATCATGAAATAAATAATTTTATTAATAATAAACCAATTAATTTCTCATATTTTCTTCAAGACTATATTGCTCTGCCAAACCAATTGTACAC
AGAGAAAACAAATTCACAAAATCATTTTCACTCAGCCACCTCCAAAGGAAAAGCTCCATAAAGGCACATGGTGAAATTGAAGCTGATTGG
GAGTAATGGCACATCTGAATTTAATGTAATGAAAAGTGGAAACGCAATTTGGAGATTGTAAAGGCCATATGCAGCCAAGGGAGTATTGT
TTAACTCTGCTGCTATGTTTGCAAAAGAGTTGGTGGGGAACCTAAATCTATTTAGTTGGCCTTTGATGTTTAAAGATACTCTCTTTTACATTG
GTTATTTTATGCATTTTGTAAGTACAAGTGGCATCAATCAAATTTATGATCTCGACATCGACAGGTTAAACAAACCTAATTTGCCAGTAG
CATCAGGAGAAATTTCAAGTTGAATTGGCATGGTTGTTGACTATAGTTTGTACAATAAGTGGCCTCACATTAACAATTATAACGAACCTCAG
GGCCATTCTTCCCTTTTCTCTACTCTGCTAGTATCTTTTTGGCTTTCTCTATTCTGCTCCTCCATTGAGATGGAAGAAGAATCCTTTTACAGC
ATGTTTCTGTAATGTTATGTTGTATGTTGGCACAAGCGTTGGTGTCTATTATGCTTGTAAAGGCTAGTCTCGGGCTTCCAGCCAACCTGGAGC
CCTGCTTTTGTGCTCTTTTGGTTTATTTTATTGTTGAGTATACCATCTCCATTGCAAAAGATCTTTCAGACATAGAAGGTGACCGCAA
GTTTGGAAATCATAACCTTCTCAACTAAATTTGGAGCAAAACCATAGCATATATTTGTCATGGACTCATGCTTCTGAATTACGTGAGTGT
ATGGCTGCAGCTATTATTTGGCCACAGTTTTCACAGTAGCGTAATATTGCTTTCTCATGCATTATGGCAATTTGGGTATTATATCAGGC
TTGGATATTGGAGAAATCAAATTACGCCACG                   <

FIG. 70

Reference CsEPSPS sequence:

LOC115705599: TTGACGAAGTTCAACTTTTCCTTGGAAATGCTGGAAAGCAATGCGTCACTCACAGCTGC (SEQ ID NO:55)

LOC115705595: TTGACGAAGTTCAACTTTTCCTTGGAAATGCTGGAAAGCAATGCGTCACTCACAGCTGC (SEQ ID NO:55)

TTGACGAAGTTCAACTTTTCCTTGGAAATGCTGGAAAGCAATGCGTCACTCACAGCTGC (SEQ ID NO:55)

XXX= spacer; XXX= PAM; █ = editing position +1

* double-nucleotide mutation: CsEPSPS_9CtoT and 20CtoT (mutation from █ to T at position 9 and from █ to T at position 20)

pegRNA: CAACTTTTCCTTGGAAA █ -scaffold- TGAGTGAACGCATTGCTATTCCAGCATTTCGAAGGA (SEQ ID NO:56)

RT + PBS

CsEPSPS-gRNA1 as oligos:

TGAAGACTTTGCACAACCTTTTCCTTGGAAA █CGTTTAAGTCTTCT (Forward) (SEQ ID NO:41)

AGAAGACTTAAACGCATTTCGAAGGAAAAGTTGTGCAAAAGTCTTCA (Reverse) (SEQ ID NO:42)

CsEPSPS-Ext1 Part for synthesis: (PBS+RT flanked by BspMI sites):

ACCTGCTATAGTGCTGAGTGAACGCATTGCTATTCCAGCATTTCGAAGGAAACATATAGCAGGT (SEQ ID NO:43)

FIG. 71

Reference CsEPSPS sequence:

LOC115705599: CTTGGAGCAACAGTTGAGGAAGGACCGATTACTGCGTGATCACTCCACCAGAGAA (SEQ ID NO:57)

LOC115705595: CTTGGAGCAACAGTTGAGGAAGGACCGATTACTGCGTGATCACTCCACCAGAGAA (SEQ ID NO:57)

CTTGGAGCAACAGTTGAGGAAGGACCGATTACTGCGTGATCACTCCACCAGAGAA (SEQ ID NO:57)

nick
↓

XXX= spacer; XXX= PAM; █ = editing position +1

* double-nucleotide mutation: CsEPSPS_9-10CCtoTT (mutation from CC to TT at position 9-10)

pegRNA: CTTGGAGCAACAGTTGAGGA-scaffold- CACGCAGTAATCAAGTCCTTCCTCAACTGTTG (SEQ ID NO:58)

RT + PBS

CsEPSPS-gRNA2 as oligos:

TGAAGACTTTGCACTTGGAGCAACAGTTGAGGAGTTTAAGTCTTCT (Forward) (SEQ ID NO:44)

AGAAGACTTAAACTCCTCAACTGTTGCTCCAAGTGCAAAGTCTTCA (Reverse) (SEQ ID NO:45)

CsEPSPS-Ext2 Part for synthesis: (PBS+RT flanked by BspMI sites):

ACCTGCTATAGTGCCACGCAGTAATCAAGTCCTTCCTCAACTGTTGAACATATAGCAGGT (SEQ ID NO:46)

FIG. 72





FIG. 73



Ln 1 = Low molecular weight ladder
Ln 2 = blank
Ln 3 = 50:50 reagent negative control
Ln 4 = Cannabis leaf negative control
Ln 5 = blank
Ln 6 = TO Cannabis WP001181-1b leaf 1
Ln 7 = TO Cannabis WP001181-1b leaf 2
Ln 8 = TO Cannabis WP001181-1b leaf 3
Ln 9 = TO Cannabis WP001181-1b leaf 4
Ln 10 = TO Cannabis WP001181-1b leaf 5
Ln 11 = TO Cannabis WP001181-1b leaf 6
Ln 12 = TO Cannabis WP001181-1b leaf 7
Ln 13 = TO Cannabis WP001181-1b leaf 8
Ln 14 = TO Cannabis WP001181-1b leaf 9
Ln 15 = TO Cannabis WP001181-1b leaf 10
Ln 16 = blank
Ln 17 = 2 ng plasmid DNA
Ln 18 = blank
Ln 19 = Low molecular weight ladder

FIG. 74




FIG. 75



FIG. 76




FIG. 77




FIG. 78



FIG. 79





FIG. 80



METHODS OF GENE EDITING AND TRANSFORMING CANNABIS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Application No. 62/875,311, filed Jul. 17, 2019, U.S. Provisional Application No. 62/906,210, filed Sep. 26, 2019, and U.S. Provisional Application No. 62/982,522, filed Feb. 27, 2020, each of which is incorporated herein by reference in their entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0002] Not Applicable.

SEQUENCE LISTING

[0003] A Sequence Listing accompanies this application and is submitted as an ASCII text file of the sequence listing named "960296_04047_ST25.txt" which is 95.4 KB in size and was created on Nov. 23, 2020. The sequence listing is electronically submitted via EFS-Web on Nov. 23, 2020 and is incorporated herein by reference in its entirety.

BACKGROUND

[0004] A key hurdle in the transformation of plants is the responsiveness of cells in tissue culture to produce embryogenic cells that go on to form clonal, intact and fertile plants. Many species and varieties of plants are recalcitrant to this type of regeneration, including *Cannabis sativa* and *Cannabis indica*. Therefore, a need in the art exists for improved *Cannabis* transformation and gene editing methods.

SUMMARY OF THE INVENTION

[0005] In a first aspect, provided herein is a method of preparing an explant, the method comprising the steps of, rehydrating a dry *Cannabis* seed in a hydration medium, and excising meristematic tissue from the rehydrated *Cannabis* seed to form an explant. In some embodiments, the hydration medium comprises one or more priming agents. In some embodiments, the seed is a *Cannabis sativa* seed.

[0006] In some embodiments, the explant preparation method additionally comprises the step of surface sterilizing the *Cannabis* seed prior to rehydration. In some embodiments, the seed is surface sterilized using bleach.

[0007] In some embodiments, the explant preparation method additionally comprises the step of drying the explant. In some embodiments, the dried explant is capable of being stored for at least 10 days. In some embodiments, the explant is dried in the presence of one or more transformation supplements. In some embodiments, the transformation supplement is selected from the group consisting of a small molecule, a nucleic acid, a polypeptide, a protein, an antibody, a transcription factor, a biological macromolecule, a nanoparticle, and a liposome.

[0008] In some embodiments, the explant preparation method additionally comprise the step of transforming the explant with a heterologous nucleic acid of interest. In some embodiments, the explant is transformed using *Agrobacterium* mediated transformation or particle bombardment.

[0009] In a second aspect, provided herein is a *Cannabis* explant generated by the explant preparation methods described herein.

[0010] In a third aspect, provided herein is a dried *Cannabis* explant generated by the methods described herein.

[0011] In a forth aspect, provided herein is a method of transforming *Cannabis* with a heterologous nucleic acid, the method comprising the steps of rehydrating a dry *Cannabis* seed in a hydration medium, excising meristematic tissue from the rehydrated *Cannabis* seed to form a *Cannabis* explant, incubating the *Cannabis* explant in a pretreatment medium, inoculating the *Cannabis* explant with *Agrobacterium* spp. comprising the heterologous nucleic acid, co-culturing the *Cannabis* explant in a co-culture medium for between about 1 day and about 6 days, and culturing the *Cannabis* explant on a selection medium to select for transformed *Cannabis* explants.

[0012] In some embodiments, the method of transforming *Cannabis* additionally comprises the step of force treating the *Cannabis* explant prior to or following inoculation. In some embodiments, the force treatment is selected from the group consisting of sonication, vortexing, centrifugation, heat-shock, and addition of chemicals.

[0013] In some embodiments, the *Cannabis* transformation method additionally comprises the step of surface sterilizing the *Cannabis* seed prior to rehydration.

[0014] In some embodiments, the heterologous nucleic acid modulates the expression or activity of an endogenous *Cannabis* gene selected from the group consisting of tetrahydrocannabinolic acid synthase (THCA synthase), cannabidiolic acid synthase (CBDA synthase), O-methyltransferase (CsOMT21), lipid transfer protein 2 (LTP2), prenyltransferase 3 (CsPT3), and prenyltransferase 1 (CsPT1). In some embodiments, the heterologous nucleic acid encodes a polypeptide at least 90% identical to SEQ ID NO:28. In some embodiments, the heterologous nucleic acid encodes a guide RNA that targets *Cannabis sativa* THCA synthase gene, *Cannabis sativa* CBDA synthase gene, or *Cannabis sativa* EPSP synthase gene.

[0015] In some embodiments, the *Cannabis* seed is a *Cannabis sativa* seed. In some embodiments, the *Cannabis* seed is a seed from a *Cannabis* plant with less than 0.3 percent THC based on dry weight.

[0016] In a fifth aspect, provided herein is a transformed *Cannabis* explant produced by the methods described herein.

[0017] In a sixth aspect, provided herein is a *Cannabis* plant grown from the *Cannabis* explant generated by the methods described herein.

[0018] In a seventh aspect, provided herein is a method of producing a transformed *Cannabis* seed, the method comprising contacting a female *Cannabis* flower with an *Agrobacterium* spp. culture, wherein the *Agrobacterium* comprises a heterologous nucleic acid and pollinating the contacted female *Cannabis* flower with male pollen from a suitable donor plant, whereby a transformed *Cannabis* seed is produced. In some embodiments, the *Agrobacterium* spp. culture comprises sucrose and a wetting agent. In some embodiments, the *Agrobacterium* comprises a vector comprising the heterologous nucleic acid.

[0019] In an eighth aspect, provided herein is a method of transforming a *Cannabis* plant, the method comprising growing a sanitized and imbibed *Cannabis* seed on a non-selective culture medium suitable for supporting the growth

and survival of the *Cannabis* seed until a *Cannabis* explant is formed, inoculating the *Cannabis* explant with a heterologous nucleic acid, co-culturing the *Cannabis* explant in a co-culture medium for between about 1 day and about 6 days, and culturing the *Cannabis* explant on a selection medium to select for transformed *Cannabis* explants. In some embodiments, the explant is selected from the group consisting of a leaf explant, a node explant, an internode explant, a petiole explant, a hypocotyl explant, and a bud explant. In some embodiments, the *Cannabis* explant is inoculated using particle bombardment, high velocity microprojection, microinjection, electroporation, direct DNA uptake, cell-penetrating peptides, silica carbide fibers, nanoparticles, and bacterially-mediated transformation. In some embodiments, *Agrobacterium* spp. is used to inoculate the *Cannabis* explant.

[0020] In some embodiments, the method additionally comprises the step of force treating the *Cannabis* explant prior to or following inoculation. In some embodiments, the force treatment is selected from the group consisting of sonication, vortexing, centrifugation, heat-shock, and addition of chemicals.

[0021] In some embodiments, the heterologous nucleic acid modulates the expression or activity of an endogenous *Cannabis* gene selected from the group consisting of tetrahydrocannabinolic acid synthase (THCA synthase), cannabidiolic acid synthase (CBDA synthase), O-methyltransferase (CsOMT21), lipid transfer protein 2 (LTP2), prenyltransferase 3 (CsPT3), and prenyltransferase 1 (CsPT1). In some embodiments, the heterologous nucleic acid encodes a polypeptide at least 90% identical to SEQ ID NO:28. In some embodiments, the heterologous nucleic acid encodes a guide RNA that targets *Cannabis sativa* THCA synthase gene, *Cannabis sativa* CBDA synthase gene, or *Cannabis sativa* EPSP synthase gene.

[0022] In some embodiments, the *Cannabis* seed is a *Cannabis sativa* seed.

BRIEF DESCRIPTION OF DRAWINGS

[0023] The patent or patent application file contains at least one drawing in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

[0024] FIG. 1 shows stable RFP expression in *Cannabis* plantlet from meristem transformation experiments outlined in Example 1.

[0025] FIG. 2 shows a non-transgenic *Cannabis* plant derived from meristem explant in tissue culture.

[0026] FIG. 3 shows transient GUS expression in *Cannabis* meristem explants varieties Honey Gold 3WS and Fiber Hemp inoculated with Ar18r12v/DICOTBINARY-19.

[0027] FIG. 4 shows the plasmid map of the DICOTBINARY-22 vector.

[0028] FIG. 5 shows flower phenotypes post inoculum application (1 day).

[0029] FIG. 6 shows GUS transient expression in *Cannabis* leaves, petioles, internodes, and nodes inoculated with Ar18r12v/DICOTBINARY-19.

[0030] FIG. 7 shows mature *Cannabis* seed and embryo.

[0031] FIG. 8 shows *Cannabis* seeds and meristem explants.

[0032] FIG. 9 shows *Cannabis* meristem explants on B5 medium and after 4 days in co-culture after inoculation with Ar18r12v/DICOTBINARY-19.

[0033] FIG. 10 shows transient GUS expression in *Cannabis* meristem explants transformed with Ar18r12v/DICOTBINARY-19.

[0034] FIG. 11 shows transient GUS expression in *Cannabis* meristem explants transformed with Ar18r12v/DICOTBINARY-19 and de-stained in 70% EtOH.

[0035] FIG. 12 shows a non-inoculated *Cannabis* seed sanitized in 20% Clorox on B5 medium (left) and a non-inoculated meristem explant on B5 medium (right).

[0036] FIG. 13 shows spectinomycin sensitive (bleaching) phenotype visible in inoculated *Cannabis* meristem explants on 150 mg/L spectinomycin B5.

[0037] FIG. 14 shows sanitized *Cannabis* seeds and axenic plantlets on B5 medium after about 6 weeks.

[0038] FIG. 15 shows greening, regenerating *Cannabis* explants that were dried, stored, and regenerated (right) against freshly excised explants (left) that were plated on Hemp Node medium (Table 7) the day of excision. Explants were imaged approximately 3 weeks after excision.

[0039] FIG. 16 shows stable GUS expression in a plantlet derived from the treatment Hemp 3/22-3B described in Example 1. One of the leaves in a plantlet transferred to non-selective BRM expressed GUS in leaf stably.

[0040] FIG. 17 shows *Cannabis* seeds imbibed at 37° C.

[0041] FIG. 18 shows embryonic parts produced by mechanical excision.

[0042] FIG. 19 shows the plasmid map of control binary construct SOYTEST-2.

[0043] FIG. 20 shows positive GUS transients in *Cannabis* meristem explants (3WS variety) using SOYTEST-2 in both the Ar18r12v (left) and GV3101 (right) strains of *Agrobacterium*, demonstrating transfection of *Cannabis* meristems using disarmed strains of both *Agrobacterium rhizogenes* (Ar18r12v) and *Agrobacterium tumefaciens* (GV3101).

[0044] FIG. 21 additional stable RFP expressing *Cannabis* (Plant WP421-1) from experiments in the Honey Gold 3WS variety. The plantlet in the center is rooting on 50 mg/L streptomycin hemp node media (after being on 50 mg/L spectinomycin hemp node media for approximately 1 month).

[0045] FIG. 22 shows the plasmid map of the DICOTBOMB-13 vector.

[0046] FIG. 23 shows GUS transient expression in *Cannabis* meristem explants bombarded with DICOTBOMB-13.

[0047] FIG. 24 shows pollen germination results.

[0048] FIG. 25 shows pollen germination results. T0=initial germination after 1 hr treatment; T1=germination after 4 hr treatment; T2=germination after overnight treatment (22 h).

[0049] FIG. 26 shows the phenotype of plant WP421-1 on the day of transfer from tissue culture to the greenhouse (right) and after approximately 4 weeks in the greenhouse (left).

[0050] FIG. 27 shows stable RFP and GUS expression in leaves of T0 *Cannabis* plant WP421-1 derived from meristem transformation.

[0051] FIG. 28 shows stable GUS expression of leaves of T0 *Cannabis* plant WP421-1 derived from meristem transformation.

[0052] FIG. 29 shows *Cannabis* VAEs after 5 weeks (left) compared to freshly excised *Cannabis* meristem explants (right).

[0053] FIG. 30 shows machine excised *Cannabis* meristem explants imaged after approximately two weeks on non-selective B5 medium.

[0054] FIG. 31 shows transgenic T0 plant WP421-1 after 7 weeks in the greenhouse.

[0055] FIG. 32 shows GUS expression in leaf samples from WP421-1. Nine out of the ten leaf samples showed positive GUS expression confirming minimal chimerism.

[0056] FIG. 33 shows the PCR amplification scheme for PCR of leaf samples from WP421-1. A 156 bp fragment within the aadA1a expression cassette of DICOTBINARY-19 was amplified using primers designated F56 and R11. The fragments and the resulting amplicon is highlighted in blue.

[0057] FIG. 34 shows the results of PCR amplification of a 156 bp fragment of the aadA1a expression cassette. All 10 leaf samples were positive.

[0058] FIG. 35 shows bleached *Cannabis* meristem explants after 1 month on 100 mg/L spectinomycin node medium. RFP negative *Cannabis* meristem explants that were previously greening on 10-50 mg/L spectinomycin were transferred to hemp node medium containing 100 mg/L spectinomycin.

[0059] FIG. 36 shows GUS expression in vascular tissue (petiole sections) of *Cannabis* T0 event WP421-1 (“Candice”).

[0060] FIG. 37 shows pollination of WP421-1 by 3WS wild-type plant.

[0061] FIG. 38 shows flower of WP421-1 pollinated by 3WS wild-type plant.

[0062] FIG. 39 shows cuttings of WP421-1.

[0063] FIG. 40 shows T1 seedlings of WP421-1 and RFP expression in T1 seedling WP421-1@2 “Carly.”

[0064] FIG. 41 shows germline confirmation of *Cannabis* meristem transformation through RFP, GUS expression, and aadA1a PCR.

[0065] FIG. 42 shows T1 *Cannabis* plants WP421-1@1 and WP421-1@2 after 2 months in greenhouse.

[0066] FIG. 43 shows T1 seed of WP421-1.

[0067] FIG. 44 shows stable RFP expression (tdTomato) in T1 seedlings of WP421-1 (germinated on germination paper).

[0068] FIG. 45 shows stable RFP expression (tdTomato) in T1 seedlings of WP421-1 (germinated in flats).

[0069] FIG. 46 shows stable RFP expression (tdTomato) in T1 seedlings of WP421-1 (germinated in flats).

[0070] FIG. 47 shows stable GUS expression in T1 seedlings of WP421-1 (germinated in flats).

[0071] FIG. 48 shows GUS expression 7-8 weeks post-inoculation in chimeric *Cannabis* shoots (3WS variety) regenerated on 75 mg/L spectinomycin.

[0072] FIG. 49 shows T0 *Cannabis* plant WP421-2 (Honey Gold 3WS+DICOTBINARY-19).

[0073] FIG. 50 shows *Cannabis* meristem explants targeted on carboxymethylcellulose media for particle bombardment.

[0074] FIG. 51 shows phenotype and stable GUS expression in transgenic 3WS *Cannabis* derived from particle-mediated transformation of meristem explants (imaged approximately 2 months post-blast).

[0075] FIG. 52 shows phenotype and stable GUS expression in T0 transgenic 3WS *Cannabis* plant WP-001181-1a (“Fernanda”) derived from particle-mediated transformation of meristem explants.

[0076] FIG. 53 shows phenotype of T0 transgenic 3WS *Cannabis* plant WP-001181-1a “Fernanda” derived from particle-mediated transformation of meristem explants after approximately 3 weeks in greenhouse.

[0077] FIG. 54 shows *Cannabis* WP001181-1a particle gun T0 event aadA1a PCR.

[0078] FIG. 55 shows *Cannabis* WP001181-1a particle gun T0 event GUS expression.

[0079] FIG. 56 shows GUS expression in vascular tissue in petiole sections (and corresponding leaf) of *Cannabis* T0 particle gun event WP001181-1a.

[0080] FIG. 57 shows phenotype and stable GUS expression in T0 transgenic 3WS *Cannabis* plant WP-001181-1b (“Hernanda”) derived from particle-mediated transformation of meristem explants.

[0081] FIG. 58 shows GUS transient expression in automated-excised *Cannabis* meristem explants of Fiber Hemp variety post co-culture.

[0082] FIG. 59 shows metal rolling pin with glass plate; optional gap can be set by adding/removing shims (metal shims shown on left, paper on right).

[0083] FIG. 60 shows transient GUS expression in *Cannabis* (Abacus variety) meristem explants mechanically excised by crushing under a rolling pin, then stored for 2 months at -20 C (left image seed crushed wet and crushed material then dried; right image seed dried first then crushed).

[0084] FIG. 61 shows GUS transient expression in *Cannabis* meristem explants of variety 3WS and Abacus post co-culture.

[0085] FIG. 62 shows RFP (tdTomato) segregation in the T2 generation of transformed *Cannabis* (derived from T0 event WP421-1).

[0086] FIG. 63 shows a schematic of the gene networks underlying cannabinoid and terpenoid accumulation in *Cannabis*. Zager et al., Plant Physiology, 2019, 180(4):1877-1897.

[0087] FIG. 64 shows a sequence alignment of various THCA synthase cDNA sequences. From top to bottom, the sequences shown are: SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:2, SEQ ID NO:5, and SEQ ID NO:59.

[0088] FIG. 65 shows a sequence alignment of various CBDA synthase cDNA sequences. From top to bottom, the sequences shown are: SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15, SEQ ID NO:11, SEQ ID NO:17, and SEQ ID NO:11.

[0089] FIG. 66 shows embodiments of a cloning strategy for BraLTP2. From top to bottom, the sequences shown are: SEQ ID NO:29, SEQ ID NO:47, and SEQ ID NO:48.

[0090] FIG. 67 shows embodiments of a cloning strategy for CsPT1. From top to bottom, the sequences shown are: SEQ ID NO:31, SEQ ID NO:49, and SEQ ID NO:50.

[0091] FIG. 68 shows embodiments of a cloning strategy for CsOMT1. From top to bottom, the sequences shown are: SEQ ID NO:33, SEQ ID NO:51, and SEQ ID NO:52.

[0092] FIG. 69 shows embodiments of a cloning strategy for CsPT3. From top to bottom, the sequences shown are: SEQ ID NO:35, SEQ ID NO:53, and SEQ ID NO:54.

[0093] FIG. 70 shows an embodiment of the CsEPSPS prime editing mutation strategy as described herein.

[0094] FIG. 71 shows an embodiment of the CsEPSPS prime editing mutation strategy as described herein.

[0095] FIG. 72 shows the phenotype of T0 transgenic 3WS *Cannabis* plant WP-001181-1b (“Hernanda”) derived from particle-mediated transformation of meristem explants after transplant to large pot.

[0096] FIG. 73 shows *Cannabis* WP001181-1b particle gun T0 event aadA1a PCR.

[0097] FIG. 74 shows *Cannabis* WP001181-1b particle gun T0 event GUS expression.

[0098] FIG. 75 shows a *Cannabis* seed (Badger variety) surface sanitized followed by incubation at 37° C. without imbibition (note radical emergence in seed on left).

[0099] FIG. 76 shows the phenotype of *Cannabis* meristem explants of Badger variety after 3-day co-culture.

[0100] FIG. 77 shows transient GUS expression in *Cannabis* meristem explants of Badger variety after 3-day co-culture.

[0101] FIG. 78 shows stable RFP (tdTomato) expression in *Cannabis* meristem explants (variety Badger).

[0102] FIG. 79 shows a process diagram timeline for *Cannabis sativa* *Agrobacterium*-mediated transformation of embodiments described herein. Stage 1 shows *Cannabis* seeds. Stage 2 shows appearance of *Cannabis* meristem explant (mature embryo) derived from imbibed seed, the explant on left has cotyledonary tissue intact; explant on right has cotyledonary tissue removed. Stage 3 shows transient GUS activity in Honey Gold 3WS *Cannabis* explants after co-culture with *Agrobacterium*. Stage 4 shows appearance of *Cannabis* meristem explants after 4 day co-culture. Stage 5 shows phenotypes of *Cannabis* meristem explants after ~3 weeks on 50 mg/L spectinomycin; note incomplete bleaching. Stage 6 shows *Cannabis* T0 event after second selection on 50 mg/L streptomycin prior to handoff to greenhouse. Finally, Stage 7 shows *Cannabis* T0 event prior to transplant in greenhouse.

[0103] FIG. 80 shows a process diagram timeline for *Cannabis sativa* Particle-mediated transformation. Stage 1 shows *Cannabis* seeds. Stage 2 shows appearance of *Cannabis* meristem explant (mature embryo) derived from imbibed seed, the explant on left has cotyledonary tissue intact; explant on right has cotyledonary tissue removed. Stage 3 shows *Cannabis* embryo target on CMC targeting media. Stage 4 shows PDS-1000 Helium gun. Stage 5 shows transient GUS activity in Honey Gold 3WS *Cannabis* explants 1-day post-bombardment. Stage 6 shows phenotypes of *Cannabis* meristem explants 2 months post-bombardment. Stage 7 shows *Cannabis* T0 event prior to handoff to greenhouse. Finally, Stage 8 shows *Cannabis* T0 event prior to transplant in greenhouse.

INCORPORATION BY REFERENCE

[0104] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, and patent application was specifically and individually indicated to be incorporated by reference.

DETAILED DESCRIPTION OF THE INVENTION

[0105] The present disclosure describes methods for preparation and transformation of meristem explants from *Cannabis*. The meristem explant preparation methods

described herein allow for pretreatment of the tissues for higher explant transformation and longer explant storage following excision.

[0106] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. All definitions, as defined and used herein, should be understood to control over dictionary definitions, definitions in documents incorporated by reference, and/or ordinary meanings of the defined terms.

[0107] The indefinite articles “a” and “an,” as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to mean “at least one.”

[0108] The phrase “and/or,” as used herein in the specification and in the claims, should be understood to mean “either or both” of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with “and/or” should be construed in the same fashion, i.e., “one or more” of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the “and/or” clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to “A and/or B,” when used in conjunction with open-ended language such as “comprising” can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other than A); in yet another embodiment, to both A and B (optionally including other elements); etc.

[0109] As used herein in the specification and in the claims, “or” should be understood to have the same meaning as “and/or” as defined above. For example, when separating items in a list, “or” or “and/or” shall be interpreted as being inclusive, i.e., the inclusion of at least one, but also including more than one, of a number or list of elements, and, optionally, additional unlisted items. Only terms clearly indicated to the contrary, such as “only one of” or “exactly one of,” or, when used in the claims, “consisting of,” will refer to the inclusion of exactly one element of a number or list of elements. In general, the term “or” as used herein shall only be interpreted as indicating exclusive alternatives (i.e. “one or the other but not both”) when preceded by terms of exclusivity, such as “either,” “one of,” “only one of,” or “exactly one of.” “Consisting essentially of,” when used in the claims, shall have its ordinary meaning as used in the field of patent law.

[0110] As used herein, the terms “approximately” or “about” in reference to a number are generally taken to include numbers that fall within a range of 5% in either direction (greater than or less than) the number unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value). Where ranges are stated, the endpoints are included within the range unless otherwise stated or otherwise evident from the context.

[0111] Provided herein are methods for preparing an explant suitable for transformation from a *Cannabis* seed. Also provided herein are methods of transforming and gene editing *Cannabis* meristem explant tissue. The explants generated by the methods described herein exhibit higher transformation efficiency with a broader capacity to customize the transformation process via pretreatment of the meri-

stematic tissue used to generate the explants. The methods described herein allow for high scale production of storable explants for more effect transformation methods. As described in further detail below, the protocols described herein allow for targeted pretreatment of the meristematic tissues used in explant preparation at various stages and with various factors to improve explant storage and transformation efficiency.

[0112] As used herein, “embryo” refers to part of a seed, consisting of precursor tissues (meristematic tissues) for the leaves, stem, and root, as well as one or more cotyledons. Once the embryo begins to grow (germinate), it becomes a seedling plant.

[0113] As used herein, “meristem” or “meristematic tissue” refers to the portion of a seed that consists of undifferentiated cells, the meristematic cells, which differentiate to produce multiple plant structures including stem, roots, leaves, germline tissues and seeds. The meristematic cells are the targets for transformation to obtain transgenic plants.

[0114] As used herein, “explant” refers to the target material for transformation.

[0115] As used herein, “germline transformation” refers to the transformation of a gene of interest into cells that give rise to pollen or ovule thus into seed.

[0116] In a first aspect, provided herein is a method for preparing an explant from the meristematic tissue of a seed, where the method generally comprises the steps of drying the seed, surface sterilizing the seed, imbibing the seed until sufficiently hydrated, excising meristematic tissue from the hydrated seed to generate an explant, and optionally drying the excised meristematic tissue to generate the storable explant for transformation. The explants generated by the methods described herein are suitable for use in any transformation method known in the art.

[0117] The methods described herein also include one or more priming steps in which one or more priming agents are added to either the hydration medium during imbibing of the seed or to the explant as it is drying to generate a value added explant. As used herein, the term “value added explant” refers to an explant prepared by the methods described herein when a priming factor has been included in the hydration medium or a transformation supplement is included during drying of the explant.

[0118] The method includes a first step of drying a seed or acquiring a dried seed from which the explant will be generated. Preferably, a dry seed for use in the methods of the present invention will have a moisture content of between 1% and 25%. Ideally seeds are grown and harvested to achieve a viable embryo and are grown and harvested and cleaned to achieve blemish-free identity preserved seeds free of plant diseases and microbes that could interfere with sterile tissue culture. It may be desirable to treat the plants with fungicides and/or natural or synthetic plant regulators to improve embryo viability, embryo storage quality, seed coat intactness, seed vigor, percent germination cell response in tissue culture and transformation.

[0119] Seeds from which explants are to be prepared may be harvested from any *Cannabis* cultivar of interest. In some embodiments, the seed is from *Cannabis sativa*. In some embodiments, the seed is from *Cannabis indica*. In some embodiments, the seed is from a *Cannabis* variety developed from cross breeding of *Cannabis sativa* and *Cannabis indica*. In some embodiments, the seed is from a *Cannabis sativa* L. plant with less than 0.3 percent tetrahydrocannabi-

nol (THC) based on the dry weight. In some embodiments, the *Cannabis sativa* cultivar is selected from the group consisting of Elektra x Chardonnay, Honey Gold 3WS (also referred to in the art at 3W1), Abacus, and Fiber Hemp.

[0120] In some embodiments of the present invention, the dry seed is surface sterilized. Any means known in the art for surface sterilization can be used. Suitable methods for surface sterilization may include, but are not limited to, exposure of the seed surface to radiation, UV light, oxidizing gasses, heat, plasma, disinfecting solvents and agents. In some embodiments, the seed is surface sterilized with a chemical agent such as sodium hypochlorite. In some embodiments, the seed is surface sterilized with an antibacterial or antifungal agent. In some embodiments, the seed is surface sterilized with ethanol (e.g., 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% ethanol). In some embodiments, the seed is surface sterilized with Clorox™ bleach.

[0121] The dry seed, which in some embodiments has undergone surface sterilization, is imbibed under conditions that support hydration of the seed. The dry seed is hydrated in a hydration medium and for a time sufficient for the seed to reach a moisture content of between 30% and 70%. In some embodiments, the seed is hydrated for at least 12 hours. In some embodiments, the seed is hydrated between 2 and 24 hours.

[0122] The hydration medium used for hydration of the seed maybe any suitable sterile hydration medium known in the art that supports survival of the meristematic tissue in the seed. In some embodiments, the hydration medium is a modified sterile water and includes antibiotics or antifungals. In some embodiments, the hydration medium is a tissue culture medium that includes natural or synthetic plant growth regulators, plant tissue culture nutrients, a carbon source or a non-nutritive osmoregulator. In one embodiment, the hydration medium is bean germination medium, which includes the components outlined in Table 1 of Example 1.

[0123] In some embodiments of the invention, the hydration medium may optionally include one or more priming factors for pretreatment of the meristematic tissue. As used herein, “priming factor” or “priming agent” references to any molecule or substance included in the hydration medium which promotes survival and storage of the prepared explant or that promotes or increases the transformation efficiency of the prepared explant. Priming factors for use in the hydration medium of the present invention may include, but are not limited to, small molecules, biological molecules such as nucleic acids, polypeptides, proteins, antibodies, transcription factors, and macromolecules or complexes thereof, nanoparticles, liposomes, and cell-penetrating peptides. In some embodiments, the priming factor is a plant growth factor including, but not limited to, thidiazuron (TDZ), 6-benzylaminopurine (BAP), polyethylene glycol (PEG), 2,4-dichlorophenoxyacetic acid (2,4-D), Paczol™, gibberellic acid (GA3), indole-3-acetic acid (IAA), indole-3-butyric acid (IBA), 1-naphthaleneacetic acid (NAA), forchlorfenuron (CPPU), spectinomycin, streptomycin, glyphosate, glufosinate, bialophos, hygromycin, amikacin, tobramycin, imazapyr, dicamba, polyvinylpyrrolidone (PVP), polyvinylpolypyrrolidone (PVPP), salicylic acid, proline, betaine, ethylene, brassinosteroids, nitrates, and gibberellins.

[0124] Following hydration of the seed, meristematic tissue is excised to form an explant. Excision of the meristematic tissue may be performed by any means known in the art in which the seed coat and cotyledons are removed

from the seed. Suitable methods for the excision of the meristematic tissue may include, but are not limited to manual processing, wet milling using a series of rollers and spray nozzles, adjustable grinding plates, rods, knives, wheels and other mechanical or machine based excision methods. These may be composed of, but are not limited to, ceramics, metals, and synthetic polymers. Induced pressure, injected gasses, vacuum and turbulence are also suitable methods. Hydrated explants may be stored in suitable storage medium for up to 7 days. Suitable storage medium for the hydrated explants may be any medium that supports survival and competence of the explant tissue. In some embodiments, the explants are stored with an excess of liquid to explants by volume. In some embodiments, the storage medium is liquid medium including MS salts.

[0125] In some embodiments, the meristematic tissue is excised from a dry seed to form an explant without first imbibing the seed. A dry seed suitable for dry excision will typically have a moisture content of between 1% and 25%. Excision of the meristematic tissue from the dry seed may be performed by any means known in the art in which the seed coat and cotyledons are removed from the seed. Suitable methods for the excision of the meristematic tissue may include, but are not limited to manual processing, wet milling using a series of rollers and spray nozzles, adjustable grinding plates, rods, knives, wheels and other mechanical or machine based excision methods. These may be composed of, but are not limited to, ceramics, metals, and synthetic polymers. Induced pressure, injected gasses, vacuum and turbulence are also suitable methods.

[0126] Following excision, the explant may be dried. Desiccation of the explant may be performed by any means known in the art such that the moisture content of the dry explant is between 1% and 25%. Suitable methods for desiccating the explant may include, but are not limited to, drying in the presence of air with and without an added dehumidifying agent. In some embodiments, the explants are dried in a laminar flow hood. In some embodiments, the explants are dried on the surface of filter paper in a laminar flow hood for about 26 hours. In some embodiments, the explants are dried in a dehumidifier. In some embodiments, the explants are dried at a temperature between 0° C. and 35° C. for at least 5 hours (e.g., at least 5, 7, 9, 12, 15, 18, 24, 30, 36, 42, 48, 72, 96 or 120 hours) and up to 2 weeks (e.g., up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 days). In some embodiments, it may be beneficial to control rates of drying by tightly controlling temperature, humidity, air flow, and time.

[0127] During desiccation of the explant, one or more transformation supplements may be added. As used herein, “transformation supplement” refers to any molecule or substance added to the explant during desiccation, which promotes survival and storage of the prepared explant or that promotes or increases the transformation efficiency of the prepared explant. Transformation supplements for use during desiccation of the explant of the present invention may include small molecules, biological molecules such as nucleic acids, polypeptides, proteins, antibodies, transcription factors, and macromolecules or complexes thereof, nanoparticles, liposomes, *Agrobacterium*, *Rhizobium*, and cell-penetrating peptides. In some embodiments, the transformation supplement is a plant growth factor, cell protectant agent including, or other agent including, but not limited to, thidiazuron (TDZ), 6-benzylaminopurine (BAP),

polyethylene glycol (PEG), alginates and alginate complexes, starches, celluloses, synthetic polymers, gums, waxes, proline, betaine, polyvinylpyrrolidone (PVP), polyvinylpyrrolidone (PVPP), salicylic acid, calcium sources, silicone sources, colchicine, 2,4-dichlorophenoxyacetic acid (2,4-D), Paczol™, gibberellic acid (GA3), gibberellin (GA) pathway inhibitors, indole-3-acetic acid (IAA), indole-3-butyric acid (IBA), 1-naphthaleneacetic acid (NAA), forchlorfenuron (CPPU), spectinomycin, streptomycin, glyphosate, glufosinate, bialophos, hygromycin, amikacin, tobramycin, imazapyr, lyophilized *agrobacterium*, lyophilized *rhizobium*, and potassium hydroxide (KOH). In some embodiments, the transformation supplement is an agent, which promotes multiplication of the meristematic tissue, such as, but not limited to, TDZ, BAP, zeatin, kinetin, and CPPU. In some embodiments, explants are mechanically wounded prior to drying and storage. This can be achieved with exposure to ultrasound energy (e.g., sonication), liquid nitrogen, centrifugation, pressure, and chemical (ex. KOH, PEG, acids, bases), enzymes, abrasives, water jets, lasers, needles, or blades.

[0128] The dried explants are suitable for storage in a variety of conditions. Dried explants may be stored at temperatures ranging from about -200° C. to 50° C. (i.e., about -190° C. to 40° C., about -170° C. to 30° C., about -150° C. to 20° C., about -130° C. to 10° C., and about -102° C. to 0° C.) for a period of time of at least 7 days (i.e., at least 10 days, at least 30 days, at least 50 days, at least 60 days, at least 75 days, at least 90 days, and at least 120 days). Storable dried explants can also be banked to create libraries of germplasms from a variety of cultivars of agronomic significance. In general, lower temperature storage under dry conditions or controlled humidity will allow for prolonged storage of the dried Cannabis explants.

[0129] Explants generated by the methods described herein can be transformed with a heterologous gene or nucleic acid of interest by any means known in the art. Various methods have been developed for transferring genes or nucleic acids into plant tissue including particle bombardment, high velocity microprojection, microinjection, electroporation, direct DNA uptake, silica carbide fibers, cell-penetrating peptides, nanoparticles, viral vectors, and bacterially-mediated transformation. Bacteria known to mediate plant cell transformation include a number of species of the Rhizobiaceae, including, but not limited to, *Agrobacterium* spp., *Sinorhizobium* spp., *Mesorhizobium* spp., *Rhizobium* spp., *Ochrobacterium* spp., and *Bradyrhizobium* spp. In some embodiments, the explant is transformed using *Agrobacterium* spp. In some embodiments, the explant is transformed using *Agrobacterium* strain Ar18r12v. In some embodiments, the explant is transformed using particle bombardment using gold microcarriers. Suitable viral vectors are known and described in the art and may include, but are not limited to, tomato yellow leaf curl virus (TYLCV), tobacco yellow dwarf virus (TobYDV), tomato golden mosaic virus (TGMV), bean pod mottle virus (BPMV). Suitable methods of plant transformation are described in the art, such as, for example, by McCabe et al. (McCabe, D. E., Swain, W. F., Martinell, B. J., Christou, P. (1988) *Nature Biotechnology* 6(8), 923-926), Chen et al. (Chen, Y., Rivlin, A. Lange, A., Ye, X., Vaghchhipawala, Z., Eisinger, E., Dersch, E., Paris, M., Martinell, B., Wan, Y. (2014) *Plant Cell Reports* 33(1), 153-164), Ye et al. (Ye, X., Williams, E. J., Shen, J., Johnson, S., Lowe, B., Radke, S., Strickland, S.

Esser, J. A., Petersen, M. W., and Gilbertson, L. A. (2011) *Transgenic Research* 20(4), 773-7860), and *Plant Transformation Technologies* (Edited by C. Neal Stewart, Alisher Touraev, Vitaly Citovsky and Tzvi Tzfira ©2011 Blackwell Publishing Ltd. ISBN: 978-0-813-82195-5.)

[0130] Prior to inoculation, dried explants to be transformed may be rehydrated using suitable tissue culture medium. In some embodiments, the rehydration medium is Soy INO medium, VAE rehydration medium, or a solid medium such as Basal MS or Gamborg's B5 medium. In some embodiments, the rehydration step is combined with the pretreatment step described below.

[0131] Embryos may be pretreated prior to inoculation and transformation. In some embodiments, the embryos are pretreated with a polyethylene glycol (PEG) solution. In some embodiments, the PEG solution includes about 20% PEG4000. In some embodiments, the embryos are pretreated in the PEG-ethanol solution for between about 1 minute and about 3 hours (e.g., 1 min., 2 min., 5 min., 15 min., 30 min., 45 min., 1 hr., 1.5 hr., 2 hr., 2.5 hr., or 3 hr.). In some embodiments, salts may be added to the PEG solution. In some embodiments, the PEG solution includes one or more fungicides (e.g., Captan, Bravo, etc.). In some embodiments, the PEG solution includes Murashige and Skoog (MS) salts. In some embodiments, the pretreatment step includes sonication, vortexing, centrifugation, heat-shock, or addition of chemicals (e.g., TDZ, glyphosate, or metolachlor).

[0132] In some embodiments, the explant transformation is supplemented with force treatments. Force treatments may include, but are not limited to, sonication, vortexing, centrifugation, increased pressure, vacuum infiltration, heat-shock, dessication or addition of chemicals (e.g., TDZ, glyphosate, or metolachlor). Force treatments may be applied prior to or during transformation. For example, prior to or concurrently with inoculation with *Agrobacterium* or particle bombardment treatment. In some embodiments, explants are sonicated for about 20 seconds at about 45 kHz. Force treatment transformation methods are described in the art. See, for example, Khanna, et al. ("Centrifugation assisted *Agrobacterium tumefaciens*-mediated transformation (CAAT) of embryogenic cell suspensions of banana (*Musa* spp. Cavendish AAA and Lady finger AAA)," Molecular Breeding, 2004, 14:239-252), Kapila et al. ("An *Agrobacterium*-mediated transient gene expression system for intact leaves," Plant Science, 1997, 122(1):101-108), Trick et al. ("SAAT: sonication-assisted *Agrobacterium*-mediated transformation," Transgenic Research, 1997, 6(5): 329-336), and Hiei et al. ("Improved frequency of transformation in rice and maize by treatment of immature embryos with centrifugation and heat prior to infection with *Agrobacterium tumefaciens*," Plant Cell, Tissue, and Organ Culture, 2006, 87(3):233-243).

[0133] Following inoculation, the explants are co-cultured for between about 1 day and about 6 days in any suitable co-culture medium that supports the growth and survival of the inoculated explant. In some embodiments, the co-culture medium is WCIC INO medium, which includes the components outlined in Table 2 of Example 1.

[0134] Following transformation, the explants or transformed tissue is regenerated using a suitable regeneration medium that supports the growth and survival of at least the positively transformed explants or transformed tissues. Suitable regeneration medium are known and described in the art. For example, the regeneration medium may be B5

medium, WPM based medium, MS salts based medium, ½×MS salts based medium, or similar medium with or without supplementation. The regeneration medium may additionally comprise nystatin, TBZ, meta-topolin (mT), naphylacetic acid (NAA) and GA3. In some embodiments, the regeneration medium includes a selection agent to select for positive transformants. Examples of suitable selection agents include, but are not limited to, RFP, GUS, aadA1a, spectinomycin, streptomycin, and imazapyr.

[0135] The heterologous gene or nucleic acid of interest may be any gene or nucleic acid that may confer a particular desirable trait or phenotype in the transformed plant. Examples of suitable genes of agronomic interest envisioned by the present invention would include but are not limited to genes for disease, insect, or pest tolerance, herbicide tolerance, genes for quality improvements such as yield, nutritional enhancements, environmental or stress tolerances, or any desirable changes in plant physiology, growth, development, morphology or plant product(s) including starch production, modified oils production, high oil production, modified fatty acid content, high protein production, fruit ripening, enhanced animal and human nutrition, and biopolymers production. Also environmental stress resistance, pharmaceutical peptides and secreted peptides, improved processing traits, industrial enzyme production, improved flavor, nitrogen fixation, hybrid seed production, and fiber production. Any of these or other genetic elements, methods, and transgenes may be used with the invention as will be appreciated by those of skill in the art in view of the instant disclosure. The heterologous gene or nucleic acid of interest may also be a sequence that can affect a phenotype of interest by encoding an RNA molecule that causes the targeted inhibition of expression on an endogenous gene via gene silencing technologies.

[0136] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis* tetrahydrocannabinolic acid synthase (THCA synthase) gene. A heterologous nucleic acid is introduced into the *Cannabis* explant to knockout or silence the THCA synthase gene. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a tetrahydrocannabinol (THC) low or THC free phenotype. The sequence and activity of the THCA synthase gene is known and described in the art. See, for example, Laverty et al. (Laverty et al., "A physical and genetic map of *Cannabis sativa* identifies extensive rearrangements at the THC/CBD acid synthase loci," Genome Research, 2018, 29:146-156).

[0137] Provided below are five cDNA sequences (SEQ ID NOs:1-5) of THCA genes from various *Cannabis sativa* strains. Single nucleotide polymorphisms (SNPs) between the active THCA synthase of SEQ ID NO:1 and each of SEQ ID NOs:2-5 are indicated in each of the respective sequences and a full sequence alignment is shown in FIG. 64.

[0138] *Cannabis sativa* cultivar Skunk #1 tetrahydrocannabinolic acid synthase (THCA1) gene cDNA (SEQ ID NO:1), Genbank: KJ469378.1

```
ATGAATTGCTCAGCATTTTCCTTTTGTTGTTTGCAAATAATATT
TTTCTTTCTCTCATTCATATCCAAATTTCATAGCTAATCCTCGAG
AAAACCTTCCTTAAATGCTTCTCAAAACATATTCACCAATGTAGCA
AATCCAAACTCGTATACACTCAACACGACCAATTGTATATGTCTAT
```

-continued

CCTGAATTCGACAATACAAAATCTTAGATTCATCTCTGATACAACCC
 CAAAACCACTCGTTATTGTCACTCCTTCAAATAACTCCCATATCCAA
 GCAACTATTTTATGCTCTAAGAAAGTTGGCTTGCGAGATTCGAACTCG
 AAGCGGTGGCCATGATGCTGAGGGTATGCTACATATCTCAAGTCC
 CATTGTGTAGTAGACTTGAGAAACATGCATTCGATCAAAATAGAT
 GTTCATAGCCAAACTGCGTGGGTGAAGCCGGAGCTACCCCTTGAGAG
 AGTTTATTATTGGATCAATGAGAAGATGAGAATCTTAGTTTCTCG
 GTGGGTATTGCCCTACTGTTGGCGTAGGTGGACACTTTAGTGGAGGA
 GGCTATGGAGCATTGATGCGAAATTATGGCCTTGCGGCTGATAATAT
 TATTGATGCACACTTAGTCAATGTTGATGGAAGTCTTAGATCGAA
 AATCCATGGGAGAAGATCTGTTTGGGCTATACGTGGTGGTGGAGGA
 GAAAACCTTGGAAATCATGCGCATGGAAGTCAAACTGGTGTGATGT
 CCCATCAAAGTCTACTATATTAGTGTGTAAGAACATGGAGATAC
 ATGGGCTTGTCAAGTTATTTAACAAATGGCAAAATATTGCTTACAAG
 TATGACAAAGATTTAGTACTCATGACTCACTTCATAACAAGAATAT
 TACAGATAATCATGGGAAGAATAAGACTACAGTACATGGTTACTTCT
 CTTC AATTTTTCATGGTGGAGTGGATAGTCTAGTCGACTTGATGAAC
 AAGAGCTTTCTGAGTTGGGTATTAAGAACTGATGCAAGAAATT
 TAGCTGGATTGATACAACCATCTTCTACAGTGGTGTGTAATTTTA
 AACTGCTAATTTTAAAAAGGAAATTTTGCTTGATAGATCAGCTGGG
 AAGAAGACGGCTTTCTCAATTAAGTTAGTATGTTAAGAAACCAAT
 TCCAGAACTGCAATGGTCAAAATTTTGAAAAATTATATGAAGAAG
 ATGTAGGAGCTGGGATGTATGTGTTGTACCCTTACGGTGGTATAATG
 GAGGAGATTTGAGAATCAGCAATTCATTCCTCATCGAGCTGGAAT
 AATGTATGAACCTTTGGTACACTGCTTCTGGGAGAAGCAAGAAGATA
 ATGAAAAGCATATAAACTGGGTTGGAAGTGTGTTATAATTTTACGACT
 CCTTATGTGTCCCAAAATCCAAGATTGGCGTATCTCAATTATAGGGA
 CCTTGATTAGGAAAACTAATCATGCGAGTCTAATAATTACACAC
 AAGCACGTATTTGGGGTGAAAAGTATTTTGGTAAAAATTTTAACAGG
 TTAGTTAAGGTGAAAACATAAGTTGATCCCAATAATTTTGTAGAAA
 CGAACAAAGTATCCACCTCTTCCACCGCATCATATTAA

[0139] *Cannabis sativa* isolate 60D2 tetrahydrocannabinolic acid synthase (THCA) gene (SEQ ID NO:2), GenBank: MG996415.1, single nucleotide polymorph (SNP) relative to SEQ ID NO:1 shown in bold underline.

atgaattgctcagcattttctcttttggtttggttgcataaatatt
 tttctttctctcattccatatacaaatctcaatagctaactcctcgag
 aaaacttccttaaatgcttctcaaaacatattcccaacaatgtagca
 aatccaaaactcgatatacactcaacacgaccaattgtatatgtctat

-continued

cctgaattcgacaatacaaaaatcttagattcatctctgatacaaccc
 caaaaccactcggtattgtcactccttcaaaataactcccatatccaa
 gcaactattttatgctctaagaaagtggcctgcagattcgaactcg
 aagcggtggccatgatgctgagggatgtcctacatatctcaagtcc
 catttggtgtagtagacttgagaacatgcattcgatcaaaatagat
 gtccatagccaaactgcgtgggttgaagccggagctacccttgagga
 agtttattattggatcaatgagaagaatgagaatcttagtttctctg
 gtgggtattgcctactgttggcgtaggtggacacttttagtggagga
 ggctatggagcattgatgcgaattatggccttgcggctgataatat
 tattgatgcacacttagtcaatgttgatggaaaagtcttagatcgaa
 aatccatgggagaagatctgttttgggctatacgtgggtggaggga
 gaaaactttggaatcattgcagcatggaaaatcaaaactgggtggtgt
 cccatcaaaagtctactatattcagtggttaaaaagaacatggagatac
 atgggcttgtcaagttatttaacaaatggcaaaatattgcttacaag
 tatgacaaagatttagtactcatgactcacttcataacaaagaatat
 tacagataatcatgggaagaataagactacagtagcatggttacttct
 ctccaattttctcaggtggagtggtatagtcagacttgatgaac
 aagagctttctgtagttgggtattaaaaaactgattgcaagaatt
 tagctggattgatacaaccatcttctacagtggtgtgtgtaatttta
 acactgctaatttttaaaaaggaaattttgcttgatagatcagctggg
 aagaagacggctttctcaattaagttagactatgttaagaaccaat
 tccgaaaactgcaatgggtcaaaattttggaaaaattatagaagaag
 atgtaggagctgggagtgatgtgtgtacccttacgggtgtataatg
 gaggagatttcagaatcagcaattccatccctcatcgagctggaat
 aatgtatgaactttggtacactgcttctgggagaagcaagaagata
 atgaaaagcatataaaactgggttcgaagtgtttataattttacgact
 ccttatgtgtcccaaaatccaagattggcgtatctcaattatagggga
 ccttgatttaggaaaaactaatcatgcgagtcctaataattacacac
 aagcacgtatttgggggtgaaaagtattttggtaaaaattttaacagg
 ttagttaagggtgaaaactaaagttgatcccaataatttttttagaaa
 cgaacaaagtatcccacctcttmcaccgcatcatcattaa

[0140] *Cannabis sativa* clone ABC67 THCA synthase gene (SEQ ID NO:3), GenBank: KT876047.1, SNPs relative to SEQ ID NO: shown in bold underline.

atgaattgctcagcattttctcttttggtttggttgcataaatatt
 tttctttctctcattccatatacaaatctcaatagctaactcctcgag
 aaaacttccttaaatgcttctcaaaacatattcccaacaatgtagca
 aatccaaaactcgatatacactcaacacgaccaattgtatatgtctat
 cctgaattcgacaatacaaaaatcttagattcatctctgatacaaccc

-continued

caaaaccactcggttattgtcactccttcaaataactcccatatccaa
gcaactattttatgctctaagaaagtggcttgagattcgaaactcg
aagcgggtggccatgatgctgagggatgtcctacatatctcaagtcc
catttggtgtagtagacttgagaaacatgcattcgatcaaaatagat
gttcatagccaaactgcgtgggtgaagccggagctacccttgaga
agtttattattggatcaatgagaagaatgagaatcttagtttcttg
gtgggtattgcctactgttggcgtaggtggacactttagtggagga
ggctatggagcattgatgcaaaattatggccttgcggtgataatat
tattgatgcacacttagtcaatgttgatggaaaagtcttagatcgaa
aatccatgggagaagatctgttttgggctatacgtggtggtggagga
gaaaactttggaatcattgcagcatggaaaatcaaaactggttgcgt
cccatcaaagtctactatattcagtggttaaaagaacatggagatac
atgggcttgtcaagttatttaacaaatggcaaaatattgcttacaag
tatgacaaagatttagtactcatgactcacttcatacaaaagaatat
tacagataatcatgggaagaataagactacagtacatggttacttct
cttcaattttcatgggtggagtggatagctagtcgacttgatgaac
aagagccttctgtagtgggtattaaaaaactgattgcaagaatt
tagctggattgatacaaccatcttctacagtggtgttgtaatttta
acactgctaattttaaaaaggaaatttggcttgatagatcagctggg
aagaagacggctttctcaattaagttagactatgttaagaaccaat
tccagaaactgcaatggtcaaaatttggaaaaatttatgaagaag
atgtaggagctgggatgtatgtgtgtacccttacgggtggtataatg
gaggagatttcagaatcagcaattccattccctcatcgagctggaat
aatgtatgaactttggtacactgcttctgggagaagcaagaagata
atgaaaagcatataaaactgggttcgaagtggttataattttacgact
ccttatgtgtcccaaaatccaagattggcgtatctcaattataggga
ccttgatttaggaaaaactaatcatgcgagtcctaataattacacac
aagcacgtatttggggtgaaaagtatttggtaaaaaattttaacagg
ttagttaagggtgaaaaactaaagttgatcccaataatttttttagaaa
cgaacaaagtatccacaccttccaccgcatcatcat

[0141] THCA synthase *Cannabis sativa* Purple Kush
(SEQ ID NO:4), SNPs relative to SEQ ID NO:1 shown in
bold underline.

ATGAATTGCTCAGCATTTTCCTTTTGGTTTGGTTGCAAAATAATAT
TTTTCTTTCTCTCATTCATATCCAAATTTCAATAGCTAATCCTCG
AGAAAACCTTCCTTAAATGCTTCTCAAAACATATTCCCAACAATGTA
GCAAATCCAAAACCTGATACACTCAACACGACCAATTGTATATGT
CTATCCTGAATTGACAATACAAAATCTTAGATTCTCTCTGATAC

-continued

AACCCCAAAACCACTCGTTATTGTCACTCCTTCAAATAACTCCCAT
ATCCAAGCAACTATTTTATGCTCTAAGAAAGTTGGCTTGAGATTCT
GAACTCGAAGCGGTGGCCATGATGCTGAGGGTATGTCTACATATC
TCAAGTCCCATTTGTTGTAGTAGACTTGAGAAACATGCATTTCGATC
AAAATAGATGTTTCATAGCCAACTGCGTGGGTGAAGCCGGAGCTA
CCCTTGGAGAAGTTTATTATTGGATCAATGAGAAGATGAGAATCT
TAGTTTTCTCGGTGGGTATTGCCCTACTGTTGGCGTAGGTGGACAC
TTTAGTGAGGAGGCTATGGAGCATTGATGCGAAATTATGGCCTTG
CGGCTGATAATATCATTGATGCACACTTAGTCAATGTTGATGGAAA
AGTTCTAGATCGAAAATCCATGGGAGAAGATCTGTTTTGGGCTATA
CGTGGTGGTGAGGAGAGAAAATTTGGAATCATTGCAGCATGGAAAA
TCAAACCTGGTTGCTGTCCCATCAAAGTCTACTATATTAGTGTAA
AAAGAACATGGAGATACATGGGCTTGTCAAGTTATTTAACAAATGG
CAAAATATTGCTTACAAGTATGACAAAGATTTAGTACTCATGACTC
ACTTCATAACAAAGAATATTACAGATAATCATGGGAAGAATAAGAC
TACAGTACATGGTTACTTCTCTTCAATTTTTCATGGTGGAGTGGAT
AGTCTAGTCGACTTGATGAACAAGAGCTTTCGTGAGTTGGGTATTA
AAAAAACTGATTGCAAGAATTAGCTGGATTGATACAACCATCTT
CTACAGTGGTGTGTAAATTACAACACTGCTAATTTAAAAAGGAA
ATTTTGTCTGATAGATCAGCTGGGAAGAAGACGGCTTTCTCAATTA
AGTTAGACTATGTTAAGAAACCAATTCAGAAACTGCAATGGTCAA
AATTTTGGAAAAATTATATGAAGAAGATGTAGGAGCTGGGATGTAT
GTGTTGTACCTTACGGTGGTATAATGGAGGAGATTTGAGAATCAG
CAATTCATTCCCTCATCGAGCTGGAATAATGTATGAACCTTTGGTA
CACTGCTTCTGGGAGAAGCAAGAAGATAATGAAAAGCATATAAAC
TGGGTTGCAAGTGTTTATAATTTTACGACTCCTTATGTGTCCCAAA
ATCCAAGATTGGCGTATCTCAATTATAGGGACCTTGATTAGGAAA
AACTAATCATGCGAGTCTAATAATTACACACAAGCACGTATTTGG
GGTGAAAAGTATTTTGGTAAAAATTTTAAACAGGTAGTTAAGGTGA
AACTAAAGTTGATCCCAATAATTTTGTAGAAACGAACAAAGTAT
CCCACCTCTTCCACCGCATCATCATTA

[0142] Inactive THCA synthase gene (SEQ ID NO:5),
SNPs relative to SEQ ID NO:1 shown in bold underline.

ATGAATTGCTCAGCATTCCTTTTGGTTTGGTTGCAAAATAATAT
TTTCTTTCTCTCATTCATATCCAAATTTCAATAGCTAATCCTCAAG
AAAACCTCCTTAAATGCTTCTCGGAATATATTCTTAACAATCCAGCA
AATCCAAAATTCATATACACTCAACACGACCAATTGTATATGTCTGT
CCTGAATTCGACAATACAAAATCTTAGATTCACTCTGATACAAACC
CAAAACCACTCGTTATTGTCACTCCTTCAAATGCTCTCCCATATCCAG

-continued

GCCAGTATTCTCTGCTCCAGAAAGTTGGTTTGCAGATTGGAACTCG
 AAGCGGTGGCCATGATGCTGAGGGTTTGTCTACATATCTCAAGTCC
 CATTGCTATAGTAGACTTGAGAAACATGCATACGGTCAAAGTAGAT
 ATTATAGCCAACTGCGTGGGTGAAGCCGAGCTACCTTGGAGA
 AGTTTATTATTGGATCAATGAGATGAATGAGAATTTAGTTTTCTCTG
 GTGGGTATTGCCCTACTGTTGGCGTAGGTGGACACTTTAGTGGAGGA
 GGCTATGGAGCATTGATGCGAAATTATGGCCTTGGCGGTGATAATAT
 CATTGATGCACACTTAGTCAATGTTGATGAAAAAGTTCTAGATCGAA
 AATCCATGGGAGAAGATCTATTTTGGGTATACGTGGTGGAGGAGGA
 GAAAACTTTGGAATCATTGCAGCATGGAATACTAACTTGTGTGTGT
 CCCATCAAAGCTACTATATTAGTGTAAAAAGAACATGGAGATAC
 ATGGGCTTGTCAAGTTATTTAAACAAATGGCAAAATATTGCTTACAAG
 TATGACAAAGATTTAAATGCTCAGACTCACTTCAGAACTAGGAATAT
 TACAGATAATCATGGGAAGAATAAGACTACAGTACATGGTTACTTCT
 CTTCCATTTTCTTGGTGGAGTGGATAGTCTAGTTGACTTGATGAAC
 AAGAGCTTTCCTGAGTTGGGTATTAATAAACTGATTGCAAGAAT
 GAGCTGGATTGATACAACCATCTTCTACAGTGGTGTGTAAATACA
 AACTGCTAATTTTAAAGAGAAATTTGCTTGATAGATCAGCTGGG
 AAGAAGACGGCTTCTCAATTAAGTTAGACTATGTTAAGAACTAAT
 ACCTGAACTGCAATGGTCAAAATTTTGGAAAAATTATGAAGAAG
 AGGTAGGAGTTGGGATGTATGTGTGTACCTTACGGTGGTATAATG
 GATGAGATTTGAGATCAGCAATTCATTCCTCATCGAGCTGGAAT
 AATGTATGAACCTTTGGTACACTGCTACCTGGGAGAAGCAAGAAGATA
 ACGAAAAGCATATAAAGTGGGTTCGAAGTGTATTAATTTACAACT
 CCTTATGTGTCCAAAAATCCAAGATTGGCGTATCTCAATTATAGGGA
 CCTTGATTTAGGAAAACTAATCTGAGAGTCTAATAATTACACAC
 AAGCAGCTATTGGGGTGAAAAGTATTTGGTAAAAATTTTAAACAGG
 TAGTTAAGGTGAAAACCAAAGCTGATCCCAATAATTTTTTAGAAA
 CGAACAAAGTATCCCACCTCTCCACCGCTCATCATTA

[0143] THCA synthase polypeptide sequence (SEQ ID NO:6)

MNCSAFSFWFVKIIFFFLSHITQISIANPRENFKCFSKHPNNVA
 NPKLVTQHDQLYMSILNSTIQNLRFISDTPKPLVIVTPSNNSHIQ
 ATILCSKKVGLQIRTRSGGHADEGMSYISQVPFVVVDLRNMHSIKID
 VHSQTAWVEAGATLGEVYVINEKENLSFPFGGYCPTVGVGGHFSG
 GYGALMRNYGLAADNIIDAHLVNVGKVLDRKSMGEDLFWAIRGGG
 GENFGIIAAWKIKLVDVPSKSTIFS VKKNMEIHGLVKLFNKWQNIAY
 KYDKDLVLMTHFITKNITDNHGKNKTTVHG YFSSIFHGGVDSLVDLM

-continued

NKSPPELGIIKKTDCKEFSWIDTTIFYSGVNFNTANFKKEILLDRSA
 GKKTAFSILKDYVKKPIPETAMVKILEKLYEEDVGAGMYVLYPYGGI
 MEEISESAIPFPHRAGIMYELWYTASWEKQEDNEKHINVVVRVSYNF
 TTPYVSQNPRLAYLNYRDLGLGKTNHASPNNTQARIWGEKYFGKNF
 NRLVKVKTKVDPNNFRNEQSIPLPPHHH

[0144] In some embodiments, the THCA synthase gene may be knocked out using Clustered Randomly Interspersed Short Palindromic Repeats (CRISPR)/Cas mediated gene editing. To knockout the THCA synthase gene using CRISPR/Cas mediated gene editing, one or more guide RNAs are designed that target the THCA synthase gene or a region adjacent there to and proximal to a Protospacer Adjacent Motif (PAM) site. Upon introduction to a *Cannabis* cell, the guide RNAs target a nuclease to induce a double strand break at the designated cut site. The cell will then undergo non-homologous end joining (NHEJ) to repair the cut site. Due to the nature of NHEJ, one or more insertions or deletions (indels) are introduced at the cut site, thereby silencing the target THCA synthase gene. In some embodiments, a homology directed repair (HDR) template oligonucleotide may be used to direct the repair at the cut site to introduce a mutation of interest. The mutation of interest may be an insertion of a stop codon, a frameshift mutation, or a nonsense mutation that disrupts expression of the THCA synthase gene. In some embodiments, the HDR oligonucleotide encodes a sequence comprising one or more of the THCA synthase gene SNPs recited herein. In some embodiments, the guide RNAs direct cleavage of the THCA gene such that all or a portion of the THCA gene is removed.

[0145] In some embodiments, the nuclease is a Cas nuclease. Suitable Cas nucleases are known and described in the art including, but not limited to, a Cas9 nuclease.

[0146] In some embodiments, the guide RNA targeting the THCA gene is selected from SEQ ID NO:7 (TGCAGCATG-GAAAATCAAAC, forward gRNA gRF743), SEQ ID NO:8 (CCCTTACGGTGGTATAATGG, forward gRNA gRF1271), SEQ ID NO:9 (TAGCTATTGAAATTTGGATA, reverse gRNA gRR63), SEQ ID NO:10 (TAGAGCAT-AAAATAGTTGCT, reverse gRNA gRR279), or combinations thereof. As used herein, the guide RNA (gRNA) nomenclature (e.g., gRF743) refers to the template direction of the gRNA, either forward (F) or reverse (R), and nucleotide position at which the 20 base pair (bp) gRNA ends. For example, gRF743 is a forward gRNA that aligns with the THCA synthase gene and ends at nucleotide 743. Other nomenclature schemes and identification methods will be known to a skilled artisan. In some embodiments, the guide RNA sequences are cloned into a vector for introduction to the cell. In some embodiments, the vector also encodes a Cas nuclease (e.g., Cas9 nuclease). In some embodiments, one or more vectors encoding the guide RNA and the Cas nuclease are introduced into the cell in the presence of an HDR oligonucleotide. Cells positive for transformation and the desired CRISPR/Cas mediated editing results may be screened and selected for using standard molecular biology and sequencing techniques known in the art.

[0147] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis* cannabidiolic-acid synthase (CBDA synthase) gene. A heterologous nucleic acid is introduced

into the *Cannabis* explant to knockout or silence the CBDA synthase gene. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a tetrahydrocannabinol (THC) low or THC free phenotype. The sequence and activity of the CBDA synthase gene is known and described in the art. See, for example, Lavery et al. (Lavery et al., "A physical and genetic map of *Cannabis sativa* identifies extensive rearrangements at the THC/CBD acid synthase loci," *Genome Research*, 2018, 29:146-156).

[0148] Provided below are four cDNA or reverse complement sequences (SEQ ID NOs:11, 13, 15, and 17) of CBDA genes from various *Cannabis sativa* strains. Single nucleotide polymorphs (SNPs) between the active CBDA synthase of SEQ ID NO:11 and each of SEQ ID NOs:13 and 15 are indicated in each of the respective sequences and a full sequence alignment is shown in FIG. 65. Additionally, mutations in the CBDA synthase sequences of SEQ ID NOs:14 and 16 are shown relative to SEQ ID NO:12.

[0149] *Cannabis sativa* cultivar Carmen cannabidiolic acid synthase (CBDA1) gene (SEQ ID NO:11), GenBank: KJ469374.1

```
atgaagtgcctcaacattctccttttggtttgttgcagataaatatt
ttctcttttctcattcaatatccaaacttcattgctaactcctcgag
aaaacttccttaaatgcttctcgcaatatattcccaataatgcaaca
aatctaaaactcgatacactcaaaacaacccattgtatatgtctgt
cctaaattcgacaatacacaaatcttagattcagctctgacacaaccc
caaaaccacttggtatcgctcactccttcacatgtctctcatatccaa
ggcactattctatgctccaagaaagtgtgctgcagattcgaactcg
aagtgggtggtcatgattctgagggcatgctacatatctcaagtcc
catttggtatagtagacttgagaacatgcgttcaatcaaaatagat
gttcatagccaaactgcattgggttgaagccggagctacccttgagaga
agtttattattgggttaatgagaaaaatgagagctcttagtttggtcg
ctgggtattgcctactgtttgcgcaggtggacactttggtggagga
ggctatggaccattgatgagaagctatggcctcgcggtgataatat
cattgatgcacacttagtcaacgttcatggaaaagtgtatagtcgaa
aatctatgggggaagatctcttttgggctttacgtggtggtggagca
gaaagcctcggaatcattgtagcatggaaaattagactggttgctgt
cccaaagtctactatgttttagtgttaaaaagatcatggagatacatg
agcttgctcaagttagttaaacaatggcaaaatattgcttacaagtat
gacaaagattttattactcatgactcacttcataactaggaacattac
agataatcaagggaagaataagacagcaatacacacttactctctt
cagttttccttggtggagtgatagtctagtcgacttgatgaacaag
agttttcctgagttgggtattaaaaaacggattgcagacaattgag
ctggattgatactatcatctctctatagtgggtgtgtaaaattacgaca
ctgataattttaacaaggaaattttgcttgatagatccgctgggcag
aacggtgctttcaagattaagtttagactacgttaagaaaccaattcc
```

-continued

```
agaatctgtatttgtccaaattttggaaaaattatatgaagaagata
taggagctggggtgatgcgttgtagccttacgggtggtataatggat
gagatttctgaatcagcaattccattccctcatcgagctggaatctt
gtatgagttatgggtacatatgttagctgggagaagcaagaagataacg
aaaagcatctaaactggattagaaatatattataaacttcagtactcct
tatgtgtcccaaatccaagattggcatatctcaattatagagacct
tgatataggaataaatgatcccaagaatccaaataattacacacaag
cacgtatttggggtgagaagtattttggtaaaaattttgacaggcta
gtaaaagtgaaaaccctgggtgatcccaataatttttttagaaacga
acaaagcatcccacctcttccacggcatcgctcattaa
```

[0150] *Cannabis sativa* CBDA synthase (SEQ ID NO:12)

```
MKCSFSPFWFVKIIFFFFSFNIQTSIANPRENFKCFSQYIPNNAT
NLKLVTYQNNPLYMSVLNSTIHNLFSSDTPPKPLVIVTPSHVSHIQ
GTILCSKKVGLQIRTRSGGHDSEGMSYISQVPFVIVDLRNMRSIKID
VHSQTAWVEAGATLGEVYYWNEKNESLSLAAGYCPTVCAGGHFGGG
GYGPLMRSYGLAADNII DAHLVNVHGKVLDRKSMGEDLFWALRGGGA
ESFGIIVAWKIRLVAVPKSTMFSVKKIMEIHELVLKLVNKWQNIAYKY
DKDLLLMTHFITRNI TDNQGKNKTAIHTYFSSVFLGGVDSLVDLMNK
SFPELGIIKKTDCRQLSWIDTII FYSGVVNYDTDNFNKEILLDRSAGQ
NGAFKIKLDYVKKPIPESVFVQILEKLYEEDIGAGMYALYPYGIMD
EISESAIPFPHRAGILYELWYICSWEKQEDNEKHLNVVIRNIYNFMT
PYVSQNPRLAYLNYRDLDIGINDPKPNPNYQARIWGEKYFGKNFDR
LVKVKTLVDPNNFFRNEQSIPPLPREIRH
```

[0151] *Cannabis sativa* cannabidiolic acid synthase (LOC115697762) (SEQ ID NO:13), GenBank: XM_030624886.1, SNPs relative to SEQ ID NO:11 shown in bold underline.

```
ATGAAGTGCTCAACATTCTCCTTTGGTTTGTGTCAGATAATATT
TTTCTTTTCTCATTCAATATCCAAACTTCCATTGCTAATCCTCGAG
AAACTTCCTTAAATGCTTCTCGCAATATATCCCAATAATGCAACA
AATCTAAAACCTCGTATACACTCAAAACAACCCATTGTATATGTCTGT
CCTAAATTCGACAATACACAATCTTAGATTCACTCTGACACAACCC
CAAAACCACTTGTATTCTGCTACTCCTTCACATGTCTCTCATATCCAA
GGCACTATTCTATGCTCCAAGAAAGTTGGCTTGCAGATTGGAACCTCG
AAGTGGTGGTCATGATTCTGAGGGCATGTCTACATATCTCAAGTCC
CATTGTATTATAGTAGACTTGAGAAACATGCGTTCAATCAAAATAGAT
GTTCATAGCCAACTGCATGGGTGGAAGCCGGAGCTACCCCTGGGAGA
AGTTTATTATTGGGTAAATGAGAAAAATGAGAACTCTTAGTTTGGCGG
CTGGGTATTGCCCTACTGTTTGCAGAGGTGGACACTTTGGTGGAGGA
```

-continued

GGCTATGGACCATTGATGAGAAACTATGGCCTCGCGGTGATAATAT
 CATTGATGCACACTTAGTCAACGTTTCATGGAAAAGTGTAGATCGAA
 AATCTATGGGGGAAGATCTCTTTTGGGCTTTACGTGGTGGTGAGCA
 GAAAGCTTCGGAATCATTGTAGCATGGAATAGACTGGTTGTGTGT
 CCCAAAGCTACTATGTTTAGTGTTAAAAAGATCATGGAGATACATG
 AGCTTGCTCAAGTTAGTTAACAAATGGCAAAATATTGCTTACAAGTAT
 GACAAAGATTATTACTCATGACTCACTTCATAACTAGGAACATTAC
 AGATAATCAAGGGAAGAATAAGACAGCAATACACACTTACTTCTCTT
 CAGTTTCTCTTGGTGGAGTGGATAGTCTAGTCGACTTGATGAACAAG
 AGTTTTCTGAGTTGGGTATTAAAAAACGGATTGCAGACAATTGAG
 CTGGATTGATACTATCATCTTCTATAGTGGTGGTTGAAATTACGACA
 CTGATAATTTTAAACAGGAAATTTTGGCTTGATAGATCCGCTGGGCAG
 AACGGTGGCTTCAAGATTAAAGTTAGACTACGTTAAGAAACCAATTCC
 AGAATCTGTATTGTCCAAATTTTGGAAAAATTATATGAAGAAGATA
 TAGGAGCTGGGATGTATGCGTTGTACCCCTACGGTGGTATAATGGAT
 GAGATTTCAGAATCAGCAATTCATTCCCTCATCGAGCTGGAATCTT
 GTATGAGTTATGGTACATATGTAGTTGGGAGAAGCAAGAAGATAACG
 AAAAGCATCTAACTGGATTAGAAATATTTATAACTTCATGACTCCT
 TATGTGTCCAAAAAATCCAGATTGGCATATCTCAATTATAGAGACCT
 TGATATAGGAATAAATGATCCCAAGATCCAAATAATTACACACAAG
 CACGTATTTGGGGTGAGAAGTATTTTGGTAAAAATTTGACAGGCTA
 GTAAAAGTGAAACCTGGTTGATCCCAATAACTTTTTTAGAAACGA
 ACAAGCATCCACCTCTTCCACGGCATCGTCATTAA

[0152] CBDA synthase protein sequence corresponding to
 SEQ ID NO:13 (SEQ ID NO:14), mutations relative to SEQ
 ID NO:12 shown in bold underline.

MKCTFSFWFVKIIFFFFSFNIQTSIANPRENFKCFSQYIPNNATN
 LKLVTQNNPLYMSVLNSTIHNLRFSTDTTPKPLVIVTPSHVSHIQGT
 ILCSKKVGLQIRTRSGGEIDSEGMSYISQVFPVIVDLRNMRSIKIDVH
 SQTAWVEAGATLGEVYVVVNEKNENLSLAAGYCPTVCAGGEIFGGGG
 YGPLMRNYGLAADNIIDAHLVNVHGKVLDRKSMGEDLFWALRGGAES
 FGIIVAWKIRLVAVPKSTMFSVKKIMEIHELVLVKNWQNIAYKYDKD
 LLLMTHFITRNI TDNQGNKTAIHTYFSSVFLGGVDSLVDLMNKSFPPE
 LGIKKTDRCQLSWIDTIIFYSGVVNYDTDNFNKEILLDRSAGQNGAFK
 IKLDYVKKPIPEFVQILEKLYEEDIGAGMYALYPYGGIMDEISESA
 IPPPHRAGILYELWYICSEWKQEDNEKHLNVVIRNIYNFMTPYVSKNP
 RLAYLNYRDLDIGINDPKNPNNYTQARIWGEKYFGKNFDRLVKVKTLV
 DPNNFFRNEQSIPPLPREIRH

[0153] *Cannabis sativa* Finola CBDA synthase gene
 reverse complement (SEQ ID NO:15), SNPs relative to both

SEQ ID NOs:11 and 13 shown in bold underline, SNPs
 relative to SEQ ID NO:13 shown in bold italics.

ATGAAGTACTCAACATTCTCCTTTTGGTTTGGTTTGCAGATAATATT
 TTTCTTTTCTCATTCAATATCCAACTTCCATTGCTAATCCTCGAG
 AAAACTTCCTTAAATGCTTCTCGCAATATATCCCAATAATGCAACA
 AATCTAAACTCGTATACACTCAAAACAACCCATTGTATATGTCTGT
 CCTAAATTCGACAATACACAATCTTAGATTCAGCTCTGACACAACC
 CCAAACCACTTGTATCGTCACTCCTTCACATGTCTCTCATATCCA
 AGGCACATTCTATGCTCCAAGAAAGTTGGCTTGAGATTGCAACTC
 GAAGNN
 NNN
 NNNNNNNNNNNNNNNNNNNNTGGGTTGAAGCCGGAGCTACCTTGGAG
 AAGTTTATTATTGGGTAAATGAGAAAAATGAGAGTCTTAGTTGGCT
 GCTGGGTATTGCCCTACTGTTTTCGCGAGGTGGACACTTTGGTGGAGG
 AGGCTATGGACCATTGATGAGAGCTATGGCTCGCGGTGATAATA
 TCATTGATGCACACTTAGTCAACGTTTCATGGAAAAGTGTAGATCGAA
 AATCTATGGGGGAAGATCTCTTTTGGGCTTTACGTGGTGGTGGAGCAG
 AAAGCTTCGGAATCATTGTAGCATGGAATTAGACTGGTTGTGTGTCC
 CAAAGTCTACTATGTTTAGTGTTAAAAAGATCATGGAGATACATGAGC
 TTGTCAAGTTAGTTAACAAATGGCAAAATATTGCTTACAAGTATGACA
 AAGATTTATTACTCATGACTCACTTCATAACTAGGAACATTACAGATA
 ATCAAGGGAAGAATAAGACAGCAATACACACTTACTTCTCTCAGTTT
 TCCTTGGTGGAGTGGATAGTCTAGTCGACTTGATGAACAAGAGTTTTC
 CTGAGTTGGGTATTAAAAAACGGATTGCAGACAATTGAGCTGGATTG
 ATACTATCATCTTCTATAGTGGTGTGTAAATTACGACACTGATAATT
 TTAACAAGGAAATTTTGGTTGATAGATCCGCTGGGCAGAACGGTGCTT
 TCAAGATTAAGTTAGACTACGTTAAGAAACCAATTCAGAATCTGTAT
 TTGTCCAAATTTTGGAAAAATTATATGAAGAAGATATAGGAGCTGGGA
 TGATGCGTTGTACCCCTACGGTGGTATAATTGGATGAGATTCT
 GAATCAGCAATTCATTCCCTCATCGAGCTGGAATCTTGATGAGTTA
 TGGTACATATGAGCTGGGAGAAGCAAGAAGATAACGAAAAGCATCTA
 AACTGGATTAGAAATATTTATAACTTCATGACTCCTTATGTGTCCCAA
 AATCCAAGATTGGCATATCTCAATTATAGAGACCTTGATATAGGAATA
 AATGATCCCAAGAATCCAAATAATTACACACAAGCACGTATTGGGGT
 GAGAAGTATTTGGTAAAAATTTTACAGGCTAGTAAAGTGAAACCC
 CTGGTTGATCCCAATAATTTTTTTAGAAACGAACAAGCATCCACCT
 CTTCCACGGCATCATCATTAA

[0154] CBDA synthase protein sequence corresponding to
 SEQ ID NO:15 (SEQ ID NO:16), mutations relative to both
 SEQ ID NOs:12 and 14 shown in bold underline, mutations
 relative to SEQ ID NO:14 shown in bold italics.

MKYSTFSFWFVCKIIFFFFSFNIQTSIANPRENPLKCFSQYIPNNATN
 LKLVTQNNPLYMSVLNSTIHLNRFSSDTPKPLVIVTPSHVSHIQGT
 ILCSKKVGLQIRTRXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX
 XXXWVEAGATLGEVYVVVNEKNE~~ES~~LSLAAGYCPTVCAGGHFGGGGY
 GPLM~~PS~~YGLAADNIIDAHLNVNHGKVLDRKSMGEDLFWALRGGAESF
 GIIVAWKIRLVAVPKSTMFSVKKIMEIHELVLVKNWQNIAYKYDKDL
 LLMTHFITRNIITDNQGNKNTAIHTYFSSVFLGGVDSLVDLMNKSFPPEL
 GIKKTDRCQLSWIDTIIIFYSGVVNYDTDNFNKEILLDRSAGQNGAFKI
 KLDYVKKPIPESVFVQILEKLYEEDIGAGMYALYPYGGIMDEISESAI
 PPHRAGILYELWYICSWEKQEDNEKHLNVVIRNIYNFMTPIVVSQNP
 LAYLNYRDLDIGINDPKPNNTQARIWGEKYPGKNFDRLVKVKTLVD
 PNNFRNEQSIPPLPRHH

[0155] Inactive CBDA synthase (SEQ ID NO:17)

ATGAAGTGCTCAACATTCCTTTGGTTGTTGCAAGATAATATTT
 TTCTTTCTCTCATTCAATATCCAACTTCAATTGCTAATCCTCGAGAA
 AACTTCTTAAATGCTTCTCGAATATATTCACCAATGTAACAAAT
 CTA~~AA~~AACTTACACCCAAACACCAATTGTATATGCCTGTCCAAAT
 CAACAATACACAATCTTAGATTCACCTCTAACACAAACCCAAACTAC
 TTGTATCGTCACTCTCATATGTCTCTCATATCCAAGGCACTATTC
 TATGTCCAAGAAATGGTTTGCAAATTCGAACTCGAAGCGGTGGTCA
 TGATTCTGAAGACATGTCTACATATCTCAAGTCCCATTTGTTATAGT
 AGACTTGAGAAACATGCATTCAATCAACATAGATGTTATAGCCAAAT
 CGCAAGGGTTGAAGCCGAGCTACCTTGGAGAAGTTTATTATGGGT
 TAATGAGAAAAATGAGAATCTTAGTTTGGCTGCTGGGTATGCCCTAC
 TGTTAGCGCAGCTGGACACTTTGGTGGAGGAGGATATGGACATTGAT
 GCAAAATTATGGCCTCGCGGCTGATAATATCGTTGATGCACACTTAGT
 CAACGTTGATGCAAAAGTGCTAGATCGAAAATCTATGGGGGAAGATCT
 CTTTGGGCTATACGTGGTGGTGGAGGAGAAAGCTTCGAATCATTGT
 AGCATGGAAAATTAGACTGGTTGTGTGTCCCAACAAAGCTACTATGTT
 TAGTGTTAAAAAGATCATGGAGATACATGAGCTTGTCAAGTGAGTTAA
 CAAATGGCAAAATATTGCTTACAAGTATGACAAAGATTATTACTCAT
 GACTCACTTCATACTAGGAATATTACAAATAATCATGGGAAGAATAA
 GACAACAATACACACTTACTTCTCTCAGTTTTCTCTGGTGGAGTGGA
 TAGTCTAGTCGACTTGATGAATAAGAGTTTCTCTGAGTTGGGTATTAA
 AAAAAACAGATTGCAAACAATTGAGCTAGATTGATATTATCATCTTTTA
 TAGCGGTGTTGTAAATTACGGCACTGATAATTTAATAAGGAAATTTT
 GCTTGATAGATCAGCTGGGCAGAACGGTCTTTAAAGATTAAGTTAGA
 CTACGTTAAGAAACCAATTCCAGAATCTGCGTTTGTCAAATTTTGGGA

-continued

AAAATTATATGAAGAAGATGAAGGAGTTGGGATGTATGCGTTGTACCC
 TTACGGTGGTATAATGGATGAGATTTCAGAATCAGCAATTCATCTCCC
 TCATTGAGCTGGAATCATGTATGAATTATGGTACATATGTAGCTGGGA
 GAAGCACGAAGATAACGAAAAAGCATCTAACTGGATTTCGAAATGTTT
 ATAGCTTCATTACTCCTTATGTGTCTCTAAATCCAAGATTGGCATATC
 TCAATTATAGAGACCTTGATACTGGAATAAATGATCCCAAGAGTCCAA
 ATAATTACACACAAGAAAGTATTTGGGGTGAGAAGTATTTTGGTAAAA
 ATTTTGACAGGGTAGTAAAGTGAAACCCCTGGTTGATCCCAATAATT
 TTTTGAAGATGAACAAAGCATCCACCTCTTCCACGGCATCGTCATT
 AA

[0156] In some embodiments, the CBDA synthase gene may be knocked out using CRISPR/Cas mediated gene editing. To knockout the CBDA synthase gene using CRISPR/Cas mediated gene editing, one or more guide RNAs are designed that target the CBDA synthase gene or a region adjacent there to and proximal to a PAM site. Upon introduction to a *Cannabis* cell, the guide RNAs target a nuclease to induce a double strand break at the designated cut site. The cell will then undergo NHEJ to repair the cut site. Due to the nature of NHEJ, one or more indels are introduced at the cut site, thereby silencing the target CBDA synthase gene. In some embodiments, an HDR template oligonucleotide may be used to direct the repair at the cut site to introduce a mutation of interest. The mutation of interest may be an insertion of a stop codon, a frameshift mutation, or a nonsense mutation that disrupts expression of the CBDA synthase gene. In some embodiments, the HDR oligonucleotide encodes a sequence comprising one or more the CBDA synthase gene SNPs recited herein or encodes a mutation in the CBDA synthase polypeptide sequence as demonstrated herein. In some embodiments, the guide RNAs direct cleavage of the CBDA gene such that all or a portion of the CBDA gene is removed. In some embodiments, the nuclease is a Cas nuclease. Suitable Cas nucleases are known and described in the art including, but not limited to, a Cas9 nuclease.

[0157] In some embodiments, the guide RNA targeting the CBDA gene is selected from SEQ ID NO:18 (GCTA-GATCGAAAATCTATGG, forward gRNA gRF668), SEQ ID NO:19 (AAAGCATCCCACCTCTTCCA, forward gRNA gRF1621), SEQ ID NO:20 (TTTAGGACAGACATATACAA, reverse gRNA gRR172), SEQ ID NO:21 (GAAAGCACCGTTCTGCCCAG, reverse gRNA gRR1118), or combinations thereof. In some embodiments, the guide RNA sequences are cloned into a vector for introduction to the cell. In some embodiments, the vector also encodes a Cas nuclease (e.g., Cas9 nuclease). In some embodiments, one or more vectors encoding the guide RNA and the Cas nuclease are introduced into the cell in the presence of an HDR oligonucleotide. Cells positive for transformation and the desired CRISPR/Cas mediated editing results may be screened and selected for using standard molecular biology and sequencing techniques known in the art.

[0158] In some embodiments, both the THCA synthase and CBDA synthase genes are knocked out to produce a THC low or THC free *Cannabis* plant. The polypeptide

sequence of the THCA synthase and CBDA synthase enzymes are approximately 84% identical and both contribute to THC production in *Cannabis*. Plants produced by knocking out both the THCA synthase and CBDA synthase genes have significantly increased production of cannabigerolic acid (CBG), which is the substrate for both THCA synthase and CBDA synthase, relative to a *Cannabis* plant with wild-type expression of THCA synthase and CBDA synthase. In some embodiments, both the THCA synthase and the CBDA synthase genes are knocked-out using CRISPR/Cas mediated gene editing as described herein. In some embodiments, guide RNAs targeting both the THCA synthase gene and the CBDA gene simultaneously are designed and used for CRISPR/Cas9 mediated gene editing. In some embodiments, the guide RNAs targeting both THCA synthase and CBDA synthase are selected from SEQ ID NO:22 (CATTTAAGGAAGTTTCTCG, reverse gRNA gRR88), SEQ ID NO:23 (AAATGGGACTTGAGATATGT, reverse gRNA gRR259), SEQ ID NO:24 (CCCTTG-GAGAAGTTTATTAT, forward gRNA gRF481), SEQ ID NO:25 (GTACCCTTACGGTGGTATAA, forward gRNA gRF1265), SEQ ID NO:26 (ATTCCAGCTCGATGAGG-GAA, reverse gRNA gRR1291), SEQ ID NO:27 (TACACACAAGCACGTATTTG, forward gRNA gRF1515), and combinations thereof. In some embodiments, the guide RNA sequences are cloned into a vector for introduction to the cell. In some embodiments, the vector also encodes a Cas nuclease (e.g., Cas9 nuclease). In some embodiments, one or more vectors encoding the guide RNA and the Cas nuclease are introduced into the cell in the presence of an HDR oligonucleotide. Cells positive for transformation and the desired CRISPR/Cas mediated editing results may be screened and selected for using standard molecular biology and sequencing techniques known in the art.

[0159] In some embodiments, the heterologous nucleic acid of interest encodes a SOLO DANCERS (SDS) and BARNASE fusion gene. The SDS gene encodes a meiosis-specific cyclin and is required for homology interaction during meiotic prophase I in Arabidopsis. The BARNASE gene encodes a ribonuclease, is driven by a tapetum-specific promoter, and, when activated is toxic to and eliminates tapetal cells to create male sterile plants. Co-expression of SDS and BARNASE from a heterologous nucleic acid will create a male and female sterile plant. See, for example, Huang et al. ("Creating completely both male and female sterile plants by specifically ablating microspore and megaspore mother cells," *Frontiers in Plant Science*, 2016, 7(30)).

[0160] SDS gene from *Arabidopsis* (SEQ ID NO:60)

ATGAAGGAGATCGCGATGAGGAATTCAAAGCGCAAGCCTGAGCCGACG
CCGTTCCGCCGGGAAGAAGCTCCGGTCGACGCGATTACGCCGAAGAGA
GCACAGATCTCTCCGTTCTTGTTCATCACCTCTCTGGAGCAACAA
ATCGGAGTCTCTGCTGCTTCTGTCGATTCTGCTCCGATTTGCTAGCT
GATGACAACGTTTCTGTGGTTCGAGCAGAGTCGAGAAGAGCTCGAAT
CCGAAGAAGACTCTAATTGAAGAGGTAGAAGTTTCTAACCTGTTAT
AATGTGAAGGAGACGATTGGTGATTGCAAAATTCGAAGGATTACGAGG
TCTTACTCTAAGCTACACAAGGAGAAGGAGGAGATGAGATCGAAGTA
AGCGAATCGTCTTGTGTGATTGCAATCTGGTGTGATTAAAGGAGA

-continued

TTGAATGTGAAGGGAAATAAAATTAACGACAACGATGAGATCTCTTTC
TCACGATCCGATGTGACCTTCGCCGGACATGTCTCCAACAGCCGGAGT
TTGAATTTGGAATCGGAGAATAAGGAGAGCGACGCTGTTTCTGTCATA
TCTGGAGTTGAGTACTGTTCCAAGTTCGGGAGCGTTACCGGAGGAGCT
GATAACGAAGAAATTGAAATCTCCAAGCCGAGCAGCTTCGTGGAAGCT
GATTCTCTCTTGGATCGGCCAAGGAATTGAAGCCGGAGCTTGAGATA
GTCCGATGCGTCTCTGATCTCGCTTGCTCTGAGAAATCTCGGAAGAG
GTTTCGGATTCTCTCGATGATGAGTCATCTGAGCAACGTTTCAGAGATA
TATTCACAGTATTCGACTTCGATTACTCGGATTACACTCCGTCCATC
TTCTTCGACTCTGGCAGCGAATTCTCTGAGAAATCTTCTCTGATTCT
CCTATTTACATTCTCGCTCTCTGTACCTCCAGTTCAAGGAACAGTTTC
TGTAAGTCCACGATTTCCCAACGATTTTGGATCTTCTTGCAGGAAGAA
ATTCACTCTGAATTGCTAAGGTTTGATGATGAGGAGGTGGAAGAGAGC
TATCTAAGGCTGAGGGAAGAGAAAGAGTATGCATATATGCGGGAC
TGTGCTAAGGCATACCTGCTCCAGGATGGACAATACTGGTCTCATCCCT
CGTCTACGCTCCATCATGTTCAATGGATTGTAAAGCAATGTTCTGAC
ATGGGGCTTCAGCAAGAGACATTGTTTCTAGGAGTTGGTCTGTTGGAT
CGATTCTGAGCAAAGGATCATTCAAAGCGAAAGGACTCTAATACTA
GTCCGGATTGCGAGTCTTACTCTGGCCACCAGAATTGAAGAAAATCAA
CCTTACAACAGCATCCGGAAGGAACCTTCACTTTCAGAACCTAAGA
TATAGCCGCATGAAGTGGTGGCAATGGAGTGGCTGGTTCAAGAAGTC
CTCAACTTCAAATGCTTCACACCCACAATCTTCAACTTCTTGTGGTTC
TACTTAAAGCTGCTCGAGCCAATCCAGAAGTTGAAAGGAAAGCCAAA
TCCTTGGCTGTTACCTCACTATCCGACCAACTCAACTCTGTTTTTGG
CCCTCAACTGTAGCAGCTGCACTCGTGGTTCTCGCTGCATCGAACAC
AACAAAATCTCTGCATACCAACGAGTCATAAAGGTCCATGTTAGAACA
ACAGATAACGAGTTGCCTGAATGCGTTAAGAGTCTGGACTGGTGTCTT
GGGCAGTAA

[0161] BARNASE gene (SEQ ID NO:61)

CTGGAACCGTCACATTGCTTCCGCATATCGGGTCAGCAACGGCTAA
AATCCGCTTGAATATGTTACACAAAGCCGCTCAAAACATGATTGACG
CCGTATACGGAAGAACGCCGAAAAACCTTACTAAGGAATTTCAATAA
GAAGAAAAATCCCGTTGGTTTCCGCGGGGTTTATTTTTCGCTAGAT
AAAAAGTACTATTTTAAATTCTTCTATTCTTTCTTTCTGTTGCTG
ATACAATGAAAAGGAATCAGCTTCACATGATGAAAATGGGAGGTATT
GCTTTGAAAAAACGATTATCGTGGATTCCGTTTGTCTTACTGGTGTCT
TGTCTCCGCGCGGGGATGCTGTTTTCAACAGCTGCCAAAACGGAAA
CATCTTCTCACAAGGCACACAGAAGCACAGGTTATCAACACGTTT

-continued

GACGGGGTTGCGGATTATCTTCAGACATATCATAAGCTACCTGATAA
 TTACATTACAAAATCAGAAGCACAAAGCCCTCGGCTGGGTGGCATCAA
 AAGGGAACCTTGACAGCTCGCTCCGGGAAAAGCATCGGCGGAGAC
 ATCTTCTCAAAACAGGGAAGGCAAACTCCCGGCAAAAGCGGACGAAC
 ATGGCGTGAAGCGGATATTAATACTATACATCAGGCTTCAGAAATTCAG
 ACCGGATTCTTTACTCAAGCGACTGGCTGATTACAAAACAACGGAC
 CATTATCAGACCTTTACAAAATCAGATAACGAAAAAACGGCTTCC
 CTGCGGAGGCCGTTTTTTTTCAGCTTTACATAAAGTGTGTAATAAATT
 TTTCTTCAAACCTCTGATCGGTCAATTCACCTT

[0162] In some embodiments, provided herein is a *Cannabis* plant with increased trichomes that has increased expression of endogenous Cannabis lipid transfer protein 2 (LTP2) or includes a heterologous nucleic acid encoding *Brassica napus* LTP2 (BraLTP2). In some embodiments, a *Cannabis* cell is transformed with a heterologous polynucleotide encoding a polypeptide at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical to SEQ ID NO:28. In some embodiments, a *Cannabis* cell is transformed with a heterologous polynucleotide encoding the polypeptide of SEQ ID NO:28. In some embodiments, the heterologous polynucleotide is at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical to SEQ ID NO:29. The heterologous nucleic acid encoding BraLTP2 may be incorporated into a construct or vector as described herein. One embodiment for cloning BraLTP2 into a vector suitable for transformation using the methods described herein is demonstrated in FIG. 66. Trichomes in the transformed *Cannabis* plant may be at least 2 fold, 5 fold, 10 fold, or 15 fold higher than trichomes in a *Cannabis* plant that does not include the heterologous polynucleotide. In some embodiments, the modulation of the LTP2 gene in other plants, such as *Brassica napus* is described in the art. See, for example, Tian et al. (Tian et al., "Overexpression of BraLTP2, a lipid transfer protein of *Brassica napus*, results in increased trichome density and altered concentration of secondary metabolites," Int. J. Mol. Sci., 2018, 19:1733).

[0163] BraLTP2 protein sequence (SEQ ID NO:28)

MATGSRVLIGLANIILIISGELLVPGQGTCCQGDIEGLMRECAVYVQR
 PGPKNPSAACCKVKRSDIPCACGRITPSVQKIDMNKVVLTVSFC
 GRPLAHGTHKCGSYIVP

[0164] BraLTP2 gene sequence GenBank: KM062522.1 (SEQ ID NO:29)

ATGGCGACAGGTTCTCGTGTCTGATCGGTCTAGCAATGATCCTCAT
 AATCTCAGGAGAACTGCTAGTTCCAGGGCAAGGAACGTGCCAAGGAG
 ACATAGAGGGTCTGATGAGAGAATGTGCGGTCTACGTCCAGCGTCCA
 GGCCCAAAGGTAAACCCATCCGCAGCGTGTGCAAAGTCGTCAAGAG
 ATCAGACATCCCTGCGCATGTGCGGTATCACACCTCGGTTCAA

-continued

AAATGATAGACATGAATAAGGTTGTTCTTGCTCACTTCCTTTGTGGG
 AGGCCTCTCGCTCATGGTACCAAGTGTGGAAGCTACATTGTGCCATG
 A

[0165] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis* LTP2 gene. A heterologous nucleic acid is introduced into the *Cannabis* explant to upregulate, overexpress, or provide multiple copies of the LTP2 gene. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a phenotype with an increase in trichomes compared to a wild-type plant as well as increased cannabidiol (CBD) production.

[0166] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis sativa* prenyltransferase 1 (CsPT1) gene. A heterologous nucleic acid is introduced into the *Cannabis* explant to upregulate, overexpress, or provide multiple copies of the CsPT1 gene. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding a polypeptide at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical to SEQ ID NO:30. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding the polypeptide SEQ ID NO:30. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid comprising SEQ ID NO:31 or a sequence at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical thereto. The heterologous nucleic acid encoding CsPT1 may be incorporated into a construct or vector as described herein. One embodiment for cloning CsPT1 into a vector suitable for transformation using the methods described herein is demonstrated in FIG. 67. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a phenotype with increased cannabigerol (CBG) production and increased cannabidiol (CBD) production. The sequence and activity of the CsPT1 gene is known and described in the art. See, for example, Luo et al. (Luo et al., "Complete biosynthesis of cannabinoids and their unnatural analogues in yeast," Nature, 2019, 567) and U.S. Pat. No. 8,884,100.

[0167] CsPT1 protein sequence (SEQ ID NO:30)

MGLSSVCTFSFQTNVHTLLNPHNNPKTSLLCYREIPKPIKYSYNN
 FPSKHCSTKSFEILQNKCSLSIAKNSIRAATTNQIEPPESDNHVS
 ATKILNFGKACWKLQRPYTIIFTSCACGLFGKELLHNTNLISWSLM
 FKAFFFLVAVLCIASFTTTINQIYDLHIDRINKPDLPLASGEISVNT
 AWIMSIIVALFGLIITIKMGGPLYIFGYCFGIFGGIVSVPPFRWK
 QNPSTAPLLNFLAIIITNTFFYYASRAALGLPFELRPSFTFLAFM
 KSMGSALALIKDASDVEGDTKFGISTLASKYGRNLTLFCSGVILLS
 YVAAILAGIIWPQAFNSNVMLLSHAILAFWLILQTRDFALTNYDPEA
 GRRFYEFMVVKLYYAEYLVYVFI

[0168] CsPT1 cDNA sequence (SEQ ID NO:31)

atgggactctcatcagtttgtaccttttcatttcaactaattaccata
 ctttattaaatcctcacaataataatcccaaacctcattattatgtta

-continued

tcgacacccccaaaacaccaattaaatactcttacaataattttcctct
 aaacattgctccaccaagagttttcatctacaaaacaatgctcagaat
 cattatcaatcgcaaaaaattccattagggcagctactacaaatcaaac
 tgagcctccagaatctgataatcattcagtagcaactaaaattttaaac
 tttgggaaggcatgttgaaacttcaaagaccatatacaatcatagcat
 ttacttcatgcgctgttggtattgtttgggaagagttgttgcataacac
 aaatttaataagttggtctctgatgttcaaggcattcttttttttggtg
 gctgtattatgcatgtcttcttttacaactaccatcaatcagatttacg
 atcttcacattgacagaataaacaagcctgatctaccactagcttcagg
 ggaaatatcagtaaacacagcttggtattatgagcataattgtggcactg
 tttggattgataataactataaaaaatgaagggtggaccactctatatat
 ttggctactgttttggatttttgggtggattgtctattctgttccacc
 atttagatggaagcaaaatccttccactgcatttcttctcaatttcctg
 gcccatattattacaaatttcacattttattatgccagcagagcagctc
 ttggcctaccatttgagttgaggtcttattactttctgctagcattt
 atgaaatcaatgggttcagctttggccttaatacaagatgcttcagacg
 ttgaaggcgacactaaatttggcatatcaacottggcaagtaaatatgg
 ttccagaaacttgacattattttgttctggaattgttctctatcctat
 gtggctgctatacttctgctgggattatctggccccaggctttcaacagta
 acgtaatgttactttctcatgcaatcttagcattttgtttaatcctcca
 gactcgagatttttgcgttaacaaattacgaccggaagcaggcagaaga
 ttttacgagttcatgtggaagctttattatgctgaatatattagtatatg
 ttttcatataa

[0169] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis sativa* O-methyltransferase (CsOMT21) gene. A heterologous nucleic acid is introduced into the *Cannabis* explant to upregulate, overexpress, or provide multiple copies of the CsOMT21 gene. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding a polypeptide at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical to SEQ ID NO:32. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding the polypeptide SEQ ID NO:32. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid comprising SEQ ID NO:33 or a sequence at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical thereto. The heterologous nucleic acid encoding CsOMT21 may be incorporated into a construct or vector as described herein. One embodiment for cloning CsOMT21 into a vector suitable for transformation using the methods described herein is demonstrated in FIG. 68. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a phenotype with increased chrysoeriol production and increased cannflavin A and cannflavin B production. The sequence and activity of the CsOMT21 gene is known and described in the art. See, for example, Rea et al. Phytochemistry 2019:

“Biosynthesis of cannflavins A and B from *Cannabis sativa* L.”

[0170] CsOMT21 (PK24150) (SEQ ID NO:32)

MGSTGIETQMTPTQISDEEANLFAMQLASASVLPMLKALELDLLEI
 TAKAGPGAFSPSDIAQQLPTQNPDPVMDRLRLLLASYNVVTYSLR
 ERETAEEEGKVERLYGLAPVSKYLTKNEDGVSIAPLCLMNQDKVLMES
 WYEILKDAVLDDGGIPFNKAYGMTAFHEYHGTQRFNKIFNRGMSDHSTI
 TMKILEITYKGFEGLSIVDVGGGTGAVNMIVSKYPTIKGINFDLPH
 VIEDAPPLTGVEHVGGDMFVSVPKGDAIFMKWICHDSDEHCLKFLKN
 CHAALPEHGKVIVAEICILPVAPDSSLATKSTVHIDIVIMLAHNPPGKKER
 TEKEFEALAKGAGFKGFKVHCNANFTHIMEFLKTI

[0171] *Cannabis sativa* PK24150.1_1.CasaPuKu, GenBank: JP459899.1 (SEQ ID NO:33)

ATGGGTTCACAGGAATAGAGACCCAAATGACCCCAACCCAAATATCC
 GACGAAGAAGCCACCTCTTCGCCATGCAATTAGCCAGTGCCTCAGTC
 TTACCCATGGTTCTCAAAGCAGCTTTAGAGCTCGACCTCTTGGAGATC
 ATAGCCAAGCCGGTCCAGGCGCGTTTCTCTCACCTTCCGACATAGCT
 CAACAGCTTCCGACTCAGAACCAGACGCCCCGGTGATGCTGGACCGG
 ATGCTGAGACTGTTGGCTAGCTACAACGTGGTGAGTACTCGCTGCGT
 GAGCGTGAGACGCGGGAAGAGGAAGGGAAGGTGGAGAGGCTTTATGGG
 TTGGCTCCGGTGAGTAAATATCTGACGAAGATGAAGATGGAGTCTCC
 ATTGCTCCTCTTTGTCTCATGAACCAGGATAAGGTTCTTATGGAGAGT
 TGGTATCACTTAAAGATGCAGTACTTGATGGAGGAATACCTTTCAAC
 AAGGCATATGGAATGACAGCATTGAATATCATGGAACCGATCAAAGG
 TTCAATAAAATCTTTAATAGAGGAATGTCCGACCACTCGACTATTACC
 ATGAAAAAATCCTCGAAACTTACAAGGGTTTCGAGGGTCTTAACTCG
 ATTGTTGATGTTGGTGGTGGTACTGGAGCTGTTGTTAACATGATCGTC
 TCTAAGTACCCTACTATTAAAGGTATTAACTTCGATTGGCTCATGTC
 ATCGAAGATGCACCTCCATTGACCGGTGTAGAGCATGTTGGAGGAGAC
 ATGTTTGTAAAGTGTACCAAAAGGAGATGCAATTTTCATGAAGTGGATT
 TGCCATGATTGGAGCGATGAACACTGCTTGAAATCTTGAAGAACTGC
 CACGCTGCAGTCCCCGAACACGGAAGGTGATCGTGGCGGAGTGCATT
 CTTCCGGTGGCACCGGACTCGAGCCTTGCCACAAAGAGTACGGTCCAC
 ATTGATGTGATCATGTTGGCCCATAAACCTGGTGGCAAGAGAGAACA
 GAGAAGAGTTTGAAGCATTGGCTAAGGGAGCTGGCTTTAAAGGCTTC
 AAAGTCCATTGCAATGCTTTCAATACCCATATCATGGAATTTCTCAAG
 ACCATTTAA

[0172] In some embodiments, the heterologous nucleic acid of interest modulates the expression or function of the endogenous *Cannabis sativa* prenyltransferase 3 (CsPT3) gene. A heterologous nucleic acid is introduced into the

Cannabis explant to upregulate, overexpress, or provide multiple copies of the CsPT3 gene. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding a polypeptide at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical to SEQ ID NO:34. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid encoding the polypeptide SEQ ID NO:34. In some embodiments, a *Cannabis* cell is transformed with a heterologous nucleic acid comprising SEQ ID NO:35 or a sequence at least 80%, 85%, 90%, 95%, 98%, 99%, or 99.9% identical thereto. The heterologous nucleic acid encoding CsPT3 may be incorporated into a construct or vector as described herein. One embodiment for cloning CsPT3 into a vector suitable for transformation using the methods described herein is demonstrated in FIG. 69. *Cannabis* plants grown from the transformed *Cannabis* explant are characterized by a phenotype with increased cannabigerol (CBG) production and increased cannabidiol (CBD) production.

[0173] CsPT3 (PK17697) (SEQ ID NO:34)

```
MVFSSVCSFPSSLGTNFKLVPRSNFKASSSHYHEINNFINNKP IKFSY
FSSRLYCSAKPIVHRENKFTKSFSLSEILQRKSSIKAHGEIADGNSG
TSEFNVMKSGNAIWRFRVRYAAKGVLFNSAAMFAKELVGNLNLFSWPL
MFKILSPTLVILCIFVSTSGINQIYDLDIRLNKPNLPVASGEISVEL
AWLLTIVCTISGLTLTIITNSGPPFPFLYSASIFFGLYSAPPFRWKK
NPFTACFCNVMLYVGTSVGVYACKASLGLPANWSPAFCLLFWFISLL
SIPISIAKDLSDIEGRKFGIITFSTKFGAKPIAYICHGLMLLNYSV
MAAAIWPQFFNSSVILLSHAFMAIWWLYQAWILEKSNYATETCQKYY
IFLWIIIFSLEHAFYLEM
```

[0174] *Cannabis sativa* PK17697.1_1.CasaPuKu, GenBank: JP460361.1, (SEQ ID NO:35)

```
atggtgttctcatcagttttagtcttccactcccttggaaactaat
tttaaattagttctcgtagtaattttaaggcatcatcttctcattat
catgaaataaataattttattaataataaaccaattaaattctcatat
ttttcttcaagactatattgctctgccaaccaattgtacacagagaa
aacaattcacaaatcattttcactcagccacctccaaaggaaaagc
tccataaaggcacatggtgaaattgaagctgatgggagtaatggcaca
tctgaatttaattgtaataaaagtggaaacgaatttgagatttgta
aggccatattgcagccaaggagattgtttaactctgctgctatgttt
gcaaaagagttggtgggaacctaattctatttagttggcattgatgt
ttaagatactctcttttacatttggtattttatgcatttttgtaagta
caagtggcatcaatcaaatttatgatctcgacatcgacaggttaaca
aacctaatttgccagtagcatcaggagaaatttcagttgaattggcat
ggttggtgactatagtttgtaataaagtggcctcacattaacaatta
taacgaactcagggccattcttcccttttctctactctgctagtatct
tttttggtcttctctattctgctcctccattcagatggaagaagaatc
```

-continued

```
cttttacagcatgtttctgtaattgtatgttggcacaagcg
ttggtgtctattatgcttgaaggctagtctcgggcttccagccaact
ggagccctgctttttgttgccttttgggttatttcattgttgagta
tacctatctccattgcaaaagatcttccagacatagaaggtagccgca
agtttggaatcataaccttctcaactaaatttggagcaaaaccatag
catatatttgcattggaactcatgcttctgaattacgtgagtggtatgg
ctgcagctattatttggccacagttttcaacagtagcgtaatttgc
tttctcatgcattcatggcaatttgggtattatcatcaggttggatatt
tggaagaatcaaattacgccacggagacgtgccaaaaatactatatat
tcttttgataatttttctcttgaaatgccttctatttggctcatgt
ag
```

[0175] In some embodiments, the heterologous nucleic acid of interest produces a glyphosate resistant *Cannabis* plant following transformation. To create a glyphosate resistant *Cannabis* plant, a *Cannabis* cell is transformed as described herein with a heterologous nucleic suitable for mutating the *Cannabis sativa* gene encoding 3-phosphoshikimate 1-carboxyvinyltransferase 2 (EPSP synthase) such that when expressed, the EPSP synthase enzyme is not inhibited by the herbicide glyphosate. Without wishing to be bound by any particular theory or embodiment, glyphosate is a competitive inhibitor of phosphoenolpyruvate (PEP) in EPSP, acting as a transition state analog that binds more tightly to the EPSPS-S3P (shikimate-3-phosphate bound EPSP synthase) complex than PEP. Upon exposure to glyphosate, the EPSP synthase enzyme is non-functional or severely inhibited resulting in plant death. However, it is possible to mutate the EPSP synthase enzyme such that it does not bind glyphosate but still catalyzes the synthesis of 5-enol-pyruvylshikimate-3-phosphate.

[0176] In some embodiments, the *Cannabis sativa* gene encoding EPSP synthase is mutated using prime editing guide RNA (pegRNA) together with a Cas9 nuclease and reverse transcriptase fusion protein. In some embodiments, the Cas9 nuclease is a mutated Cas9 nuclease that only cleaves a single strand of the target DNA. In some embodiments, the mutated Cas9 nuclease is a Cas9 H840A nickase or a Cas9 D10A nickase (Cas9n). pegRNA are designed with the desired genetic mutation and to target the *Cannabis sativa* EPSPS gene loci of interest. A nucleic acid encoding the pegRNA and the Cas9 nuclease/reverse transcriptase fusion protein is introduced into a *Cannabis sativa* cell. Cells positive for transformation and the desired prime editing mutations may be screened and selected for using standard molecular biology and sequencing techniques known in the art.

[0177] In some embodiments, the *Cannabis* EPSPS gene (SEQ ID NO:40) is mutated at positions 1790, 1801, 3620, and 3621 to encode an EPSP synthase that is glyphosate resistant and includes T181I, P185S, and P460L mutations relative to the wild-type sequence. In some embodiments, the pegRNA for the mutations at positions 1790 and 1801 comprises the gRNA sequences tgaagactt tgcaCAACTTTTCCTTGGAAATGCgttaagtcttct (forward, SEQ ID NO:41) and AGAAGACTT AAACGCATTTCCAAGGAAAAGTTG

TGCAAAGTCTTCA (reverse, SEQ ID NO:42) and the sequence ACCTGCTATA GTGCTGAGTGAACGCATTGCTATTCCAGCATTCCA AGGAAACATAT AGCAGGT (SEQ ID NO:43) encoding the two C→T mutations at positions 1790 and 1801. See, FIG. 70. In some embodiments, the pegRNA for the mutations at positions 3620 and 3621 comprises the gRNA sequences tgaagactttgcaCTTGAGCAACAGTTGAGGA gtttaagtcttct (forward, SEQ ID NO: 44) and AGAAGACTT AAACCTCCTCAACTGTTGCTCCAAG TGCAAAGTCTTCA (reverse, SEQ ID NO:45) and the sequence ACCTGCTATA GTGCCACGCAGTAATCAAGTCCTTCTCAACTGTTG AACATATAGCA GGT (SEQ ID NO:46) encoding the two C→T mutations at positions 3620 and 3621. See, FIG. 71. **[0178]** Cannabis EPSP synthase (SEQ ID NO:36)

MAQVSKICSNGAQITILTLPNISKSHTPRSLNSVSLRSPFLGSSNSLSL
KIGTEFGGCS TVGKAMAGPVMASAVTAEKPSKVPEIVLQPIKDISGTV
KLPGSKSLSNRILLALLAALSEGTTVDNLLSDDIHYMLGALETGLRLV
EADKESKRAIVEGCAGQFPAGKESVDEVQLFLGNAGTAMRPLTAAVTV
AGGNASYVLDGVPRMRERPIGDLVTGLKQLGADVDFHGTDCPPVRVL
GKGGLPGGKVKLSGSISSQYL TALLMAAPLALGDVEIEIIDKLISVPY
VDMTLKLMARFGVTVVEHSDSWDRFLVKGGQKYKSPGNAYVEGDASSAS
YFLAGAAVTGGTVTVVEGCGTSSLQGDVKFAEVLEKMGAKVSWTENSVT
VTGP PRDSVKSKHLKAIDVNMNKMPPDVAMTLAVVALFADGPTAIRDVA
SWRVKETERMIAICTELRKL GATVEEGPDYCVITPPEKLNITAITD TYD
DEIRMAMAFSLAACSDVPVTIKDPGCTRKTFFPDYFEVLERFTKH

[0179] Glyphosate resistant Cannabis EPSP synthase (SEQ ID NO:37), mutations relative to SEQ ID NO:36 shown in bold underline including T181I, P185S, P460L

MAQVSKICSNGAQITILTLPNISKSHTPRSLNSVSLRSPFLGSSNSLSL
KIGTEFGGCS TVGKAMAGPVMASAVTAEKPSKVPEIVLQPIKDISGTV
KLPGSKSLSNRILLALLAALSEGTTVDNLLSDDIHYMLGALETGLRLV
EADKESKRAIVEGCAGQFPAGKESVDEVQLFLGNAG**I**AMRSLTAAVTV
AGGNASYVLDGVPRMRERPIGDLVTGLKQLGADVDFHGTDCPPVRVL
GKGGLPGGKVKLSGSISSQYL TALLMAAPLALGDVEIEIIDKLISVPY
VDMTLKLMARFGVTVVEHSDSWDRFLVKGGQKYKSPGNAYVEGDASSAS
YFLAGAAVTGGTVTVVEGCGTSSLQGDVKFAEVLEKMGAKVSWIENSVT
VTGP PRDSVKSKHLKAIDVNMNKMPPDVAMTLAVVALFADGPTAIRDVA
SWRVKETERMIAICTELRKL GATVEEG**L**DYCVITPPEKLNITAITD TYD
DEIRMAMAFSLAACSDVPVTIKDPGCTRKTFFPDYFEVLERFTKH

[0180] Cannabis EPSP synthase cDNA (SEQ ID NO:38)

ATGGCCCAAGTGAGCAAAATCTGTAGCAATGGAGCTCAAACATATCCTTA
CTCTCCCAAATATATCTAAGTCTCATACACCAAGATCCCTAAATTCAGT

-continued

TTCGTTGAGATCACCGTTTTTGGGTT CATCTAACTCTTTGAGTTTGAAG
ATTGGAACTGAATTTGGGGTTGTTCTACGGTTGGTAAAGCTATGGCTG
GTCCAGTCATGGCTTCAGCTGTCACAGCGGAGAAAGCCTTCAAAGGTACC
GGAGATTGTGTTGCAGCCATTAAAGATATCTCTGGCACTGTCAAGTTG
CCGGGTTCCAAGTCACTATCGAATCGGATTCTACTCCTGGCTGCTCTTT
CTGAGGGGACAACTGTTGTGGACAACTGTTAGATAGTGATGACATTCA
CTACATGCTTGGTGCCTTGGAACCCCTGGTCTTCGTGTGAAGCAGAC
AAGGAAAGCAAACGAGCAATTGTGGAAGGTTGTGCGGGTCAGTTTCCTG
CAGGTAAAGAACTCTGTTGACGAAGTTCAACTTTTCCTTGGAATGCTGG
AACAGCAATGCGTCCACTCACAGCTGCGGTGACTGTTGCTGGTGGAAT
GCTAGCTACGTACTTGATGGTGTTCCTCGAATGAGAGAAAGACCAATTG
GAGATTGGTGACTGGTCTTAAGCAGCTTGGTGCAGATGTTGATTGTTT
TCATGGTACGGATTGTCCCCCTGTTCTGTGCTTGGAAAGGAGGCCTT
CCTGGGGCAAGGTGAAACTTTCTGGATCAATTAGCAGTCAATATTTGA
CAGCCTTGCTTATGGCAGCTCCCTTGCTCTTGGAGATGTTGAAATCGA
GATAATTGATAAATTGATCTCGGTTCCCTATGTTGATGACTTTGAAG
TTGATGGCACGTTTTTGGGGTTACTGTTGAACACAGTGATAGCTGGGATC
GATTTTTAGTTAAAGGAGGTCAAAGTACAAATCTCTGGAACAGCTTA
TGTTGAAGGTGATGCTTCAAGTGCTAGTTACTTCTAGCTGGTGTGCA
GTCACTGGTGGTACAGTCAACCGTAGAAGGTTGTGGGACTAGTAGTTTAC
AGGGAGACGTAAATTTGCTGAAGTTCTTGAGAAAATGGGTGCTAAAGT
TAGCTGGACAGAGAACAGTGTCACGGTCACTGGACCACCACAGATTCT
GTAAAAAGTAAACACTTGAAAGCCATTGATGTCAACATGAACAAAATGC
CTGATGTTGCCATGACTCTTGCTGTAGTTGCTCTTTTTGCTGATGGCCC
CACTGCTATAAGAGATGTGGCAAGTTGGAGAGTCAAGGAGACAGAGAGA
ATGATTGCCATCTGCACTGAACTCAGAAAGCTTGAGCAACAGTTGAGG
AAGACCCGATTACTGCGTGATCACTCCACGAGAGAACTAAATATCAC
AGCAATAGACACATACGACGACCACAGGATGGCTATGGCGTTCTCTCTT
GCAGCTTGTTGAGATGTGCCAGTTACCATTAAAGGATCCTGGTTGCACCC
GAAAAAAGTTCCAGATTACTTTGAAGTCTTGAGAGATTTACAAAGCA
CTGA

[0181] Glyphosate resistant Cannabis EPSP synthase cDNA (SEQ ID NO:39), mutations relative to SEQ ID NO:38 shown in bold underline, including C542T, C553T, C1379T, and C1380T

ATGGCCCAAGTGAGCAAAATCTGTAGCAATGGAGCTCAAACATATCCTTA
CTCTCCCAAATATATCTAAGTCTCATACACCAAGATCCCTAAATTCAGT
TTCGTTGAGATCACCGTTTTTGGGTT CATCTAACTCTTTGAGTTTGAAG
ATTGGAACTGAATTTGGGGTTGTTCTACGGTTGGTAAAGCTATGGCTG

-continued

GTCCAGTCATGGCTTCAGCTGTCACAGCGGAGAAGCCTCAAAGGTACC
 GGAGATTGTGTTGCAGCCATTAAAGATATCTCTGGCACTGTCAAGTTG
 CCGGGTTCCAAGTCACTATCGAATCGGATTCTACTCCTGGCTGCTCTTT
 CTGAGGGGACAACCTGTTGAGCAACTTGTAGATAGTGATGACATTCA
 CTACATGCTTGGTGCCTTGGAACCCCTGGTCTTCGTGTTGAAGCAGAC
 AAGGAAAGCAAACGAGCAATTGTGGAAGGTTGTGCGGGTCAGTTTCCTG
 CAGGTAAAGAATCTGTTGACGAAGTTCAACTTTTCTTGGAATGCTGG
 AATAGCAATGCGTTCACTCACAGCTGCGGTGACTGTTGCTGGTGGAAAT
 GCTAGCTACGTACTTGATGGTGTCTCTCGAATGAGAGAAAGACCAATTG
 GAGATTGGTGACTGGTCTTAAGCAGCTTGGTGCAGATGTTGATTGTTT
 TCATGGTACGGATTGTCCCTGTTTCGTGTGCTTGGAAGGAGGCTTT
 CCTGGGGCAAGGTGAACTTTCTGGATCAATTAGCAGTCAATATTTGA
 CAGCCTTGCTTATGGCAGCTCCCTTGGCTCTTGGAGATGTTGAAATCGA
 GATAATTGATAAATGATCTCGGTTCCCTATGTTGATATGACTTTGAAG
 TTGATGGCAGCTTTTGGGGTACTGTTGAACACAGTGATAGCTGGGATC
 GATTTTGTAGTTAAAGGAGGTCAAAGTACAAATCTCCTGGAACGCTTA

-continued

TGTTGAAGGTGATGCTTCAAGTGCTAGTTACTTCTAGCTGGTGCTGCA
 GTCCTGGTGGTACAGTCACCGTAGAAGGTTGTGGGACTAGTAGTTTAC
 AGGGAGACGTAAAATTTGCTGAAGTTCTTGAGAAAATGGGTGCTAAAGT
 TAGCTGGACAGAGAACAGTGTACGGTCACTGGACCACCACGAGATTCT
 GTAAAAAGTAAACACTTGAAAGCCATTGATGTCAACATGAACAAAATGC
 CTGATGTTGCCATGACTCTTGCTGTAGTTGCTCTTTTGCTGATGGCCC
 CACTGCTATAAGAGATGTGGCAAGTTGGAGAGTCAAGGAGACAGAGAGA
 ATGATTGCCATCTGCACCTGAAGTCTGAGAAAGCTTGGAGCAACAGTTGAGG
 AAGGACTTGATTACTGCGTGATCACTCCACCAGAGAACTAAATATCAC
 AGCAATAGACACATACGACGACCACAGGATGGCTATGGCGTTCTCTCTT
 GCAGCTTGTTCAGATGTGCCAGTTACCATTAAAGATCCTGGTTGCACCC
 GAAAACTTTCCAGATTACTTTGAAGTCCTTGAGAGATTTACAAAGCA
 CTGA

[0182] *Cannabis* EPSP synthase gene, *Cannabis sativa* chromosome 2, cs10, whole genome shotgun sequence, *Cannabis sativa* 3-phosphoshikimate 1-carboxyvinyltransferase 2 (LOC115705599), (SEQ ID NO:40), possible glyphosate resistant mutation target loci indicated in bold underline italics including positions 1790, 1801, 3620, and 3621.

GGTTGGTAAGCCCTCTACCTCTTTGAAAATTGAAAGAGAGTCAATGTGACCTACAGCAG
 CAGCATCCATTAACTGTTACCATGGCCACCAAAATCCAACCTTTATTTGTATAGAGAGAATC
 AGAGAAGGTTTGGGTTTTCAGAGAGAGAGAGAAGAAGAACAAAAAATGGCCCAAGTGAGCAA
 AATCTGTAGCAATGGAGCTCAAACATATCCTTACTCTCCCAATATATCTAAGTCTCATACA
 CCAAGATCCCTAAATTGAGTTTCGTTGAGATCACCGTTTTTGGGTTTCACTAACTCTTTGAG
 TTTGAAGATTGGAACGAAATTTGGGGTGTGTTCTACGGTTGGTAAAGCTATGGCTGGTCCAG
 TCATGGCTTCAGCTGTACAGCGGAGAAGCCTTCAAAGGTACCGGAGATTGTGTTGCAGCCC
 ATTAAAGATATCTCTGGCACTGTCAAGTTGCCGGGTTCCAAGTCACTATCGAATCGGATTCT
 ACTCCTGGCTGCTCTTTCTGAGGTATATTTCAATTTTTTTTAAACGTCAAACATGTATTTTT
 GTCGAGGAAGTTTTCTGTATATACAAGATAAGAGAGTAAAAATATGGAACATCAATACCAA
 AATGAACCAAACTAGGCTAAGCTATCAAATCATGTATGGTATGCCACTCTACTTTTCCT
 ATCTCAAGCTCCACAGCTATAAAATACTATATCGTAATATTTTGTCAACTGCTTTTCATATT
 CCTTGTAATTTCCCTCATTTCCCACTAAACTAGTTCCAATGGATTGTGTGGCTGGAACTGT
 AGTTAGTTACATTAGCTAGATCTGAACCATGATCAGCATCGACTGCCCACTGGTAAACCAT
 GTAATTGCATGGAATTTCTCCTTTGTTATCCACAAATTTGAAAAGTATTTTGGAGGTATACA
 AAGATTGTGCTTTTTATGAGCAATTTCTTTTAGTTTATGTTAAGAGTTTGTAGCGATGGG
 ATGTTTTTTTTCTAGAAAATGGACAGTAAAGCTTAGCATTTTTACTTTTATTGTTGTAATGA
 ATAGTGTTCATTGAAGCTGAACTCATGCCCTTAATTGGGAGGAAAATTGAGAGAAATGGAGT
 AAAGTAATATGATATTTTGGTTAAATTCGTAAGAATATGATGGAAATAAAAAATGCAACTCA
 ACTGGGTTACTGAAGTTATATTTCTGGTCTCAGTTGTGCTTTTACAACCTTAGTCTAGAGCT
 CCACGCTGCGGAGAGATTCGGAGTCCTTACAGTTTATTTTGATAATGATTTATGAGAATTTT

- continued

ATAACTCTACGCTTTTGGTTACATTATATATGAGGTGTTTCGTTGGTGCAATTGCTCACCTGA
ACTCCCTAAATTTTAGAATGTGGGATTTAGAATGAAGTTATACTATTAGTGTGAGTCATC
TAGAATTTGTAGTGCTCATTCTCCATATACTCTTCTCTATTTCTCCCATATTTTGGCG
CTACTACTTATCTTTACAGTTCATGTTATTTTCATGTACTTGAGTTTTTGGCCATAAAAT
ATTTTGAGCGGTGGGAAGTAACTGTTTTTTTTGTTATAATTATCCAGGGGACAACTGTTGTG
GACAACTTGTTAGATAGTGATGACATTCACTACATGCTTGGTGCCTTGGAAACCCCTGGTCT
TCGTGTGAAGCAGACAAGGAAAGCAAACGAGCAATTGTGGAAGGTTGTGCGGGTCAGTTTC
CTGCAGGTAAAGAATCTGTTGACGAAGTTCAACTTTTCCTTGGAATGCTGGAAAGCAATG
CGTCCACTCACAGCTGCGGTGACTGTTGCTGGTGGAATGCTAGGTTTGTCTTCATTGCAAT
TGCTTTGAATATAAAGTACTTCTAATGCAGTGAATTTATGCTCTGTTTTCTTACTGGCC
GAGTAGCTCTTACATTTTAGGTAAAGAAAGTCATTTTGCTAACACATCACCATTTATACT
TCCCCTTTTACTTTGATGTGTTATGCTAGAAATTACATGTTGGAATGAAGTAGCACATAT
CATAAATTATTTGTATGCTGTTATTACATTTTCTCAGTAACCTCTTAACTTCTATATCTCA
GCTACGTACTTGATGGTGTTCCTCGAATGAGAGAAAGACCAATTGGAGATTGGTGACTGGT
CTTAAGCAGCTTGGTGAGATGTTGATTGTTTTCATGGTACGGATTGTCCCTGTTGCTGT
GCTTGGAAGGAGGCTTCTGCGGGCAAGGTGAGGCTTGCAATGCTTCTTCTTATTCTTT
TTGGCCATAAAACATCATTGTAATAGTGGTTTTATGTTATGAAATCCATTGACTGGTTATT
TTTAGGTTGTTGTTTTGCTTTTAAATAAAACAATATTGTCAATGATGCATAAGTAGTGAT
TACATCTACATCATTTAATTTATTTATCTTAAATGATGACAAACTTCATCATTTGACTCAGA
ATTATGTAATATTACCCCTTGCAGGTGAAACTTTCTGGATCAATTAGCAGTCAATATTGAC
AGCCTTGCTTATGCGCAGCTCCCTTGGCTCTTGAGATGTTGAAATCGAGATAATTGATAAAT
TGATCTCGGTTCCCTATGTTGATATGACTTTGAAGTTGATGGCACGTTTTGGGGTACTGTT
GAACACAGTGATAGCTGGGATCGATTTTTAGTTAAAGGAGGTCAAAGTACAAGTAGGTTTC
TTCTGAATATAGTTGATAGTATTGTTACATTACATCTGTTTATGTCAAAGAGTAATAAATTG
AAAAATAAAATCTGTCAGATCTCCTGGAAACGCTTATGTTGAAGGTGATGCTTCAAGTGCT
AGTTACTTCTAGCTGGTGCTGCAGTCACTGGTGGTACAGTCACCGTAGAAGGTTGTGGGAC
TAGTAGTTTACAGGTATTTTGCTTAGACCTTGAAATCTCTTATTCTGTACTTGTGTTTACA
TAGAATCTAAGATTAAGTGATTTTACATACATTAAGTGGTGTAAATAAGGGAGACGTAAA
ATTTGCTGAAGTTCTTGAGAAAATGGGTGCTAAAGTTAGCTGGACAGAGAACAGTGTCACGG
TCACTGGACCACCAGAGATTCTGTAAAAAGTAAACACTTGAAAGCCATTGATGTCAACATG
AACAAAATGCCTGATGTTGCCATGACTCTGCTGTAGTTGCTCTTTTGTGATGGCCCCAC
TGCTATAAGAGATGGTATGTTTTCTTAAATTTGTGAGATGGTAAATGGGGCAGTCGGTT
TGGGTTGGGTAGATTATCGGTTCTCGTCTGCCATAATAAAAAATAATCTGCTCATTGCAA
TAAATTTACAGACACAAAAATGAAACCAATAAAATATTATTTGTTTAGAGGATTAAATA
CTCATTTCTTGCCTTTCCTAATCCCAGTGGCAAGTTGGAGAGTCAAGGAGACAGAGAGAAT
GATTGCCATCTGCACTGAACTCAGAAAGGTTAGTTTTATGCTGTTTTATGTACTTGTATG
TCATGCGCTTGGAATGTAATGGCTGATAGCTATCTGTTCTTATGGGAACAAACATTTGAGC
TTGGAGCAACAGTTGAGGAAGGACCGGATTACTGCGTGATCACTCCACCAGAGAACTAAAT
ATCACAGCAATAGACATACGACGACCACAGGATGGCTATGGCGTTCTCTTGCAGCTTG

- continued

TTCAGATGTGCCAGTTACCATTAAAGGATCCTGGTTGCACCCGAAAACTTTCCAGATTACT
 TTGAAGTCCTTGAGAGATTACAAAGCACTGAATGAGTATTTATTAAGTGGATAGAGAACAA
 TAGCATCGGCTACTGTCATTACAATAAGCAGTTGGGAGGCAGGCAATCCTTTTCAATTAT
 CATGTGTTTGAATTTTGGTCGACTGTATTGCAAGTTGAGCTTCTCATTATTATTAAGACTGTA
 ATCGTAGTTATTTGTTGTAACCTTCTGCAAACCCCTTCATGTTATTTTTTACCCTTCTAATAA
 GCCAGTGGGGCAAATTCATATCTGCTATATGAGCTTGAGTGTAGAGAGAACTTTTGTCAAT
 GTATAGGTTTCTAGCAGAAGCACCATCCCTAATATGCTTTATTATAAGAGTTGCTGTGATCG
 TGTAGTGTATTTTATTGAAAGTAACCGACGCATCTCATATTA

[0183] As used herein, the terms “polynucleotide,” “polynucleotide sequence,” “nucleic acid” and “nucleic acid sequence” refer to a nucleotide, oligonucleotide, polynucleotide (which terms may be used interchangeably), or any fragment thereof. These phrases also refer to DNA or RNA of natural or synthetic origin (which may be single-stranded or double-stranded and may represent the sense or the antisense strand). The polynucleotides may be cDNA or genomic DNA.

[0184] As used herein, the term “construct” refers to recombinant polynucleotides including, without limitation, DNA and RNA, which may be single-stranded or double-stranded and may represent the sense or the antisense strand. Recombinant polynucleotides are polynucleotides formed by laboratory methods that include polynucleotide sequences derived from at least two different natural sources or they may be synthetic. Constructs thus may include new modifications to endogenous genes introduced by, for example, genome-editing technologies. Constructs may also include recombinant polynucleotides created using, for example, recombinant DNA methodologies.

[0185] The constructs provided herein may be prepared by methods available to those of skill in the art. Notably each of the constructs used or claimed are recombinant molecules and as such do not occur in nature. Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, and recombinant DNA techniques that are well known and commonly employed in the art. Standard techniques available to those skilled in the art may be used for cloning, DNA and RNA isolation, amplification and purification. Such techniques are thoroughly explained in the literature.

[0186] The constructs provided herein may include a promoter operably linked to any one of the polynucleotides described herein. The promoter may be a heterologous promoter or an endogenous promoter associated with the heterologous gene, nucleic acid, or polypeptide of interest.

[0187] As used herein, the terms “heterologous promoter,” “promoter,” “promoter region,” or “promoter sequence” refer generally to transcriptional regulatory regions of a gene, which may be found at the 5' or 3' side of the polynucleotides described herein, or within the coding region of the polynucleotides, or within introns in the polynucleotides. Typically, a promoter is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. The typical 5' promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of

bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence is a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

[0188] In some embodiments, the heterologous polynucleotides of interest are operably connected to the promoter. As used herein, a polynucleotide is “operably connected” or “operably linked” when it is placed into a functional relationship with a second polynucleotide sequence. For instance, a promoter is operably linked to a polynucleotide if the promoter is connected to the polynucleotide such that it may effect transcription of polynucleotides. In various embodiments, the polynucleotides may be operably linked to at least 1, at least 2, at least 3, at least 4, at least 5, or at least 10 promoters.

[0189] Heterologous promoters useful in the practice of the present invention include, but are not limited to, constitutive, inducible, temporally-regulated, developmentally regulated, chemically regulated, tissue-preferred and tissue-specific promoters. The heterologous promoter may be a plant, animal, bacterial, fungal, or synthetic promoter. Suitable promoters for expression in plants include, without limitation, the 35S promoter of the cauliflower mosaic virus, ubiquitin, tCUP cryptic constitutive promoter, actin, the Rsyn7 promoter, pathogen-inducible promoters, the maize In2-2 promoter, the tobacco PR-1a promoter, glucocorticoid-inducible promoters, estrogen-inducible promoters and tetracycline-inducible and tetracycline-repressible promoters. Other promoters include the T3, T7 and SP6 promoter sequences, which are often used for in vitro transcription of RNA. Those of skill in the art are familiar with a wide variety of additional promoters for use in various cell types. In some embodiments, the heterologous promoter includes a plant promoter, either endogenous to the plant host or heterologous.

[0190] Vectors including any of the constructs or polynucleotides described herein are provided. The term “vector” is intended to refer to a polynucleotide capable of transporting another polynucleotide to which it has been linked. In some embodiments, the vector may be a “plasmid,” which refers to a circular double-stranded DNA loop into which additional DNA segments may be ligated. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors can be integrated into the genome of a host cell upon introduction into the host cell,

and thereby are replicated along with the host genome, such as some viral vectors or transposons. Plant mini-chromosomes are also included as vectors. Any suitable vector design known in the art may be used with the explants of the present invention.

[0191] Vectors may carry genetic elements, such as those that confer resistance to certain drugs or chemicals. In some embodiments, the vector will additionally include one or more selectable or screenable markers. The selectable or screenable marker may confer upon the plant tissue resistance to an otherwise toxic compound. A number of screenable or selectable marker are known in the art and can be used in the present invention. The screenable marker may be fluorescent (e.g., RFP) or non-fluorescent (e.g., GUS). More than 20 selectable marker genes have been reported in the transformation of higher plants (Komari T, Takakura Y, Ueki J, Kato N, Ishida Y, Hiei Y (2006) Binary vectors and super-binary vectors. In: KanWang (ed.), and *Methods in Molecular Biology*, vol. 343: *Agrobacterium* Protocols, Vol. 1, Second Edition. Humana Press Inc., Totowa, N.J., pp. 15-41). In some embodiments, the selectable or screenable marker is selected from the group consisting of RFP, GUS, aadA, spectinomycin, streptomycin, and imazapyr. In some embodiments, the vector is a DICOTBINARY-19 plasmid. In some embodiments, the vector is a DICOTBINARY-22 plasmid.

[0192] In some aspects, provided herein are methods for transforming *Cannabis* using floral dip transformation. Female *Cannabis* flowers are exposed to *Agrobacterium* cultures suitable for flower dip transformation. In some embodiments, Cannabis flowers are submerged in an *Agrobacterium* culture under vacuum to induce transformation. In some embodiments, the *Agrobacterium* culture comprises *Agrobacterium* comprising the heterologous gene or nucleic acid of interest, a wetting agent, and a carrier. In some embodiments, the heterologous gene or nucleic acid of interest is included on a vector such as the DICOTBINARY-22 vector described herein. Other suitable vectors are known in the art. In some embodiments, the *Agrobacterium* strain is Ar18r12v, although other suitable *Agrobacterium* strains are known and used in the art. In some embodiments, the *Agrobacterium* used is a constitutively active variant or virG mutant of *Agrobacterium* (e.g., N45D mutant *Agrobacterium*). In some embodiments, the carrier is a cell culture medium suitable for the survival of the *Agrobacterium*. In some embodiments, the carrier is 5% sucrose. In some embodiments, the wetting agent is 0.05% silwet L-77. In some embodiments, the *Agrobacterium* culture includes Triton X-100, NP-40, Tween, or combinations thereof. In some embodiments, the *Agrobacterium* culture additionally comprises acetosyringone, galacturonic acid, cinnamic acid, coumarin, vanillin, other phenolic compounds, or combinations thereof. Suitable surfactants for plant floral dip transformations are known in the art. Following application of the *Agrobacterium* culture to the female *Cannabis* flower, the flowers are pollinated with male pollen from a suitable donor plant. In some embodiments, female *Cannabis* flowers are exposed to the *Agrobacterium* culture for at least 1, at least 2, at least 5, at least 10, at least 12, at least 15, at least 16, or at least 18 days before pollination. In some embodiments, the female *Cannabis* flowers are pollinated using a paintbrush.

[0193] In some embodiments of the floral dip transformations, DNA is delivered directly to the *Cannabis* ovules. In

some embodiments, DNA is complexed with cell penetrating peptides prior to administration to *Cannabis* ovules.

[0194] In some aspects, provided herein are methods for transforming *Cannabis* nodes, internodes, leafs, petioles, hypocotyls, and buds. Sanitized and imbibed seeds are plated on non-selective medium (e.g., B5 medium) and grown for approximately 6 weeks or until explants suitable for transformation are formed. Resulting explants are inoculated with a heterologous gene or nucleic acid of interest. In some embodiments, the explants are inoculated using a force treatment as described herein. In some embodiments, the explants are inoculated by sonication with a vector comprising the heterologous gene or nucleic acid of interest. In some embodiment, the vector additionally comprises a selectable markers.

[0195] Following inoculation, node and bud explants are co-cultured in a culture medium that supports growth and survival of the node or bud for at least about 4 days. In some embodiments, the culture medium that supports the growth and survival of the node or bud is WCIC INO medium. In some embodiments, the medium additionally comprises nystatin, TBZ, and meta-topolin (mT). Following co-culture of at least about 4 days, the nodes and buds are transferred to a second culture medium suitable for the growth and survival of the nodes or buds. In some embodiments, the second culture medium is hemp node medium described herein in Table 7. In some embodiments, the hemp node medium additionally comprises a selection agent. In some embodiments, the hemp node medium additionally comprises activated charcoal.

[0196] Following inoculation, internode, leaf, hypocotyl, and petiole explants are co-cultured in a culture medium that supports growth and survival of the internode, leaf, hypocotyl, or petiole for at least about 4 days. In some embodiments, the culture medium that supports the growth and survival of the internode, leaf, hypocotyl, or petiole is WCIC INO medium. In some embodiments, the medium additionally comprises nystatin, TBZ, meta-topolin (mT), naphthalacetic acid (NAA) and GA3. Following co-culture for at least about 4 days, leaf, petiole, hypocotyl, and internode explants are transferred to a second culture medium suitable for the growth and survival of the leaf, petiole, or internode. In some embodiments, the second culture medium is hemp internode medium described herein in Table 12. In some embodiments, the medium additionally includes a selection agent.

[0197] In some aspects, provided herein are methods for transforming *Cannabis* pollen or anther cultures. *Cannabis* pollen is harvested by shaking branches of male plants and collecting the pollen. In some embodiments, after harvest and collection, the pollen may be sized by passing the pollen through a sieve (e.g., a #80 sieve). Pollen may be used immediately for transformation, or may be stored prior to use. Pollen may be stored at a temperature between about 4° C. and about 20° C. The pollen may be stored in the presence of a storage medium. In some embodiments, the storage medium is medium suitable for pollen germination. In some embodiments, the storage medium includes boric acid, calcium chloride, potassium phosphate, and water. In some embodiments, the storage medium additionally includes glycerol.

[0198] Pollen may be transformed using particle bombardment, high velocity microprojection, microinjection, electroporation, direct DNA uptake, cell-penetrating peptides,

silica carbide fibers, nanoparticles, and bacterially-mediated transformation. In some embodiments, pollen is transformed using silica carbide fibers. In some embodiments, pollen is transformed using cell-penetrating peptides. Following transformation, pollen is co-cultured and germinated on medium suitable for the survival of the pollen. The medium may include a suitable selection agent. In some embodiments, transformed pollen is used to pollenate female flowers for rapid generation of transgenic T1 progeny.

[0199] In some aspects, provided herein are methods for transforming Cannabis callus tissue or embryogenic suspension cells. In general, Cannabis leaf tissue is cultured on plant medium containing hormones suitable to induce embryogenic calli formation. Embryogenic calli are then transformed using any of the suitable transformation methods as described herein. Calli are then grown a suitable plant medium containing hormones suitable for induction of shooting in the plant. Suspension cells may similarly be transformed in suspension as single cells then subsequently grown into plants using suitable medium including suitable hormones. Suitable medium and hormones are known and described in the art.

[0200] The present invention has been described in terms of one or more preferred embodiments, and it should be appreciated that many equivalents, alternatives, variations, and modifications, aside from those expressly stated, are possible and within the scope of the invention.

Example 1

[0201] The embodiment described here demonstrates *Cannabis* meristem explant transformation.

[0202] Seeds of variety Elektra x Chardonnay were surface sanitized with 20% Clorox for 5 minutes, rinsed, and sat for ~2 hrs before overnight imbibition with WCIC Bean Germination Media (BGM).

TABLE 1

WCIC Bean Germination Media (BGM)	
Ingredients and Notes	Amount to add per liter (grams)
Phytotechnology Laboratories WPM L449	2.41
Sucrose	20
pH to 5.8 with 1N KOH and autoclave	
Add the following prior to use:	
Captan fungicide (50WP)	0.06
Bravo fungicide (Daconil) (82DP)	0.03
Cefotaxime (100 mg/ml)	1.25 ml

[0203] The next day, meristem explants were aseptically excised from seed and incubated for approximately 2 hrs in 20% PEG4000 with 60 mg/L Captan and 30 mg/L Bravo fungicides. Explants were then rinsed and inoculated with Agrobacterium strain Ar18r12v harboring the binary plasmid DICOTBINARY-19. During inoculation, explants were exposed to 20 seconds of sonication at ~45 kHz. Explants

were co-cultured in either 1.75 ml or 2.0 ml WCIC INO media with 100 uM acetosyringone, 50 mg/L nystatin, 10 mg/L TBZ, and 95 uM lipoic acid for 4 days at 23 C 16/8 photoperiod. Thidiazuron (TDZ) was added to some co-culture treatments at 1 mg/L.

TABLE 2

WCIC INO media	
Ingredients and Notes	Amount to add per liter (grams)
Gamborg B5 Phytotechnology Laboratories G398	1.284
Glucose	30
MES	2.8
pH to 5.4 with 1N KOH and autoclave	

[0204] After 4 days of co-culture, transient GUS expression was evaluated in explants. Explants were then transferred to either 10 mg/L spectinomycin or 150 mg/L spectinomycin WCIC Gamborg B5 medium (Table 3) for selection. Explants on 10 mg/L spectinomycin B5 were transferred to 150 mg/L spectinomycin B5 approximately 1 week later, and then explants from all treatments were transferred to 50 mg/L spectinomycin B5 1 month later. Additional transfers have been made with explants remaining green on spectinomycin. A summary of this experiment is given in Table 4.

TABLE 3

WCIC Gamborg B5 Medium	
Ingredients and Notes	Amount to add per liter (grams)
Phytotechnology Laboratories B5 salts G398	2.41
Sucrose	20
Cleary's 3336 (50WP)	0.06
Ca Gluconate	1.29
pH to 5.8 with 1N KOH	
Phytigel	3.50
autoclave	
Add the following fresh before use:	
Timetin (150 mg/ml stock)	Use 1 mL per Liter (150 mg/L)
Cefotaxime (100 mg/ml stock)	Use at 2 ml per Liter (200 mg/L)
Carbenicillin (100 mg/ml stock)	Use at 4 ml per Liter (400 mg/L)
Selective Agent	as needed

TABLE 4

Description and summary of <i>Cannabis</i> meristem explant transformation experiments.							
Co-Culture conditions	Experiment ID	# embryos to Selection	Selection Media	2nd media	3rd media	Notes	Notes
Filter paper in plantcon with 1.75 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid; 23 C. 16/8 photoperiod	Hemp 3/22-1A	7	10 ppm spec B5; 28 C. 16/8 photoperiod;	150 ppm spec B5; 28 C. 16/8 photoperiod;	50 ppm spec B5; 28 C. 16/8 photoperiod;	1 greening explant transferred to non-selective BRM	
Filter paper in plantcon with 1.75 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid; 23 C. 16/8 photoperiod	Hemp 3/22-1B	4	150 ppm spec B5; 28 C. 16/8 photoperiod;	NA	50 ppm spec B5; 28 C. 16/8 photoperiod;	1 greening explant transferred to non-selective BRM	
Filter paper in plantcon with 2 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid; 23 C. 16/8 photoperiod	Hemp 3/22-2A	9	10 ppm spec B5; 28 C. 16/8 photoperiod;	150 ppm spec B5; 28 C. 16/8 photoperiod;	50 ppm spec B5; 28 C. 16/8 photoperiod;	1 greening explant transferred to 10 ppm spec B5	
Filter paper in plantcon with 2 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid; 23 C. 16/8 photoperiod	Hemp 3/22-2B	10	150 ppm spec B5; 28 C. 16/8 photoperiod;	NA	50 ppm spec B5; 28 C. 16/8 photoperiod;	1 chimeric RFP+ explant (primary leaves) transferred to B5	1 greening explant transferred to non-selective BRM
Filter paper in plantcon with 1.75 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 3/22-3A	9	10 ppm spec B5; 28 C. 16/8 photoperiod;	150 ppm spec B5; 28 C. 16/8 photoperiod;	50 ppm spec B5; 28 C. 16/8 photoperiod;		
Filter paper in plantcon with 1.75 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 3/22-3B	10	150 ppm spec B5; 28 C. 16/8 photoperiod;	NA	50 ppm spec B5; 28 C. 16/8 photoperiod;	1 chimeric RFP + explant (primary leaves) transferred to B5	4 greening explant transferred to non-selective BRM; GUS positive leaf imaged
Filter paper in plantcon with 2 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 3/22-4A	9	10 ppm spec B5; 28 C. 16/8 photoperiod;	150 ppm spec B5; 28 C. 16/8 photoperiod;	50 ppm spec B5; 28 C. 16/8 photoperiod;	RFP positive shoot imaged	
Filter paper in plantcon with 2 ml INO + 50 ppm nystatin + 10 ppm TBZ + lipoic acid + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 3/22-4B	12	150 ppm spec B5; 28 C. 16/8 photoperiod;	NA	50 ppm spec B5; 28 C. 16/8 photoperiod;	5 greening explants transferred to 10 ppm spec B5	

[0205] Stable RFP (tdTomato) was imaged several explants 3 weeks after inoculation, but was confined primarily to primary leaves and cotyledonary remnants. The stable RFP signal in the chimeric plantlet that appeared to be from new growth was from treatment 4A (treatment bolded in Table 4—2 ml co-culture volume with TDZ, and initial transfer to 10 mg/L spectinomycin B5), shown again in FIG. 1. This plant did not survive tissue culture, which may have been result of its chimerism in response to selection, or possibly non-optimal media conditions.

[0206] We have also observed stable GUS expression in a plantlet derived from this experiment in treatment Hemp 3/22-3B. One of the leaves in a plantlet transferred to non-selective BRM expressed GUS in leaf stably. The leaf shown in FIG. 16 was imaged 2 months after inoculation after clearing with 70% ethanol.

[0207] The non-inoculated control meristem explant was sent to the greenhouse as a proof of concept of tissue culture (TC) regeneration of a plant from a meristem explant (rooted on non-selective B5 media). See FIG. 2.

[0208] Additional experiments with additional varieties of *Cannabis* inoculated with Ar18r12v/DICOTBINARY-19 and a 4 day co-culture are outlined in Table 5. The co-culture volume was increased as we noted in the prior tests explants were very dry post co-culture. Some of these experiments used meta-topolin in co-culture, which has been demonstrated to encourage propagation in *Cannabis* nodal cultures (H. Lata et al./Journal of Applied Research on Medicinal and Aromatic Plants 3 (2016) 18-26); some used a full-strength formulation of B5 media for selection; and some used an MS-based selection media with meta-topolin (mT) based on Lata 2016 but without activated charcoal.

TABLE 5

Description and summary of follow-up experiments with <i>Cannabis</i> meristem explants.							
<i>Cannabis sativa</i> Genotype/ Line		Experi- ment ID	Strain	Binary	# embryos to Selection	Notes	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 5/9-1	Ar18r12v	DICOT- BINARY- 19	35	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm meta-topolin (mT); 23 C. 16/8 photoperiod	Hemp 5/9-2	Ar18r12v	DICOT- BINARY- 19	25	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Fiber Hemp	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 5/9-3	Ar18r12v	DICOT- BINARY- 19	21	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Fiber Hemp	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm meta-topolin (mT); 23 C. 16/8 photoperiod	Hemp 5/9-4	Ar18r12v	DICOT- BINARY- 19	12	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 5/16-1	Ar18r12v	DICOT- BINARY- 19	59	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper on top of semisolid INO (8 g/L agarose I) + 60 ppm Cleary's + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 5/16-2	Ar18r12v	DICOT- BINARY- 19	19	10 ppm spec 100% B5; 28 C. 16/8 photoperiod;	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 5/30-1	Ar18r12v	DICOT- BINARY- 19	5	non- selective hemp node media (minus activated charcoal)	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 5/30-2	Ar18r12v	DICOT- BINARY- 19	36	10 ppm sec hemp node media (minus activated charcoal)	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 5/30-3	Ar18r12v	DICOT- BINARY- 19	43	50 ppm sec hemp node media (minus activated charcoal)	
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/6-1	Ar18r12v	SOYTEST-2	36	50 ppm sec hemp node media (minus activated charcoal)	

TABLE 5-continued

Description and summary of follow-up experiments with <i>Cannabis</i> meristem explants.						
<i>Cannabis sativa</i> Genotype/ Line	Comments	Experi- ment ID	Strain	Binary	# embryos to Selection	Notes
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 2 min 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/6-2	Ar18r12v	SOYTEST-2	36	50 ppm sec hemp node media (minus activated charcoal)
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/13-1	Ar18r12v	SOYTEST-2	44	50 ppm sec hemp node media (minus activated charcoal)
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 2 min 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/13-2	Ar18r12v	SOYTEST-2	45	50 ppm sec hemp node media (minus activated charcoal)
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-1A	Ar18r12v	SOYTEST-2	20	10 ppm sec B5; 28 C. 16/8 photocopied;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-1B	Ar18r12v	SOYTEST-2	10	10 ppm sec node-AC; 28 C. 16/8 photocopied;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-1C	Ar18r12v	SOYTEST-2	10	50 ppm sec B5; 28 C. 16/8 photocopied;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-1D	Ar18r12v	SOYTEST-2	10	50 ppm strep node-AC; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 1 ppm TDZ, 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-2A	Ar18r12v	SOYTEST-2	10	10 ppm spec B5; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 1 ppm TDZ, 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-2B	Ar18r12v	SOYTEST-2	10	10 ppm spec node-AC; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 1 ppm TDZ, 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-2C	Ar18r12v	SOYTEST-2	10	50 ppm spec B5; 28 C. 16/8 photoperiod;

TABLE 5-continued

Description and summary of follow-up experiments with <i>Cannabis</i> meristem explants.						
<i>Cannabis sativa</i> Genotype/ Line	Comments	Experiment ID	Strain	Binary	# embryos to Selection	Notes
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.5 ml INO + 1 ppm TDZ, 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ; 23 C. 16/8 photoperiod	Hemp 6/20-2D	Ar18r12v	SOYTEST-2	10	50 ppm strep node-AC; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.25 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/27-1	GV3101	SOYTEST-2	48	50 ppm spec meristem regeneration; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.25 ml INO + 60 ppm Cleary's + 50 ppm nystatin + 10 ppm TBZ + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 6/27-2	GV3101	SOYTEST-2	36	50 ppm spec meristem regeneration with 0.5 ppm MT; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper in plantcon with 2.25 ml INO + 60 ppm Cleary's + 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 7/3-1	Ar18r12v	SOYTEST-2	83	10 ppm spec meristem regeneration; 28 C. 16/8 photoperiod;
Honey Gold 3WS	Hand excised from seed surface sanitized in 20% Clorox 5 min; rinsed; imbibed for ~20 h in BGM at 37 C.; explants placed in 20% PEG4000 with Captan/Bravo for 1.5-2.5 h, rinsed, inoculated and sonicated 20 s 45 kHz; incubated 30 min, inoculum removed; explants co-cultured on filter paper on semisolid INO (8 g/L agarose I) + 60 ppm Cleary's 1 ppm TDZ; 23 C. 16/8 photoperiod	Hemp 7/8-1	Ar18r12v	SOYTEST-2	48	50 ppm spec meristem regeneration with 0.5 ppm mT; 28 C. 16/8 photoperiod;

TABLE 6

WCIC TRUE Gamborg B5 Media	
Ingredients and Notes	Amount to add per liter (grams)
Phytotechnology Laboratories B5 salts G398	3.21
Sucrose	20
Cleary's 3336 (50WP)	0.06
Ca Gluconate	1.29
pH to 5.8 with 1 N KOH	
Phytigel	3.50
autoclave	
Add the following fresh before use:	
Timetin (150 mg/ml stock)	Use 1 mL per Liter (150 mg/L)
Cefotaxime (100 mg/ml stock)	Use at 2 ml per Liter (200 mg/L)
Carbenicillin (100 mg/ml stock)	Use at 4 ml per Liter (400 mg/L)
Selective Agent	as needed

TABLE 7

WCIC Hemp Node Media (modified from Lata 2016)	
Ingredients and Notes	Amount to add per liter (grams)
MS Salts complete with vitamins (PhytoTech M519)	4.43

TABLE 7-continued

WCIC Hemp Node Media (modified from Lata 2016)	
Ingredients and Notes	Amount to add per liter (grams)
Sucrose	30
Cleary's 3336	0.06
pH to 5.7 with 1 N KOH	
Agar (Sigma A7921)	8
autoclave	
Meta-topolin (mT) (1 mg/ml)	0.5 ml
Carbenicillin (200 mg/ml)	1.25 ml
Cefotaxime (100 mg/ml)	2 ml
Selection	as needed

[0209] We did obtain strong GUS transients using both *Cannabis* varieties Honey Gold 3WS and the Fiber Hemp (in addition to the transients shown for Elektra x Chardon-nay in initial disclosure). See FIG. 3.

[0210] Additionally, *Cannabis* seeds were also imbibed at 37° C. to facilitate excision of the meristem explants. With an overnight imbibition at 37° C. the radical begins to emerge from the seed which makes a natural crack in the hard seed coat. This makes isolating the *Cannabis* mature embryo/meristem explant easier. See FIG. 17.

[0211] It is also possible to mechanically excise the *Cannabis* meristem explants from the seen. 10 g *Cannabis* seed from the Fiber variety were surface sterilized with 20%

bleach solution for 5 minutes, then rinsed with sterile distilled water for 2 minutes. Seed were then imbibed in sterile distilled water at 37 degrees C. for approximately 24 hours. Seed was then rinsed with sterile distilled water for 2 minutes and weighed (collected rehydrated weight was 18 grams). Seed was then split into two 9 gram samples and spread on sterile filter paper in petri dishes and dried in laminar flow hood. One sample was removed 24 hrs later, the other 49 hours later. The collected dry weight of each sample was 4.8 grams.

[0212] A portion of this dry hemp seed from the 49 hour dried material was used for machine excision experiments. Seed were placed through a Perten Instruments Laboratory Mill 3310 using seven different gap settings (0 to 6, smallest to largest gap) with 20 seeds per gap setting. Ground material was collected and embryonic parts were counted under a microscope. While the Perten Lab mill was used for these examples, excision of embryos and explants can be performed using and dry mill or equivalent instrumentation known in the art, for example, roller mills, hammer mills, and bladed mills or other suitable means described herein. Equivalent wet mill processing is also envisioned.

TABLE 8

Embryonic parts produced using various gap setting on the Perten Instruments Laboratory Mill 3310	
Gap Setting	Embryonic Parts Produced
6	20
5	14
4	14
3	5
2	3
1	4
0	0

[0213] Further experimentation will be carried out using gap setting 6 for grinding hemp varieties. Following embryonic part production, regeneration of the embryonic material will be checked on non-selective medium. Embryonic parts generated using the gap 6 setting on Perten is shown in FIG. 18.

[0214] Additionally, MS-based medium with or without meta-topolin may be used with meristem explants. When TDZ is used in co-culture with *Agrobacterium*, the use of meta-topolin during selection/regeneration may not be necessary.

TABLE 9

Hemp Meristem Regeneration Medium	
Ingredients and Notes	Amount to add per liter (grams)
MS Salts complete with vitamins (PhytoTech M519)	4.43
Sucrose	30
Cleary's 3336	0.06
pH to 5.7 with 1 N KOH	
Agar (Sigma A7921)	8
autoclave	
Meta-topolin (mT) (1 mg/ml)	as needed
Carbenicillin (100 mg/ml)	2 ml
Cefotaxime (100 mg/ml)	2 ml
Timetin (150 mg/ml)	1 ml
Selection	as needed

[0215] Additional control binary constructs have also been testing including SOYTEST-2. SOYTEST-2 has a different promoter driving tdTomato to test impacts on RFP visualization (DICOTBINARY-19 uses the *Glycine max* Ubiquitin 3 XL promoter driving tdTomato, where SOYTEST-2 uses *Pinus radiata* Super Ubiquitin promoter driving tdTomato). See FIG. 19.

[0216] We have obtained positive GUS transients in *Cannabis* meristem explants (3WS variety) using SOYTEST-2 in both the Ar18r12v and GV3101 strains of *Agrobacterium* (see FIG. 20), demonstrating transfection of *Cannabis* meristems using disarmed strains of both *Agrobacterium rhizogenes* (Ar18r12v) and *Agrobacterium tumefaciens* (GV3101).

[0217] We have obtained additional stable RFP expressing *Cannabis* from experiments in the Honey Gold 3WS variety. The plantlet in the center of FIG. 21 (*Cannabis* plant WP421-1) is rooting on 50 mg/L streptomycin hemp node media (after being on 50 mg/L spectinomycin hemp node media for approximately 1 month).

[0218] Plant WP421-1 was transferred to the greenhouse and imaged the day of transfer, approximately 4 weeks later (FIG. 26), and approximately 7 weeks later (FIG. 31). Three leaves from WP421-1 were sampled after it had been in greenhouse for 4 weeks. Two of the three leaves were stably expressing RFP (FIG. 27), while all three were stable expressing GUS (FIGS. 27 and 28). RFP expression was relatively weaker than when we initially sent the plant, which is likely due to tissue age. That all three randomly selected leaves are stably expressing GUS is a good indicator WP421-1 is not overly chimeric.

[0219] Additional data in FIGS. 32-34 confirms the chimerism in the WP421-1 plant. 10 leaf samples were taken from EP421-1 and divided for PCR (FIGS. 33-34) and GUS expression (FIG. 32) analysis. For PCR, we amplified a 156 bp fragment within the aadA expression cassette of DICOTBINARY-19 using primers designated F56 and R11 (fragments and amplicon highlighted in blue in FIG. 33). Leaf DNA from WP421-1 was extracted using the REDExtract-N-Amp™ Plant PCR Kit (Sigma-Aldrich XNAP-1KT) following manufacturer's instructions. PCR reaction was run with following:

[0220] 1. 3 minutes at 94 C for initial denaturation
[0221] 2. 30 seconds at 94 C for denaturation
[0222] 3. 30 seconds at 55 C for annealing
[0223] 4. 1 minute at 72 C for primer extension
[0224] 5. Cycle steps 2-4 34 more times (35 total cycles)
[0225] 6. 10 minutes at 72 C for final primer extension
[0226] PCR products were run on 1.5% agarose gel in SB buffer. All 10 leaf samples gave the expected 156 bp product, confirming minimal chimerism in this event

[0227] We have also blasted DNA into *Cannabis* meristem explants using PDS-1000 Helium gun with a plasmid designated DICOTBOMB-13. For particle bombardment experiments, gold-DNA "bead prep" was prepared by first washing 50 mg 0.6 um gold microcarriers (BioRad part #1652262) in 1 ml 100% ethanol and sonicating for 1 min 45 kHz. Gold was pelleted by centrifugation at 5000 rpm in microfuge (~2300xg) and ethanol removed. Gold was then resuspended in 1 ml 100% ethanol and stored at -20 C until use. To precipitate DNA onto beads, the 50 mg gold/l ml ethanol stock was sonicated for 1 min 45 kHz. 42 ul of this stock was transferred to an Eppendorf tube, then pelleted by centrifugation at 2500 rpm for 10 seconds, after which

ethanol was removed. 500 uL sterile water was added and mixture sonicated 1 min 45 kHz. Gold was again pelleted by centrifugation at 2500 rpm for 10 seconds and water removed. 25 ul sterile water was then added, followed by sonication for 1 min 45 kHz. 2.6 ug DICOTBOMB-13 DNA was added, then sterile water to bring volume up to 245 ul. 250 ul cold 2.5 M CaCl₂ was added, followed by 50 ul 0.1 M spermidine. Solution was mixed by low speed vortexing. Tube was incubated on ice for approximately 1 hour with gentle inversions every 5-10 minutes. DNA/gold was pelleted at 1000 rpm (~100×g) for 2 min and supernatant removed. Pellet was then washed with 1 ml 100% EtOH w/ pipette tip, then pelleted again at 1000 rpm (~100×g) for 2 min and supernatant removed 36 ul 100% EtOH was added

Example 2

[0229] The embodiments described herein demonstrate *Cannabis* floral dip experiments.

[0230] We used strain Ar18r12v based on positive GUS transients in meristem explants, harboring DICOTBINARY-22, which has the aadA1a protein targeted to both plastid and mitochondria in the plants. *Agrobacterium* cultures were resuspended in 5% sucrose with 0.05% silwet L-77 as a wetting agent. One of the cultures was induced with 100 uM acetosyringone and one was not. These cultures were applied directly to female flowers (some flowers received only 5% sucrose+0.05% silwet L-77 “blank”). Experiment is summarized in Table 11.

TABLE 11

Description and summary of experiments with Cannabis floral dip				
Experiment ID	Comments	AS	Co-Culture Duration (Days)	Observations and Comments
Hrmp 4/17-1	Agro washed once with none sterile water; spun 10 min, resuspended in 5% sucrose; Silwet L-77 added to 0.05%		Agro applied directly to flowers; 1d dark in high humidity LEDA; then moved to GH7 with 3 males	5 female plants; 3 hermaphrodites (red/white twist tie below flower for inoculated, white for blank)
Hemp 4/17-2	Agro washed once with 100 uM AS sterile water; spun 10 min, resuspended in 5% sucrose; Silwet L-77 added to 0.05%		Agro applied directly to flowers; 1d dark in high humidity LEDA; then moved to GH7 with 3 males	5 female plants; 3 hermaphrodites (green twist tie below flower for inoculated, white for blank)

to tube and gold completely resuspended with low-speed vortexing. Bead prep was stored at -20 C until used, with 5 ul used per bombardment. This corresponds to 360 ng DNA per blast; 290 ug gold per blast (1.2 ng DNA per ug gold).

[0228] *Cannabis* meristem explants were excised from seed and precultured on EJW1 media overnight at 28 C 16/8 photoperiod, arranged on 12% xanthan gum targeting plates (with 60 mg/L Cleary's 3336 fungicide), blasted at 6 cm from the launch assembly using 5 uL bead prep per target and a range of rupture disks (650 psi-1550 psi), then allowed to rest on EJW1 media overnight. Transients were taken the next day. GUS transient expression in *Cannabis* meristem explants bombarded with DICOTBOMB-13 is shown in FIG. 23.

TABLE 10

Preculture Medium EJW1	
Ingredients and Notes	Amount to add per liter (grams)
MS salts no vitamins	4.3
Sucrose	30
2,4-D (1 mg/ml stock)	0.2 ml
MES	2
Cleary's 3336	.03
pH	5.6
Agarose	4
Autoclave	
Carbenicillin (100 mg/ml)	.25
TDZ (1 mg/ml stock)	1 ml

[0231] We noted no obvious ill effects of the inoculum application to the flowers one-day post-application. See FIG. 5. Female flowers were then pollinated with male pollen from donor plants by dusting the plants with pollen; as well as using a paint brush to apply pollen earlier.

Example 3

[0232] The embodiments described herein demonstrate *Cannabis* node, internode, leaf, and petiole transformation experiments.

[0233] We inoculated node, internode, leaf, and petiole explants of *Cannabis* variety Elektra x Chardonnay. We used aseptically grown plantlets from sanitized and imbibed seed plated on non-selective B5 media for approximately 6 weeks. Explants were sonicated for 20 seconds at ~45 kHz in the presence of Ar18r12v/DICOTBINARY-19. Nodal explants were co-cultured on 2.5 ml WCIC INO media with 50 mg/L nystatin, 10 mg/L TBZ, and 0.5 mg/L meta-topolin (mT). Internode, leaf, and petiole explants were co-cultured on 2.5 ml WCIC INO media with 50 mg/L nystatin, 10 mg/L TBZ, 1 mg/L meta-topolin (mT), 1 mg/L naphylacetic acid (NAA), and 0.2 mg/L GA3. After 4 days of co-culture at 23 C 16/8 photoperiod, GUS transient expression was observed in all explant types. See FIG. 6.

[0234] After co-culture, nodal explants were transferred to 100 mg/L spectinomycin hemp node media (Table 7) supplemented with 500 mg/L activated charcoal. Leaf, petiole, and internode explants were transferred to 100 mg/L spectinomycin hemp internode media, which is a modification of potato ZIG media (Cearley J A, Bolyard M G: Regeneration of *Solanum tuberosum* cv. Katandin from leaf explants in

vitro. Am Potato J 74: 125-129 (1997)). A summary of these experiments is provided in Table 13.

TABLE 12

WCIC Hemp Internode Media (modified from Cearley 1997)	
Ingredients and Notes	Amount to add per liter (grams)
MS Salts complete with vitamins (PhytoTech M519)	4.43
Sucrose	20
Cleary's 3336	0.06
pH to 5.7 with 1 N KOH	
Gelrite (or Phytigel), then autoclave	2
Meta-topolin (mT) (1 mg/ml)	1 ml
Naphthylacetic acid (NAA) (1 mg/ml)	1 ml
GA3 (FS) Sigma Prod G7645 (1 mg/ml)	0.2 ml
Carbenicillin (200 mg/ml)	1.25 ml
Cefotaxime (100 mg/ml)	2 ml
Selection	as needed

pollen from male plants can be stored. *Cannabis* pollen was harvested in by shaking branches of male plants onto a creased sheet of blue paper (high visibility of yellow on blue). Pollen was poured through a #80 sieve into a 50 mL conical tube. A small, visible amount of pollen was added to 7 1.5 mL microfuge tubes comprising the following treatments:

- [0237] 1. 25% glycerol (w/v) PGM, inverted several times to mix and stored at -20 C.
- [0238] 2. 25% glycerol (w/v) PGM, inverted several times to mix and stored at 4 C
- [0239] 3. Untreated, fresh pollen, stored at 4 C
- [0240] 4. Untreated pollen, stored at -20 C
- [0241] 5. Pollen placed in 1.5 mL microfuge tube, placed in sweater box with MgNO3 solution. Moved to -20 C on 5/3 at 11 am.
- [0242] 6. 25% w/g glycerol PGM, 1 hour incubation at RT, followed by storage at -80 C
- [0243] 7. One hour untreated at room temperature in 50 mL tube.

TABLE 13

Description and summary of experiments with Cannabis nodes, leaves, petioles, and internodes.					
Experiment ID	Comments	# embryos to Selection	Notes	# explants to second media	2nd media
Hemp 5/2-1	Nodes from hemp seed sanitized and imbibed; germinated on B5 3/22/19; harvested in INO; inoculated and sonicated 20s 45 kHz; co-cultured in plantcons with filter paper + 2.5 ml INO with 50 ppm nystatin +10 ppm TBZ + .5 ppm meta-topolin (mT); 23 C. 16/8 photoperiod	11	100 ppm spec hemp node media; 28 C. 16/8/photoperiod	NA	NA
Hemp 5/2-2	Leaves from hemp seed sanitized and imbibed; germinated on B5 3/22/19; harvested in INO; inoculated and sonicated 20s 45 kHz; co-cultured in plantcons with filter paper + 2.5 ml INO with 50 ppm nystatin + 10 ppm TBZ + 1 ppm meta-topolin (mT) + 1 ppm NAA + 0.2 ppm GA3; 23 C. 16/8 photoperiod	68	100 ppm spec hemp internode media; 28 C. 16/8/photoperiod	18	non selective hemp node media; 28 C. 16/8/photoperiod
Hemp 5/2-2	Petioles from hemp seed sanitized and imbibed; germinated on B5 3/22/19; harvested in INO; inoculated and sonicated 20s 45 kHz; co-cultured in plantcons with filter paper + 2.5 ml INO with 50 ppm nystatin + 10 ppm TBZ + 1 ppm meta-topolin (mT) + 1 ppm NAA + 0.2 ppm GA3; 23 C. 16/8 photoperiod	68	100 ppm spec hemp internode media; 28 C. 16/8/photoperiod	3	non selective hemp node media; 28 C. 16/8/photoperiod
Hemp 5/2-2	Internodes from hemp seed sanitized and imbibed; germinated on B5 3/22/19; harvested in INO; inoculated and sonicated 20s 45 kHz; co-cultured in plantcons with filter paper + 2.5 ml INO with 50 ppm nystatin + 10 ppm TBZ + 1 ppm meta-topolin (mT) + 1 ppm NAA + 0.2 ppm GA3; 23 C. 16/8 photoperiod	68	100 ppm spec hemp internode media; 28 C. 16/8/photoperiod	3	non selective hemp node media; 28 C. 16/8/photoperiod

Example 4

[0235] *Cannabis* meristem explants were excised, dried, and stored at -20 C. *Cannabis* meristem explants of variety 3WS were excised from seed and then dried on the surface of filter paper in a laminar flow hood for 26 hours. Dried *Cannabis* meristem explants were then stored at -20 C for 3 days. Explants were then rehydrated in 20% PEG4000 with 60 mg/L Captan and 30 mg/L Bravo fungicides, rinsed, and plated on Hemp Node media without activated charcoal (Table 7).

Example 5

[0236] This embodiment describes the transformation of *Cannabis* pollen, in particular the advantages provided when

[0244] The same day pollen was harvested, pollen from tubes #6,7 were placed onto solid PGM after one hour of incubation at room temperature. Plates were then wrapped in parafilm and placed in dark room and examined under microscope for germination in the form of pollen tube expansion. Germination of tubes was visible on both. Tube emerged more quickly from pollen from tube #6, which may be due to the liquid PGM matrix. Germination continue onto the next day, 22 hours after plating. See FIG. 24.

[0245] To test for overnight storage, we assayed pollen for germination the next day. Pollen from tubes 1,4 and 6 were placed on ice to thaw. After half an hour, a small amount of pollen from each was added to a PGM plate partition. 50 uL of pollen was taken from 1 & 6. Tube 6 was noticeably more opaque than tube 1. Tube germination was visible only in

treatment #1. Multiple, branched tube-like structures were visible in treatment #4 at T2 (overnight storage) which may have been aberrant germination or the first indications of fungal hyphae. Following 24H of incubation, tube germination was visible in the -20° C. stored pollen that was stored in 25% (w/v) glycerol PGM. No survival was observed in either -80° C. treatment, which perhaps indicates that a slower rate of freezing is important for retaining viability. See FIG. 35.

TABLE 14

Pollen Germination Medium - Salts II Solution (PGM2 Salts)				
Pollen Germination Medium Salts II Solution (PGM2 Salts)				
Stock (100X) Reagent	Volume: 10 mL CAS	g/10 mL	Original Units	100 mL stock grams added
Boric Acid (H ₃ BO ₃)	0043-35-3	0.05	0.005%	0.5
CaCl ₂ *2H ₂ O	10035-04-8	1.47	10 mM	14.7
Potassium Phosphate, monobasic (KH ₂ PO ₄)	7778-77-0	0.00680	0.05 mM	0.068
ddH ₂ O (dissolve salts)		Add 8 mL		Add 80 mL
dissolve by stirring and/or sonication				
ddH ₂ O		Fill to 10 mL w/ graduated cylinder		Fill to 100 mL w/ graduated cylinder

TABLE 15

2X PGM (Pollen Germination Medium)		
2X Pollen Germination Medium		
Volume	1 L	
	amount	units
PGM Salts	10 mL	
Sucrose	100 g	
PEG 4000	600 g	
ddH ₂ O	Fill to 1 L	
Heat to 70 C. for 10 minutes on stir plate.		
Filter sterilize.		
Store at 4 C.		

TABLE 16

Solid Pollen Germination Medium		
Solid Pollen Germination Medium		
Volume	1 L	
	amount	units
2X PGM	500 mL	
0.6% Noble Agar	500 mL	

TABLE 17

25% glycerol PGM		
25% glycerol PGM		
Volume	1 L	
	amount	units
2X PGM	500 mL	
50% (w/y) glycerol	500 mL	

Example 6

[0246] *Cannabis* value added explants (VAEs) were generated by hand excising meristem tissue from seeds surface sanitized in 20% Clorox for 5 min, rinsed, and imbibed for ~20 h in BGM. *Cannabis* VAEs of variety 3WS were excised from seed, then dried on the surface of a filter paper in a laminar flow hood for 26 hours. Dried *Cannabis* meristem explants were then stored at -20° C for 3 days.

Explants were then rehydrated in 20% PEG4000 with 60 mg/L Captan and 30 mg/L Bravo fungicides, rinsed, and plated on Hemp Node media without activated charcoal. FIG. 29 shows dried *Cannabis* VAEs after 5 weeks plated on non-selective medium. FIG. 30 shows machine excised *Cannabis* meristem explants on non-selective medium for 2 weeks.

[0247] *Cannabis* VAEs may also be primed with WCIC INO or other medium prior to drying. Examples of priming and drying conditions are outlined below. These examples are for soybean transformations but are applicable to *Cannabis* VAE priming and drying and the same or similar conditions may be used with *Cannabis* explants.

TABLE 18

Priming and Drying of Explants (Soybean genotype W28, strain GV3101)					
VAE Batch/Priming	Co-culture	Ex-plants	Shoots har-vested	T0 plants to GH	TF
Fresh explants giving rise to S92 series	1 ppm TDZ	75	6	3	4.0%
Fresh explants giving rise to S92 series	no TDZ	71	3	3	4.2%
S92A (std dry)	1 ppm TDZ	75	8	4	5.3%
S92B (3 hr prime in INO, then dry)	1 ppm TDZ	50	10	10	20.0%
S92C (3 hr prime in INO with 1 ppm TDZ, then dry)	1 ppm TDZ	75	9	6	8.0%
S92D (3 hr prime in INO with 10 ppm TDZ, then dry)	1 ppm TDZ	50	0	0	0.0%
S92E (3 hr prime in INO with 50 ppm TDZ, then dry)	1 ppm TDZ	63	0	0	0.0%

TABLE 18-continued

Priming and Drying of Explants (Soybean genotype W28, strain GV3101)					
VAE Batch/Priming	Co-culture	Ex-plants	Shoots har-vested	T0 plants to GH	TF
S92E (3 hr prime in INO with 100 ppm TDZ, then dry)	1 ppm TDZ	50	0	0	0.0%

Example 7

[0248] This example demonstrates additional embodiments of the transformation methods and transgenic *Cannabis* plants described herein.

[0249] T1 data/POC of germline transformation of *Cannabis* using meristem explants: A leaf sample of *Cannabis* T0 event WP421-1 (plant “Candice”) was taken for long-term storage at -80 for further genetic analysis, and at this time we took sections of the petiole as well for GUS expression analysis. FIG. 36 shows GUS expression in the vascular tissue of WP421-1, which is an indicator of germline status for the transgene.

[0250] WP421-1 was moved to short days in GH15 approximately 9 weeks after plant handoff to greenhouse. 10 cuttings were taken 2 days later. Some cuttings had roots approximately 2 weeks after cuttings were taken. First flowers (female) observed 5 days after moving plant to short days. WP421-1 was move into GH9 2 weeks after being put in short-day to be exposed to pollen from the 3WS 2-5 auto-flowering male. FIG. 37 shows WP421-1 adjacent to the male pollen donor wild-type 3WS plant.

[0251] 2.5 weeks after pollination with male 3WS plant, immature T1 seed of WP421-1 was planted. We noted RFP+ expression (tdTomato) in one of the seedlings 3.5 weeks after this (in plantlet “Carly”), confirming germline transmission of transgene. (FIG. 40)

[0252] We followed this observation up with additional RFP expression (tdTomato) observations (eliminating background in a null segregant and in a wild-type leaf), GUS expression, and an aadA1a PCR that produced the expected 156 bp amplicon. (FIGS. 41 and 42). Approximately 5250 T1 seed were harvested from WP421-1 approximately 2 months after pollination was initiated (FIG. 43). We germinated T1 seed of WP421-1 on germination paper and planted a flat of T1 seed to examine more plants for RFP expression (tdTomato) and segregation. RFP expression (tdTomato) was apparent along with null segregants (FIG. 44)

[0253] Of the 48 T1 WP421-1 seed planted in flats, 44 germinated and were designated WP421-1@3 through WP421-1@46. 22 of these 44 expressed RFP (tdTomato) and GUS, giving the expected 1:1 segregation ratio for this cross. (FIG. 45)

[0254] Additional *Cannabis* transformation experiments using meristem explants: Additional experiments with *Cannabis* meristem explants inoculated with Ar18r12v/DICOT-BINARY-19 using 75 mg/L spectinomycin selection have yielded plantlets chimeric for GUS as well as greening plantlets that are GUS negative. These chimeric plantlets did not root on the subsequent 50 mg/L streptomycin media and were instead transferred to selection-free BRM (FIG. 48). One chimeric GUS+ plant (center photo in FIG. 48) from

this set rooted on selection-free BRM and was sent to the greenhouse as the T0 plant WP421-2, but we could not detect positive GUS or RFP expression or transgene integration by PCR in subsequent tests of WP421-2, possibly indicating negative sectors of the plant outcompeted transgenic sector (FIG. 49).

[0255] We have regenerated a transgenic *Cannabis* plantlet from particle-mediated transformation that is stably expressing GUS in all it leaves. *Cannabis* explants prepared from seed germinated at 37 C overnight of variety 3WS were precultured on EJW1 overnight and blasted on carboxymethylcellulose media using DICOTBOMB-13 (at 1.2 ng DNA/ug gold; 0.6 um gold; 1 cm gap, 6 cm distance, and 1350 psi).

[0256] Explants were rested on EJW1 overnight, then transferred to 100 mg/L spectinomycin hemp node media without activated charcoal for 1 month. Explants were then transferred to 50 mg/L streptomycin hemp regeneration media with 1 mg/L meta-topolin (mT). Leaves were sampled from greening explant and incubated in X-gluc at 37 C. Stable GUS expression was present in every leaf, demonstrating POC of particle-mediated transformation of *Cannabis* meristem explants (FIG. 51).

Example 8

[0257] Transgenic T0 *Cannabis* Plant Generation Using Particle-Mediated Transformation of *Cannabis* Meristem Explants

[0258] The regenerating transgenic *Cannabis* plantlet from particle-mediated transformation stably expressing GUS in all it leaves from last report has rooted and been sent to GH as T0 plant WP-001181-1. This T0 plant was derived from *Cannabis* meristem explants prepared from seed germinated at 37 C overnight of variety 3WS. Meristem explants were precultured on EJW1 overnight and blasted on carboxymethylcellulose media using DICOTBOMB-13 (at 1.2 ng DNA/ug gold; 0.6 um gold; 1 cm gap, 6 cm distance, and 1350 psi). Explants were rested on EJW1 overnight, then transferred to 100 mg/L spectinomycin hemp node media without activated charcoal for 1 month (media previously described). Explants were then transferred to 50 mg/L streptomycin hemp regeneration media with 1 mg/L meta-topolin (mT) for 6 weeks (media previously described). We then attempted to root this explant by transferring it to: (i) 4 weeks on 50 mg/L streptomycin hemp node media after cutting hypocotyl to remove necrotic tissue; (ii) 2 weeks on non-selective B5 media; (iii) 6 weeks on non-selective dicot BRM media (½ MS salts with 0.1 mg/L IAA); (iv) Explant split into two; hypocotyl cut on each piece to remove necrotic tissue; and (v); 4 weeks (plant “A”) to 7 weeks (plant “B”) on non-selective dicot BRM media. We re-assayed leaves for GUS and found all leaves remained GUS+. (FIG. 52).

[0259] After the WP-001181a event had been in the greenhouse for 3 weeks, we used the previously described primer set and PCR conditions to amplify a 156 bp fragment of the aadA1a gene from 10 separate leaves. At the same time, we assayed these leaves for GUS expression (each leaf sample was divided in 2, one sample for PCR, one for GUS assay). All 10 leaves tested positive for aadA1a by PCR, and all 10 were GUS positive (FIG. 53).

[0260] Previous examples described taking petiole sections of the *Cannabis* T0 event WP421-1 to examine GUS expression in vascular tissue, which appeared to be an early

indicator of germline status in the *Cannabis* WP421-1 T0 event, and found GUS expression to be present in vascular tissue of WP001181-1a as well (FIG. 56). The second piece of this explant rooted approximately 3 weeks after the first, and was sent to greenhouse after finding all leaves GUS+ (FIG. 57). The ability to split the event into two T0 plants in the meristem transformation system may offer some downstream advantages. It may be possible to feminize either the WP-001181-1a or WP-001181-1b to make subsequent flow-ers male (process described later). Crosses from -1a onto -1b would then yield homozygotes in the T1 progeny, which may offer timing advantages for transgenic *Cannabis* breeding programs (and advantages to attempting to make hermaphroditic plants for same purpose).

[0261] DNA complexed with gold “bead prep” was made by these steps:

[0262] Sonicate 50 mg/1 ml EtOH gold suspension for 3 min 45-55 kHz to resuspend gold.

[0263] Transfer 42 ul to a new tube. Pellet by centrifugation at 2500 rpm for 10 seconds, remove EtOH.

[0264] Add 500 uL sterile water, sonicate 3 min 45-55 kHz. Pellet by centrifugation at 2500 rpm for 10 seconds, remove water.

[0265] Add 25 ul sterile water (wash sides of tube with pipette tip). Sonicate 3 min 45-55 kHz.

[0266] Add 2.6 ug DNA

[0267] Add cold sterile water to bring volume up to 245 ul.

[0268] Add 250 ul cold 2.5 M CaCl₂.

[0269] Add 50 ul 0.1 M spermidine

[0270] Mix solution by low speed vortexing. Incubate tube on ice for at least 45 min (I usually go 1.5 hrs). Invert tube every 5-10 minutes.

[0271] Once coating is done, pellet the DNA/gold at 1000 rpm (~100xg) for 2 min. Remove supernatant.

[0272] Wash pellet with 1 ml 100% EtOH w/ pipette tip. Pellet DNA/gold at 1000 rpm (~100xg) for 2 min. Remove EtOH.

[0273] Add 36 ul 100% EtOH, completely resuspend gold with low-speed vortexing.

[0274] Store at -20 C, we generally use 5 ul of this prep per bombardment.

[0275] This DNA prep gives a DNA Loading Rate of 1.2 ng DNA/ug gold (360 ng DNA per blast, 290 ug gold per blast).

[0276] CMC targeting media includes 8% low viscosity carboxymethylcellulose, 2% medium viscosity carboxymethylcellulose, and 0.4% washed agar. CMC was made by adding washed agar to water, autoclaving, pouring into blender, adding carboxymethylcellulose, blending, pouring into bottle, then re-autoclaving, then poured into plates or stored.

[0277] EJW1 media is shown above in Table 10.

TABLE 19

BRM media	
Ingredients and Notes	Amount to add per liter (grams)
MS Salts no vitamins M524	2.15
myo-inositol	0.1

TABLE 19-continued

BRM media	
Ingredients and Notes	Amount to add per liter (grams)
sucrose	30
pH 5.8 with KOH	
Agar, Sigma A7921	8
Autoclave	
Add after autoclaving	
Cysteine (100 mg/ml) - make stock fresh	Use at 1.0 ml per Liter (100 mg/L)
Cefotaxime (100 mg/ml)	Use at 2.0 ml per Liter (200 mg/L)
IAA (1 mg/ml)	Use at 0.1 ml per Liter (0.1 mg/L)
MS Vitamins (1000X)	Use at 1.0 ml per Liter
Selection	as needed

[0278] It should also be possible to transform *Cannabis* meristem explants with free DNA methods where the explants do not require targeting (i.e. nanotechnology such as cell penetrating peptides, silica carbide fibers or others; as well as use of lipid and/or cationic lipid compounds).

[0279] Transient transformation of storable *Cannabis* meristem explants excised from seed using automation: We have demonstrated transient expression of GUS in meristem explants of the Fiber Hemp variety using meristem explants excised from seed using automation (described in comments to application sent in July 2019) with H1-H5 fractions pooled. These explants had been stored at -20 C for x weeks before rehydration in INO+60 mg/L Cleary's and inoculation with Ar18r12v/DICOTBINARY-19. Presence of GUS also indicates cells remained viable during this ~8 month storage time. (FIG. 58).

[0280] *Cannabis* seeds can also be crushed to extract mature embryos using a metal rolling pin on a glass plate, with an adjustable gap made using shims (FIG. 59). *Cannabis* seed of the Abacus variety was sanitized and imbibed overnight at 23 C. Seed was then either crushed under the rolling pin, with crushed material allowed to dry in a laminar flow hood for 72 hours; or seed was dried first for 72 hours in laminar flow hood and then crushed. Crushed material was stored at -20 C for 2 months, then rehydrated in INO media with 60 mg/L Cleary's fungicide. Explants were inoculated with Ar18r12v/DICOTBINARY-19 and co-cultured as previously described. We noticed transient GUS expression in meristem explants, again indicating cells were capable of surviving storage as well as maintaining some competency for *Agrobacterium* transformation (FIG. 60). This crushing technology and roller technology can be applied to wet or dry *Cannabis* seed, and can be used in conjunction with other treatments to compromise seed coat (ex. seed placed in paint shaker with a metal bead or beads).

[0281] Transient transformation of elite Abacus *Cannabis* variety using *Agrobacterium*-mediated transformation of meristem explants: We have also demonstrated transient expression of GUS in meristem explants of the elite Abacus variety using meristem explants inoculated with Ar18r12v/DICOTBINARY-19 (derived from seed imbibed overnight at 23 C) (FIG. 61).

[0282] Transmission of transgenes into T2 generation of *Cannabis* 3WS variety (from initial T0 event WP421-1): The previously described GUS positive, RFP (tdTomato) positive, and aadA positive T1 plant WP421-1@2 and cuttings from this plant (WP421-1-08C; WP421-1-10C; WP421-1-07C; WP421-1-02C) were used to examine transmission of transgenes into the T2 generation. This T1 plant and its cuttings were pollinated with pollen from a separate

GUS positive and RFP positive feminized WP421-1 T1 plant. Feminization stresses the female plant out enough that it is making male structures that have female genetic pollen. Seed is feminized from the subsequent crosses (all female). Three main substances can be used: Colloidal Silver Solution (~120 ppm); Silver Thiosulfate/Silver Nitrate (STS); or GA3. These chemicals reduce ethylene production that ordinarily helps ripening and the production of female flowers. Temperature or other stresses can cause feminization as well. This T1 plant was feminized by the following procedure. 5 days before moving plants to short days (i.e., 12 hour light periods), all branches and leaves of the plants were sprayed to saturation with either the STS or Colloidal Silver solution. The spraying was repeated every 5 days until major male flower formation has begun. Usually male flowers begin showing at about 16-20 days after initial spraying and spraying finished after 25-30 days. Resulting T2 seeds were harvested and germinated and imaged for RFP expression. These crosses gave rise to the expected 3:1 segregation ratio (from both parents segregating 1:1), and the 3:1 ratio should include 1 homozygote, 2 hemizygotes and 1 null at T2.

TABLE 20

RFP segregation ratios in T2 Cannabis lines derived from transformed meristem explants.	
Cannabis line	Ratio
WP421-1@2	10/15 positive
WP421-1-08C	13/18 positive
WP421-1-10C	16/18 positive
WP421-1-07C	13/18 positive
WP421-1-02C	12/16 positive
Total	64/85 positives

Example 9

[0283] The reduction or elimination of THC will be accomplished by knocking out the THCA synthase gene using a CRISPR/Cas9 gene editing approach. *Cannabis* has two genes that contribute to THC production, namely THCA synthase (primary) and CBDA synthase (secondary). It is postulated that in high CBD varieties of *Cannabis*, the THCA synthase gene(s) are either inactive or highly suppressed. Since CBDA synthase has about an 84% amino acid similarity to THCA synthase, it is also possible that it also plays a role in producing the low amounts of THC seen in these varieties. The proposed CRISPR approach includes a gRNAs designed to target both the THCA synthase as well as the CBDA synthase resulting in plants with single gene knockouts as well as knockouts for both genes. Plants produced from this editing approach would be analyzed with our UPLC to determine the content of 13 different cannabinoids including THCA, CBDA, and CBG. In general, high CBD varieties of cannabis have a CBDA:THCA ratio of 20:1 to 27:1 and these compounds are tightly related to each other (i.e. as CBDA goes up so does THCA). A potential outcome of this editing approach may also be a plant that is skewed with a higher ratio of CBDA:THCA, again reducing the risk to the grower of cultivating plants producing THC as well as producing more CBDA. Another possible outcome of knocking out both THCA synthase and CBD synthase would be the accumulation of the precursor com-

pound CBG. A plant high in CBG is also a highly valued product due to the fact that the plant would normally make very little CBG and thus is a cannabinoid in higher demand. [0284] A big risk to farmers trying to maximize their CBD amounts in their *Cannabis* plants, is pollination from male plants in the population or from adjacent fields. Pollination of female plants substantially reduces the amount of CBD made by the plants due to the reduction of trichome producing pistils and energy going toward making seed, which is low in CBD. Huang et al. successfully expressed the Solo Dancers and Barnase genes using a fusion gene, which resulted in fully sterile male/female flowers in *Arabidopsis* and tobacco without affecting growth or development. Since *Cannabis* can be easily cloned from a “mother” plant as demonstrated herein, a female sterile plant could be of high value to the grower in eliminating the risk of pollination to the crop.

[0285] Increasing trichome numbers: Cannabinoids are mainly made and secreted in the trichomes of the cannabis plant, and more specifically the trichomes associated with female flowers (pistils) of the plant. Increasing the number of trichomes will increase the total cannabinoid production in the plant. Tian, et al. successfully overexpressed the BraLTP2, a lipid transfer protein from *Brassica napus*, resulting in a 10-fold increase of trichomes in *B. napus*. The transgenic lines also produce elevated levels of 43 different secondary metabolites, which could also prove interesting to the amount and type of cannabinoids made in a transgenic cannabis plant with overexpression of this type of gene. LTPs belong to a large multigene family with many complex physiological functions. We have identified similar genes in the cannabis genome and propose to overexpress both the *B. napus* BraLTP2 as well as some homologues from cannabis. These genes could be driven by either a constitutive or flower/pistil specific promoter to drive these potential outcomes. Transgenic plants would be visually screened for an increase in trichomes as well as tested for cannabinoid content using our UPLC.

[0286] High CBGA plants: The CsPT1 ((geranylpyrophosphate:olivetolate geranyltransferase (GOT)) gene is involved in the production of CBGA, the precursor to THCA and CBDA. By overexpressing this gene, it may be possible to increase the amount of CBGA in the plant resulting in the increase of the downstream cannabinoids CBDA and THCA. This approach could also be used in conjunction with knocking out the THCA synthase gene and ultimately making a very high CBDA plant with little or no THCA.

[0287] Herbicide resistant plants: Like many other plants, cannabis has an EPSP Synthase gene. It is known that changes to 1-3 amino acids in this gene tend to confer glyphosate resistance to the plant. Since weed pressure in cannabis is high due to the competition of weeds before its canopy is established, this would be a good target. Currently there are no federally registered pesticides for cannabis. A gene editing is used to make the required amino acid substitutions, and produce a glyphosate resistant cannabis plant.

Example 10

[0288] Transgenic T0 *Cannabis* plant generation using particle-mediated transformation of *Cannabis* meristem explants—We tested 10 leaves of the twin particle gun event described in Example 8, WP-001181-1b “Hernanda” for aadA1a by PCR and GUS by expression and all 10 leaves

were positive. (FIGS. 72-74). As described herein (Example 2), it is also possible to transform plastids and/or proplastids of *Cannabis* meristem explants using plastid specific promoters driving aadA (and GOI) such as the large subunit promoter of RUBISCO; or a suitable prokaryotic promoter. This would give advantages of plastid transformation (maternal inheritance with null pollen, whole operon engineering, greater protein expression, etc.) in *Cannabis* and *Cannabis* breeding.

[0289] Alternate seed conditioning without imbibition—We were able to get radical emergence from *Cannabis* seed of the Badger variety using seed sanitization followed by

incubation at 37° C. without imbibition. This enables new forms of both hand and machine excision and storage (as seed coat is split but at relatively low moisture compared with imbibed seed). (FIG. 75)

[0290] Transient transformation and Stable expression of RFP (tdTomato) in Badger variety using *Agrobacterium*-mediated transformation of meristem explants—We have also demonstrated transient expression of GUS in meristem explants of the elite Badger variety using meristem explants inoculated with Ar18r12v/DICOTBINARY-19 (derived from seed imbibed overnight at 23° C. or 37° C.) under a variety of co-culture conditions (FIGS. 76-78).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 61

<210> SEQ ID NO 1

<211> LENGTH: 1638

<212> TYPE: DNA

<213> ORGANISM: *Cannabis sativa*

<400> SEQUENCE: 1

```

atgaattgct cagcattttc cttttgggtt gtttgcaaaa taatatTTTT ctttctctca    60
ttccatatcc aaatttcaat agctaactct cgagaaaact tccttaaatg cttctcaaaa    120
catattccca acaatgtagc aaatccaaaa ctcgatataca ctcaacacga ccaattgtat    180
atgtctatcc tgaattcgac aatacaaaat cttagattca tctctgatac aacccccaaa    240
ccactcgtaa ttgtcactcc ttcaataaac tcccatatcc aagcaactat tttatgctct    300
aagaaaagttg gcttgcagat tcgaactcga agcgggtggc atgatgctga gggatgtcc    360
tacatatctc aagtccattt tgtttagtagt gacttgagaa acatgcattc gatcaaaata    420
gatgttcata gccaaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat    480
tggatcaatg agaagaatga gaatcttagt tttcctgggt ggtattgcc tactgttggc    540
gtagggtggc acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg    600
gctgataata ttattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa    660
tccatgggag aagatctgtt ttgggctata cgtgggtggt gaggagaaaa ctttggaaatc    720
attgcagcat ggaatacaa actggttgat gtcccatcaa agtctactat attcagtgtt    780
aaaaagaaca tggagataca tgggcttgtc aagttattta acaaatggca aaatattgct    840
tacaagtatg acaaagattt agtactcatg actcacttca taacaagaa tattacagat    900
aatcatggga agaataagac tacagtacat ggttacttct cttcaatttt tcatggtgga    960
gtggatagtc tagtcgactt gatgaacaag agctttcctg agttgggtat taaaaaaact   1020
gattgcaaag aatttagctg gattgatata accatcttct acagtgggtg tgtaaatTTT   1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct   1140
ttctcaatta agttagacta tgtaagaaa ccaattccag aaactgcaat ggtcaaaatt   1200
ttggaaaaat tatatgaaga agatgtagga gctgggatgt atgtgttgta cccttacggt   1260
ggtataatgg aggagatttc agaatcagca attccattcc ctcatcgagc tggaataatg   1320
tatgaacttt ggtacactgc ttcctgggag aagcaagaag ataataaaaa gcataataac   1380
tgggttcgaa gtgtttataa ttttacgact ccttatgtgt cccaaaatcc aagattggcg   1440
tatctcaatt atagggacct tgatttagga aaaactaatc atgcgagtc taataattac   1500

```

-continued

acacaagcac gtatttgggg tgaagagtat ttgtgtaaaa attttaacag gttagttaag	1560
gtgaaaacta aagttgatcc caataatttt tttagaaacg aacaaagtat cccacctctt	1620
ccaccgcac atcattaa	1638

<210> SEQ ID NO 2
 <211> LENGTH: 1638
 <212> TYPE: DNA
 <213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 2

atgaattgct cagcattttc cttttggttt gtttgcaaaa taatattttt cttctctca	60
ttccatatcc aaatttcaat agctaactc cgagaaaact tccttaaatg cttctcaaaa	120
catattccca acaatgtagc aaatccaaa ctcgtatata ctcaacacga ccaattgtat	180
atgtctatcc tgaattcgac aatacaaaat cttagattca tctctgatac aacccccaaa	240
ccactcgta ttgtcactcc ttcaaataac tcccatatcc aagcaactat tttatgctct	300
aagaaagttg gcttgcagat tcgaactcga agcgggtggc atgatgctga gggatgtcc	360
tacatatctc aagtcctatt tgtttagta gacttgagaa acatgcattc gatcaaaata	420
gatgttcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat	480
tggatcaatg agaagaatga gaacttagt tttcctggtg ggtattgcc tactgttggc	540
gtaggtggac acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg	600
gctgataata ttattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa	660
tccatgggag aagatctgtt ttgggtcata cgtggtggtg gaggagaaaa ctttggatc	720
attgcagcat ggaaaatcaa actggttgct gtcccatcaa agtctactat attcagtggt	780
aaaaagaaca tggagataca tgggctgtc aagttattta acaaatggca aaatattgct	840
tacaagtatg acaagatgtt agtactcatg actcaattca taacaaagaa tattacagat	900
aatcatggga agaataagac tacagtacat ggttacttct cttcaatttt tcatggtgga	960
gtggatagtc tagtcgactt gatgaacaag agctttcgtg agttgggtat taaaaaaact	1020
gattgcaaag aatttagctg gattgatata accatcttct acagtgggtg tgtaaatatt	1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct	1140
ttctcaatta agttagacta tgttaagaaa ccaattccag aaactgcaat ggtcaaaatt	1200
ttggaaaaat tatatgaaga agatgtagga gctgggatgt atgtgttgta cccttacggg	1260
ggtataatgg aggagatttc agaactagca attccattcc ctcatcgagc tggaaataatg	1320
tatgaacttt ggtacactgc ttctgggag aagcaagaag ataataaaaa gcatataaac	1380
tgggttcgaa gtgtttataa ttttacgact ccttatgtgt cccaaaatcc aagattggcg	1440
tatctcaatt atagggacct tgatttagga aaaactaatc atgcgagtc taataattac	1500
acacaagcac gtatttgggg tgaagagtat ttgtgtaaaa attttaacag gttagttaag	1560
gtgaaaacta aagttgatcc caataatttt tttagaaacg aacaaagtat cccacctctt	1620
mcaccgcac atcattaa	1638

<210> SEQ ID NO 3
 <211> LENGTH: 1635
 <212> TYPE: DNA
 <213> ORGANISM: Cannabis sativa

-continued

<400> SEQUENCE: 3

```

atgaattgct cagcattttc cttttgggtt gtttgcaaaa taatattttt ctttctctca    60
ttccatatcc aaatttcaat agctaactc cgagaaaact tccttaaatg cttctcaaaa    120
catattccca acaatgtagc aaatccaaaa ctcgatatca ctcaacacga ccaattgtat    180
atgtctatcc tgaattcgac aatacaaaaat cttagattca tctctgatac aacccccaaa    240
ccactcgta ttgtcactcc ttcaaataac tcccatatcc aagcaactat tttatgctct    300
aagaaagttg gcttgcagat tcgaactcga agcgggtggc atgatgctga gggatgtgcc    360
tacatatctc aagccccatt tgtttagta gacttgagaa acatgcattc gatcaaaata    420
gatgttcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat    480
tggatcaatg agaagaatga gaacttagt tttcctggtg ggtattgcc tactgttggc    540
gtaggtggac acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg    600
gctgataata ttattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa    660
tccatgggag aagatctggt ttgggctata cgtggtggtg gaggagaaaa ctttggaatc    720
attgcagcat ggaaaatcaa actggttgct gtcccatcaa agtctactat attcagtggt    780
aaaaagaaca tggagataca tgggcttgc aagttattta acaaatggca aaatattgct    840
tacaagtatg acaagatgtt agtactcatg actcaattca taacaaagaa tattacagat    900
aatcatggga agaataagac tacagtacat ggttacttct cttcaatttt tcatggtgga    960
gtgtagatgc tagtcgactt gatgaacaag agctttcctg agttgggtat taaaaaaact   1020
gattgcaaag aatttagctg gattgataca accatcttct acagtgggtg tgtaaatatt   1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct   1140
ttctcaatta agttagacta tgtaagaaa ccaattccag aaactgcaat ggtcaaaatt   1200
ttggaaaaat tatatgaaga agatgtagga gctgggatgt atgtgttgta cccttacggt   1260
gggtataatg aggagatttc agaatcagca attccattcc ctcatcgagc tggaaataatg   1320
tatgaacttt ggtacactgc ttctgggag aagcaagaag ataataaaaa gcatataaac   1380
tgggttcgaa gtgtttataa ttttacgact cttatgtgt cccaaaatcc aagattggcg   1440
tatctcaatt atagggacct tgatttagga aaaactaatc atgcgagtc taataattac   1500
acacaagcac gtatttgggg tgaagatgt tttggtaaaa attttaacag gttagttaag   1560
gtgaaaacta aagttgatcc caataatttt ttagaaacg aacaaagtat cccacctctt   1620
ccaccgcac atcat                                     1635

```

<210> SEQ ID NO 4

<211> LENGTH: 1638

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 4

```

atgaattgct cagcattttc cttttgggtt gtttgcaaaa taatattttt ctttctctca    60
ttccatatcc aaatttcaat agctaactc cgagaaaact tccttaaatg cttctcaaaa    120
catattccca acaatgtagc aaatccaaaa ctcgatatca ctcaacacga ccaattgtat    180
atgtctatcc tgaattcgac aatacaaaaat cttagattca tctctgatac aacccccaaa    240
ccactcgta ttgtcactcc ttcaaataac tcccatatcc aagcaactat tttatgctct    300

```

-continued

aagaaagttg gcttcagat tcgaactcga agcgggtggc atgatgctga gggatatgtcc	360
tacatatctc aagtoccatt tgtttagta gacttgagaa acatgcattc gatcaaaata	420
gatgttcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat	480
tggatcaatg agaagaatga gaattctagt ttctctggtg ggtattgccc tactgttggc	540
gtaggtggac acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg	600
gctgataata tcattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa	660
tccatgggag aagatctgtt ttgggtctata cgtggtggtg gaggagaaaa ctttgaatc	720
attgcagcat ggaaaatcaa actggttgct gtcccatcaa agtctactat attcagtgtt	780
aaaaagaaca tggagataca tgggcttgtc aagttattta acaaatggca aaatattgct	840
tacaagtatg acaagatgtt agtactcatg actcaattca taacaaagaa tattacagat	900
aatcatggga agaataagac tacagtacat ggttacttct cttcaatttt tcatggtgga	960
gtggatagtc tagtcgactt gatgaacaag agctttctgt agttgggtat taaaaaaact	1020
gattgcaaag aattgagctg gattgataca accatcttct acagtgggtg tgtaaattac	1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct	1140
ttctcaatta agtttagacta tgttaagaaa ccaattccag aaactgcaat ggtcaaaatt	1200
ttggaaaaat tatatgaaga agatgttaga gctgggatgt atgtgttgta cccttacggt	1260
ggtataatgg aggagatttc agaatcagca attccattcc ctcatcgagc tggaaataatg	1320
tatgaacttt ggtacactgc ttctctggag aagcaagaag ataataaaaa gcatataaac	1380
tgggttcgaa gtgtttataa ttttacgact cttatgtgt cccaaaatcc aagattggcg	1440
tatctcaatt atagggacct tgatttagga aaaactaatc atgcgagtc taataattac	1500
acacaagcac gtatttgggg tgaagatgt tttggtaaaa attttaacag gttagttaag	1560
tgaaaaacta aagttgatcc caataatttt tttagaaacg aacaaagtat cccacctctt	1620
ccaccgcac atcattaa	1638

<210> SEQ ID NO 5

<211> LENGTH: 1638

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 5

atgaattgct cagcattctc cttttggttt gtttgcaaaa taatattttt cttctctca	60
ttcaatatcc aaatttcaat agctaactc caagaaaact tccttaaatg cttctcgga	120
tatattccta acaatccagc aaatccaaaa ttcatataca ctcaacacga ccaattgtat	180
atgtctgtcc tgaattcgac aatacaaaat cttagattca cctctgatac aacccaaaa	240
ccactcgta ttgtcactcc ttcaaatgtc tcccatatcc aggcagtat tctctgtcc	300
aagaaagttg gtttcagat tcgaactcga agcgggtggc atgatgctga gggtttgtcc	360
tacatatctc aagtoccatt tgotatagta gacttgagaa acatgcatac ggtcaaagta	420
gatattcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat	480
tggatcaatg agatgaatga gaattttagt ttctctggtg ggtattgccc tactgttggc	540
gtaggtggac acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg	600
gctgataata tcattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa	660

-continued

tccatgggag aagatctatt ttgggtata cgtggtggag gaggagaaaa ctttgaatc	720
attgcagcat ggaaatcaa acttgttgtt gtccatcaa aggtactat attcagtgtt	780
aaaaagaaca tggagataca tgggcttgc aagttattta acaaatggca aaatattgct	840
tacaagtatg acaagatatt aatgctcag actcacttca gaactaggaa tattacagat	900
aatcatggga agaataagac tacagtacat ggttacttct cttccatttt tcttggtgga	960
gtggatagtc tagttgactt gatgaacaag agctttcctg agttgggtat taaaaaaact	1020
gattgcaaag aattgagctg gattgataca accatcttct acagtgggtg tgtaaattac	1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct	1140
ttctcaatta agttgacta tgtaagaaa ctaatacctg aaactgcaat ggtcaaaatt	1200
ttggaaaaat tatatgaaga agaggttagga gttgggatgt atgtgttgta cccttacggt	1260
ggtataatgg atgagatttc agaatacagc attccattcc ctcatcgagc tggaataatg	1320
tatgaacttt ggtacactgc tacctgggag aagcaagaag ataacgaaaa gcatataaac	1380
tgggttcgaa gtgtttataa tttcacaact ccttatgtgt cccaaaatcc aagattggcg	1440
tatctcaatt atagggacct tgatttagga aaaactaatc ctgagagtcc taataattac	1500
acacaagcac gtatttgggg tgaagatg tttggtaaaa attttaacag gttagttaag	1560
gtgaaaacca aagctgatcc caataatttt ttagaaacg aacaaagtat cccacctctt	1620
ccaccgctc atcattaa	1638

<210> SEQ ID NO 6

<211> LENGTH: 545

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 6

Met Asn Cys Ser Ala Phe Ser Phe Trp Phe Val Cys Lys Ile Ile Phe	
1 5 10 15	
Phe Phe Leu Ser Phe His Ile Gln Ile Ser Ile Ala Asn Pro Arg Glu	
20 25 30	
Asn Phe Leu Lys Cys Phe Ser Lys His Ile Pro Asn Asn Val Ala Asn	
35 40 45	
Pro Lys Leu Val Tyr Thr Gln His Asp Gln Leu Tyr Met Ser Ile Leu	
50 55 60	
Asn Ser Thr Ile Gln Asn Leu Arg Phe Ile Ser Asp Thr Thr Pro Lys	
65 70 75 80	
Pro Leu Val Ile Val Thr Pro Ser Asn Asn Ser His Ile Gln Ala Thr	
85 90 95	
Ile Leu Cys Ser Lys Lys Val Gly Leu Gln Ile Arg Thr Arg Ser Gly	
100 105 110	
Gly His Asp Ala Glu Gly Met Ser Tyr Ile Ser Gln Val Pro Phe Val	
115 120 125	
Val Val Asp Leu Arg Asn Met His Ser Ile Lys Ile Asp Val His Ser	
130 135 140	
Gln Thr Ala Trp Val Glu Ala Gly Ala Thr Leu Gly Glu Val Tyr Tyr	
145 150 155 160	
Trp Ile Asn Glu Lys Asn Glu Asn Leu Ser Phe Pro Gly Gly Tyr Cys	
165 170 175	
Pro Thr Val Gly Val Gly Gly His Phe Ser Gly Gly Gly Tyr Gly Ala	

-continued

180						185						190					
Leu	Met	Arg	Asn	Tyr	Gly	Leu	Ala	Ala	Asp	Asn	Ile	Ile	Asp	Ala	His		
		195					200					205					
Leu	Val	Asn	Val	Asp	Gly	Lys	Val	Leu	Asp	Arg	Lys	Ser	Met	Gly	Glu		
	210					215					220						
Asp	Leu	Phe	Trp	Ala	Ile	Arg	Gly	Gly	Gly	Gly	Glu	Asn	Phe	Gly	Ile		
225					230					235					240		
Ile	Ala	Ala	Trp	Lys	Ile	Lys	Leu	Val	Asp	Val	Pro	Ser	Lys	Ser	Thr		
				245					250					255			
Ile	Phe	Ser	Val	Lys	Lys	Asn	Met	Glu	Ile	His	Gly	Leu	Val	Lys	Leu		
			260					265					270				
Phe	Asn	Lys	Trp	Gln	Asn	Ile	Ala	Tyr	Lys	Tyr	Asp	Lys	Asp	Leu	Val		
		275					280					285					
Leu	Met	Thr	His	Phe	Ile	Thr	Lys	Asn	Ile	Thr	Asp	Asn	His	Gly	Lys		
	290					295					300						
Asn	Lys	Thr	Thr	Val	His	Gly	Tyr	Phe	Ser	Ser	Ile	Phe	His	Gly	Gly		
305					310					315					320		
Val	Asp	Ser	Leu	Val	Asp	Leu	Met	Asn	Lys	Ser	Phe	Pro	Glu	Leu	Gly		
				325					330					335			
Ile	Lys	Lys	Thr	Asp	Cys	Lys	Glu	Phe	Ser	Trp	Ile	Asp	Thr	Thr	Ile		
			340					345					350				
Phe	Tyr	Ser	Gly	Val	Val	Asn	Phe	Asn	Thr	Ala	Asn	Phe	Lys	Lys	Glu		
		355					360					365					
Ile	Leu	Leu	Asp	Arg	Ser	Ala	Gly	Lys	Lys	Thr	Ala	Phe	Ser	Ile	Lys		
	370					375					380						
Leu	Asp	Tyr	Val	Lys	Lys	Pro	Ile	Pro	Glu	Thr	Ala	Met	Val	Lys	Ile		
385					390					395					400		
Leu	Glu	Lys	Leu	Tyr	Glu	Glu	Asp	Val	Gly	Ala	Gly	Met	Tyr	Val	Leu		
			405						410					415			
Tyr	Pro	Tyr	Gly	Gly	Ile	Met	Glu	Glu	Ile	Ser	Glu	Ser	Ala	Ile	Pro		
			420					425					430				
Phe	Pro	His	Arg	Ala	Gly	Ile	Met	Tyr	Glu	Leu	Trp	Tyr	Thr	Ala	Ser		
		435					440					445					
Trp	Glu	Lys	Gln	Glu	Asp	Asn	Glu	Lys	His	Ile	Asn	Trp	Val	Arg	Ser		
	450					455					460						
Val	Tyr	Asn	Phe	Thr	Thr	Pro	Tyr	Val	Ser	Gln	Asn	Pro	Arg	Leu	Ala		
465					470					475					480		
Tyr	Leu	Asn	Tyr	Arg	Asp	Leu	Asp	Leu	Gly	Lys	Thr	Asn	His	Ala	Ser		
			485						490					495			
Pro	Asn	Asn	Tyr	Thr	Gln	Ala	Arg	Ile	Trp	Gly	Glu	Lys	Tyr	Phe	Gly		
		500						505					510				
Lys	Asn	Phe	Asn	Arg	Leu	Val	Lys	Val	Lys	Thr	Lys	Val	Asp	Pro	Asn		
		515					520					525					
Asn	Phe	Phe	Arg	Asn	Glu	Gln	Ser	Ile	Pro	Pro	Leu	Pro	Pro	His	His		
	530					535					540						
His																	
545																	

<210> SEQ ID NO 7

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 7

tgacgcatgg aaaatcaaac 20

<210> SEQ ID NO 8
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 8

cccttacggt ggtataatgg 20

<210> SEQ ID NO 9
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 9

tagctattga aatttgata 20

<210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 10

tagagcataa aatagttgct 20

<210> SEQ ID NO 11
 <211> LENGTH: 1635
 <212> TYPE: DNA
 <213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 11

atgaagtgtc caacattctc cttttggttt gtttgcaaga taatattttt cttttttctca 60

ttcaatatcc aaacttccat tgctaatacct cgagaaaact tccttaaatg cttctcgcaa 120

tatattccca ataatgcaac aaatctaaaa ctggtatata ctcaaaaaca cccattgtat 180

atgtctgtcc taaattcgac aatacacaat cttagattca gctctgacac aacccccaaa 240

ccacttgta tcgtcactcc ttcacatgct tctcatatcc aaggcactat tctatgctcc 300

aagaaagttg gcttgagat tcgaactcga agtggtggtc atgattctga gggcattgctc 360

tacatatctc aagtcccatt tggtatagta gacttgagaa acatgcgttc aatcaaaata 420

gatgttcata gccaaactgc atgggttgaa gccggagcta cccttgaga agtttattat 480

tgggttaatg agaaaaatga gagtcttagt ttggctgctg ggtattgccc tactgtttgc 540

gcaggtggac acttttggtg aggaggctat ggaccattga tgagaagcta tggcctcgcg 600

gctgataata tcattgatgc acacttagtc aacgttcatt gaaaagtgtc agatcgaaaa 660

tctatggggg aagatctctt ttgggtctta cgtggtggtg gagcagaaag cttcggaatc 720

attgtagcat ggaaaattag actggttgct gtcccaaagt ctactatgtt tagtggttaa 780

-continued

```

aagatcatgg agatacatga gcttgtaag ttagttaaca aatggcaaaa tattgcttac      840
aagtatgaca aagatttatt actcatgact cacttcataa ctaggaacat tacagataat      900
caagggaaga ataagacagc aatacacact tacttctctt cagttttcct tgggtggagt      960
gatatgttag tcgacttgat gaacaagagt ttctctgagt tgggtattaa aaaaacggat     1020
tgcagacaat tgagctggat tgatactatc atcttctata gtggtgttgt aaattacgac     1080
actgataatt ttaacaagga aattttgctt gatagatccg ctgggcagaa cgggtgctttc     1140
aagattaagt tagactacgt taagaaacca attccagaat ctgtatttgt ccaaattttg     1200
gaaaaattat atgaagaaga tataggagct gggatgtatg cgttgtaacc ttacggtggg     1260
ataatggatg agatttttga atcagcaatt ccattccctc atcgagctgg aatcttgat      1320
gagttatggg acatatgtag ctggggagaag caagaagata acgaaaagca tctaaactgg     1380
attagaaata ttataactt catgactcct tatgtgtccc aaaatccaag attggcatat      1440
ctcaattata gagaccttga tataggaata aatgatccca agaatccaaa taattacaca     1500
caagcacgta tttgggtgga gaagtatttt ggtaaaaatt ttgacaggct agtaaaagt      1560
aaaaccctgg ttgatcccaa taattttttt agaaacgaac aaagcatccc acctcttcca     1620
cggcatcgtc attaa                                           1635

```

<210> SEQ ID NO 12

<211> LENGTH: 544

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 12

```

Met Lys Cys Ser Thr Phe Ser Phe Trp Phe Val Cys Lys Ile Ile Phe
1             5             10            15

Phe Phe Phe Ser Phe Asn Ile Gln Thr Ser Ile Ala Asn Pro Arg Glu
20            25            30

Asn Phe Leu Lys Cys Phe Ser Gln Tyr Ile Pro Asn Asn Ala Thr Asn
35            40            45

Leu Lys Leu Val Tyr Thr Gln Asn Asn Pro Leu Tyr Met Ser Val Leu
50            55            60

Asn Ser Thr Ile His Asn Leu Arg Phe Ser Ser Asp Thr Thr Pro Lys
65            70            75            80

Pro Leu Val Ile Val Thr Pro Ser His Val Ser His Ile Gln Gly Thr
85            90            95

Ile Leu Cys Ser Lys Lys Val Gly Leu Gln Ile Arg Thr Arg Ser Gly
100           105           110

Gly His Asp Ser Glu Gly Met Ser Tyr Ile Ser Gln Val Pro Phe Val
115           120           125

Ile Val Asp Leu Arg Asn Met Arg Ser Ile Lys Ile Asp Val His Ser
130           135           140

Gln Thr Ala Trp Val Glu Ala Gly Ala Thr Leu Gly Glu Val Tyr Tyr
145           150           155           160

Trp Val Asn Glu Lys Asn Glu Ser Leu Ser Leu Ala Ala Gly Tyr Cys
165           170           175

Pro Thr Val Cys Ala Gly Gly His Phe Gly Gly Gly Gly Tyr Gly Pro
180           185           190

Leu Met Arg Ser Tyr Gly Leu Ala Ala Asp Asn Ile Ile Asp Ala His

```

-continued

195	200	205
Leu Val Asn Val His Gly	Lys Val Leu Asp Arg	Lys Ser Met Gly Glu
210	215	220
Asp Leu Phe Trp Ala Leu Arg Gly Gly Gly Ala Glu Ser Phe Gly Ile		
225	230	235 240
Ile Val Ala Trp Lys Ile Arg Leu Val Ala Val Pro Lys Ser Thr Met		
	245	250 255
Phe Ser Val Lys Lys Ile Met Glu Ile His Glu Leu Val Lys Leu Val		
	260	265 270
Asn Lys Trp Gln Asn Ile Ala Tyr Lys Tyr Asp Lys Asp Leu Leu Leu		
	275	280 285
Met Thr His Phe Ile Thr Arg Asn Ile Thr Asp Asn Gln Gly Lys Asn		
	290	295 300
Lys Thr Ala Ile His Thr Tyr Phe Ser Ser Val Phe Leu Gly Gly Val		
	305	310 315 320
Asp Ser Leu Val Asp Leu Met Asn Lys Ser Phe Pro Glu Leu Gly Ile		
	325	330 335
Lys Lys Thr Asp Cys Arg Gln Leu Ser Trp Ile Asp Thr Ile Ile Phe		
	340	345 350
Tyr Ser Gly Val Val Asn Tyr Asp Thr Asp Asn Phe Asn Lys Glu Ile		
	355	360 365
Leu Leu Asp Arg Ser Ala Gly Gln Asn Gly Ala Phe Lys Ile Lys Leu		
	370	375 380
Asp Tyr Val Lys Lys Pro Ile Pro Glu Ser Val Phe Val Gln Ile Leu		
	385	390 395 400
Glu Lys Leu Tyr Glu Glu Asp Ile Gly Ala Gly Met Tyr Ala Leu Tyr		
	405	410 415
Pro Tyr Gly Gly Ile Met Asp Glu Ile Ser Glu Ser Ala Ile Pro Phe		
	420	425 430
Pro His Arg Ala Gly Ile Leu Tyr Glu Leu Trp Tyr Ile Cys Ser Trp		
	435	440 445
Glu Lys Gln Glu Asp Asn Glu Lys His Leu Asn Trp Ile Arg Asn Ile		
	450	455 460
Tyr Asn Phe Met Thr Pro Tyr Val Ser Gln Asn Pro Arg Leu Ala Tyr		
	465	470 475 480
Leu Asn Tyr Arg Asp Leu Asp Ile Gly Ile Asn Asp Pro Lys Asn Pro		
	485	490 495
Asn Asn Tyr Thr Gln Ala Arg Ile Trp Gly Glu Lys Tyr Phe Gly Lys		
	500	505 510
Asn Phe Asp Arg Leu Val Lys Val Lys Thr Leu Val Asp Pro Asn Asn		
	515	520 525
Phe Phe Arg Asn Glu Gln Ser Ile Pro Pro Leu Pro Arg His Arg His		
	530	535 540

<210> SEQ ID NO 13

<211> LENGTH: 1635

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 13

atgaagtgcct caacattctc cttttgggtt gtttgcaaga taatattttt ctttttctca 60

ttcaatatcc aaacttccat tgctaatacct cgagaaaact tccttaaagt cttctcgcaa 120

-continued

tatattccca ataatgcaac aaatctaaaa ctcgatatca ctcaaaaacaa cccattgtat	180
atgtctgtcc taaattcgac aatacacaaat cttagattca cctctgacac aacccccaaaa	240
ccacttggtta tcgtcactcc ttccacatgtc tctcatatcc aaggcactat tctatgctcc	300
aagaaagtgt gcttgcagat tcgaactcga agtgggtggtc atgattctga gggcatgtcc	360
tacatatctc aagtccatt tgttatagta gacttgagaa acatgcgttc aatcaaaata	420
gatgttcata gccaaactgc atgggttgaa gccggagcta cccttgaga agtttattat	480
tgggttaatg agaaaaatga gaattctagt ttggcggctg ggtattgccc tactgtttgc	540
gcaggtggac actttggtgg aggaggtat ggaccattga tgagaaacta tggcctcgcg	600
gctgataata tcattgatgc acacttagtc aacgttcagt gaaaagtgt agatcgaaaa	660
tctatggggg aagatctctt ttgggcttta cgtggtggtg gagcagaaag cttcggaatc	720
attgtagcat ggaaaattag actggttgct gtcccaaagt ctactatgtt tagtgtaaa	780
aagatcatgg agatacatga gcttgtcaag ttagttaaca aatggcaaaa tattgcttac	840
aagtatgaca aagatttatt actcatgact cacttcataa ctaggaaact tacagataat	900
caagggaaga ataagacagc aatacacact tacttctctt cagttttcct tgggtgagtg	960
gatagtctag tcgacttgat gaacaagagt ttctctgagt tgggtattaa aaaaacggat	1020
tgcagacaat tgagctggat tgatactatc atcttctata gtggtgttgt aaattacgac	1080
actgataatt ttaacaagga aattttgctt gatagatccg ctgggcagaa cgggtgcttc	1140
aagattaagt tagactacgt taagaaacca attccagaat ctgtatttgt ccaaattttg	1200
gaaaaattat atgaagaaga tataggagct gggatgtatg cgttgtacct ttacggtggt	1260
ataatggatg agatttcaga atcagcaatt ccattccctc atcgagctgg aatcttgat	1320
gagttatggt acatatgtag ttggggagaag caagaagata acgaaaagca tctaaactgg	1380
attagaaata ttataactt catgactcct tatgtgtcca aaaatccaag attggcatat	1440
ctcaattata gagacctga tataggaata aatgatccca agaatccaaa taattacaca	1500
caagcacgta tttgggtgga gaagtattt ggtaaaaatt ttgacaggct agtaaaagt	1560
aaaaccctgg ttgatcccaa taactttttt agaaacgaac aaagcatccc acctcttcca	1620
cggcatcgtc attaa	1635

<210> SEQ ID NO 14

<211> LENGTH: 544

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 14

Met	Lys	Cys	Ser	Thr	Phe	Ser	Phe	Trp	Phe	Val	Cys	Lys	Ile	Ile	Phe
1				5				10					15		
Phe	Phe	Phe	Ser	Phe	Asn	Ile	Gln	Thr	Ser	Ile	Ala	Asn	Pro	Arg	Glu
			20					25					30		
Asn	Phe	Leu	Lys	Cys	Phe	Ser	Gln	Tyr	Ile	Pro	Asn	Asn	Ala	Thr	Asn
			35					40					45		
Leu	Lys	Leu	Val	Tyr	Thr	Gln	Asn	Asn	Pro	Leu	Tyr	Met	Ser	Val	Leu
			50				55					60			
Asn	Ser	Thr	Ile	His	Asn	Leu	Arg	Phe	Thr	Ser	Asp	Thr	Thr	Pro	Lys
					70					75					80

-continued

Pro	Leu	Val	Ile	Val	Thr	Pro	Ser	His	Val	Ser	His	Ile	Gln	Gly	Thr	85	90	95
Ile	Leu	Cys	Ser	Lys	Lys	Val	Gly	Leu	Gln	Ile	Arg	Thr	Arg	Ser	Gly	100	105	110
Gly	His	Asp	Ser	Glu	Gly	Met	Ser	Tyr	Ile	Ser	Gln	Val	Pro	Phe	Val	115	120	125
Ile	Val	Asp	Leu	Arg	Asn	Met	Arg	Ser	Ile	Lys	Ile	Asp	Val	His	Ser	130	135	140
Gln	Thr	Ala	Trp	Val	Glu	Ala	Gly	Ala	Thr	Leu	Gly	Glu	Val	Tyr	Tyr	145	150	155
Trp	Val	Asn	Glu	Lys	Asn	Glu	Asn	Leu	Ser	Leu	Ala	Ala	Gly	Tyr	Cys	165	170	175
Pro	Thr	Val	Cys	Ala	Gly	Gly	His	Phe	Gly	Gly	Gly	Gly	Tyr	Gly	Pro	180	185	190
Leu	Met	Arg	Asn	Tyr	Gly	Leu	Ala	Ala	Asp	Asn	Ile	Ile	Asp	Ala	His	195	200	205
Leu	Val	Asn	Val	His	Gly	Lys	Val	Leu	Asp	Arg	Lys	Ser	Met	Gly	Glu	210	215	220
Asp	Leu	Phe	Trp	Ala	Leu	Arg	Gly	Gly	Gly	Ala	Glu	Ser	Phe	Gly	Ile	225	230	235
Ile	Val	Ala	Trp	Lys	Ile	Arg	Leu	Val	Ala	Val	Pro	Lys	Ser	Thr	Met	245	250	255
Phe	Ser	Val	Lys	Lys	Ile	Met	Glu	Ile	His	Glu	Leu	Val	Lys	Leu	Val	260	265	270
Asn	Lys	Trp	Gln	Asn	Ile	Ala	Tyr	Lys	Tyr	Asp	Lys	Asp	Leu	Leu	Leu	275	280	285
Met	Thr	His	Phe	Ile	Thr	Arg	Asn	Ile	Thr	Asp	Asn	Gln	Gly	Lys	Asn	290	295	300
Lys	Thr	Ala	Ile	His	Thr	Tyr	Phe	Ser	Ser	Val	Phe	Leu	Gly	Gly	Val	305	310	315
Asp	Ser	Leu	Val	Asp	Leu	Met	Asn	Lys	Ser	Phe	Pro	Glu	Leu	Gly	Ile	325	330	335
Lys	Lys	Thr	Asp	Cys	Arg	Gln	Leu	Ser	Trp	Ile	Asp	Thr	Ile	Ile	Phe	340	345	350
Tyr	Ser	Gly	Val	Val	Asn	Tyr	Asp	Thr	Asp	Asn	Phe	Asn	Lys	Glu	Ile	355	360	365
Leu	Leu	Asp	Arg	Ser	Ala	Gly	Gln	Asn	Gly	Ala	Phe	Lys	Ile	Lys	Leu	370	375	380
Asp	Tyr	Val	Lys	Lys	Pro	Ile	Pro	Glu	Ser	Val	Phe	Val	Gln	Ile	Leu	385	390	395
Glu	Lys	Leu	Tyr	Glu	Glu	Asp	Ile	Gly	Ala	Gly	Met	Tyr	Ala	Leu	Tyr	405	410	415
Pro	Tyr	Gly	Gly	Ile	Met	Asp	Glu	Ile	Ser	Glu	Ser	Ala	Ile	Pro	Phe	420	425	430
Pro	His	Arg	Ala	Gly	Ile	Leu	Tyr	Glu	Leu	Trp	Tyr	Ile	Cys	Ser	Trp	435	440	445
Glu	Lys	Gln	Glu	Asp	Asn	Glu	Lys	His	Leu	Asn	Trp	Ile	Arg	Asn	Ile	450	455	460
Tyr	Asn	Phe	Met	Thr	Pro	Tyr	Val	Ser	Lys	Asn	Pro	Arg	Leu	Ala	Tyr	465	470	475
Leu	Asn	Tyr	Arg	Asp	Leu	Asp	Ile	Gly	Ile	Asn	Asp	Pro	Lys	Asn	Pro			

-continued

485					490					495					
Asn	Asn	Tyr	Thr	Gln	Ala	Arg	Ile	Trp	Gly	Glu	Lys	Tyr	Phe	Gly	Lys
			500					505					510		
Asn	Phe	Asp	Arg	Leu	Val	Lys	Val	Lys	Thr	Leu	Val	Asp	Pro	Asn	Asn
		515					520					525			
Phe	Phe	Arg	Asn	Glu	Gln	Ser	Ile	Pro	Pro	Leu	Pro	Arg	His	Arg	His
		530					535					540			

<210> SEQ ID NO 15
 <211> LENGTH: 1635
 <212> TYPE: DNA
 <213> ORGANISM: Cannabis sativa
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (333)..(441)
 <223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 15

atgaagtact caacattctc cttttggttt gtttgcaaga taatattttt ctttttctca	60
ttcaatatcc aaacttccat tgctaactct cgagaaaact tccttaaatg cttctcgcaa	120
tatattccca ataatgcaac aaatctaaaa ctcgtataca ctcaaaacaa cccattgtat	180
atgtctgtcc taaattcgac aatacacaaat cttagattca gctctgacac aacccccaaa	240
ccacttgta tcgtcactcc ttcacatgtc tctcatatcc aaggcaactat tctatgctcc	300
aagaaagttg gcttgcagat tcgaactcga agnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	360
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	420
nnnnnnnnnn nnnnnnnnnn ntgggttgaa gccggagcta cccttgaga agtttattat	480
tgggttaatg agaaaaatga gagtcttagt ttggctgctg ggtattgcc tactgtttgc	540
gcaggtggac actttgtgtg aggaggctat ggaccattga tgagaagcta tggcctcgcg	600
gctgataata tcattgatgc acacttagtc aacgttcattg gaaaagtgt agatcgaaaa	660
tctatggggg aagatctctt ttgggcttta cgtgggtgtg gaggagaaag cttcggaatc	720
attgtagcat ggaaaattag actggttgct gtcccaaagt ctactatgtt tagtgtaaaa	780
aagatcatgg agatacatga gcttgcgaag ttagttaaca aatggcaaaa tattgcttac	840
aagtatgaca aagatttatt actcatgact cacttcataa ctaggaaacac tacagataat	900
caagggaaga ataagacagc aatacacact tacttctctt cagttttcct tgggtggagt	960
gatagtctag tcgacttgat gaacaagagt tttcctgagt tgggtattaa aaaaacggat	1020
tcgagacaat tgagctggat tgatactatc atcttctata gtggtgttgt aaattacgac	1080
actgataatt ttaacaagga aattttgctt gatagatccg ctgggcagaa cggtgctttc	1140
aagattaagt tagactacgt taagaaacca attccagaat ctgtatttgt ccaaattttg	1200
gaaaaattat atgaagaaga tataggagct gggatgtatg cggtgtaccc ttacggtggt	1260
ataatggatg agattttctga atcagcaatt ccattccctc atcgagctgg aatcttgtat	1320
gagttatggt acatatgtag ctgggagaag caagaagata acgaaaagca tctaaactgg	1380
attagaaata tttataactt catgactcct tatgtgtccc aaaatccaag attggcatat	1440
ctcaattata gagaccttga tataggaata aatgatccca agaattccaaa taattacaca	1500
caagcacgta tttgggttga gaagtatttt ggtaaaaatt ttgacaggct agtaaaagt	1560
aaaaccctgg ttgatcccaa taattttttt agaaacgaac aaagcatccc acctcttcca	1620

-continued

cggcacatc attaa

1635

<210> SEQ ID NO 16

<211> LENGTH: 544

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (111)..(147)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 16

```

Met Lys Tyr Ser Thr Phe Ser Phe Trp Phe Val Cys Lys Ile Ile Phe
 1             5             10             15

Phe Phe Phe Ser Phe Asn Ile Gln Thr Ser Ile Ala Asn Pro Arg Glu
 20             25             30

Asn Phe Leu Lys Cys Phe Ser Gln Tyr Ile Pro Asn Asn Ala Thr Asn
 35             40             45

Leu Lys Leu Val Tyr Thr Gln Asn Asn Pro Leu Tyr Met Ser Val Leu
 50             55             60

Asn Ser Thr Ile His Asn Leu Arg Phe Ser Ser Asp Thr Thr Pro Lys
 65             70             75             80

Pro Leu Val Ile Val Thr Pro Ser His Val Ser His Ile Gln Gly Thr
 85             90             95

Ile Leu Cys Ser Lys Lys Val Gly Leu Gln Ile Arg Thr Arg Xaa Xaa
100             105             110

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
115             120             125

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
130             135             140

Xaa Xaa Xaa Trp Val Glu Ala Gly Ala Thr Leu Gly Glu Val Tyr Tyr
145             150             155             160

Trp Val Asn Glu Lys Asn Glu Ser Leu Ser Leu Ala Ala Gly Tyr Cys
165             170             175

Pro Thr Val Cys Ala Gly Gly His Phe Gly Gly Gly Gly Tyr Gly Pro
180             185             190

Leu Met Arg Ser Tyr Gly Leu Ala Ala Asp Asn Ile Ile Asp Ala His
195             200             205

Leu Val Asn Val His Gly Lys Val Leu Asp Arg Lys Ser Met Gly Glu
210             215             220

Asp Leu Phe Trp Ala Leu Arg Gly Gly Gly Ala Glu Ser Phe Gly Ile
225             230             235             240

Ile Val Ala Trp Lys Ile Arg Leu Val Ala Val Pro Lys Ser Thr Met
245             250             255

Phe Ser Val Lys Lys Ile Met Glu Ile His Glu Leu Val Lys Leu Val
260             265             270

Asn Lys Trp Gln Asn Ile Ala Tyr Lys Tyr Asp Lys Asp Leu Leu Leu
275             280             285

Met Thr His Phe Ile Thr Arg Asn Ile Thr Asp Asn Gln Gly Lys Asn
290             295             300

Lys Thr Ala Ile His Thr Tyr Phe Ser Ser Val Phe Leu Gly Gly Val
305             310             315             320

Asp Ser Leu Val Asp Leu Met Asn Lys Ser Phe Pro Glu Leu Gly Ile

```

-continued

325						330						335			
Lys	Lys	Thr	Asp	Cys	Arg	Gln	Leu	Ser	Trp	Ile	Asp	Thr	Ile	Ile	Phe
			340					345					350		
Tyr	Ser	Gly	Val	Val	Asn	Tyr	Asp	Thr	Asp	Asn	Phe	Asn	Lys	Glu	Ile
		355					360					365			
Leu	Leu	Asp	Arg	Ser	Ala	Gly	Gln	Asn	Gly	Ala	Phe	Lys	Ile	Lys	Leu
	370					375					380				
Asp	Tyr	Val	Lys	Lys	Pro	Ile	Pro	Glu	Ser	Val	Phe	Val	Gln	Ile	Leu
385					390					395					400
Glu	Lys	Leu	Tyr	Glu	Glu	Asp	Ile	Gly	Ala	Gly	Met	Tyr	Ala	Leu	Tyr
				405					410					415	
Pro	Tyr	Gly	Gly	Ile	Met	Asp	Glu	Ile	Ser	Glu	Ser	Ala	Ile	Pro	Phe
			420					425					430		
Pro	His	Arg	Ala	Gly	Ile	Leu	Tyr	Glu	Leu	Trp	Tyr	Ile	Cys	Ser	Trp
		435					440					445			
Glu	Lys	Gln	Glu	Asp	Asn	Glu	Lys	His	Leu	Asn	Trp	Ile	Arg	Asn	Ile
	450					455					460				
Tyr	Asn	Phe	Met	Thr	Pro	Tyr	Val	Ser	Gln	Asn	Pro	Arg	Leu	Ala	Tyr
465					470					475					480
Leu	Asn	Tyr	Arg	Asp	Leu	Asp	Ile	Gly	Ile	Asn	Asp	Pro	Lys	Asn	Pro
				485					490					495	
Asn	Asn	Tyr	Thr	Gln	Ala	Arg	Ile	Trp	Gly	Glu	Lys	Tyr	Phe	Gly	Lys
			500					505					510		
Asn	Phe	Asp	Arg	Leu	Val	Lys	Val	Lys	Thr	Leu	Val	Asp	Pro	Asn	Asn
		515					520					525			
Phe	Phe	Arg	Asn	Glu	Gln	Ser	Ile	Pro	Pro	Leu	Pro	Arg	His	His	His
	530					535					540				

<210> SEQ ID NO 17

<211> LENGTH: 1634

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 17

```

atgaagtgtct caacattccc cttttggttt gtttgcaaga taatatTTTT ctttctctca    60
ttcaatatcc aaacttcaat tgctaactct cgagaaaact tccttaaatg cttctcgcaa    120
tatattccca ccaatgtaac aaatctaaaa cttacaccca aaacaaccaa ttgtatatgc    180
ctgtccaaaa ttcaacaata cacaatctta gattcacctc taacacaacc ccaaaactac    240
ttgttatcgt cactccttca tatgtctctc atatccaagg cactattcta tgtccaagaa    300
aattggtttg caaattcgaa ctggaagcgg tgggtcatgat tctgaagaca tgcctcatat    360
atctcaagtc ccatttggtta tagtagactt gagaacatg cattcaatca acatagatgt    420
tcatagccaa atcgcaaggg ttgaagccgg agctaccctt ggagaagttt attattgggt    480
taatgagaaa aatgagaatc ttagtgtggc tgctgggtat tgccctactg ttagcgcagc    540
tggacacttt ggtggaggag gatattggacc attgatgcaa aattatggcc tcgcggtgga    600
taatatcggt gatgcacact tagtcaacgt tgatgcaaaa gtgctagatc gaaaaatctat    660
gggggaagat ctcttttggg ctatacgtgg tgggtggagga gaaagcttcg gaatcattgt    720
agcatggaaa attagactgg ttgctgtccc aacaaagtct actatgttta gtgttaaaaa    780
gatcatggag atacatgagc ttgtcaagtg agttaacaaa tggcaaaaata ttgcttacia    840

```

-continued

```

gtatgacaaa gatttattac tcatgactca cttcataact aggaatatta caaataatca    900
tggaagaagt aagacaacaa tacacactta cttctcttca gtttccttg gtggagtgga    960
tagtctagtc gacttgatga ataagagttt tcctgagttg ggtattaaaa aaacagattg   1020
caaacaattg agctagattg atattatcat cttttatagc ggtgttgtaa attacggcac   1080
tgataatfff aataaggaaa ttttgcttga tagatcagct gggcagaacg gttctttaa    1140
gattaagtta gactacgtta agaaccaat tccagaatct gcgtttgtca aaattttgga    1200
aaaattatat gaagaagatg aaggagttgg gatgtatgcg ttgtaccctt acggtgggat   1260
aatggatgag atttcagaat cagcaattcc attccctcat tgagctggaa tcatgtatga   1320
attatggtac atatgtagct gggagaagca cgaagataac gaaaaagcat ctaaactgga   1380
ttcgaaatgt ttatagcttc attactcctt atgtgtccta aaatccaaga ttggcatatc   1440
tcaattatag agaccttgat actggaataa atgatcccaa gagtccaaat aattacacac   1500
aagaaagtat ttgggggtgag aagtattttg gtaaaaaatt tgacagggtg gtaaaagtga   1560
aaacctgggt tgatcccaat aattttttta gaaatgaaca aagcatccca cctcttccac   1620
ggcatcgtca ttaa                                     1634

```

```

<210> SEQ ID NO 18
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

```

```

<400> SEQUENCE: 18

```

```

gctagatcga aaatctatgg                                     20

```

```

<210> SEQ ID NO 19
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

```

```

<400> SEQUENCE: 19

```

```

aaagcatccc acctcttcca                                     20

```

```

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

```

```

<400> SEQUENCE: 20

```

```

tttaggacag acatatacaa                                     20

```

```

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

```

```

<400> SEQUENCE: 21

```

```

gaaagcaccg ttctgcccag                                     20

```

-continued

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 22

catttaagga agttttctcg 20

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 23

aaatgggact tgagatatgt 20

<210> SEQ ID NO 24
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 24

cccttggaga agtttattat 20

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 25

gtacccttac ggtggtataa 20

<210> SEQ ID NO 26
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 26

attccagctc gatgaggaa 20

<210> SEQ ID NO 27
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- guide RNA

<400> SEQUENCE: 27

tacacacaag cacgtatttg 20

<210> SEQ ID NO 28
<211> LENGTH: 109

-continued

<212> TYPE: PRT

<213> ORGANISM: Brassica napus

<400> SEQUENCE: 28

```

Met Ala Thr Gly Ser Arg Val Leu Ile Gly Leu Ala Met Ile Leu Ile
1           5           10           15
Ile Ser Gly Glu Leu Leu Val Pro Gly Gln Gly Thr Cys Gln Gly Asp
20           25           30
Ile Glu Gly Leu Met Arg Glu Cys Ala Val Tyr Val Gln Arg Pro Gly
35           40           45
Pro Lys Val Asn Pro Ser Ala Ala Cys Cys Lys Val Val Lys Arg Ser
50           55           60
Asp Ile Pro Cys Ala Cys Gly Arg Ile Thr Pro Ser Val Gln Lys Met
65           70           75           80
Ile Asp Met Asn Lys Val Val Leu Val Thr Ser Phe Cys Gly Arg Pro
85           90           95
Leu Ala His Gly Thr Lys Cys Gly Ser Tyr Ile Val Pro
100          105

```

<210> SEQ ID NO 29

<211> LENGTH: 330

<212> TYPE: DNA

<213> ORGANISM: Brassica napus

<400> SEQUENCE: 29

```

atggcgacag gttctcgtgt tctgatcggt ctagcaatga tcctcataat ctcaggagaa    60
ctgctagttc cagggcaagg aacgtgccaa ggagacatag agggctctgat gagagaatgt    120
gcggtctacg tccagcgtec agggcccaaag gtaaaccat cgcagcgtg ttgcaaagtc    180
gtcaagagat cagacatccc ctgcgcatgt ggccgtatca caccctcggt tcaaaaaatg    240
atagacatga ataaggttgt tcttgtcact tccttttgtg ggaggcctct cgctcatggt    300
accaagtgtg gaagctacat tgtgccatga    330

```

<210> SEQ ID NO 30

<211> LENGTH: 395

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 30

```

Met Gly Leu Ser Ser Val Cys Thr Phe Ser Phe Gln Thr Asn Tyr His
1           5           10           15
Thr Leu Leu Asn Pro His Asn Asn Asn Pro Lys Thr Ser Leu Leu Cys
20           25           30
Tyr Arg His Pro Lys Thr Pro Ile Lys Tyr Ser Tyr Asn Asn Phe Pro
35           40           45
Ser Lys His Cys Ser Thr Lys Ser Phe His Leu Gln Asn Lys Cys Ser
50           55           60
Glu Ser Leu Ser Ile Ala Lys Asn Ser Ile Arg Ala Ala Thr Thr Asn
65           70           75           80
Gln Thr Glu Pro Pro Glu Ser Asp Asn His Ser Val Ala Thr Lys Ile
85           90           95
Leu Asn Phe Gly Lys Ala Cys Trp Lys Leu Gln Arg Pro Tyr Thr Ile
100          105          110
Ile Ala Phe Thr Ser Cys Ala Cys Gly Leu Phe Gly Lys Glu Leu Leu

```

-continued

115	120	125
His Asn Thr Asn Leu Ile Ser Trp Ser Leu Met Phe Lys Ala Phe Phe 130 135 140		
Phe Leu Val Ala Val Leu Cys Ile Ala Ser Phe Thr Thr Thr Ile Asn 145 150 155 160		
Gln Ile Tyr Asp Leu His Ile Asp Arg Ile Asn Lys Pro Asp Leu Pro 165 170 175		
Leu Ala Ser Gly Glu Ile Ser Val Asn Thr Ala Trp Ile Met Ser Ile 180 185 190		
Ile Val Ala Leu Phe Gly Leu Ile Ile Thr Ile Lys Met Lys Gly Gly 195 200 205		
Pro Leu Tyr Ile Phe Gly Tyr Cys Phe Gly Ile Phe Gly Gly Ile Val 210 215 220		
Tyr Ser Val Pro Pro Phe Arg Trp Lys Gln Asn Pro Ser Thr Ala Phe 225 230 235 240		
Leu Leu Asn Phe Leu Ala His Ile Ile Thr Asn Phe Thr Phe Tyr Tyr 245 250 255		
Ala Ser Arg Ala Ala Leu Gly Leu Pro Phe Glu Leu Arg Pro Ser Phe 260 265 270		
Thr Phe Leu Leu Ala Phe Met Lys Ser Met Gly Ser Ala Leu Ala Leu 275 280 285		
Ile Lys Asp Ala Ser Asp Val Glu Gly Asp Thr Lys Phe Gly Ile Ser 290 295 300		
Thr Leu Ala Ser Lys Tyr Gly Ser Arg Asn Leu Thr Leu Phe Cys Ser 305 310 315 320		
Gly Ile Val Leu Leu Ser Tyr Val Ala Ala Ile Leu Ala Gly Ile Ile 325 330 335		
Trp Pro Gln Ala Phe Asn Ser Asn Val Met Leu Leu Ser His Ala Ile 340 345 350		
Leu Ala Phe Trp Leu Ile Leu Gln Thr Arg Asp Phe Ala Leu Thr Asn 355 360 365		
Tyr Asp Pro Glu Ala Gly Arg Arg Phe Tyr Glu Phe Met Trp Lys Leu 370 375 380		
Tyr Tyr Ala Glu Tyr Leu Val Tyr Val Phe Ile 385 390 395		

<210> SEQ ID NO 31

<211> LENGTH: 1188

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 31

```

atgggactct catcagtttg taccttttca tttcaaacta attaccatac ttatttaaatt    60
cctcacaata ataatcccaa aacctcatta ttatgttatc gacaccccaa aacaccaatt    120
aaatactctt acaataatth tccctctaaa cattgctcca ccaagagttt tcatctacaa    180
aacaaatgct cagaatcatt atcaatcgca aaaaattcca ttagggcagc tactacaaat    240
caaaatgagc ctccagaatc tgataatcat tcagtagcaa ctaaaatttt aaactttggg    300
aaggcatggt ggaaacttca aagaccatat acaatcatag catttacttc atgcgcttgt    360
ggattgtttg ggaaagagtt gttgcataac acaaatthta taagttgggc tctgatgttc    420
aaggcattct ttttttgggt ggctgtatta tgcattgctt cttttacaac taccatcaat    480

```


-continued

```

cagatttacg atcttcacat tgacagaata aacaagcctg atctaccact agcttcaggg 540
gaaatatcag taaacacagc ttggattatg agcataattg tggcactgtt tggattgata 600
ataactataa aaatgaaggg tggaccactc tatatatttg gctactgttt tggatatttt 660
ggtgggattg tctattctgt tccaccattt agatggaagc aaaatccttc cactgcattt 720
cttctcaatt tcctggccca tattattaca aatttcacat tttattatgc cagcagagca 780
gctcttggcc taccatttga gttgaggcct tcttttactt tcctgctagc atttatgaaa 840
tcaatggggtt cagctttggc tttaatcaaa gatgcttcag acgttgaagg cgacactaaa 900
tttggcatat caaccttggc aagtaaatat ggttccagaa acttgacatt attttgttct 960
ggaattgttc tcctatccta tgtggctgct atacttgctg ggattatctg gccccaggct 1020
ttcaacagta acgtaatgtt actttctcat gcaatcttag cattttgggtt aatcctccag 1080
actcgagatt ttgcgttaac aaattacgac ccggaagcag gcagaagatt ttacgagttc 1140
atgtggaagc tttattatgc tgaatattta gtatatgttt tcatataa 1188

```

<210> SEQ ID NO 32

<211> LENGTH: 370

<212> TYPE: PRT

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 32

```

Met Gly Ser Thr Gly Ile Glu Thr Gln Met Thr Pro Thr Gln Ile Ser
1           5           10           15

Asp Glu Glu Ala Asn Leu Phe Ala Met Gln Leu Ala Ser Ala Ser Val
20          25          30

Leu Pro Met Val Leu Lys Ala Ala Leu Glu Leu Asp Leu Leu Glu Ile
35          40          45

Ile Ala Lys Ala Gly Pro Gly Ala Phe Leu Ser Pro Ser Asp Ile Ala
50          55          60

Gln Gln Leu Pro Thr Gln Asn Pro Asp Ala Pro Val Met Leu Asp Arg
65          70          75          80

Met Leu Arg Leu Leu Ala Ser Tyr Asn Val Val Thr Tyr Ser Leu Arg
85          90          95

Glu Arg Glu Thr Ala Glu Glu Glu Gly Lys Val Glu Arg Leu Tyr Gly
100         105         110

Leu Ala Pro Val Ser Lys Tyr Leu Thr Lys Asn Glu Asp Gly Val Ser
115         120         125

Ile Ala Pro Leu Cys Leu Met Asn Gln Asp Lys Val Leu Met Glu Ser
130         135         140

Trp Tyr His Leu Lys Asp Ala Val Leu Asp Gly Gly Ile Pro Phe Asn
145         150         155         160

Lys Ala Tyr Gly Met Thr Ala Phe Glu Tyr His Gly Thr Asp Gln Arg
165         170         175

Phe Asn Lys Ile Phe Asn Arg Gly Met Ser Asp His Ser Thr Ile Thr
180         185         190

Met Lys Lys Ile Leu Glu Thr Tyr Lys Gly Phe Glu Gly Leu Asn Ser
195         200         205

Ile Val Asp Val Gly Gly Gly Thr Gly Ala Val Val Asn Met Ile Val
210         215         220

Ser Lys Tyr Pro Thr Ile Lys Gly Ile Asn Phe Asp Leu Pro His Val

```

-continued

225	230	235	240
Ile Glu Asp Ala Pro Pro Leu Thr Gly Val Glu His Val Gly Gly Asp			
	245	250	255
Met Phe Val Ser Val Pro Lys Gly Asp Ala Ile Phe Met Lys Trp Ile			
	260	265	270
Cys His Asp Trp Ser Asp Glu His Cys Leu Lys Phe Leu Lys Asn Cys			
	275	280	285
His Ala Ala Leu Pro Glu His Gly Lys Val Ile Val Ala Glu Cys Ile			
	290	295	300
Leu Pro Val Ala Pro Asp Ser Ser Leu Ala Thr Lys Ser Thr Val His			
	305	310	315
Ile Asp Val Ile Met Leu Ala His Asn Pro Gly Gly Lys Glu Arg Thr			
	325	330	335
Glu Lys Glu Phe Glu Ala Leu Ala Lys Gly Ala Gly Phe Lys Gly Phe			
	340	345	350
Lys Val His Cys Asn Ala Phe Asn Thr His Ile Met Glu Phe Leu Lys			
	355	360	365
Thr Ile			
	370		

<210> SEQ ID NO 33

<211> LENGTH: 1113

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 33

```

atgggttcaa caggaataga gacccaaatg accccaaccc aaatatccga cgaagaagcc      60
aacctcttcg ccattgcaatt agccagtgcc tcagttctac ccattggttct caaagcagct      120
ttagagctcg acctcttgga gatcatagcc aaggccggtc caggcgcgtt tctctcacct      180
tccgacatag ctcaacagct tccgactcag aaccagacg ccccggtgat gctggaccgg      240
atgctgagac tgttggttag ctacaacgtg gtgacgtact cgctgcgtga gcgtgagacg      300
gcggaagagg aagggaaggt ggagaggctt tatgggttgg ctccggtgag taaatatctg      360
acgaagaatg aagatggagt ctccattgct cctctttgtc tcatgaacca ggataaggtt      420
cttatggaga gttggtatca cttaaagat gcagtacttg atggaggaat acctttcaac      480
aaggcatatg gaatgacagc atttgaatat catggaacgg atcaaagggt caataaaatc      540
tttaatatag gaatgtccga ccactcgact attaccatga aaaaaatcct cgaaacttac      600
aagggtttcg aggttcttaa ctcgattgtt gatgttggtg gtggtactgg agctgttggt      660
aacatgatcg tctctaagta cctactatt aagggtatta acttcgattt gcctcatgtc      720
atcgaagatg cacctccatt gaccggtgta gagcatgttg gaggagacat gtttgaagt      780
gtacaaaaag gagatgcaat tttcatgaag tggatttgcc atgattggag cgatgaacac      840
tgcttgaaat tcttgaagaa ctgccacgct gcactgcccg aacacggaaa agtgatcgtg      900
gcggagtgca ttcttccggt ggccaccggac tcgagccttg ccacaaagag tacggtcac      960
attgatgtga tcattgttgc ccataaccct ggtggcaaa agagaacaga gaaagagttt     1020
gaggcattgg ctaaggagc tggtctttaa ggcttcaaag tccattgcaa tgccttcaat     1080
accatataca tggaaatttct caagaccatt taa                                1113

```

-continued

```

<210> SEQ ID NO 34
<211> LENGTH: 400
<212> TYPE: PRT
<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 34

Met Val Phe Ser Ser Val Cys Ser Phe Pro Ser Ser Leu Gly Thr Asn
1          5          10          15
Phe Lys Leu Val Pro Arg Ser Asn Phe Lys Ala Ser Ser Ser His Tyr
20          25          30
His Glu Ile Asn Asn Phe Ile Asn Asn Lys Pro Ile Lys Phe Ser Tyr
35          40          45
Phe Ser Ser Arg Leu Tyr Cys Ser Ala Lys Pro Ile Val His Arg Glu
50          55          60
Asn Lys Phe Thr Lys Ser Phe Ser Leu Ser His Leu Gln Arg Lys Ser
65          70          75          80
Ser Ile Lys Ala His Gly Glu Ile Glu Ala Asp Gly Ser Asn Gly Thr
85          90          95
Ser Glu Phe Asn Val Met Lys Ser Gly Asn Ala Ile Trp Arg Phe Val
100         105         110
Arg Pro Tyr Ala Ala Lys Gly Val Leu Phe Asn Ser Ala Ala Met Phe
115         120         125
Ala Lys Glu Leu Val Gly Asn Leu Asn Leu Phe Ser Trp Pro Leu Met
130         135         140
Phe Lys Ile Leu Ser Phe Thr Leu Val Ile Leu Cys Ile Phe Val Ser
145         150         155         160
Thr Ser Gly Ile Asn Gln Ile Tyr Asp Leu Asp Ile Asp Arg Leu Asn
165         170         175
Lys Pro Asn Leu Pro Val Ala Ser Gly Glu Ile Ser Val Glu Leu Ala
180         185         190
Trp Leu Leu Thr Ile Val Cys Thr Ile Ser Gly Leu Thr Leu Thr Ile
195         200         205
Ile Thr Asn Ser Gly Pro Phe Phe Pro Phe Leu Tyr Ser Ala Ser Ile
210         215         220
Phe Phe Gly Phe Leu Tyr Ser Ala Pro Pro Phe Arg Trp Lys Lys Asn
225         230         235         240
Pro Phe Thr Ala Cys Phe Cys Asn Val Met Leu Tyr Val Gly Thr Ser
245         250         255
Val Gly Val Tyr Tyr Ala Cys Lys Ala Ser Leu Gly Leu Pro Ala Asn
260         265         270
Trp Ser Pro Ala Phe Cys Leu Leu Phe Trp Phe Ile Ser Leu Leu Ser
275         280         285
Ile Pro Ile Ser Ile Ala Lys Asp Leu Ser Asp Ile Glu Gly Asp Arg
290         295         300
Lys Phe Gly Ile Ile Thr Phe Ser Thr Lys Phe Gly Ala Lys Pro Ile
305         310         315         320
Ala Tyr Ile Cys His Gly Leu Met Leu Leu Asn Tyr Val Ser Val Met
325         330         335
Ala Ala Ala Ile Ile Trp Pro Gln Phe Phe Asn Ser Ser Val Ile Leu
340         345         350
Leu Ser His Ala Phe Met Ala Ile Trp Val Leu Tyr Gln Ala Trp Ile
355         360         365

```

-continued

Leu Glu Lys Ser Asn Tyr Ala Thr Glu Thr Cys Gln Lys Tyr Tyr Ile
370 375 380

Phe Leu Trp Ile Ile Phe Ser Leu Glu His Ala Phe Tyr Leu Phe Met
385 390 395 400

<210> SEQ ID NO 35
<211> LENGTH: 1203
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 35

```

atggtgttct catcagtttg tagttttcca tctcccttg gaactaattt taaattagtt      60
cctcgtagta attttaaggc atcatcttct cattatcatg aaataaataa ttttattaat    120
aataaaccaa ttaaattctc atatttttct tcaagactat attgctctgc caaaccaatt    180
gtacacagag aaaacaaatt cacaaaatca ttttactca gccacctcca aaggaaaagc    240
tccataaagg cacatggtga aattgaagct gatgggagta atggcacatc tgaatttaat    300
gtaatgaaaa gtggaaagc aatttgaga tttgtaaggc catatgcagc caaggagta      360
ttgtttaact ctgctgctat gtttgcaaaa gagttggagg ggaacctaaa tctatttagt    420
tggcctttga tgtttaagat actctctttt acattggtta ttttatgcac ttttgaagt     480
acaagtggca tcaatcaaat ttatgatctc gacatcgaca ggtaaaciaa acctaatttg    540
ccagtagcat caggagaaat ttcagttgaa ttggcatggt tgttgactat agtttgataga    600
ataagtggcc tcacattaac aattataacg aactcagggc cattcttccc tttctctac     660
tctgctagta tcttttttgg ctttctctat tctgctctc cattcagatg gaagaagaat    720
ccttttacag catgtttctg taatgttatg ttgtatgttg gcacaagcgt tgggtgtctat    780
tatgcttgta aggctagtct cgggcttcca gccaaactgga gccctgcttt ttgtttgctc    840
ttttggttta tttcattggt gagtatacc atctccattg caaaagatct ttcagacata     900
gaagtgacc gcaagtttgg aatcataacc ttctcaacta aatttgagc aaaaccata     960
gcataatatt gtcattgact catgctctg aattacgtga gtgttatggc tgcagctatt   1020
atttgccac agtttttcaa cagtagcgta atattgcttt ctcatgcatt catggcaatt   1080
tgggtattat atcaggcttg gatattggag aaatcaaatt acgccacgga gacgtgccaa   1140
aaatactata tattccttgg gataattttt tctcttgaac atgccttcta ttgttcatg   1200
tag                                              1203

```

<210> SEQ ID NO 36
<211> LENGTH: 523
<212> TYPE: PRT
<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 36

Met Ala Gln Val Ser Lys Ile Cys Ser Asn Gly Ala Gln Thr Ile Leu
1 5 10 15

Thr Leu Pro Asn Ile Ser Lys Ser His Thr Pro Arg Ser Leu Asn Ser
20 25 30

Val Ser Leu Arg Ser Pro Phe Leu Gly Ser Ser Asn Ser Leu Ser Leu
35 40 45

Lys Ile Gly Thr Glu Phe Gly Gly Cys Ser Thr Val Gly Lys Ala Met
50 55 60

-continued

Ala	Gly	Pro	Val	Met	Ala	Ser	Ala	Val	Thr	Ala	Glu	Lys	Pro	Ser	Lys	65	70	75	80
Val	Pro	Glu	Ile	Val	Leu	Gln	Pro	Ile	Lys	Asp	Ile	Ser	Gly	Thr	Val	85	90	95	
Lys	Leu	Pro	Gly	Ser	Lys	Ser	Leu	Ser	Asn	Arg	Ile	Leu	Leu	Leu	Ala	100	105	110	
Ala	Leu	Ser	Glu	Gly	Thr	Thr	Val	Val	Asp	Asn	Leu	Leu	Asp	Ser	Asp	115	120	125	
Asp	Ile	His	Tyr	Met	Leu	Gly	Ala	Leu	Glu	Thr	Leu	Gly	Leu	Arg	Val	130	135	140	
Glu	Ala	Asp	Lys	Glu	Ser	Lys	Arg	Ala	Ile	Val	Glu	Gly	Cys	Ala	Gly	145	150	155	160
Gln	Phe	Pro	Ala	Gly	Lys	Glu	Ser	Val	Asp	Glu	Val	Gln	Leu	Phe	Leu	165	170	175	
Gly	Asn	Ala	Gly	Thr	Ala	Met	Arg	Pro	Leu	Thr	Ala	Ala	Val	Thr	Val	180	185	190	
Ala	Gly	Gly	Asn	Ala	Ser	Tyr	Val	Leu	Asp	Gly	Val	Pro	Arg	Met	Arg	195	200	205	
Glu	Arg	Pro	Ile	Gly	Asp	Leu	Val	Thr	Gly	Leu	Lys	Gln	Leu	Gly	Ala	210	215	220	
Asp	Val	Asp	Cys	Phe	His	Gly	Thr	Asp	Cys	Pro	Pro	Val	Arg	Val	Leu	225	230	235	240
Gly	Lys	Gly	Gly	Leu	Pro	Gly	Gly	Lys	Val	Lys	Leu	Ser	Gly	Ser	Ile	245	250	255	
Ser	Ser	Gln	Tyr	Leu	Thr	Ala	Leu	Leu	Met	Ala	Ala	Pro	Leu	Ala	Leu	260	265	270	
Gly	Asp	Val	Glu	Ile	Glu	Ile	Ile	Asp	Lys	Leu	Ile	Ser	Val	Pro	Tyr	275	280	285	
Val	Asp	Met	Thr	Leu	Lys	Leu	Met	Ala	Arg	Phe	Gly	Val	Thr	Val	Glu	290	295	300	
His	Ser	Asp	Ser	Trp	Asp	Arg	Phe	Leu	Val	Lys	Gly	Gly	Gln	Lys	Tyr	305	310	315	320
Lys	Ser	Pro	Gly	Asn	Ala	Tyr	Val	Glu	Gly	Asp	Ala	Ser	Ser	Ala	Ser	325	330	335	
Tyr	Phe	Leu	Ala	Gly	Ala	Ala	Val	Thr	Gly	Gly	Thr	Val	Thr	Val	Glu	340	345	350	
Gly	Cys	Gly	Thr	Ser	Ser	Leu	Gln	Gly	Asp	Val	Lys	Phe	Ala	Glu	Val	355	360	365	
Leu	Glu	Lys	Met	Gly	Ala	Lys	Val	Ser	Trp	Thr	Glu	Asn	Ser	Val	Thr	370	375	380	
Val	Thr	Gly	Pro	Pro	Arg	Asp	Ser	Val	Lys	Ser	Lys	His	Leu	Lys	Ala	385	390	395	400
Ile	Asp	Val	Asn	Met	Asn	Lys	Met	Pro	Asp	Val	Ala	Met	Thr	Leu	Ala	405	410	415	
Val	Val	Ala	Leu	Phe	Ala	Asp	Gly	Pro	Thr	Ala	Ile	Arg	Asp	Val	Ala	420	425	430	
Ser	Trp	Arg	Val	Lys	Glu	Thr	Glu	Arg	Met	Ile	Ala	Ile	Cys	Thr	Glu	435	440	445	
Leu	Arg	Lys	Leu	Gly	Ala	Thr	Val	Glu	Glu	Gly	Pro	Asp	Tyr	Cys	Val	450	455	460	
Ile	Thr	Pro	Pro	Glu	Lys	Leu	Asn	Ile	Thr	Ala	Ile	Asp	Thr	Tyr	Asp				

-continued

465	470	475	480
Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ser Asp Val			
	485	490	495
Pro Val Thr Ile Lys Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro Asp			
	500	505	510
Tyr Phe Glu Val Leu Glu Arg Phe Thr Lys His			
	515	520	

<210> SEQ ID NO 37
 <211> LENGTH: 523
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- engineered protein

<400> SEQUENCE: 37

Met Ala Gln Val Ser Lys Ile Cys Ser Asn Gly Ala Gln Thr Ile Leu			
1	5	10	15
Thr Leu Pro Asn Ile Ser Lys Ser His Thr Pro Arg Ser Leu Asn Ser			
	20	25	30
Val Ser Leu Arg Ser Pro Phe Leu Gly Ser Ser Asn Ser Leu Ser Leu			
	35	40	45
Lys Ile Gly Thr Glu Phe Gly Gly Cys Ser Thr Val Gly Lys Ala Met			
	50	55	60
Ala Gly Pro Val Met Ala Ser Ala Val Thr Ala Glu Lys Pro Ser Lys			
	65	70	75
Val Pro Glu Ile Val Leu Gln Pro Ile Lys Asp Ile Ser Gly Thr Val			
	85	90	95
Lys Leu Pro Gly Ser Lys Ser Leu Ser Asn Arg Ile Leu Leu Leu Ala			
	100	105	110
Ala Leu Ser Glu Gly Thr Thr Val Val Asp Asn Leu Leu Asp Ser Asp			
	115	120	125
Asp Ile His Tyr Met Leu Gly Ala Leu Glu Thr Leu Gly Leu Arg Val			
	130	135	140
Glu Ala Asp Lys Glu Ser Lys Arg Ala Ile Val Glu Gly Cys Ala Gly			
	145	150	155
Gln Phe Pro Ala Gly Lys Glu Ser Val Asp Glu Val Gln Leu Phe Leu			
	165	170	175
Gly Asn Ala Gly Ile Ala Met Arg Ser Leu Thr Ala Ala Val Thr Val			
	180	185	190
Ala Gly Gly Asn Ala Ser Tyr Val Leu Asp Gly Val Pro Arg Met Arg			
	195	200	205
Glu Arg Pro Ile Gly Asp Leu Val Thr Gly Leu Lys Gln Leu Gly Ala			
	210	215	220
Asp Val Asp Cys Phe His Gly Thr Asp Cys Pro Pro Val Arg Val Leu			
	225	230	235
Gly Lys Gly Gly Leu Pro Gly Gly Lys Val Lys Leu Ser Gly Ser Ile			
	245	250	255
Ser Ser Gln Tyr Leu Thr Ala Leu Leu Met Ala Ala Pro Leu Ala Leu			
	260	265	270
Gly Asp Val Glu Ile Glu Ile Ile Asp Lys Leu Ile Ser Val Pro Tyr			
	275	280	285
Val Asp Met Thr Leu Lys Leu Met Ala Arg Phe Gly Val Thr Val Glu			

-continued

290	295	300
His Ser Asp Ser Trp Asp Arg Phe Leu Val Lys Gly Gly Gln Lys Tyr		
305	310	315 320
Lys Ser Pro Gly Asn Ala Tyr Val Glu Gly Asp Ala Ser Ser Ala Ser		
	325	330 335
Tyr Phe Leu Ala Gly Ala Ala Val Thr Gly Gly Thr Val Thr Val Glu		
	340	345 350
Gly Cys Gly Thr Ser Ser Leu Gln Gly Asp Val Lys Phe Ala Glu Val		
	355	360 365
Leu Glu Lys Met Gly Ala Lys Val Ser Trp Thr Glu Asn Ser Val Thr		
	370	375 380
Val Thr Gly Pro Pro Arg Asp Ser Val Lys Ser Lys His Leu Lys Ala		
385	390	395 400
Ile Asp Val Asn Met Asn Lys Met Pro Asp Val Ala Met Thr Leu Ala		
	405	410 415
Val Val Ala Leu Phe Ala Asp Gly Pro Thr Ala Ile Arg Asp Val Ala		
	420	425 430
Ser Trp Arg Val Lys Glu Thr Glu Arg Met Ile Ala Ile Cys Thr Glu		
	435	440 445
Leu Arg Lys Leu Gly Ala Thr Val Glu Glu Gly Leu Asp Tyr Cys Val		
	450	455 460
Ile Thr Pro Pro Glu Lys Leu Asn Ile Thr Ala Ile Asp Thr Tyr Asp		
465	470	475 480
Asp His Arg Met Ala Met Ala Phe Ser Leu Ala Ala Cys Ser Asp Val		
	485	490 495
Pro Val Thr Ile Lys Asp Pro Gly Cys Thr Arg Lys Thr Phe Pro Asp		
	500	505 510
Tyr Phe Glu Val Leu Glu Arg Phe Thr Lys His		
	515	520

<210> SEQ ID NO 38

<211> LENGTH: 1572

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 38

```

atggcccaag tgagcaaaat ctgtagcaat ggagctcaaa ctatccttac tctcccaaat    60
atatctaagt ctcatacacc aagatcccta aattcagttt cgttgagatc accgtttttg    120
ggttcatcta actctttgag tttgaagatt ggaactgaat ttgggggttg ttctacgggt    180
ggtaaagcta tggtggtgcc agtcatggct tcagctgtca cagcggagaa gccttcaaag    240
gtaccggaga ttgtgttgca gccattaaa gatattcttg gcaactgtcaa gttgccgggt    300
tccaagtcac tatcgaatcg gattctactc ctggctgctc tttctgaggg gacaactgtt    360
gtggacaact tgtagatag tgatgacatt cactacatgc ttggtgcctt ggaaaccctt    420
ggtcttcctg ttgaagcaga caaggaaagc aaacgagcaa ttgtggaagg ttgtgcgggt    480
cagtttctcg caggtaaaga atctgttgac gaagttcaac ttttcttgg aaatgctgga    540
acagcaatgc gtccactcac agctgcgggt actgttgctg gtggaaatgc tagctacgta    600
cttgatggtg ttctctgaat gagagaaaga ccaattggag atttggtgac tggctttaag    660
cagcttggtg cagatgttga ttgttttcat ggtacggatt gtccccctgt tcgtgtgctt    720

```

-continued

ggaaaaggag gccttcctgg gggcaagggtg aaactttctg gatcaattag cagtcaatat	780
ttgacagcct tgcttatggc agtcaccttg gctcttgag atgttgaaat cgagataatt	840
gataaattga tctcggttcc ctatgttgat atgactttga agttgatggc acgttttggg	900
gttactgttg aacacagtga tagctgggat cgatttttag ttaaaggagg tcaaaagtac	960
aaatctcctg gaaacgctta tgttgaaggt gatgcttcaa gtgctagtta cttcctagct	1020
ggtgctgcag tcaactggtg tacagtcacc gtagaagggt gtgggactag tagtttacag	1080
ggagacgtaa aatttgctga agttcttgag aaaatgggtg cttaaagttag ctggacagag	1140
aacagtgtca cggtcactgg accaccacga gattctgtaa aaagtaaaca cttgaaagcc	1200
attgatgtca acatgaacaa aatgcctgat gttgccatga ctcttgctgt agttgctctt	1260
tttgctgatg gcccactgac tataagagat gtggcaagtt ggagagtcaa ggagacagag	1320
agaatgattg ccactctcac tgaactcaga aagcttgagg caacagtga ggaaggaccc	1380
gattactgag tgatcactcc accagagaaa ctaaatatca cagcaataga cacatacgac	1440
gaccacagga tggctatggc gttctctctt gcagcttggt cagatgtgcc agttaccatt	1500
aaggatcctg gttgcaccgc aaaaactttc ccagattact ttgaagtcct tgagagattt	1560
acaaagcact ga	1572

<210> SEQ ID NO 39

<211> LENGTH: 1572

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 39

atggcccaag tgagcaaaat ctgtagcaat ggagctcaaa ctatccttac tctcccaaat	60
atatctaagt ctcatacacc aagatcccta aattcagttt cgttgagatc accgtttttg	120
ggttcaccta actctttgag tttgaagatt ggaactgaat ttgggggttg ttctacgggt	180
ggtaaagcta tggctggtcc agtcatggct tcagctgtca cagcggagaa gccttcaaag	240
gtaccggaga ttgtgttgca gccattaaa gatatctctg gcaactgtcaa gttgcgggt	300
tccaagtcac tatcgaatcg gattctactc ctggctgtct tttctgaggg gacaactgtt	360
gtggacaact tgtagatag tgatgacatt cactacatgc ttggtgcctt ggaaccctt	420
ggtcttcctg ttgaagcaga caaggaaagc aaacgagcaa ttgtggaagg ttgtgcgggt	480
cagtttcctg caggtaaaga atctgttgac gaagttcaac tttccttggt aaatgctgga	540
atagcaatgc gttcactcac agctgcgggt actgttgctg gtggaaatgc tagctacgta	600
cttgatggtg ttcctcgaat gagagaaaga ccaattggag atttggtgac tggctttaag	660
cagcttggtg cagatgttga ttgttttcat ggtacggatt gtccccctgt tcgtgtgctt	720
ggaaaaggag gccttcctgg gggcaagggtg aaactttctg gatcaattag cagtcaatat	780
ttgacagcct tgcttatggc agtcaccttg gctcttgag atgttgaaat cgagataatt	840
gataaattga tctcggttcc ctatgttgat atgactttga agttgatggc acgttttggg	900
gttactgttg aacacagtga tagctgggat cgatttttag ttaaaggagg tcaaaagtac	960
aaatctcctg gaaacgctta tgttgaaggt gatgcttcaa gtgctagtta cttcctagct	1020
ggtgctgcag tcaactggtg tacagtcacc gtagaagggt gtgggactag tagtttacag	1080

-continued

ggagacgtaa aatttgctga agttcttgag aaaatgggtg cttaaagttag ctggacagag	1140
aacagtgtca cggtcactgg accaccacga gattctgtaa aaagtaaaca cttgaaagcc	1200
attgatgtca acatgaacaa aatgctgat gttgccatga ctcttgctgt agttgctctt	1260
tttgctgatg gccccactgc tataagagat gtggcaagtt ggagagtcaa ggagacagag	1320
agaatgattg ccatctgcac tgaactcaga aagcttggag caacagttga ggaaggactt	1380
gattactgcg tgatcactcc accagagaaa ctaaatatca cagcaataga cacatacgac	1440
gaccacagga tggctatggc gttctctctt gcagcttggt cagatgtgcc agttaccatt	1500
aaggatcctg gttgcaccgc aaaaactttc ccagattact ttgaagtcct tgagagattt	1560
acaaagcact ga	1572

<210> SEQ ID NO 40

<211> LENGTH: 4196

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 40

ggttggtaag ccctcctacc ctctttgaaa attgaaagag agtcaatgtc gacctacagc	60
agcagcatcc attaacgtta ccattgccac caaaaatcca acctttatct gtatagagag	120
aatcagagaa ggtttgggtt tcagagagag agagaagaag aacaaaaaaa tggccaagt	180
gagcaaaatc tgtagcaatg gagctcaaac tatccttact ctcccaaata tatctaagtc	240
tcatacacca agatccctaa attcagtttc gttgagatca ccgttttttg gttcatctaa	300
ctctttgagt ttgaagattg gaactgaatt tgggggttgt tctacggttg gtaaagctat	360
ggctgggtcca gtcattggct cagctgtcac agcggagaag ccttcaaagg tacggagat	420
tgtgttcag cccattaaag atatctctgg cactgtcaag ttgccgggtt ccaagtcact	480
atcgaatcgg attctactcc tggctgctct ttctgaggtt tatttcattt tttttaaac	540
gtcaaacatg tattttgtgc gaggaagttt tctgtatata caaagataag agagtaaaaa	600
tatggaacat caataccaaa atgaacaaa actaggctaa gctatcaaat catgtcatgg	660
tatgccatcc tctactttcc tatctcaagc tccacagcta taaaatacta tatcgtaatt	720
atcttgtcaa ctgcttccat attccttgta atttccctca ttcccaacta aactagttcc	780
aatggattgt gtggctggaa actgtagtta gttacattag ctagatctga accatgatca	840
gcacgcactg cccaactggg aaacctatga attgcatgga attcttcctt tggtatccac	900
aaatttgaaa agtatttttg aggtatacaa agattgtgct ttttatgagc aattttcttt	960
tagttttatg ttaagagttt gtagcgatgg gatgtttttt ttctagaaaa tggacagtaa	1020
agcttagcat ttttacttta ttggtgtaaa tgaatagtgt tcattgaagc tgaactcatg	1080
cccttaattg ggaggaaaat tgagagaaat ggagtaaagt aatatgatat ttgggttaaa	1140
ttcgtaaaga tatgatggaa ataaaaaatg caactcaact gggttactga agttatattt	1200
ctggtctcag ttgtgctttt acaactttag tctagagctc cacgctcggg agagattcgg	1260
agtccttaca gtttattttg ataagtattt atgagaattt cataactcta cgcttttggt	1320
acattatata tgagggtgtt cgttgggtga ttgtctcacc tgaactccct aaattttaga	1380
atgtgggatt tagaatgaag ttatactatt agtgtttgag tcatctagaa ttgttagctg	1440
ctcattctcc atatactctt tctctatttc ctccccatat ttggcgcta ctacttatct	1500

-continued

ttacagttca	tggtattttc	atgtacttga	gtttttttgcc	ctataaaata	ttttgagcgg	1560
tgggaagtaa	ctgttttttt	tggtataaatt	atccagggga	caactgttgt	ggacaacttg	1620
ttagatagtg	atgacattca	ctacatgctt	ggtgccttgg	aaacccttgg	tcttcgtgtt	1680
gaagcagaca	aggaaagcaa	acgagcaatt	gtggaaggtt	gtgcgggtca	gtttcctgca	1740
ggtaagaat	ctgttgacga	agttcaactt	ttccttggaa	atgctggaac	agcaatgcgt	1800
ccactcacag	ctgcggtgac	tggtgctgg	ggaaatgcta	ggtttgtctt	cattgcaatt	1860
gcttttgaat	ataaagtact	tctaatagcag	tgaatttatg	ctcttgtttt	tcttactggc	1920
cgagtagctc	ttacatttta	ggtaaagaaa	gtcacttttg	ctaacaacat	caccatttat	1980
acttccctct	ttactttgat	gtgggtatgc	tagaaattac	atgttggaac	tgaactagca	2040
catatcataa	attattttgt	atgctgttat	tacattttct	cagtaacctc	ttacttcta	2100
tatctcagct	acgtacttga	tggtgttcct	cgaatgagag	aaagaccaat	tggagatttg	2160
gtgactggtc	ttaagcagct	tggtgcagat	gttgattgtt	ttcatggtag	ggattgtccc	2220
cctgttcgtg	tgcttggaac	aggaggcctt	cctgggggca	aggtgaggct	tgcattgctt	2280
cttcttattc	tttttgccca	taaaacatca	ttgtaaatgt	ggttttatgt	tatgaaatcc	2340
attgactggg	ttatttttag	gttggtgttt	tgcttttaaa	taaaaacaat	attgtcaaat	2400
gatgcataag	tagtgattac	atctacatca	tttaatttat	tatcttaaat	gatgacaaac	2460
ttcatcattt	tgactcagaa	ttatgtaata	ttaccctttg	caggtgaaac	tttctggatc	2520
aattagcagt	caatatattga	cagccttgct	tatggcagct	cccttggtct	ttggagatgt	2580
tgaaatcgag	ataattgata	aattgatctc	ggttccttat	gttgatatga	ctttgaagtt	2640
gatggcacgt	tttgggggta	ctgttgaaca	cagtgatagc	tggtgatcga	ttttagttaa	2700
aggagggtcaa	aagtacaagt	aggtttcttc	tgaatatagt	tgatagtatt	gttacattac	2760
atctgggttat	gtcaaagagt	aataaattga	aaaataaaaa	tctgtcagat	ctcctggaaa	2820
cgcttatgtt	gaagggtgatg	cttcaagtgc	tagttacttc	ctagctgggtg	ctgcagtcac	2880
tggttggtaca	gtcaccttag	aagggtgtgg	gactagtagt	ttacaggtag	tttgcttaga	2940
ccttgaaatc	tcttattctt	gtacttgtgt	ttacatagaa	tctaagatta	agtgtattta	3000
catacattaa	ctggtgttta	ataaaggag	acgtaaaatt	tgctgaagtt	cttgagaaaa	3060
tgggtgctaa	agttagctgg	acagagaaca	gtgtcacggt	cactggacca	ccacgagatt	3120
ctgtaaaaag	taaacacttg	aaagccattg	atgtcaacat	gaacaaaatg	cctgatgttg	3180
ccatgactct	tgctgtagtt	gctctttttg	ctgatggccc	cactgctata	agagatggta	3240
tgtttttctt	taaatttggtg	agatggtaaa	atggggcag	cggtttgggt	tggggtagat	3300
tatcgggttct	cgtctgccat	aataaaaaat	aatctgctca	tttgcaataa	atttcacaga	3360
cacaaaaatg	aaaaccaata	aaatattatt	ttgttttagag	gattaaatac	tcatttcttg	3420
cctttcctaa	ttccagtggtg	caagttggag	agtcaggag	acagagagaa	tgattgccat	3480
ctgcactgaa	ctcagaaagg	ttagttttta	tgctgtttta	tgtacttgtt	atgtcatgctg	3540
ccttggaatg	taattggctga	tagctatctg	ttcttatggg	aacaaacatt	tcagcttgga	3600
gcaacagttg	aggaaggacc	cgattactgc	gtgatcactc	caccagagaa	actaaatctc	3660
acagcaatag	acacatacga	cgaccacagg	atggctatgg	cggtctctct	tgcagcttgt	3720
tcagatgtgc	cagttacat	taaggatcct	ggttgcaccc	gaaaaacttt	cccagattac	3780

-continued

```

tttgaagtcc ttgagagatt tacaaagcac tgaatgagta tttattaact ggatagagaa 3840
caatagcatc ggctactgtc attacaacta aagcagttgg gaggcaggca atccttttca 3900
attatcatgt gtttgatttt ggtcgactgt attgcaagtt gagcttctca ttattattaa 3960
gactgtaatc gtagttatct gttgtaactt ctgcaaacc cttcatgttat tttttcacc 4020
ttctaataag ccagtggggc aaattctata tctgctatat gagcttgagt gtagagagaa 4080
cttttgtcaa tgtataggtt tctagcagaa gcaccatccc taatatgctt tattataaga 4140
gttgctgtga tcgtgtagtg ttattttatt gaaagtaacc gacgcatctc atatta 4196

```

```

<210> SEQ ID NO 41
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- prime editing guide RNA

```

```

<400> SEQUENCE: 41

```

```

tgaagacttt gcacaacttt tccttggaaa tgcgtttaag tcttct 46

```

```

<210> SEQ ID NO 42
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- prime editing guide RNA

```

```

<400> SEQUENCE: 42

```

```

agaagactta aacgcatttc caaggaaaag ttgtgcaaag tcttca 46

```

```

<210> SEQ ID NO 43
<211> LENGTH: 64
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

```

```

<400> SEQUENCE: 43

```

```

acctgctata gtgctgagtg aacgcattgc tattccagca tttccaagga aacatatagc 60
aggt 64

```

```

<210> SEQ ID NO 44
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- prime editing guide RNA

```

```

<400> SEQUENCE: 44

```

```

tgaagacttt gcacttggag caacagttga ggagttaag tcttct 46

```

```

<210> SEQ ID NO 45
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- prime editing guide RNA

```

```

<400> SEQUENCE: 45

```

```

agaagactta aactcctcaa ctgttgctcc aagtgcaaag tcttca 46

```

-continued

```

<210> SEQ ID NO 46
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 46

```

```

acctgctata gtgccacgca gtaatcaagt ccttcctcaa ctgttgaaca tatagcaggt      60

```

```

<210> SEQ ID NO 47
<211> LENGTH: 349
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 47

```

```

ggtctcaaat ggcgacaggt tctcgtgttc tgatcgggtc agcaatgac ctcataatct      60
caggagaact gctagttcca gggcaaggaa cgtgccaaagg agacatagag ggtctgatga    120
gagaatgtgc ggtctacgtc cagcgtccag gcccaaagg aaacccatcc gcagcgtgtt    180
gcaaagtcgt caagagatca gacatccct gcgcgtgtgg ccgtatcaca ccctcggttc    240
aaaaaatgat agacatgaat aagggtgttc ttgtcacttc cttttgtggg aggcctctcg    300
ctcatggtac caagtgtgga agctacattg tgccatgagc ttagagacc                349

```

```

<210> SEQ ID NO 48
<211> LENGTH: 348
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 48

```

```

ggtctcaaat ggcgacaggt tctcgtgttc tgatcgggtc agcaatgac ctcataatct      60
caggagaact gctagttcca gggcaaggaa cgtgccaaagg agacatagag ggtctgatga    120
gagaatgtgc ggtctacgtc cagcgtccag gcccaaagg aaacccatcc gcagcgtgtt    180
gcaaagtcgt caagagatca gacatccct gcgcgtgtgg ccgtatcaca ccctcggttc    240
aaaaaatgat agacatgaat aagggtgttc ttgtcacttc cttttgtggg aggcctctcg    300
ctcatggtac caagtgtgga agctacattg tgccaagttc gagagacc                348

```

```

<210> SEQ ID NO 49
<211> LENGTH: 1207
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 49

```

```

ggtctcaaat gggactctca tcagtttgta ccttttcatt tcaaactaat taccatactt      60
tattaaatcc tcacaataat aatcccaaaa cctcattatt atgttatcga cccccaaaa    120
caccaattaa atactcttac aataattttc cctctaaaca ttgctccacc aagagttttc    180
atctacaaaa caaatgtcga gaatcattat caatcgcaa aaattccatt agggcagcta    240
ctacaaatca aactgagcct ccagaatctg ataatcattc agtagcaact aaaattttaa    300

```

-continued

actttgggaa ggcattgttg aaacttcaaa gaccatatac aatcatagca tttacttcat	360
gcgcttgttg attgtttggg aaagagttgt tgcataaac aaatttaata agttggtcac	420
tgatgttcaa ggcattcttt tttttggttg ctgtattatg cattgcttct tttacaacta	480
ccatcaatca gatttacgat cttcacattg acagaataaa caagcctgat ctaccactag	540
cttcagggga aatatcagta aacacagctt ggattatgag cataattgtg gcactgtttg	600
gattgataat aactataaaa atgaagggtg gaccactcta tatatttggc tactgttttg	660
gtatttttgg tgggattgtc tattctgttc caccatttag atggaagcaa aatccttcca	720
ctgcatttct tctcaatttc ctggcccata ttattacaaa tttcacattt tattatgcca	780
gcagagcagc tcttggccta ccatttgagt tgaggccttc ttttacttcc ctgctagcat	840
ttatgaaatc aatgggttca gctttggctt taatcaaaga tgcttcagac gttgaaggcg	900
acactaaatt tggcatatca accttggcaa gttaaattgg ttccagaaac ttgacattat	960
tttgttctgg aattgttctc ctatcctatg tggtgtctat acttgcctgg attatctggc	1020
cccaggcttt caacagtaac gtaattgtac tttctcatgc aatcttagca ttttggttaa	1080
tcctccagac tcgagatttt gcgttaacaa attacgacct ggaagcaggc agaagatttt	1140
acgagttcat gtggaagctt tattatgtct aatatttagt atatgttttc atataagctt	1200
agagacc	1207

<210> SEQ ID NO 50

<211> LENGTH: 1206

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 50

ggtctcaaat gggactctca tcagtttgta ccttttcatt tcaaactaat taccatactt	60
tattaaatcc tcacaataat aatcccaaaa cctcattatt atgttatcga cccccaaaa	120
caccaattaa atactcttac aataattttc cctctaaaca ttgtccacc aagagttttc	180
atctacaaaa caaatgtcga gaatcattat caatcgcaaa aaattccatt agggcagcta	240
ctacaaatca aactgagcct ccagaatctg ataatcattc agtagcaact aaaattttaa	300
actttgggaa ggcattgttg aaacttcaaa gaccatatac aatcatagca tttacttcat	360
gcgcttgttg attgtttggg aaagagttgt tgcataaac aaatttaata agttggtcac	420
tgatgttcaa ggcattcttt tttttggttg ctgtattatg cattgcttct tttacaacta	480
ccatcaatca gatttacgat cttcacattg acagaataaa caagcctgat ctaccactag	540
cttcagggga aatatcagta aacacagctt ggattatgag cataattgtg gcactgtttg	600
gattgataat aactataaaa atgaagggtg gaccactcta tatatttggc tactgttttg	660
gtatttttgg tgggattgtc tattctgttc caccatttag atggaagcaa aatccttcca	720
ctgcatttct tctcaatttc ctggcccata ttattacaaa tttcacattt tattatgcca	780
gcagagcagc tcttggccta ccatttgagt tgaggccttc ttttacttcc ctgctagcat	840
ttatgaaatc aatgggttca gctttggctt taatcaaaga tgcttcagac gttgaaggcg	900
acactaaatt tggcatatca accttggcaa gttaaattgg ttccagaaac ttgacattat	960
tttgttctgg aattgttctc ctatcctatg tggtgtctat acttgcctgg attatctggc	1020

-continued

```

cccaggcttt caacagtaac gtaatgttac tttctcatgc aatcttagca ttttggttaa 1080
tctccagac tcgagatttt gcgttaacaa attacgaccc ggaagcaggc agaagatttt 1140
acgagttcat gtggaagcgt tattatgctg aatatttagt atatgttttc ataagttcga 1200
gagacc                                           1206

```

```

<210> SEQ ID NO 51
<211> LENGTH: 1132
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

```

```

<400> SEQUENCE: 51

```

```

ggtctcaaat gggttcaaca ggaatagaaa cccaaatgac cccaaccaa atatccgacg 60
aagaagccaa cctcttcgcc atgcaattag ccagtgcctc agtcttacc atggtttctca 120
aagcagcttt agagctcgac ctcttgaga tcatagccaa ggccggtcca ggcgcgtttc 180
tctcaccttc cgacatagct caacagcttc cgactcagaa ccagacgcc ccggtgatgc 240
tggaccggat gctgagactg ttggctagct acaacgtggt gacgtactcg ctgcgtgagc 300
gtgaaacggc ggaagaggaa ggaaggtgg agaggcttta tgggttggt ccggtgagta 360
aatatctgac gaagaatgaa gatggagtct ccattgctcc tctttgtctc atgaaccagg 420
ataaggttct tatggagagt tggatcact taaaagatgc agtacttgat ggaggaatac 480
ctttcaacaa ggcataatga atgacagcat ttgaatatca tggaaacgat caaaggttca 540
ataaaatctt taatagagga atgtccgacc actcgactat taccatgaaa aaaatcctcg 600
aaacttacaa gggtttcgag ggtcttaact cgattgtga tgggttggt ggtactggag 660
ctgttggtta catgatcgtt tctaagtacc ctactattaa ggggtattaac ttcgatttgc 720
ctcatgtcat cgaagatgca cctccattga ccggtgtaga gcatgttgga ggagacatgt 780
ttgtaagtgt accaaaagga gatgcaattt tcatgaagtg gatttgccat gattggagcg 840
atgaacactg cttgaaatc ttgaagaact gccacgtgc actgccgaa cacggaaaag 900
tgatcgtggc ggagtgcatt cttccggtgg caccggactc gagccttgcc acaaagagta 960
cggccacat tgatgtgatc atgttgccc ataaccctgg tggcaaagag agaacagaga 1020
aagagtttga ggcattggct aagggagctg gctttaaagg cttcaaagtc cattgcaatg 1080
ctttcaatac ccatatcatg gaatttctca agaccattta agcttagaga cc 1132

```

```

<210> SEQ ID NO 52
<211> LENGTH: 1131
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene

```

```

<400> SEQUENCE: 52

```

```

ggtctcaaat gggttcaaca ggaatagaaa cccaaatgac cccaaccaa atatccgacg 60
aagaagccaa cctcttcgcc atgcaattag ccagtgcctc agtcttacc atggtttctca 120
aagcagcttt agagctcgac ctcttgaga tcatagccaa ggccggtcca ggcgcgtttc 180
tctcaccttc cgacatagct caacagcttc cgactcagaa ccagacgcc ccggtgatgc 240
tggaccggat gctgagactg ttggctagct acaacgtggt gacgtactcg ctgcgtgagc 300

```

-continued

gtgaaacggc ggaagaggaa ggaagagtg agaggttta tgggttggt ccggtgagta	360
aatatctgac gaagaatgaa gatggagtct ccattgctcc tctttgtctc atgaaccagg	420
ataaggttct tatggagagt tggatcact taaaagatgc agtacttgat ggaggaatac	480
ctttcaacaa ggcatatgga atgacagcat ttgaatatca tggaaccgat caaaggttca	540
ataaaatctt taatagagga atgtccgacc actcgactat taccatgaaa aaaatcctcg	600
aaacttacaa gggtttcgag ggtcttaact cgattgttga tgttggtggt ggtactggag	660
ctgttgtaa catgatcgtt tctaagtacc ctactattaa gggatttaac ttcgatttgc	720
ctcatgtcat cgaagatgca cctccattga ccggtgtaga gcatgttga ggagacatgt	780
ttgtaagtgt accaaaagga gatgcaattt tcatgaagtg gatttgccat gattggagcg	840
atgaacactg cttgaaattc ttgaagaact gccacgctgc actgccgaa cacggaaaag	900
tgatcgtggc ggagtgcatt ctcccggtgg caccggactc gagccttgcc acaaagagta	960
cgggtccacat tgatgtgatc atgttggccc ataaccctgg tggcaaagag agaacagaga	1020
aagagtttga ggcattggct aaggagctg gctttaagg cttcaaagtc cattgcaatg	1080
ctttcaatc ccatatcatg gaatttctca agaccattag ttcgagagac c	1131

<210> SEQ ID NO 53

<211> LENGTH: 1222

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 53

ggtctcaaat ggtgttctca tcagtttgta gttttccatc ctcccttgga actaatTTTA	60
aattagttcc tcgtagtaat ttaaggcat catcttctca ttatcatgaa ataaataatt	120
ttattaataa taaaccaatt aaattctcat atttttcttc aagactatat tgctctgcca	180
aaccaattgt acacagagaa aacaaattca caaatcatt ttcactcagc cacctccaaa	240
ggaaaagctc cataaaggca catggtgaaa ttgaagctga tgggagtaat ggcacatctg	300
aatttaattg aatgaaaagt ggaacgcaa tttggagatt tgtaaggcca tatgcagcca	360
agggagtatt gtttaactct gctgctatgt ttgcaaaaga gttgggtggg aacctaaatc	420
tatttagttg gcctttgatg ttaagatac tctcttttac attggttatt ttatgcattt	480
ttgtaagtac aagtggcatc aatcaaatTT atgatctcga catcgacagg ttaaacaac	540
ctaatttgcc agtagcatca ggagaaattt cagttgaatt ggcattggtt ttgactatag	600
ttgtacaat aagtggctc acattaacaa ttataacgaa ctgagggcca ttcttccctt	660
ttctctactc tgctagtatc ttttttggt tctctattc tgctcctcca ttcagatgga	720
agaagaatcc ttttacagca tgtttctgta atgttatgtt gtatgttggc acaagcgttg	780
gtgtctatta tgcttgtaag gctagtctcg ggcttcacgc caactggagc cctgcttttt	840
gtttgctctt ttggtttatt tcattgttga gtatacccat ctccattgca aaagatcttt	900
cagacataga aggtgaccgc aagtttgga tcataacctt ctcaactaaa tttggagcaa	960
aacctatagc atatatttgt catggactca tgcttctgaa ttacgtgagt gttatggctg	1020
cagctattat ttggccacag tttttcaaca gtacgtaat attgctttct catgcattca	1080
tggcaatttg ggtattatat caggcttga tattggagaa atcaaatTAC gccacggaaa	1140

-continued

cgtgccaaaa atactatata ttcccttgga taatttttcc tcttgaacat gccttctatt	1200
tgttcatgta ggcttagaga cc	1222

<210> SEQ ID NO 54
 <211> LENGTH: 1221
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- engineered gene

<400> SEQUENCE: 54

ggtctcaaat ggtgttctca tcagtttgta gttttccatc ctcccttgga actaatTTTA	60
aattagttcc tcgtagtaat ttaaggcat catcttctca ttatcatgaa ataaataatt	120
ttattaataa taaaccaatt aaattctcat atttttcttc aagactatat tgctctgcca	180
aaccaattgt acacagagaa aacaaattca caaaatcatt ttcactcagc cacctccaaa	240
ggaaaagctc cataaaggca catggtgaaa ttgaagctga tgggagtaat ggcacatctg	300
aatttaattgt aatgaaaagt ggaaacgcaa ttggagatt tgtaaggcca tatgcagcca	360
agggagtatt gtttaactct gctgctatgt ttgcaaaaga gttgggtggg aacctaaatc	420
tatttagttg gcctttgatg ttaagatac tctcttttac attggttatt ttatgcattt	480
ttgtaagtac aagtggcatc aatcaaattt atgatctcga catcgacagg ttaaacaac	540
ctaatttgcc agtagcatca ggagaaattt cagttgaatt ggcatggttg ttgactatag	600
tttgtaaat aagtggcctc acattaacaa ttataacgaa ctcaggggcca ttcttccctt	660
ttctctactc tgctagtatc ttttttggtt ttctctatc tgctctcca ttcagatgga	720
agaagaatcc ttttacagca tgtttctgta atgttatgtt gtatggtggc acaagcgttg	780
gtgtctatta tgcttgtaag gctagtctcg ggcttccagc caactggagc cctgcttttt	840
gtttgctctt ttggtttatt tcattgttga gtatacccat ctccattgca aaagatcttt	900
cagacataga aggtgaccgc aagtttgga tcataacctt ctcaactaaa ttgggagcaa	960
aaccatagc atatatattgt catggactca tgcttctgaa ttacgtgagt gttatggctg	1020
cagctattat ttggccacag tttttcaaca gtagcgtaat attgctttct catgcattca	1080
tggcaatttg ggtattatat caggcttgga tattggagaa atcaaattac gccacggaaa	1140
cgtgccaaaa atactatata ttcccttgga taatttttcc tcttgaacat gccttctatt	1200
tgttcatgag ttcgagagac c	1221

<210> SEQ ID NO 55
 <211> LENGTH: 61
 <212> TYPE: DNA
 <213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 55

ttgacgaagt tcaacttttc cttggaaatg ctggaacagc aatgcgtcca ctacagctg	60
c	61

<210> SEQ ID NO 56
 <211> LENGTH: 57
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic- engineered gene
 <220> FEATURE:

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: N represents a scaffold sequence

<400> SEQUENCE: 56

caacttttcc ttggaaatgc ntgagtgaac gcattgctat tccagcattt ccaagga      57

<210> SEQ ID NO 57
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 57

cttgagcaaa cagttgagga aggacccgat tactgcgtga tcaactccacc agagaa      56

<210> SEQ ID NO 58
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic- engineered gene
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: N represents a scaffold sequence

<400> SEQUENCE: 58

cttgagcaaa cagttgagga ncacgcagta atcaagtcct tcctcaactg ttg      53

<210> SEQ ID NO 59
<211> LENGTH: 1638
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: In silico- consensus sequence

<400> SEQUENCE: 59

atgaattgct cagcattttc cttttggttt gtttgcaaaa taatattttt ctttctctca      60
ttccatatcc aaatttcaat agctaactct cgagaaaact tccttaaatg cttctcaaaa      120
catattccca acaatgtagc aaatccaaaa ctcgatataca ctcaacacga ccaattgtat      180
atgtctatcc tgaattcgac aatacaaaaat cttagattca tctctgatac aacccccaaa      240
ccactcgtaa ttgtcactcc ttcaaataac tcccatatcc aagcaactat tttatgctct      300
aagaaagttg gcttgcagat tcgaactcga agcgggtggc atgatgctga gggatgtgcc      360
tacatatctc aagtcgccatt tgtttagta gacttgagaa acatgcattc gatcaaaata      420
gatgttcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat      480
tggatcaatg agaagaatga gaatcttagt ttctctggtg ggtattgccc tactgttggc      540
gtaggtggac acttttagtg aggaggctat ggagcattga tgcgaaatta tggccttgcg      600
gctgataata ttattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa      660
tccatgggag aagatctggt ttgggtctata cgtggtggtg gaggagaaaa ctttggaaac      720
attgcagcat ggaaaatcaa actggttgct gtcccatcaa agtctactat attcagtgtt      780
aaaaagaaca tggagatata tgggcttgtc aagttattta acaaatggca aaatattgct      840
tacaagtatg acaagatgtt agtactcatg actcacttca taacaaagaa tattacagat      900
aatcatggga agaataagac tacagtacat ggttacttct cttcaatttt tcatgggtga      960

```

-continued

gtggatagtc tagtcgactt gatgaacaag agctttcctg agttgggtat taaaaaaact	1020
gattgcaaag aatttagctg gattgataca accatcttct acagtgggtg tgtaaatTTT	1080
aacactgcta attttaaaaa ggaaatTTTg cttgatagat cagctgggaa gaagacggct	1140
ttctcaatta agtttagacta tgtaagaaa ccaattccag aaactgcaat ggtcaaaatt	1200
ttggaaaaat tatatgaaga agatgttaga gctgggagt atgtgttgta cccttacggg	1260
ggtataatgg aggagatttc agaactcagca attccattcc ctcacgagc tggaataatg	1320
tatgaacttt ggtacactgc ttcctgggag aagcaagaag ataataaaaa gcatataaac	1380
tgggttcgaa gtgtttataa ttttacgact ccttatgtgt cccaaaatcc aagattggcg	1440
tatctcaatt atagggaact tgatttagga aaaactaatc atgcgagtc taataattac	1500
acacaagcac gtatttgggg tgaaaagtat tttggtaaaa attttaacag gttagttaag	1560
gtgaaaacta aagttgatcc caataatttt tttgaaaacg aacaaagtat cccacctctt	1620
ccaccgcac atcattaa	1638

<210> SEQ ID NO 60

<211> LENGTH: 1737

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 60

atgaaggaga tcgcgatgag gaattcaaag cgcaagcctg agccgacgcc gttcgccggg	60
aagaagctcc ggtcgacgag attacgccg aagagagcac agatctctcc cgttcttgtt	120
caatcacctc tctggagcaa acaaatcgga gtctctgctg cttctgtcga ttcctgctcc	180
gatttgctag ctgatgacaa cgtttcctgt ggttcgagca gagtcgagaa gagctcgaat	240
cgaagaaga ctctaattga agaggtagaa gtttctaacc ctggttataa tgtgaaggag	300
acgattgggt attcgaaatt tcgaaggatt acgaggtctt actctaagct acacaaggag	360
aaggaggag atgagatcga agtaagcgaa tcgtcttggt ttgattcgaa ttcgtggtgct	420
ggattaagga gattgaatgt gaagggaat aaaattaacg acaacgatga gatctcttcc	480
tcacgatccg atgtgacctt cgccgacat gtctccaaca gccggagtgt gaatttcgaa	540
tcggagaata aggagagcga cgtcgtttct gtcatatctg gagttgagta ctgttccaag	600
ttcggggagc ttaccggagg agctgataac gaagaaattg aaatctccaa gccgagcagc	660
ttcgtggaag ctgattcctc tcttgatcg gccaaaggaat tgaagccgga gcttgagata	720
gtcggatgag tctctgatct cgttgctctt gagaaattct cggaagaggt ttcggattct	780
ctcgatgatg agtcacatga gcaacgttca gagatatatt cacagtattc cgacttcgat	840
tactcggatt aactccgctc catcttcttc gactctggca gcgaattctc tgagaaatct	900
tcctctgatt ctctatttcc acattctcgc tctctgtacc tcaggttcaa ggaacagttc	960
tgtagatcca cgattcccaa cgattttgga tcttcttgag aggaagaaat tcaactcgaa	1020
ttgctaagggt ttgatgatga ggagggtgaa gagagctatc taaggctgag ggaaagagaa	1080
agaagtcatg catatatgag ggactgtgct aaggcatact gctccaggat ggacaatact	1140
ggtctcatcc ctcgtctacg ctccatcatg gttcaatgga ttgtaaagca atgttctgac	1200
atggggcttc agcaagagac attgtttcta ggagttggct tggtagatcg attcctgagc	1260
aaaggatcat tcaaaagcga aaggactcta atactagtcg ggattgcgag tcttactctg	1320

-continued

```

gccaccagaa ttgaagaaaa tcaaccttac aacagcatcc ggaaaaggaa ctccaccatt 1380
cagaacctaa gatatagccg gcatgaagtg gtggcaatgg agtggctggg tcaagaagtc 1440
ctcaacttca aatgcttcac acccacaatc ttcaacttct tgtggttcta cttaaaagct 1500
gctcgagcca atccagaagt tgaaggaaa gccaaatcct tggctgttac ctactatcc 1560
gaccaaactc aactctgttt ttggccctca actgtagcag ctgcactcgt gggtctcgcc 1620
tgcacgaac acaacaaaat ctctgcatac caacgagtc taaaggcca tgtagaaca 1680
acagataacg agttgcctga atgcgttaag agtctggact gggtgcttgg gcagtaa 1737

```

<210> SEQ ID NO 61

<211> LENGTH: 832

<212> TYPE: DNA

<213> ORGANISM: *Bacillus amyloliquefaciens*

<400> SEQUENCE: 61

```

ctggaaaacg tcacattgct tccgcatatc gggtcagcaa cggctaaaat ccgcttgaat 60
atgttcacac aagccgctca aaacatgatt gacgcgctat acggaagaac gccgaaaaac 120
cttactaagg aatttcaata agaagaaaaa tcccggttgg ttcagccggg gtttattttt 180
cgctagataa aaagtactat ttttaaatc tttctattcc tttcttctgt tgetgataca 240
atgaaaagga atcagcttca catgatgaaa atgggaggtg ttgctttgaa aaacgatta 300
tcgtggattt ccgtttgttt actggtgctt gtctccggc cggggatgct gttttcaaca 360
gtgccccaaa cggaaacatc ttctcacaag gcacacacag aagcacaggt tatcaacacg 420
tttgacgggg ttgcggtgta tcttcagaca taccataagc tacctgataa ttacattaca 480
aaatcagaag cacaagccct cggctgggtg gcatcaaaag ggaaccttgc agacgtcgct 540
ccggggaaaa gcatcgccgg agacatcttc tcaaacaggg aaggcaaaact cccgggcaaa 600
agcggacgaa catggcgtga agcggatatt aactatacat caggcttcag aaattcagac 660
cggattcttt actcaagcga ctggctgatt taaaaacaa cggaccatta tcagaccttt 720
acaaaaatca gataacgaaa aaaacggctt cctgcggag gccgtttttt tcagctttac 780
ataaagtgtg taataaatTT ttcttcaaac tctgatcggc caatttcact tt 832

```

We claim:

1. A method of preparing an explant, the method comprising the steps of,

rehydrating a dry *Cannabis* seed in a hydration medium, and

excising meristematic tissue from the rehydrated *Cannabis* seed to form an explant.

2. The method of claim 1, wherein the hydration medium comprises one or more priming agents.

3. The method of claim 1, wherein the seed is a *Cannabis sativa* seed.

4. The method of claim 1, additionally comprising the step of surface sterilizing the *Cannabis* seed prior to rehydration.

5. The method of claim 1, wherein the method additionally comprises the step of drying the explant.

6. The method of claim 5, wherein the dried explant is capable of being stored for at least 10 days.

7. The method of claim 5, wherein the explant is dried in the presence of one or more transformation supplements.

8. The method of claim 1, wherein the method additionally comprises the step of transforming the explant with a heterologous nucleic acid of interest.

9. The method of claim 8, wherein the explant is transformed using *Agrobacterium* mediated transformation or particle bombardment.

10. A *Cannabis* explant generated by the method of claim 1.

11. A dried *Cannabis* explant generated by the method of claim 5.

12. A method of transforming *Cannabis* with a heterologous nucleic acid, the method comprising the steps of, rehydrating a dry *Cannabis* seed in a hydration medium, excising meristematic tissue from the rehydrated *Cannabis* seed to form a *Cannabis* explant, incubating the *Cannabis* explant in a pretreatment medium, inoculating the *Cannabis* explant with *Agrobacterium* spp. comprising the heterologous nucleic acid, co-culturing the *Cannabis* explant in a co-culture medium for between about 1 day and about 6 days,

culturing the *Cannabis* explant on a selection medium to select for transformed *Cannabis* explants.

13. The method of claim 12, additionally comprising the step of force treating the *Cannabis* explant prior to or following inoculation.

14. The method of claim 13, wherein the force treatment is selected from the group consisting of sonication, vortexing, centrifugation, heat-shock, increase pressure, vacuum infiltration, and addition of chemicals.

15. The method of claim 12, additionally comprising the step of surface sterilizing the *Cannabis* seed prior to rehydration.

16. The method of claim 12, wherein the heterologous nucleic acid modulates the expression or activity of an endogenous *Cannabis* gene selected from the group consisting of tetrahydrocannabinolic acid synthase (THCA synthase), cannabidiolic acid synthase (CBDA synthase), O-methyltransferase (CsOMT21), lipid transfer protein 2 (LTP2), prenyltransferase 3 (CsPT3), and prenyltransferase 1 (CsPT1).

17. The method of claim 12, wherein the *Cannabis* seed is a *Cannabis sativa* seed.

18. A transformed *Cannabis* explant produced by the method of claim 12.

19. A *Cannabis* plant grown from the *Cannabis* explant of claim 18.

20. A method of producing a transformed *Cannabis* seed, the method comprising:

contacting a female *Cannabis* flower with an *Agrobacterium* spp. culture, wherein the *Agrobacterium* comprises a heterologous nucleic acid; and
pollinating the contacted female *Cannabis* flower with male pollen from a suitable donor plant, whereby a transformed *Cannabis* seed is produced.

21. The method of claim 20, wherein the *Agrobacterium* spp. culture comprises sucrose and a wetting agent.

22. The method of claim 20, wherein the *Agrobacterium* comprises a vector comprising the heterologous nucleic acid.

23. A method of transforming a *Cannabis* plant, the method comprising:

growing a sanitized and imbibed *Cannabis* seed on a non-selective culture medium suitable for supporting the growth and survival of the *Cannabis* seed until a *Cannabis* explant is formed;
inoculating the *Cannabis* explant with a heterologous nucleic acid;

co-culturing the *Cannabis* explant in a co-culture medium for between about 1 day and about 6 days,

culturing the *Cannabis* explant on a selection medium to select for transformed *Cannabis* explants.

24. The method of claim 23, wherein the explant is selected from the group consisting of a leaf explant, a node explant, an internode explant, a petiole explant, a hypocotyl explant, and a bud explant.

25. The method of claim 23, wherein the *Cannabis* explant is inoculated using particle bombardment, high velocity microprojection, microinjection, electroporation, direct DNA uptake, cell-penetrating peptides, silica carbide fibers, nanoparticles, and bacterially-mediated transformation.

26. The method of claim 23, wherein *Agrobacterium* spp. is used to inoculate the *Cannabis* explant.

27. The method of claim 23, additionally comprising the step of force treating the *Cannabis* explant prior to or following inoculation.

28. The method of claim 27, wherein the force treatment is selected from the group consisting of sonication, vortexing, centrifugation, heat-shock, increased pressure, vacuum infiltration, and addition of chemicals.

29. The method of claim 23, wherein the heterologous nucleic acid modulates the expression or activity of an endogenous *Cannabis* gene selected from the group consisting of tetrahydrocannabinolic acid synthase (THCA synthase), cannabidiolic acid synthase (CBDA synthase), O-methyltransferase (CsOMT21), lipid transfer protein 2 (LTP2), prenyltransferase 3 (CsPT3), and prenyltransferase 1 (CsPT1).

30. The method of claim 23, wherein the *Cannabis* seed is a *Cannabis sativa* seed.

31. The method of claim 12, wherein the heterologous nucleic acid encodes a polypeptide at least 90% identical to SEQ ID NO:28.

32. The method of claim 23, wherein the heterologous nucleic acid encodes a polypeptide at least 90% identical to SEQ ID NO:28.

33. The method of claim 12, wherein the heterologous nucleic acid encodes a guide RNA that targets *Cannabis sativa* THCA synthase gene, *Cannabis sativa* CBDA synthase gene, or *Cannabis sativa* EPSP synthase gene.

34. The method of claim 23, wherein the heterologous nucleic acid encodes a guide RNA that targets *Cannabis sativa* THCA synthase gene, *Cannabis sativa* CBDA synthase gene, or *Cannabis sativa* EPSP synthase gene.

* * * * *