



US 20240026466A1

(19) **United States**

(12) **Patent Application Publication**
Venturelli et al.

(10) **Pub. No.: US 2024/0026466 A1**

(43) **Pub. Date: Jan. 25, 2024**

(54) **CELL-BASED DNA SENSORS AND METHODS OF USING SAME**

(71) Applicant: **Wisconsin Alumni Research Foundation, Madison, WI (US)**

(72) Inventors: **Ophelia Venturelli, Madison, WI (US); Yu-Yu Cheng, Madison, WI (US); Zhengyi Chen, Madison, WI (US)**

(73) Assignee: **Wisconsin Alumni Research Foundation, Madison, WI (US)**

(21) Appl. No.: **18/067,194**

(22) Filed: **Dec. 16, 2022**

Related U.S. Application Data

(60) Provisional application No. 63/290,442, filed on Dec. 16, 2021.

Publication Classification

(51) **Int. Cl.**

C12Q 1/689 (2006.01)

C12Q 1/6897 (2006.01)

C12N 15/90 (2006.01)

(52) **U.S. Cl.**


CPC **C12Q 1/689** (2013.01); **C12Q 1/6897** (2013.01); **C12N 15/902** (2013.01)

(57)

ABSTRACT

Cell-based DNA sensors, compositions comprising the cell-based DNA sensors, and methods of detecting DNA and cells. The cell-based DNA sensors include competent cells that include genetic circuits. Each genetic circuit includes homology arms separated by an interstitial region that comprises at least one element of a reporter switch and/or a kill switch. The compositions include one or more cell-based DNA sensors. The cell-based DNA sensors can be used to detect DNA and cells by the genetic circuits undergoing homologous recombination with target regions of target DNA to activate or deactivate the reporter switch and/or kill switch.

Specification includes a Sequence Listing.






FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 2G




FIG. 2H




FIG. 2I




FIG. 2J




FIG. 2K




FIG. 2L




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 5A




FIG. 5B






FIG. 5C



+gDNA

Fluorescent colonies
Recombination: 12

FIG. 6A



-gDNA

Fluorescent colonies
txpA deletion: 6
Non-fluorescent colonies
lacI mutation: 6

FIG. 6B

Transformed *E. coli* sensor: left junction




FIG. 6C

Transformed *E. coli* sensor: right junction




FIG. 6D

Fluorescent *E. coli* sensor mutant: *txpA* deletion

SEQ ID NO:3




FIG. 6E

Non-fluorescent of *E. coli* sensor mutant: *lacI* R195 and L196 deletion

SEQ ID NO:4




FIG. 6F




FIG. 7A







FIG. 7B

**FIG. 8A****FIG. 8B****FIG. 8C**




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 10E




FIG. 10F




FIG. 10G




FIG. 10H




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 14A




FIG. 14B

Coupling GFP expression to DNA detection Repressor-based toxin expression




FIG. 15A

Toxin x	2nd repressor y	Promoter P_y
<i>txpA</i>	<i>xylR</i>	P_{xylA}
<i>mazF</i>	<i>cl</i>	P_R
<i>hwl</i>	CRISPRi	P_{const}

Activator-based toxin expression




FIG. 15B

Toxin <i>x</i>	Activator <i>y</i>	Promoter P_y	Inducer
<i>txpA</i>	<i>spaR</i>	P_{spaS}	subtilin
<i>mazF</i>	<i>liaR</i>	P_{liaI}	bacitracin
<i>hewI</i>	<i>ccaR</i>	P_{cpcG2}	green light

Counter-selectable marker-based DNA detection




FIG. 15C

Counter-selectable marker <i>x</i>	Sensitive compound <i>X</i>
<i>upp</i>	5-fluorouracil
<i>pheS</i> [*]	p-chloro-phenylalanine
<i>ysbC</i>	fluoro-orotate

Selectable marker-based DNA detection




FIG. 15D

Selectable marker x	Essential for
<i>neo</i>	neomycin resistance
<i>lysA</i>	lysine production




FIG. 15E

Toxin x	Quorum-sensing gene y	Promoter P_y
<i>txpA</i>	<i>agrBDCA</i>	P_3
<i>mazF</i>	<i>comA</i>	P_{ertA}
<i>hewI</i>		




FIG. 15F




FIG. 15G




FIG. 15H




FIG. 15I

CELL-BASED DNA SENSORS AND METHODS OF USING SAME

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

[0001] This invention was made with government support under HR-0011-18-2-0002 awarded by the DOD/DARPA. The government has certain rights in the invention.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been electronically submitted in XML format and is hereby incorporated by reference in its entirety. The ASCII copy was created on Dec. 2, 2022, is named Seq_List-P210411US02.xml and is 54,692 bytes in size.

FIELD OF THE INVENTION

[0003] The invention is directed to cell-based DNA sensors and methods of using same for detecting DNA and cells.

BACKGROUND

[0004] Chemical and electrical signaling in microbial communities play key roles in biofilm development, activation of virulence pathways, and symbioses with multicellular organisms⁵³. These signals can be exploited to control the collective growth or gene expression of the population or mediate interactions between constituent community members^{5,55}. For example, circuits have been designed in engineered organisms to sense specific signals produced by pathogens for selective inhibition of growth^{7,8}. However, there are limited well-characterized and orthogonal chemical signals systems for building communication networks between strains due to signal crosstalk^{9,10}. In addition, there are challenges to engineering these chemical signals for inter-species communication^{55,56}. Therefore, new versatile mechanisms are needed for sensing diverse species in microbial communities.

[0005] Towards this goal, sensing of bacterial pathogens is a critical and unsolved challenge, as new pathogens can emerge⁵⁷. Current methods for pathogen detection include quantitative PCR (qPCR), immunology-based testing, selective culturing, and Next-Generation Sequencing (NGS)^{58,59}. Further, diagnostic tools based on CRISPR-Cas nucleases have also been developed^{60,61}. While qPCR is sensitive to the concentration of the target sequence, this method requires specialized equipment and trained personnel which may limit its broad deployment. Immunological detection methods have lower sensitivity and specificity than PCR-based techniques, but have a faster turnaround time. Due to these limitations, new cost-effective, sensitive, generalizable and easy to implement pathogen detection methods are needed.

SUMMARY OF THE INVENTION

[0006] The invention provides cell-based DNA sensors. One version of a cell-based DNA sensor comprises a competent cell that comprises a genetic circuit. The genetic circuit preferably comprises a pair of homology arms and at least one of a reporter switch and a kill switch. The pair of homology arms are preferably comprised in a DNA strand. The homology arms preferably comprise a first homology arm and a second homology arm. The first homology arm is

preferably homologous to a first portion of a target DNA. The second homology arm is preferably homologous to a second portion of the target DNA. The first portion and the second portion in some versions are contiguous in the target DNA. The first homology arm and the second homology arm are preferably separated within the DNA strand by an interstitial region of the DNA strand. The reporter switch preferably comprises a reporter gene and a negative regulator of the reporter gene. The reporter gene preferably comprises a promoter and a coding sequence that are not comprised within the interstitial region of the DNA strand. The negative regulator of the reporter gene is preferably comprised within the interstitial region of the DNA strand. The kill switch preferably comprises one or more genetic elements effective to inhibit growth of the competent cell. At least one of the one or more genetic elements is preferably comprised within the interstitial region of the DNA strand.

[0007] The invention also provides compositions comprising one or more cell-based DNA sensors of the invention. One version of a composition of the invention comprises two or more cell-based DNA sensors. The pairs of homology arms in the two or more cell-based sensors are preferably each homologous to different target DNA sequences. Each of the two or more cell-based DNA sensors preferably comprises the reporter switch. The reporter genes in the two or more cell-based DNA sensors preferably express reporters that are each detectably different from each other.

[0008] The invention also provides methods of detecting target DNA with a cell-based DNA sensor of the invention. One version of such methods comprises culturing the DNA sensor in a culture medium comprising the target DNA for a time effective to transform the DNA sensor with the target DNA and detecting the transformed DNA sensor.

[0009] The invention also provides methods of detecting a target cell comprising target DNA with a cell-based DNA sensor of the invention. One version of such methods comprises culturing the DNA sensor in a culture medium with the target cell for a time effective to transform the DNA sensor with the target DNA and detecting the transformed DNA sensor.

[0010] The objects and advantages of the invention will appear more fully from the following detailed description of the preferred embodiment of the invention made in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0011] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[0012] FIGS. 1A-1C. Construction and characterization of a living cell-based DNA sensor. (FIG. 1A) Schematic of a synthetic genetic circuit constructed in *B. subtilis* that allows the recognition of specific extracellular DNA. The xylose-inducible master regulator of competence comK can induce *B. subtilis* to take up eDNA and undergo homologous recombination. The toxin-antitoxin system txpA-ratA and a distal fluorescent reporter gfp are regulated by the repressor lacI. The target sequences are split into two and introduced to the flanking region of toxin and repressor as landing pad. In the presence of target DNA, homologous recombination can remove the toxin and the repressor, so a sub-population of transformed cells can express GFP. To select the trans-

formed cells, IPTG can induce the toxin to kill the non-transformed cells, so the fluorescence can be enhanced by the growth of transformed cells. (FIG. 1B) Line plot of transformation efficiency versus transformation time for the *E. coli* DNA sensor in the presence (solid lines) or absence (dashed lines) of 100 ng/mL purified *E. coli* gDNA as input. The colonies on agar plate can be fluorescent (green) or not (blue), coming from background escape mutation or transformation. Transformation efficiency as output is optimal at 10 hr. Data points represent three biological replicates and lines are the average of the replicates. (FIG. 1C) Line plot of transformation efficiency versus homology length of *E. coli* xdhABC operon on each side of flanking region in the circuit in the presence (solid lines) or absence (dashed lines) of 100 ng/mL purified *E. coli* gDNA as input. The colonies on agar plate can be fluorescent (green) or not (blue). The transformation efficiency as output is above the background escape mutation when the homology length is equal or greater than 1 kbp and increases as the homology increases. Data points represent three biological replicates and lines are the average of replicates.

[0013] FIGS. 2A-2L. Cell-based sensors can detect DNA from diverse bacteria with high sensitivity and specificity. (FIG. 2A) Transformation efficiency of DNA sensors that can detect *E. coli* (EC sensor), *S. typhimurium* (ST sensor), *S. aureus* (SA sensor), or *C. difficile* (CD sensor) in the presence of 100 ng/mL extracted gDNA (triangles) or no gDNA (circles) at 10 hr. Bar represents the average of three biological replicates. (FIG. 2B) Schematic of experimental procedure for quantifying GFP expression after transformation. (FIG. 2C) Time-series measurements of GFP expression of EC sensor in liquid medium after the transformation of varying *E. coli* gDNA concentrations (ng/mL). A threshold of GFP 400 was used to determine the detection time for each gDNA concentration. Line is the average of four technical replicates and the shaded region represents one standard deviation from the average. Detection time versus gDNA concentration for (FIG. 2D) EC sensor, (FIG. 2E) ST sensor, (FIG. 2F) SA sensor, and (FIG. 2G) CD sensor. Horizontal line (pale color) is the background GFP fluorescence in the absence of gDNA. Unpaired t-test was performed to determine if the detection time with specific DNA concentration is different from the background fluorescence, and *, **, and *** denote p-values <0.05, 0.01 and 0.001, respectively. A straight line was fitted to the transformation efficiencies versus logarithmic gDNA concentrations with statistical differences. The slope of the fitted line was determined by the cell growth of *B. subtilis*, while the intercept was determined by the background escape mutation. The coefficient of determination R^2 shows the goodness of the fit. (FIG. 2H) Detection time of each DNA sensor after the transformation of 100 ng/mL gDNA of a given donor species or no gDNA representing the negative control (NC). Each sensor expressed GFP 3–4 hours earlier in the presence of gDNA from its target strain than gDNA from other strains. Data are the average of four technical replicates. (FIG. 2I) Nucleotide BLAST search of 5000 bp *S. aureus* hemEH in the NCBI database. Each circle represents a homolog found in species other than *S. aureus* and its coverage and identity similarity. A closely related human commensal strain *S. epidermidis* with high coverage and similarity was selected for specificity test using SA sensor. The region within dashed lines indicates *Staphylococcus* species that could be recognized by the SA sensor based on their high similarity

of homologous sequences. (FIG. 2J) Comparison of colony numbers of SA sensor (with GFP expression) on selective agar plate after the transformation of 100 ng/mL *S. aureus* gDNA, 100 ng/mL *S. epidermidis* gDNA or no gDNA. Bar represents the average of three technical replicates. SA sensor can distinguish between *S. aureus* and closely related *S. epidermidis*. (FIG. 2K) Nucleotide BLAST search of 5000 bp *C. difficile* pheST in the NCBI database. Each circle represents a homolog found in species other than *C. difficile* and its coverage and identity similarity. A closely related human commensal strain *C. hiranonis* with high coverage and similarity was selected for specificity test using CD sensor. The region within dashed lines indicates *Clostridium* species that could be recognized by the CD sensor based on their high similarity of homologous sequences. (FIG. 2L) Comparison of colony numbers of CD sensor (with GFP expression) on selective agar plate after the transformation of 100 ng/mL *C. difficile* gDNA, 100 ng/mL *C. hiranonis* gDNA or no gDNA. Bar represents the average of three technical replicates. CD sensor can distinguish between *C. difficile* and closely related *C. hiranonis*.

[0014] FIGS. 3A-3F. DNA sensors can perform multiplexed detection in complex DNA samples. (FIG. 3A) Schematic of experimental procedure using the living cell-based DNA sensors for multiplexed detection. *E. coli*, *S. typhimurium*, and *S. aureus* DNA sensors were labeled GFP (EC-G), RFP (ST-R) and BFP (SA-B) for detection, respectively. Three sensors and gDNA extracted from different strains were mixed into liquid medium for transformation. Transformed cells were selected on agar and colonies expressing GFP, RFP and BFP can indicate the presence of target DNA. (FIG. 3B) Numbers of GFP, RFP, or BFP-expressing colonies on agar plate after the transformation of different combinations of gDNA extracted from *E. coli*, *S. typhimurium*, or *S. aureus*. Sensors can report the presence of target DNA for all 8 different combinations. Bar represents the average of three technical replicates. (FIG. 3C) Relative abundance of six species in a synthetic gut microbial community composed of *S. aureus* (SA), *S. typhimurium* (ST), *Bifidobacterium longum* (BL), *Bacteroides thetaiotaomicron* (BT), *Anaerostipes caccae* (AC), and *Clostridium asparagiforme* (CG) over two days of growth. The six bacteria were co-cultured in liquid medium anaerobically for 24 hours and cell culture was diluted in fresh medium once for species to continue their competition. 16S rRNA gene of each strain was PCR amplified and sequenced using NGS to determine the relative abundance over time. Bar represents the average of three technical replicates of 16S rRNA sequencing. *S. typhimurium* abundance remained similar while *S. aureus* abundance decreased over time. (FIG. 3D) Representative fluorescence images of transformed SA-G and ST-R sensors on selective agar plate after the transformation of gDNA extracted from the bacterial community in a mixture of SA-G and ST-R sensors. The numbers of green and red colonies indicate the abundance of *S. aureus* and *S. typhimurium* in gut microbial community, respectively. (FIG. 3E) Numbers of GFP or RFP-expressing colonies on agar plate after the transformation of gDNA extracted from the community at different time. Green colonies (SA-G) decreased over time while red colonies (ST-R) remained similar numbers. Bar represents the average of three technical replicates. (FIG. 3F) Fold change of *S. aureus* and *S. typhimurium* abundance compared to day 1 characterized by NGS (dashed line) or cell-based detection

method (solid line). Measurements of *S. aureus* (orange) by the two methods were similar, while characterization of *S. typhimurium* (green) by the cell-based detection method had larger variability than NGS and the data at Day 1 were statistically different determined by Unpaired t-test (p-values=0.0015).

[0015] FIGS. 4A-4D. DNA sensors can directly detect target species without DNA extraction. (FIG. 4A) Schematic of experimental design for the detection in the co-culture. Sensor and target strain were co-cultured in liquid medium with or without selective antibiotics (ABX) to determine the effect of antibiotic for detection. Heat-killed target strain was also tested without the use of antibiotics in the co-culture. The cell culture was plated on IPTG and ABX (1 μ g/mL erythromycin and 25 μ g/mL lincomycin) agar plates to select for transformed *B. subtilis*. (FIG. 4B) Colony numbers of transformed EC, ST, SA, and CD sensors co-cultured with (1) target cell, (2) target cell and 100 μ g/mL spectinomycin (spec), (3) target cell, spec, and 1 unit/mL DNase I, and (4) spec only. Spectinomycin can enhance the detection of *E. coli*, *S. typhimurium*, and *S. aureus*, but it is not required for *C. difficile* detection. Addition of DNase I in the co-culture reduced the numbers of transformed sensors significantly. (FIG. 4C) Colony number of transformed EC sensor co-cultured with *E. coli* using spectinomycin or co-cultured with heat-treated *E. coli*. Heat treatment of *E. coli* can significantly increase the number of transformed EC sensor. (FIG. 4D) Colony number of transformed EC-G and ST-R sensors co-cultured with heat-treated mice cecal samples spiked in with different amounts of *E. coli* and *S. typhimurium*. Both sensors can detect the presence of target strains in cecal samples. Unpaired t-test was performed to determine if the colony number is different from no cell condition, and *, **, and *** denote p-values <0.05, 0.01, and 0.001, respectively. Bar represents the average of three technical replicates.

[0016] FIGS. 5A-5C. Plasmid maps for the construction of DNA-sensing *B. subtilis*. (FIG. 5A) Plasmid constructed for the DNA detection via homologous recombination. The toxin-antitoxin system *txpA-ratA* is regulated by the repressor *lad*, both of which are flanked by 2.5 kbp target DNA sequence on each side. The plasmid was integrated into the *amyE* locus on the *B. subtilis* PY79 genome by spectinomycin selection. (FIG. 5B) Plasmid constructed for the fluorescent reporter after DNA detection. The green fluorescent protein *gfp(Sp)* is regulated by the distal repressor *lad*. The plasmid was integrated into the *ycgO* locus on *B. subtilis* PY79 genome by chloramphenicol selection. The green fluorescent protein *gfp(Sp)* was codon-optimized for *Streptococcus pneumoniae* and displayed high fluorescence signal in *B. subtilis*⁵¹. (FIG. 5C) Modularity of the synthetic genetic circuit allows customized target DNA sequence as input and gene expression as output.

[0017] FIGS. 6A-6F. Sequencing of transformed *E. coli* sensor and escape mutants. Representative fluorescence images of colonies of *E. coli* DNA sensor on selective agar plate after the transformation of (FIG. 6A) 100 ng/mL purified *E. coli* gDNA or (FIG. 6B) no DNA. DNA of the circuit in *B. subtilis* genome was PCR amplified from different colonies and sequenced to confirm the homologous recombination for the transformed sensor (N=12) or mutations for the escape mutants (N=12). (FIGS. 6C and 6D) Sanger sequencing of transformed *E. coli* sensor confirmed the homologous recombination at the predicted region that

removed the whole cassette of *txpA-ratA* and *lad*. (FIG. 6E) Sanger sequencing of a GFP-expressing escape mutant shows deletion in the toxin *txpA*. The mutant can grow and express GFP in the presence of IPTG due to the non-functional toxin. (FIG. 6F) Sanger sequencing of a non-fluorescent escape mutant shows deletion of R195 and L196 in the repressor *lad*, which has been shown to affect IPTG binding⁵². The mutant can grow but cannot express GFP in the presence of IPTG due to the non-functional *LacI*.

[0018] FIGS. 7A and 7B. Nucleotide BLAST search of homology sequences in EC sensor and ST sensor. (FIG. 7A) Nucleotide BLAST search of 5000 bp *S. typhimurium* sipBCDA in the NCBI database. Each circle represents a homolog found in species other than *S. typhimurium* and its coverage and identity similarity. Homologs were found mostly in the *Salmonella enterica* (*S. enterica*) species but were rarely found in other species. (FIG. 7B) Nucleotide BLAST search of 5000 bp *E. coli* MG1655 xdhABC in the NCBI database. Each circle represents a homolog found in species other than *E. coli* and its coverage and identity similarity. Homologs were found in closely related *Shigella* and *Escherichia* species. Highly similar homologs were found in the closely related *Shigella* species.

[0019] FIGS. 8A-8C. Time-series measurements of GFP expression of ST, SA, CD sensors in liquid medium after transformation. Time-series measurements of GFP expression of (FIG. 8A) ST sensor, (FIG. 8B) SA sensor, and (FIG. 8C) CD sensor in liquid medium after the transformation of varying target gDNA concentrations (ng/mL). A threshold of GFP 400 was used to determine the detection time for each gDNA concentration. Line is the average of four technical replicates and the shaded region represents one standard deviation from the average.

[0020] FIGS. 9A-9D. Time-series OD measurements of DNA sensors in liquid medium after transformation. Time-series measurements of OD600 absorbance of (FIG. 9A) EC sensor, (FIG. 9B) ST sensor, (FIG. 9C) SA sensor, and (FIG. 9D) CD sensor in liquid medium after the transformation of varying target gDNA concentrations (ng/mL). Line is the average of four technical replicates and the shaded region represents one standard deviation from the average. Cell growth correlated with DNA concentration during transformation.

[0021] FIGS. 10A-10H. Orthogonality test of the four constructed DNA sensors. Time-series measurements of GFP expression of (FIG. 10A) EC sensor, (FIG. 10C) ST sensor, (FIG. 10E) SA sensor, and (FIG. 10G) CD sensor in liquid medium after the transformation of gDNA extracted from different strains or no gDNA. Line is the average of four technical replicates and the shaded region represents one standard deviation from the average. A threshold of GFP 400 was used to determine the detection time for the target gDNA and non-target gDNA or no gDNA for (FIG. 10B) EC sensor, (FIG. 10D) ST sensor, (FIG. 10F) SA sensor, and (FIG. 10H) CD sensor. Unpaired t-test was performed to determine if the detection time for the target gDNA is different from non-target gDNA or no gDNA. Based on the calculated p-values, sensors expressed GFP hours earlier only in the presence of target gDNA.

[0022] FIGS. 11A and 11B. Detection efficiency based on number of transformed cells. (FIG. 11A) Colony number per 5 μ L for transformed *B. subtilis* with or without 100 ng/mL target gDNA. (FIG. 11B) Colony number per 10⁻⁴ μ L for total *B. subtilis* with or without 100 ng/mL target gDNA. Bar

represents the average of three biological replicates. The transformation efficiency in FIG. 2A was calculated by the ratio of the density of transformed *B. subtilis* to the density of total *B. subtilis* shown here.

[0023] FIG. 12. Representative fluorescence images of transformed EC-G, ST-R, and SA-B sensors for multiplexed detection. Colonies with different fluorescence on agar plate after the transformation of combinations of extracted gDNA in a mixture of EC-G, ST-R, and SA-G sensors. Sensors were transformed with (A) *E. coli*, *S. aureus* and *S. typhimurium* gDNA, (B) *E. coli* and *S. typhimurium* gDNA, (C) *E. coli* and *S. aureus* gDNA, (D) *S. typhimurium* and *S. aureus* gDNA, (E) *E. coli* gDNA, (F) *S. typhimurium* gDNA, (G) *S. aureus* gDNA, and (H) no gDNA.

[0024] FIGS. 13A and 13B. Negative control for the multiplexed detection in complex DNA samples. (FIG. 13A) Relative abundance of six species in a synthetic gut microbial community composed of *Bifidobacterium longum* (BL), *Bacteroides thetaiotaomicron* (BT), *Anaerostipes caccae* (AC), and *Clostridium asparagiforme* (CG). The four bacteria were co-cultured in liquid medium anaerobically for 24 hours and cell culture was diluted in fresh medium once for species to continue their competition. 16S rRNA gene of each strain was PCR amplified and sequenced using NGS to determine the relative abundance over time. (FIG. 13B) Numbers of GFP or RFP-expressing colonies on agar plate after the transformation of gDNA extracted from the community at different time in a mixture of SA-G and ST-R sensors. Bar represents the average of three technical replicates of 16S rRNA sequencing. Only a few colonies appeared on agar plates, indicating that sensors did not detect the gDNA of *S. aureus* or *S. typhimurium*. Bar represents the average of three technical replicates.

[0025] FIGS. 14A and 14B. Detection of *E. coli* or *S. typhimurium* in cecal samples using mixed EC-G and ST-R sensors. Colony number of transformed EC-G and ST-R sensors co-cultured with heat-treated cecal samples spiked in with different amounts of (FIG. 14A) *E. coli* or (FIG. 14B) *S. typhimurium*. High density (10^8 CFU/mL) of *E. coli* and *S. typhimurium* can lead to false positive results for ST-R and EC-G, respectively. Unpaired t-test was performed to determine if the colony number is different from no cell condition, and *, **, and *** denote p-values <0.05, 0.01, and 0.001, respectively. Bar represents the average of three technical replicates.

[0026] FIGS. 15A-15I. Additional exemplary synthetic genetic circuits coupling GFP expression and growth regulation to DNA detection. (FIG. 15A) Synthetic genetic circuit for repressor-based toxin expression. (CRISPRi is listed as a repressor in FIG. 15A for convenience but is properly categorized as a “negative regulator” using the terminology employed herein.) (FIG. 15B) Synthetic genetic circuit for activator-based toxin expression. (FIG. 15C) Synthetic genetic circuit for counter-selectable marker-based DNA detection. (FIG. 15D) Synthetic genetic circuit for selectable marker-based DNA detection. (FIG. 15E) Synthetic genetic circuit for autonomous DNA detection (toxin expresses as cell density increases). (FIGS. 15F and 15G) Synthetic genetic circuits with a negative regulator (inducible regulator) comprised within the interstitial region (FIG. 15F) and outside the interstitial region (FIG. 15G) (Y is a growth inhibitor gene, and X is a positive regulator of the growth inhibitor gene). (FIGS. 15H and 15I) Synthetic genetic circuits with a negative regulator (inducible regula-

tor) comprised within the interstitial region (FIG. 15H) and outside the interstitial region (FIG. 15I).

DETAILED DESCRIPTION OF THE INVENTION

[0027] An aspect of the invention is directed to cell-based DNA sensors (also referred to herein as “DNA sensors”). The DNA sensors comprise competent cells. “Competent cell” refers to a cell capable of taking up extracellular DNA from its surrounding environment. The competent cells can be naturally competent cells or artificially competent cells.

[0028] Naturally competent cells are cells that, in their unmodified state, are capable of taking up external DNA. Naturally competent cells are not necessarily in a constant state of competence. Natural regulation of competence is common among naturally competent cells. *Streptococcus pneumoniae*, for example, is a naturally competent bacterium, but its competence at a given point in time is prompted by quorum sensing (detecting and responding to cell population density through gene expression). More than more than 80 species of bacteria are known to be naturally competent, including both Gram-positive and Gram-negative bacteria (Johnston, C., Martin, B., Fichant, G., Polard, P., and Claverys, J. P. (2014). Bacterial transformation: distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.* 12, 181-196). Exemplary naturally competent cells include members of the *Bacillus* genus, such as *Bacillus subtilis*; members of the *Streptococcus* genus, such as *Streptococcus pneumoniae*; members of the *Neisseria* genus, including *Neisseria gonorrhoeae* and *Neisseria meningitidis*; members of the *Haemophilus* genus, such as *Haemophilus influenzae*; members of the *Helicobacter* genus, such as *Helicobacter pylori*; members of the *Acinetobacter* genus, such as *Acinetobacter baylyi*; members of the *Vibrio* genus, such as *Vibrio cholerae*; members of the *Thermus* genus, such as *Thermus thermophilus*; and *Synechococcus* sp., among others.

[0029] Naturally competent cells can be modified to increase competency. Such cells are still considered to be naturally competent in light of their competency in their natural, unmodified state. Naturally competent cells, for example can be modified to express genes that increase competence. An example of such a gene is *comK*²⁵. Other genes or modifications that increase competency are known in the art.

[0030] Artificially competent cells are cells that are not capable of taking up external DNA in their unmodified state but are modified to do so. Artificially competent cells include chemically competent cells and electrocompetent cells. Chemically competent cells are cells that are made competent with a chemical treatment. Common treatments include a salt treatment followed by a heat-shock step. This process permeabilizes the cell membrane, allowing entry of DNA. Protocols using CaCl_2 or MgCl_2 are the most common method for making chemically competent cells, but other salts and chemicals can be used. These include dimethyl sulfoxide (DMSO), polyethylene glycol (PEG), and rubidium chloride (RbCl). Electrocompetent cells are made competent using an electrical pulse from an electroporator to create temporary pores (poration) in the cell membrane. An exemplary artificially competent cell is *Saccharomyces cerevisiae*, which can be artificially induced into competence and have stronger homologous recombination and may detect short DNA sequences.

[0031] The competent cells of the invention can be prokaryotic cells or eukaryotic cells. In exemplary versions of the invention, the competent cells are prokaryotic cells, such as naturally competent prokaryotic cells, such as *Bacillus subtilis*.

[0032] The competent cells of the invention can comprise a genetic circuit. A genetic circuit is a combination of genetic elements that enable a cell to perform a logical function. As used herein, “genetic element” refers to any sequence of DNA that confers a genetic function. Exemplary genetic elements, include genes, promoters, exons, introns, enhancers, silencers, 5' untranslated regions, 3' untranslated regions, open reading frames, coding regions, codons, terminators, etc. In the context of the present invention, an exemplary logical function is the detection of DNA.

[0033] The genetic circuits of the invention can comprise a pair of homology arms on a DNA strand. Each homology arm is a portion of DNA in the genetic circuit that has a sequence homologous to target DNA. The homology arms in each pair are configured to excise a portion of DNA from the DNA strand through homologous recombination with a target DNA. To perform this function, the homology arms are spaced apart on the DNA strand to form an interstitial region therebetween and together have homology to one or more homologous sequences in the target DNA. Homologous recombination of the homology arms with the target DNA thereby excises the interstitial region from the DNA strand. “Strand” in this context refers to a contiguous, connected string of nucleic acid bases. The DNA strand can be a portion of the competent cell's chromosome or a portion of an extra-chromosomal DNA, such as a plasmid. In some versions, the homology arms are homologous to a single, continuous sequence in the target DNA (contiguous portions of the target DNA). In some versions, the homology arms are homologous to two separated sequences in the target DNA. The number of bases separating the homologous sequences in the target DNA can vary depending on the length of the homology arms, with longer homology arms permitting further separation (separation being defined as the number of bases between the homologous sequences). In various versions, the homologous sequences are contiguous or separated by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 50, 100, 250, 500, 1000, 2,500, 5,000, 7,500, 10,000 or more bases, or any range therebetween any two of the foregoing values.

[0034] The terms “homologous” and “homology,” used herein with reference to sequences or portions of DNA, refer to having a sufficient length and sequence identity to be exchanged by homologous recombination in a competent cell of the invention.

[0035] Each homology arm in the genetic circuits of the invention preferably comprises a length of at least 0.025 kbp (kilobase pairs), such as at least 0.050 kbp, at least 0.070 kbp, at least 0.075 kbp, at least 0.1 kbp, at least 0.5 kbp, at least 1 kbp, at least 1.5 kbp, at least 2 kbp, or at least 2.5 kbp. Each homology arm in the genetic circuits of the invention can comprises a length up to 3 kbp, up to 5 kbp, up to 10 kbp, up to 25 kbp, up to 50 kbp, up to 75 kbp, up to 100 kbp, up to 250 kbp, up to 500 kbp, up to 750 kbp, up to 1,000 kbp, up to 1,250 kbp, up to 1,500 kbp, up to 1,750 kbp, up to 2,000 kbp, up to 2,250 kbp, up to 2,500 kbp, up to 2,750 kbp, up to 3,000 kbp, or more.

[0036] Each homology arm in the genetic circuits of the invention preferably comprises at least 75% sequence identity, at least 77% sequence identity, at least 80% sequence

identity, at least 85% sequence identity, at least 90% sequence identity, at least 91% identity, at least 92% identity, at least 93% identity, at least 94% identity, at least 95% identity, at least 96% identity, at least 97% identity, at least 98% identity, at least 99% identity, or 100% identity, to a portion of a target DNA. Such sequence identity is preferably assessed over the entire length of the homology arm, such as a length as defined in the preceding paragraph.

[0037] The term “sequence identity” (or “identity”), in the context of percent sequence identity, refers to the percentage of bases (or residues) in two sequences that are the same when aligned for maximum correspondence, as measured using a sequence comparison or analysis algorithm such as those described herein. For example, if when properly aligned, the corresponding segments of two sequences have identical bases (or residues) at 5 positions out of 10, the two sequences have a 50% identity. Most bioinformatic programs report percent identity over aligned sequence regions, which are typically not the entire molecules. If an alignment is long enough and contains enough identical residues, an expectation value can be calculated, which indicates that the level of identity in the alignment is unlikely to occur by random chance. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2008)). One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity for purposes of defining homologs is the BLAST algorithm, which is described in Altschul et al., *J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always>0) and N (penalty score for mismatching residues; always<0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X

determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

[0038] The homology arm pairs in the genetic circuits are preferably spaced on the DNA strand such that the interstitial region has a length of at least 0.001 kbp, at least 0.01 kbp, at least 0.05 kbp, at least 0.1 kbp, at least 0.5 kbp, at least 1 kbp, at least 1.25 kbp, at least 1.5 kbp, at least 1.75 kbp, at least 2 kbp, at least 2.5 kbp, at least 3 kbp, at least 3.5 kbp, at least 4 kbp, at least 4.5 kbp, or at least 5 kbp. In various versions of the invention, the interstitial region has a length up to 5 kbp, up to 10 kbp, up to 15 kbp, up to 20 kbp, up to 25 kbp, up to 30 kbp, up to 35 kbp, up to 40 kbp, up to 45 kbp, up to 50 kbp, up to 55 kbp, up to 60 kbp, up to 65 kbp, up to 70 kbp, up to 75 kbp, up to 80 kbp, up to 85 kbp, up to 90 kbp, up to 95 kbp, up to 100 kbp, or more.

[0039] The genetic circuits of the invention can comprise genes. “Gene” as used herein refers to the combination of genetic elements effective for the expression of a gene product such as RNA (e.g., mRNA, microRNA, etc.) and/or a protein. Genes typically minimally include a promoter and a coding sequence. Genes can also include other genetic elements, including enhancers, silencers, etc.

[0040] The genetic circuits of the invention can comprise a reporter gene. A reporter gene is a gene that expresses a reporter. The reporter can be any detectable gene product, such as any protein or RNA (e.g., mRNA, microRNA, etc.). Depending on the reporter, the reporter can be detected by visual inspection, Western blotting, Northern blotting, and mRNA sequencing and quantitation, among other methods. Reporters and reporter genes are well known in the art. Examples of common reporters include fluorescent proteins (e.g., green fluorescent protein (GFP), yellow fluorescent protein (YFP), blue fluorescent protein (BFP), red fluorescent protein (RFP), etc.), enzymes or colorimetric reporter proteins (e.g., β -galactosidase, β -D-galactopyranoside (lacZ)), luminescent proteins (e.g., luciferase), and fluorescent RNA aptamers (Bai, J., Luo, Y., Wang, X. et al. A protein-independent fluorescent RNA aptamer reporter system for plant genetic engineering. *Nat Commun* 11, 3847 (2020)), among others.

[0041] The genetic circuits of the invention can comprise a growth inhibitor gene. A growth inhibitor gene is a gene that expresses a gene product that kills or otherwise inhibits the growth and proliferation of a cell in which it is expressed. Examples of growth inhibitor genes include toxin genes and counter-selectable marker genes. Toxin genes are genes that express a gene product that kills a cell in which it is expressed in a non-regulatable manner. Toxin genes are well-known in the art. Examples of common toxin genes include txpA⁴⁹, mazF⁵⁰, and hewl⁵¹, among others. Counter-selectable marker genes are genes that express a gene product that kills a cell in which it is expressed in the presence or absence of a particular condition. In some cases, the condition is the presence or absence of a particular compound or chemical. Counter-selectable marker genes are well known in the art (see, e.g., Reyrat, J M et al. “Counterselectable markers: untapped tools for bacterial genetics and pathogenesis.” *Infection and immunity* vol. 66,9 (1998):

4011-7). Exemplary counter-selectable marker gene/chemical combinations include upp/5-fluorouracil⁵², pheS*/p-chloro-phenylalanine⁵³, and ysbC/fluoro-orotate⁵⁴. Counter-selectable marker genes are sometimes referred to as negative selectable marker genes.

[0042] The genetic circuits of the invention can comprise selectable marker genes. “Selectable marker gene” as used herein refers to a gene that confers a trait suitable for positive selection. Selectable marker genes are well known in the art. Selectable marker genes are often antibiotic resistance genes, which typically produce a protein that provides cells expressing the protein with resistance to an antibiotic. Normally, the genes confer resistance to antibiotics such as ampicillin, chloramphenicol, tetracycline, neomycin, or kanamycin, among others. Other common selectable marker genes are genes that are necessary for an organism to synthesize a particular compound required for survival or growth, e.g., for use in auxotrophic selection. Exemplary selectable marker/function combinations include neo/neomycin resistance⁵⁵ and lysA/lysine production⁵⁶, among others.

[0043] The genetic circuits of the invention can comprise negative regulators of genes. The term “negative regulator” used in reference to regulating a gene refers to any genetic element or combination of elements capable of negatively regulating a gene. “Negatively regulating” (and grammatical variants thereof) as used herein refers to inhibiting, reducing, or repressing expression of a gene or reducing the abundance of a gene product of a gene. The negative regulation of a gene can operate at any step of the expression of the gene, including the transcription of the gene to mRNA and/or the translation of the mRNA to a protein. The negative regulation can also or alternatively operate on the gene products themselves, including reducing the abundance of expressed mRNA or protein (e.g., by specific degradation). Exemplary systems or methods of negatively regulating a gene include siRNA, RNAi, and CRISPRi⁵⁹. Exemplary proteins capable of negatively regulating a gene include repressors (otherwise known as repressor proteins). Repressors are proteins that repress the expression of a gene, typically by binding to an element of the gene, such as its promoter. Some repressors are inducible repressors. Inducible repressors are repressors that repress the expression of a gene in an inducible manner depending on the absence or presence of an inducer, whereby the inducer either activates repression or inhibits repression. A number of inducible repressor/promoter/inducer combinations are known in the art. These include lacI/P_{hyperspank}/IPTG⁵⁷, xylR/P_{xylA}/xylose⁵⁸, and cI/P_R promoter/temperature increase⁵⁵, among others. A gene that expresses a repressor is referred to herein as a “repressor gene.”

[0044] The genetic circuits of the invention can comprise positive regulators of genes. The term “positive regulator” used in reference to regulating a gene refers to any genetic element or combination of elements capable of positively regulating a gene. “Positively regulating” (and grammatical variants thereof) as used herein refers to stimulating expression of a gene or protecting the abundance of any gene product. The positive regulation of a gene can operate at any step of the expression of the gene, including the transcription of the gene to mRNA and/or the translation of the mRNA to a protein. The positive regulation can also or alternatively operate on the gene products themselves, including protecting or maintaining the abundance of expressed mRNA or

protein (e.g., from specific degradation). Exemplary systems or methods of positively regulating a gene include CRISPRa (Liu, Y., Wan, X. & Wang, B. Engineered CRISPRa enables programmable eukaryote-like gene activation in bacteria. *Nat Commun* 10, 3693 (2019)). Exemplary proteins capable of negatively regulating a gene include activators (otherwise known activator proteins). Activators are proteins that activate the expression of a gene, typically by binding to an element of the gene, such as its promoter. Some activators are inducible activators. Inducible activators are activators that activate the expression of a gene in an inducible manner depending on the absence or presence of an inducer, whereby the inducer either stimulates activation or represses activation. A number of inducible activator/promoter/inducer combinations are known in the art. These include exemplary activator/promoter/inducer combinations include *spaR/P_{spaS}/subtilin*⁶⁰, *liaR/P_{liaI}/bacitracin*⁶¹, and *ccaR/P_{cpcG2}/green light*⁶², among others. A gene that expresses an activator is referred to herein as an “activator gene.”

[0045] Quorum-sensing genes can also be employed as positive regulators of genes. Quorum sensing is a specific type of regulation of gene expression in bacteria that is dependent on population density. Quorum-sensing systems include two components: a regulator (autoinducer) and a regulatory receptor protein that interacts with the regulator. The regulator is typically of low-molecular-weight and readily diffuses through the cytoplasmic membrane. As the bacterial population reaches a critical density level, autoinducers accumulate to a threshold value, the regulatory receptor becomes activated, and activation (induction) of genes comprising response elements to the regulatory receptor protein occurs. A gene intending to be positively regulated according to the present invention can comprise or be modified to comprise a promoter sensitive to a regulatory receptor protein of a quorum-sensing system. Autoinducers produced in response to cell density thereby stimulate the regulatory receptor protein to activate expression of such a gene. See e.g., FIG. 15E. A gene that expresses a regulatory receptor protein of a quorum-sensing system is referred to herein as a “quorum-sensing gene.”

[0046] The genetic circuits of the invention can comprise a reporter switch. A reporter switch is a combination of genetic elements configured to express a reporter from a reporter gene in response to a certain condition. The condition in the reporter switches of the present invention is preferably the presence of a target DNA.

[0047] In preferred versions of the invention, the reporter switch comprises a reporter gene in combination with a negative regulator of the reporter gene. The reporter gene preferably comprises a promoter and coding sequence that is not comprised within the interstitial region of the DNA strand. In some versions, no part of the reporter gene is comprised within the interstitial region of the DNA strand. “Not comprised within the interstitial region” as used herein with reference to a particular element means that no portion of that particular element is included within the interstitial region. The negative regulator of the reporter gene is preferably comprised within the interstitial region of the DNA strand. “Comprised within the interstitial region” as used herein with reference to a particular element means that at least some portion of the element is included within the interstitial region. In reporter switches comprising a reporter gene with a promoter and coding sequence not comprised within the interstitial region of the DNA strand and a

negative regulator of the reporter gene comprised within the interstitial region of the DNA strand, the negative regulator of the reporter gene remains intact in the genetic circuit in the absence of target DNA and thereby inhibits expression of the reporter under such conditions. In the presence of target DNA, however, the negative regulator of the reporter gene is excised from the DNA strand, thereby permitting expression of the reporter gene for detection.

[0048] The genetic circuits of the invention can comprise a kill switch. The kill switch comprises one or more genetic elements configured to inhibit growth of the competent cell in response to a certain condition. “Inhibit growth of the competent cell” in this context refers to killing the cell or otherwise inhibiting its proliferation. The condition in the reporter switches of the present invention preferably comprises the absence of a target DNA. The condition may comprise other aspects, such as the presence or absence of an inducer. In preferred versions, at least one of the one or more genetic elements is comprised within the interstitial region of the DNA strand. In such a configuration, the kill switch remains intact in the absence of target DNA, thereby maintaining the ability to inhibit growth of the competent cell with the kill switch. In the presence of target DNA, however, the interstitial region and at least a portion of the kill switch is excised from the DNA strand, thereby removing the ability to inhibit growth of the cell with the kill switch and thereby permitting growth of the cell.

[0049] In some versions, the interstitial region comprises at least one of a growth inhibitor gene, a positive regulator of a growth inhibitor gene, and a negative regulator of a selectable marker gene. In such configurations, the interstitial region the growth inhibitor gene, the positive regulator of the growth inhibitor gene, and/or the negative regulator of the selectable marker gene remain(s) intact in the kill switch in the absence of target DNA, thereby maintaining the ability to inhibit growth of the competent cell with these elements. In the presence of target DNA, however, the growth inhibitor gene, the positive regulator of the growth inhibitor gene, and/or the negative regulator of the selectable marker gene comprised within the interstitial region is excised from the DNA strand, thereby removing the ability to inhibit growth of the cell and thereby permitting growth of the cell with these elements. In some versions, the growth inhibitor gene included in the interstitial region comprises at least one of a toxin gene and a counter-selectable marker gene. In some versions, the positive regulator of the growth inhibitor gene included in the interstitial region comprises at least one of a quorum-sensing gene and an activator gene. In some versions, the negative regulator of the selectable marker gene included in the interstitial region comprises a repressor gene.

[0050] In some versions, the kill switch comprises a toxin gene comprised within the interstitial region and a repressor gene that expresses a repressor of the toxin gene. The repressor of the toxin gene preferably expresses an inducible repressor. An exemplary genetic circuit comprising such a kill switch is shown in FIGS. 1A and 14A. The repressor gene can be comprised within the interstitial region or elsewhere within the competent cell. In such configurations, the toxin gene and, optionally, the repressor of the toxin gene remain intact in the kill switch in the absence of target DNA, thereby maintaining the ability of the toxin gene to inhibit growth of the competent cell. In the presence of target DNA, however, the toxin gene and, optionally, the repressor of the

toxin gene is/are excised from the DNA strand, thereby removing the ability of the toxin gene to inhibit growth of the cell and thereby permitting growth of the cell.

[0051] In some versions, the kill switch comprises a toxin gene and a positive regulator of the toxin gene, wherein one or both of the toxin gene and the positive regulator of the toxin gene is comprised within the interstitial region. An exemplary genetic circuit comprising such a kill switch is shown in FIGS. 15B and 15E. In such a configuration, the toxin gene and the positive regulator of the toxin gene remain intact in the kill switch in the absence of target DNA, thereby maintaining the ability of the toxin gene and the positive regulator of the toxin gene to inhibit growth of the competent cell. In the presence of target DNA, however, the toxin gene and/or the positive regulator of the toxin gene are/is excised from the DNA strand, thereby removing the ability of the toxin gene and the positive regulator of the toxin gene to inhibit growth of the cell and thereby permitting growth of the cell. In some versions, the positive regulator of the toxin gene comprises at least one of an activator gene that expresses an inducible activator of the toxin gene and a quorum-sensing gene that expresses regulatory receptor protein capable of activating the toxin gene.

[0052] In some versions, the kill switch comprises a counter-selectable marker gene comprised within the interstitial region. An exemplary genetic circuit comprising such a kill switch is shown in FIG. 15C. In such a configuration, the counter-selectable marker gene remains intact in the kill switch in the absence of target DNA, thereby maintaining the ability of the counter-selectable marker gene to inhibit growth of the competent cell. In the presence of target DNA, however, the counter-selectable marker gene is excised from the DNA strand, thereby removing the ability of the selectable marker gene to inhibit growth of the cell and thereby permitting growth of the cell.

[0053] In some versions, the kill switch comprises a selectable marker gene not comprised within the interstitial region of the DNA strand and a negative regulator of the selectable marker gene comprised within the interstitial region of the DNA strand. An exemplary genetic circuit with such a kill switch is shown in FIG. 15D. In such a configuration, the negative regulator of the selectable marker gene remains intact in the kill switch in the absence of target DNA, thereby maintaining the ability of the counter-selectable marker to inhibit growth of the competent cell. In the presence of target DNA, however, the negative regulator of the selectable marker gene is excised from the DNA strand, thereby removing the ability of the negative regulator of the selectable marker gene to inhibit growth of the cell and thereby permitting growth of the cell. In some versions, as shown in FIG. 15D, the negative regulator of the selectable marker gene is also the negative regulator of the reporter gene.

[0054] In some versions, the genetic circuit comprises a negative regulator that functions in both the reporter switch and the kill switch (FIGS. 15F-15I). The negative regulator is preferably inducible (shown is *lacI*, but others can be used). The reporter switch comprises a reporter gene (shown is *gfp*, but others can be used) with a promoter and a coding sequence that are not comprised within the interstitial region of the DNA strand, as well as the negative regulator, which negatively regulates the reporter gene.

[0055] In FIGS. 15F and 15G, the kill switch comprises a growth inhibitor gene within the interstitial region, a posi-

tive regulator of the growth inhibitor gene (which can be inside or outside the interstitial region), and the negative regulator, which, in addition to negatively regulating the reporter gene, also negatively regulates the positive regulator of the growth inhibitor gene. In some versions, the negative regulator is comprised within the interstitial region (FIG. 15F). In other versions, the negative regulator is not comprised within the interstitial region (FIG. 15G). In the versions of FIGS. 15F and 15G, cells can be grown in the absence of inducer (such as IPTG) so that the positive regulator of the growth inhibitor gene is inhibited. After exposing to target DNA (or target cells), inducer can be added to stimulate expression of the positive regulator of the growth inhibitor gene and the reporter gene. If the growth inhibitor gene is present, the cells will die. If the growth inhibitor gene is not present, the cells will grow and the reporter will be expressed. In FIGS. 15H and 15I, the kill switch in comprises a growth inhibitor gene (toxin as shown, but others can be used) within the interstitial region and the negative regulator, which, in addition to negatively regulating the reporter gene, also negatively regulates the growth inhibitor gene. In some versions, the first negative regulator is comprised within the interstitial region (FIG. 15H). In other versions, the first negative regulator is not comprised within the interstitial region (FIG. 15I). In the versions of FIGS. 15H and 15I, cells can be grown in the absence of inducer (such as IPTG) so that the growth inhibitor gene is inhibited. After exposing to target DNA (or target cells), inducer can be added to stimulate expression of the growth inhibitor gene and the reporter gene. If the growth inhibitor gene is present, the cells will die. If the growth inhibitor gene is not present, the cells will grow and the reporter will be expressed.

[0056] The target DNA detected with the DNA sensors of the invention can comprise any sequence. In some versions, the target DNA is native DNA. "Native DNA" as used herein is DNA that consists of native sequence, wherein "native sequence" refers to a natural DNA sequence (a sequence found in nature). In some versions, the target DNA is recombinant DNA. "Recombinant DNA" as used herein is DNA that comprises a recombinant sequence, wherein "recombinant sequence" refers to non-natural sequence (a sequence not found in nature.) In some versions, the portion(s) of the target DNA homologous to the pair of homology arms (referred to herein as the first and second portions of the target DNA) consists of native sequence. In some versions, the portion(s) of the target DNA homologous to the pair of homology arms (referred to herein as the first and second portions of the target DNA) comprises recombinant sequence. In some versions, the target DNA is cellular DNA. "Cellular DNA" refers to DNA presently or formerly comprised within a cell. Cellular DNA can comprise genomic DNA, chromosomal DNA, plasmid DNA, etc. In some versions, the target DNA is non-isolated cellular DNA. "Non-isolated cellular DNA" refers to cellular DNA that is not isolated, purified, released, or removed (e.g., by cell lysis) from the cell using chemical or physical treatment. Free cellular DNA present in a culture medium that has not been isolated, purified, released, or removed (e.g., by cell lysis) from the cell using chemical or physical treatment, for example, constitutes non-isolated cellular DNA. In some versions, the non-isolated cellular DNA is from eukaryotic cells. In some versions, the non-isolated DNA is from mammalian cells (e.g., cancer cells). In some versions, the

non-isolated cellular DNA is from prokaryotic cells. In some versions, the non-isolated cellular DNA is from bacterial cells (referred to herein as non-isolated bacterial DNA).

[0057] The target DNA can be DNA from any target organism or target cell. Exemplary organisms comprise bacteria, viruses, fungi, and animals such as humans. For the purposes herein, viruses are considered to be organisms. “Target organism” and “target cell” refer to an organism or cell, respectively, in which target DNA is comprised or from which target DNA is derived.

[0058] Exemplary target bacteria include members of the genus *Bacillus*, such as *Bacillus anthracis* and *Bacillus cereus*; members of the genus *Bordetella*, such as *Bordetella pertussis*; members of the genus *Borrelia*, such as *Borrelia burgdorferi*; members of the genus *Brucella*, such as *Brucella abortus*, *Brucella canis*, *Brucella melitensis*, and *Brucella suis*; members of the genus *Campylobacter*, such as *Campylobacter jejuni*; members of the genus *Chlamydia*, such as *Chlamydia pneumoniae*, *Chlamydia trachomatis*, and *Chlamydia psittaci*; members of the genus *Clostridium*, such as *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, and *Clostridium tetani*; members of the genus *Corynebacterium*, such as *Corynebacterium diphtheriae*; members of the genus *Clostridioides*, such as *Clostridioides difficile* and *Clostridioides mangenotii*; members of the genus *Enterococcus*, such as *Enterococcus faecalis* and *Enterococcus faecium*; members of the genus *Escherichia*, such as *Escherichia coli*; members of the genus *Francisella*, such as *Francisella tularensis*; members of the genus *Haemophilus*, such as *Haemophilus influenzae*; members of the genus *Helicobacter*, such as *Helicobacter pylori*; members of the genus *Legionella*, such as *Legionella pneumophila*; members of the genus *Leptospira*, such as *Leptospira interrogans*; members of the genus *Listeria*, such as *Listeria monocytogenes*; members of the genus *Mycobacterium*, such as *Mycobacterium leprae*, *Mycobacterium tuberculosis*, and *Mycobacterium ulcerans*; members of the genus *Mycoplasma*, such as *Mycoplasma pneumoniae*; members of the genus *Neisseria*, such as *Neisseria gonorrhoeae* and *Neisseria meningitidis*; members of the genus *Pseudomonas*, such as *Pseudomonas aeruginosa*; members of the genus *Rickettsia*, such as *Rickettsia rickettsii*; members of the genus *Salmonella*, such as *Salmonella typhi* and *Salmonella typhimurium*; members of the genus *Shigella*, such as *Shigella sonnei*; members of the genus *Staphylococcus*, such as *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Staphylococcus saprophyticus*; members of the genus *Streptococcus*, such as *Streptococcus agalactiae*, *Streptococcus pneumoniae*, and *Streptococcus pyogenes*; members of the genus *Treponema*, such as *Treponema pallidum*; members of the genus *Vibrio*, such as *Vibrio cholerae*; members of the genus *Yersinia*, such as *Yersinia pestis*, *Yersinia enterocolitica*, and *Yersinia pseudotuberculosis*, among others.

[0059] Exemplary target viruses include viruses in the family adenoviridae, such as adenovirus; viruses in the family herpesviridae such as herpes simplex, type 1, herpes simplex, type 2, varicella-zoster virus, epstein-barr virus, human cytomegalovirus, human herpesvirus, and type 8; viruses in the family papillomaviridae such as human papillomavirus; viruses in the family polyomaviridae such as BK virus and JC virus; viruses in the family poxviridae such as smallpox; viruses in the family hepadnaviridae such as hepatitis B virus; viruses in the family parvoviridae such as

human bocavirus and parvovirus B19; viruses in the family astroviridae such as human astrovirus; viruses in the family caliciviridae such as norwalk virus; viruses in the family picornaviridae such as coxsackievirus, hepatitis A virus, poliovirus, and rhinovirus; viruses in the family coronaviridae such as severe acute respiratory syndrome (SARS) viruses, including SARS-CoV-2; viruses in the family flaviviridae such as hepatitis C virus, yellow fever virus, dengue virus, and West Nile virus; viruses in the family togaviridae such as rubella virus; viruses in the family hepeviridae such as hepatitis E virus; viruses in the family retroviridae such as human immunodeficiency virus (HIV); viruses in the family orthomyxoviridae such as influenza virus; viruses in the family arenaviridae such as guarnarito virus, junin virus, lassa virus, machupo virus, and sabia virus; viruses in the family bunyaviridae such as Crimean-Congo hemorrhagic fever virus; viruses in the family filoviridae such as ebola virus and marburg virus; viruses in the family paramyxoviridae such as measles virus, mumps virus, parainfluenza virus, respiratory syncytial virus, human metapneumovirus, hendra virus, and nipah virus; viruses in the family rhabdoviridae such as rabies virus; unassigned viruses such as hepatitis D virus; and viruses in the family reoviridae such as rotavirus, orbivirus, coltivirus, and banna virus, among others.

[0060] Exemplary target fungi include fungi of the genus *Aspergillus*, such as *Aspergillus fumigatus*, which cause aspergillosis; fungi of the genus *Blastomyces*, such as *Blastomyces dermatitidis*, which cause blastomycosis; fungi of the genus *Candida*, such as *Candida albicans*, which cause candidiasis; fungi of the genus *Coccidioides*, which cause coccidioidomycosis (valley fever); fungi of the genus *Cryptococcus*, such as *Cryptococcus neoformans* and *Cryptococcus gattii*, which cause cryptococcosis; dermatophytes fungi, which cause ringworm; fungi that cause fungal keratitis, such as *Fusarium* species, *Aspergillus* species, and *Candida* species; fungi of the genus *Histoplasma*, such as *Histoplasma capsulatum*, which cause histoplasmosis; fungi of the order Mucorales, which cause mucormycosis; fungi of the genus *Saccharomyces*, such as *Saccharomyces cerevisiae*; fungi of the genus *Pneumocystis*, such as *Pneumocystis jirovecii*, which cause *pneumocystis* pneumonia; and fungi of the genus *Sporothrix*, such as *Sporothrix schenckii*, which cause sporotrichosis.

[0061] In some versions, the target DNA sequence is an antibiotic resistance gene or a regulator gene, wherein homologous recombination corrects the point mutation in the kill switch or the reporter to thereby activate the kill switch or reporter.

[0062] The DNA sensors of the invention can be comprised in compositions. In some versions, the compositions comprise one or more DNA sensors in a culture medium. The culture medium is preferably capable of supporting growth of the DNA sensor under certain conditions. Such conditions preferably include the presence of target DNA. In some versions, the compositions comprise one or more DNA sensors in combination with a target DNA. In some versions, the compositions comprise two or more DNA sensors, such as three or more, four or more, five or more, or six or more DNA sensors. The two or more DNA sensors can be configured to detect different target DNA sequences, such as from different target organisms or target cells, and express different reporters. In one exemplary configuration, the pairs of homology arms in the two or more cell-based

sensors are each homologous to different target DNA sequences, and the reporter genes in the two or more cell-based DNA sensors express reporters that are each detectably different from each other. In some versions, the detectably different reporters comprise fluorescent reporters that emit fluorescence at different wavelengths. The compositions comprising the two or more DNA sensors can be used in multiplex detection methods.

[0063] The DNA sensors of the invention can be used in methods of detecting target DNA. The methods can comprise culturing the DNA sensor in a culture medium comprising the target DNA for a time effective to transform the DNA sensor with the target DNA and then detecting the transformed DNA sensor. “Transformation” (and grammatical variants) as used herein refers to the uptake of DNA into a cell. The genetic circuits of the transformed DNA sensors can then undergo homologous recombination with the transformed DNA, thereby excising the interstitial region from the DNA strand and altering the reporter switch (if present), the kill switch (if present), or both (if present). In some versions, the DNA sensor comprises a reporter switch, and detection of the target DNA occurs by detecting growth of the DNA sensor. In some versions, the DNA sensor comprises a kill switch, and detection of the target DNA occurs by detecting a reporter expressed from the reporter gene. In some versions, the DNA sensor comprises a reporter switch and a kill switch, and detection of the target DNA occurs by detecting a reporter expressed from the reporter gene and/or growth of the DNA sensor. It has been found that including both a reporter switch and a kill switch in a DNA sensor of the invention can enhance detection by amplifying the signal to noise (background) ratio of the reporter with respect to including a reporter switch alone.

[0064] It has been surprisingly found that target cells such as bacteria release sufficient DNA during culture to transform the DNA sensors of the invention and undergo homologous recombination for detection thereof without isolating, purifying, releasing, or removing (e.g., by cell lysis) target DNA from the target cell using chemical or physical treatment. Accordingly, some methods of the invention are directed to detecting a target cell comprising target DNA. Such methods can comprise culturing the DNA sensor in a culture medium with the target cell for a time effective to transform the DNA sensor with the target DNA and then detecting the transformed DNA sensor. In some methods, the culturing is performed without lysing the target cell. In some methods, the method is performed without isolating, purifying, releasing, or removing (e.g., by cell lysis) the target DNA from the target cell using chemical or physical treatment. In some methods, the target DNA is non-isolated cellular DNA. In some methods the target cell comprises a bacterium. In some methods, the target DNA is non-isolated bacterial DNA. In some methods, the cell-based DNA sensor comprises two or more cell-based DNA sensors, wherein: the pairs of homology arms the two or more cell-based sensors are each homologous to different target DNA sequences from different target cells; each of the two or more cell-based DNA sensors comprises the reporter switch; and the reporter genes in the two or more cell-based DNA sensors express reporters that are each detectably different from each other. In some methods, the two or more cell-based sensors comprise three or more cell-based sensors.

[0065] Aspects pertaining to homologous recombination such as homology length and sequence identity are known in the art. 66,67,68,69

[0066] Unless the context indicates otherwise, any gene described herein as “expressing” or being one that “expresses” a particular gene product can constitutively express the gene product (e.g., via a constitutive promoter) or inducibly express the gene product (e.g., via an inducible promoter or a promoter sensitive to a regulatory protein, such as a repressor or an activator). The unmodified term “expresses” therefore encompasses but does not necessarily require constitutive expression. The phrases “expresses” and “configured to express” are used interchangeably herein.

[0067] Terms used herein pertaining to genetic manipulation are defined as follows.

[0068] Endogenous: As used herein with reference to a polynucleotide molecule and a particular cell, “endogenous” refers to a polynucleotide sequence or polypeptide that is in the cell and was not introduced into the cell using recombinant engineering techniques. For example, an endogenous gene is a gene that was present in a cell when the cell was originally isolated from nature.

[0069] Exogenous: As used herein with reference to a polynucleotide molecule or polypeptide in a particular cell, “exogenous” refers to any polynucleotide molecule or polypeptide that does not originate from that particular cell as found in nature. Thus, a non-naturally-occurring polynucleotide molecule or protein is considered to be exogenous to a cell once introduced into the cell. A polynucleotide molecule or protein that is naturally occurring also can be exogenous to a particular cell. For example, an entire coding sequence isolated from cell X is an exogenous polynucleotide with respect to cell Y once that coding sequence is introduced into cell Y. The term “heterologous” is used herein interchangeably with “exogenous.”

[0070] Expression: The process by which a gene’s coded information is converted into the structures and functions of a cell, such as a protein, transfer RNA, or ribosomal RNA. Expressed genes include those that are transcribed into mRNA and then translated into protein and those that are transcribed into RNA but not translated into protein (for example, transfer and ribosomal RNAs).

[0071] Isolated: Except as otherwise defined herein, an “isolated” biological component (such as a polynucleotide molecule, polypeptide, or cell) has been substantially separated or purified away from other biological components in which the component naturally occurs, such as other chromosomal and extrachromosomal DNA and RNA and proteins. Polynucleotide molecules and polypeptides that have been “isolated” include polynucleotide molecules and polypeptides purified by standard purification methods. The term also includes polynucleotide molecules and polypeptides prepared by recombinant expression in a cell as well as chemically synthesized polynucleotide molecules and polypeptides.

[0072] Polynucleotide: Encompasses both RNA and DNA molecules including, without limitation, cDNA, genomic DNA, and mRNA. Polynucleotides also include synthetic polynucleotide molecules, such as those that are chemically synthesized or recombinantly produced. The polynucleotide can be double-stranded or single-stranded. Where single-stranded, the polynucleotide molecule can be the sense strand, the antisense strand, or both. In addition, the polynucleotide can be circular or linear.

[0073] Operably linked: A first polynucleotide sequence is operably linked with a second polynucleotide sequence when the first polynucleotide sequence is placed in a functional relationship with the second polynucleotide sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. An origin of replication is operably linked to a coding sequence if the origin of replication controls the replication or copy number of the polynucleotide in the cell. Operably linked polynucleotides may or may not be contiguous.

[0074] Operon: Configurations of separate genes that are transcribed in tandem as a single messenger RNA are denoted as operons. Thus, a set of in-frame genes in close proximity under the transcriptional regulation of a single promoter constitutes an operon. Operons may be synthetically generated.

[0075] Overexpress: When a gene is caused to be transcribed at an elevated rate compared to the endogenous or basal transcription rate for that gene. In some examples, overexpression additionally includes an elevated rate of translation of the gene compared to the endogenous translation rate for that gene. Methods of testing for overexpression are well known in the art, for example transcribed RNA levels can be assessed using rtPCR and protein levels can be assessed using SDS page gel analysis.

[0076] Recombinant cell: A cell that comprises a recombinant polynucleotide.

[0077] Unless explained otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this disclosure belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below.

[0078] A coding sequence can be operably linked to an appropriate expression control sequence (promoters, enhancers, and the like) to direct synthesis of the encoded gene product. Such promoters can be derived from microbial or viral sources, including CMV and SV40. Depending on the cell/vector system utilized, any of a number of suitable transcription and translation control elements, including constitutive and inducible promoters, transcription enhancer elements, transcription terminators, etc. can be used in the expression vector (see e.g., Bitter et al. (1987) *Methods in Enzymology*, 153:516-544).

[0079] Suitable promoters for use in prokaryotic cells include but are not limited to: promoters capable of recognizing the T4, T3, Sp6, and T7 polymerases; the P_R and P_L promoters of bacteriophage lambda; the trp, recA, heat shock, and lacZ promoters of *E. coli*; the alpha-amylase and the sigma-specific promoters of *B. subtilis*; the promoters of the bacteriophages of *Bacillus*; *Streptomyces* promoters; the int promoter of bacteriophage lambda; the bla promoter of the beta-lactamase gene of pBR322; and the CAT promoter of the chloramphenicol acetyl transferase gene. Prokaryotic promoters are reviewed by Glick, *J. Ind. Microbiol.* 1:277 (1987); Watson et al, *Molecular Biology of the Gene*, 4th Ed., Benjamin Cummins (1987); and Sambrook et al., In: *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press (2001).

[0080] Non-limiting examples of suitable promoters for use within a eukaryotic cell are typically viral in origin and include the promoter of the mouse metallothionein I gene

(Hamer et al. (1982) *J. Mol. Appl. Genet.* 1:273); the TK promoter of Herpes virus (McKnight (1982) *Cell* 31:355); the SV40 early promoter (Benoist et al. (1981) *Nature* (London) 290:304); the Rous sarcoma virus promoter; the cytomegalovirus promoter (Foecking et al. (1980) *Gene* 45:101); the yeast gal4 gene promoter (Johnston et al. (1982) *PNAS* (USA) 79:6971; Silver et al. (1984) *PNAS* (USA) 81:5951); and the IgG promoter (Orlandi et al. (1989) *PNAS* (USA) 86:3833).

[0081] Coding sequences can be operably linked to an inducible promoter. Inducible promoters are those wherein addition of an effector affects expression. Suitable effectors include proteins, metabolites, chemicals, or culture conditions capable of affecting expression. Suitable inducible promoters include but are not limited to the lac promoter (regulated by IPTG or analogs thereof), the lacUV5 promoter (regulated by IPTG or analogs thereof), the tac promoter (regulated by IPTG or analogs thereof), the trc promoter (regulated by IPTG or analogs thereof), the araBAD promoter (regulated by L-arabinose), the phoA promoter (regulated by phosphate starvation), the recA promoter (regulated by nalidixic acid), the proU promoter (regulated by osmolarity changes), the cst-1 promoter (regulated by glucose starvation), the tetA promoter (regulated by tetracycline), the cadA promoter (regulated by pH), the nar promoter (regulated by anaerobic conditions), the p_L promoter (regulated by thermal shift), the cspA promoter (regulated by thermal shift), the T7 promoter (regulated by thermal shift), the T7-lac promoter (regulated by IPTG), the T3-lac promoter (regulated by IPTG), the T5-lac promoter (regulated by IPTG), the T4 gene 32 promoter (regulated by T4 infection), the nprM-lac promoter (regulated by IPTG), the VHb promoter (regulated by oxygen), the metallothionein promoter (regulated by heavy metals), the MMTV promoter (regulated by steroids such as dexamethasone) and variants thereof.

[0082] In some versions, the promoter is a constitutive promoter. Suitable constitutive promoters are known in the art and include constitutive adenovirus major late promoter, a constitutive MPSV promoter, and a constitutive CMV promoter.

[0083] The elements of the genetic circuits of the invention can be integrated in the competent cell's chromosome or an extra-chromosomal DNA, such as a plasmid, or a combination thereof.

[0084] Polynucleotides encoding enzymes desired to be expressed in a cell may be codon-optimized for that particular type of cell. Codon optimization can be performed for any polynucleotide by "OPTIMUMGENE"-brand gene design system by GenScript (Piscataway, NJ).

[0085] The elements and method steps described herein can be used in any combination whether explicitly described or not.

[0086] All combinations of method steps as used herein can be performed in any order, unless otherwise specified or clearly implied to the contrary by the context in which the referenced combination is made.

[0087] As used herein, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise.

[0088] Numerical ranges as used herein are intended to include every number and subset of numbers contained within that range, whether specifically disclosed or not. Further, these numerical ranges should be construed as

providing support for a claim directed to any number or subset of numbers in that range. For example, a disclosure of from 1 to 10 should be construed as supporting a range of from 2 to 8, from 3 to 7, from 5 to 6, from 1 to 9, from 3.6 to 4.6, from 3.5 to 9.9, and so forth.

[0089] All patents, patent publications, and peer-reviewed publications (i.e., “references”) cited herein are expressly incorporated by reference to the same extent as if each individual reference were specifically and individually indicated as being incorporated by reference. In case of conflict between the present disclosure and the incorporated references, the present disclosure controls.

[0090] It is understood that the invention is not confined to the particular construction and arrangement of parts herein illustrated and described, but embraces such modified forms thereof as come within the scope of the claims.

EXAMPLES

[0091] Programming Bacteria for Multiplexed DNA Detection Summary

[0092] DNA is a universal and programmable signal of living organisms. Here we developed cell-based DNA sensors by engineering the naturally competent bacterium *Bacillus subtilis* (*B. subtilis*) to detect specific DNA sequences in the environment. The DNA sensor strains can identify diverse bacterial species including major human pathogens with high specificity and sensitivity. Multiplexed detection of genomic DNA from different species in complex samples can be achieved by coupling the sensing mechanism to orthogonal fluorescent reporters. We also demonstrate that the DNA sensors can detect the presence of species in the complex samples without requiring DNA extraction. The modularity of the living cell-based DNA sensing mechanism and simple detection procedure enables programmable DNA sensing for broad applications.

INTRODUCTION

[0093] Next-generation engineered bacteria hold tremendous promise for a wide range of applications in human health, environment and agriculture by sensing key environmental signals, performing computation on these signals to regulate a response that modulates specific environmental parameters¹. Developing specific and selective sensors of key environmental signals is a critical feature of next-generation engineered bacteria. For example, bacteria have been engineered to detect physical and chemical signals such as light, ultrasound, and quorum-sensing molecules²⁻⁴. These signals can be exploited to control the collective growth or gene expression of the bacterial population or mediate interactions between constituent community members⁴⁻⁶. In addition, we can exploit their sensing ability to achieve real-time monitoring of natural environments. For example, synthetic genetic circuits have been designed in *Escherichia coli* (*E. coli*) to sense signals produced by pathogens and use this information to regulate the production of antimicrobials that inhibit the target pathogen^{7,8}. However, there are limited well-characterized and orthogonal signals that can be exploited to sense different bacterial species in a microbial community^{9,10}.

[0094] DNA provides the blueprint for living organisms and is prevalent in natural environments¹¹. Therefore, extracellular DNA (eDNA) could be exploited as a biomarker for identifying different species. Naturally competent bacteria

have the ability to take up DNA from the environment and integrate imported sequences onto the genome based on sequence homology requirements. Horizontal gene transfer (HGT) via natural transformation has been shown to have a variety of benefits such as nutrient utilization, DNA repair, or acquisition of genes¹². Since homologous recombination of imported DNA requires sequences of sufficient length and homology¹³, natural transformation could be exploited to build a selective cell-based DNA sensor.

[0095] We constructed a living cell-based DNA sensor by engineering the naturally competent bacterium *B. subtilis*. This circuit controls *B. subtilis* growth and fluorescence reporter genes in response to specific input DNA sequences. We demonstrate that the cell-based DNA sensor is sensitive and highly specific to species harboring the target DNA sequence. In addition, we demonstrate that our cell-based DNA sensor can perform multiplexed DNA detection in complex samples. The cell-based DNA sensors can detect DNA released from pre-treated donor cells (i.e. crude samples). Our detailed characterization of the cell-based DNA sensors in vitro provides a foundation for future in vitro DNA detection and in situ sense-and-respond DNA applications.

Results

Construction of a Living DNA Sensor Strain

[0096] To build the living cell-based DNA sensor, we exploited the natural competence ability of the well-characterized soil bacterium *B. subtilis*¹⁴. The natural competence ability of *B. subtilis* enables uptake of environmental DNA and integration of specific sequences with sufficient homology into genome via homologous recombination¹⁵. The efficiency of homologous recombination depends stringently on the sequence percent identity and length^{13,16}, which can be exploited to build a highly specific DNA sensor.

[0097] To detect eDNA sequences in a programmable fashion, we constructed a synthetic genetic circuit in *B. subtilis* that implements a growth selection function based on the presence of target DNA sequence in the environment. The circuit consists of a xylose-inducible master regulator of competence *comK*¹⁷ and IPTG-inducible toxin-antitoxin system *txpA-ratA*¹⁸ and GFP regulated by the repressor *Lad* (FIGS. 1A and 5A-5C). The target sequences were introduced to the flanking regions (upstream and downstream) of *txpA-ratA* and *lad* and referred to as landing pads for homologous recombination.

[0098] In the presence of xylose, *ComK* activates competence genes for DNA uptake and homologous recombination (FIG. 1A). Bistability and stochastic processes in the regulation of natural competence can yield a sub-population that can be transformed with extracellular DNA¹⁹. This naturally competent sub-population forms competence pili which bind to double stranded DNA outside the cell. The DNA is cleaved into single-stranded DNA (ssDNA) outside the cell membrane, and transported into the cell¹². Inside the cell, *RecA* binds the ssDNA sequences and searches the *B. subtilis* genome for a region with sufficient homology. If the target DNA sequence is present, homologous recombination removes the toxin-antitoxin *txpA-ratA* and repressor *lad*. In the presence of the chemical inducer Isopropyl β -D-1-thiogalactopyranoside (IPTG), cell growth and GFP expression is enabled in the transformed sub-population. Growth

of the non-transformed subpopulation is inhibited by the activity of TxpA, which blocks cell wall synthesis¹⁸ (FIG. 1A).

[0099] We constructed a sensor for *E. coli* (EC sensor) by introducing the xdhABC operon onto the *B. subtilis* genome (landing pad region), which encodes genes for purine catabolism²⁰. The xdhABC operon is a representative sequence that can detect a wide range of *E. coli* strains. This sequence is highly conserved such that 99% of 5000 *E. coli* genomes in the NCBI database contain this sequence with >95% coverage (the degree of alignment of the query sequence with a reference sequence) and >95% identity similarity (the percentage of bases that are identical to the target sequence within the aligned region).

[0100] To characterize the homology length needed for robust DNA sensing, we varied the homology length of the xdhABC operon in each landing pad (0.5 to 2.5 kb). We performed time-series measurements of transformation efficiency (number of colonies for transformed *B. subtilis* cells divided by the number total *B. subtilis* colonies) with 100 ng/mL *E. coli* genomic DNA (gDNA). The transformation efficiency is defined as the ratio of the number of transformed *B. subtilis* to the total number of *B. subtilis* based on colony forming units (CFU). Transformation efficiency plateaued at approximately 10 hr and the colonies expressed GFP (FIG. 1B). In addition, transformation efficiency increased with homology length at 10 hr (FIG. 1C). A homology length of 1 kb or greater was required to robustly sense the target sequence over the background frequency of escape mutants (10^{-7} - 10^{-6} frequency) that displayed heterogeneous GFP expression (FIGS. 6A and 6B). To achieve high performance of the DNA sensor ($>10^2$ increase in transformation efficiency above background), we used a landing pad homology length of 2.5 kb (transformation efficiency of 10^{-5} - 10^{-4}). In the transformed sub-population, homologous recombination was confirmed by sequencing to occur at the expected location with the elimination of txpA-ratA and lad (FIGS. 6C and 6D). The moderate number of escape mutants that displayed growth in the absence of gDNA had mutations in txpA or lacI, which reduced the growth inhibitory activity of TxpA (FIGS. 6E and 6F). In sum, the synthetic genetic circuit enabled *B. subtilis* to sense specific DNA sequences present in the environment.

Building Living DNA Sensors to Sense Human Pathogens

[0101] Exploiting the modularity of the DNA sensing circuit, we replaced the landing pad region with specific sequences targeting different bacterial strains (FIG. 5C). To this end, we constructed DNA sensors to detect sequences harbored in human intestinal pathogens *Salmonella typhimurium*²¹ (*S. typhimurium*), *Clostridium difficile*²² (*C. difficile*), or the skin pathogen *Staphylococcus aureus*²³ (*S. aureus*). We selected two 2.5 kb sequences in the pathogenicity island sipBCDA of *S. typhimurium* (ST sensor), the heme biosynthesis pathway hemEH in *S. aureus* (SA sensor), and the phenylalanyl-tRNA synthetase pheST in *C. difficile* (CD sensor)²⁴⁻²⁶.

[0102] The selected set of target DNA sequences are highly conserved within a given species (ST sensor: 94%, SA sensor: 96%, and CD sensor: 96% all with >95% coverage and >95% identity similarity). In addition, some of the sequences are linked to virulence activities of the pathogen or encode enzymes that are critical for fitness²⁴⁻²⁶. To further explore the conservation of the target sequences

across different strains, we performed nucleotide BLAST using the NCBI Database to quantify the homology coverage and sequence similarity across species. The pathogenicity island sipBCDA in *S. typhimurium* was found only in *Salmonella enterica* species and infrequently observed in other species (FIG. 7A). Homologs in other species have low coverage and identity similarity, suggesting that the pathogenicity island could be a good target sequence for this species (FIG. 7A). The heme biosynthesis pathway hemEH in *S. aureus* and phenylalanyl-tRNA synthetase pheST in *C. difficile* are conserved in some closely related strains with varying degrees of similarity and coverage (FIGS. 2I and 2K). The *E. coli* MG1655 xdhABC purine catabolism operon is found in other closely related bacteria such as *Shigella* with high coverage and identity similarity (FIG. 7B). Although the target sequences for building the different DNA sensor strains varied in the degree of specificity based on bioinformatic analyses, a detailed characterization of circuit performance could guide the design of optimized cell-based DNA sensors for future applications.

[0103] The four sensors robustly detected the presence of 100 ng/mL target gDNA over background (0 ng/mL gDNA) based on transformation efficiency (FIG. 2A). We evaluated the sensitivity of each DNA sensor strain by performing time-series GFP measurements in liquid culture after being transformed with a wide range of gDNA concentrations (0-1500 ng/mL gDNA) from single species (FIGS. 2B, 2C, and 8A-8C). We evaluated the time required for each culture to display a fluorescence level higher than a threshold (i.e. detection time) (FIGS. 2C and 8A-8C). The sensitivity of the circuit was evaluated as the lowest gDNA concentration that yielded a statistically significant difference in the detection time in the presence versus absence of gDNA (FIGS. 2D-2G).

[0104] The relationship between the log transformed gDNA concentration and detection time is linear due to the exponential growth of the fluorescent *B. subtilis* sub-population successfully transformed with the input target sequence (FIG. 9A-9D). Therefore, to assess the range of gDNA concentrations that can be accurately sensed, we inferred the parameters of a linear function fit to the log transformed gDNA concentration versus detection time (FIGS. 2D-2G). The inferred slope of the linear function is determined by the cell doubling time (~0.5 hour) and intercept is determined by the background mutation frequency (FIGS. 2D-2G). The EC, SA, and CD DNA sensor strains displayed high sensitivity of 1-16 ng/mL (10^5 - 10^6 chromosome copy number/mL), whereas the ST sensor displayed a lower sensitivity (62.5 ng/mL, 10^7 chromosome copy number/mL). While the DNA sensor strains displayed lower sensitivities than quantitative real-time polymerase chain reaction (qPCR) reported for *E. coli* (3.5×10^3 CFU/mL in pure culture²⁷), the observed sensitivities are within the range of the sensitivities reported for the lateral flow immunoassay (1.8×10^5 CFU/mL for a pure culture of *E. coli*²⁸).

[0105] To characterize the specificity of each DNA sensor strain to the target sequence, we performed time-series fluorescence measurements in liquid culture in response to all individual species gDNA. The fluorescence signal was observed at a substantially earlier time (6.1-7.1 hr) in the presence of the corresponding species' gDNA than in the presence of a non-target species' gDNA or in the absence of

DNA (9.1-10.7 hr) (FIGS. 2H and 10A-10H). This demonstrates that the cell-based DNA sensors were highly specific to the target sequence.

[0106] In the context of microbial communities, the cell-based DNA sensors may need to distinguish between closely related species. Therefore, we evaluated the ability of the DNA sensors to distinguish between closely related species with similar target sequences. To this end, we measured the transformation frequency of the SA sensor in the presence of gDNA (100 ng/mL) derived from *S. epidermidis*. *S. epidermidis* is a closely related human skin commensal bacterium that harbors a similar hemeH sequence to the SA sensor landing pad region (89% coverage and 77% identity similarity) (FIG. 2I). Since the number of total *B. subtilis* colonies was similar in the presence and absence of gDNA, we quantified the number of transformed colonies as opposed to transformation efficiency (FIGS. 2A, 11A, and 11B). The number of colonies in the presence of *S. epidermidis* gDNA (SE) was substantially lower than in the presence of *S. aureus* gDNA (SA) and similar to the absence of DNA (FIG. 2J). Similarly, we characterized the ability of the CD sensor to detect gDNA derived from a human gut commensal bacterium *C. hiranonis*, a close relative of *C. difficile* that contains a similar pheST sequence in its genome to the landing pad region in CD sensor (87% coverage and 75% identity similarity). The number of colonies in the presence of 100 ng/mL *C. hiranonis* gDNA (CH) was substantially lower than in the presence of *C. difficile* gDNA (CD) and also similar to the absence of DNA (FIGS. 2K and 2L). These data demonstrate that the cell-based DNA sensors are highly specific to species that harbor an exact match to the target sequence. Therefore, the sensors do not display false positives in the presence of closely related species that harbor similar target sequences. The high specificity of the sensors is due to the stringent requirements for homologous recombination in *B. subtilis*^{13,29}.

Multiplexed Detection of Pathogen DNA in Complex Samples

[0107] Since certain future applications may require sensing of more than one organism, we tested the ability of the DNA sensors to detect more than one species within mixed DNA samples. To this end, we constructed individual sensors with orthogonal fluorescent reporters to achieve multiplexed DNA detection. Exploiting the modularity of the circuit, we constructed an RFP-labeled ST sensor (ST-R) and a BFP-labeled SA sensor (SA-B), in addition to the GFP-labeled EC sensor (EC-G) (FIG. 5C). We introduced gDNA (200 ng/mL) extracted from each of the three target strains into a culture containing EC-G, ST-R, and SA-B sensors and determined the number of fluorescent colonies for each reporter (FIG. 3A). The sensors accurately reported the presence/absence of all combinations of species' gDNA reliably (FIGS. 3B and 12). Therefore, a mixture of DNA sensor strains each individually labeled with a unique fluorescent reporter enabled multiplexed detection of gDNA derived from different species. To investigate if multiplexed detection can be achieved for samples derived from a complex microbial community, we constructed a four-member human gut community composed of diverse commensal bacteria from three major phyla in human gut—*Anaerostipes caccae* (AC, Firmicutes), *Bacteroides thetaiotaomicron* (BT, Bacteroidetes), *Bifidobacterium longum* (BL, Actinobacteria), and *Clostridium asparagiforme* (CG, Firmicutes). This community also contained the target pathogens *S. typhimurium* (ST) and *S. aureus* (SA). We tested whether the

DNA sensors could accurately report the relative abundance of the two pathogens during community assembly. The 6-member community was inoculated in equal initial species proportions based on absorbance at 600 nm (OD₆₀₀, Day 0) and cultured anaerobically for 24 hr (Day 1). An aliquot of the community was transferred to fresh media and community composition was characterized following an additional 24 hr (Day 2). Based on 16S rRNA gene sequencing, the abundance of *S. typhimurium* was similar as a function of time whereas the abundance of *S. aureus* decreased over time (FIG. 3C).

[0108] We characterized the ability of the DNA sensors to accurately track the temporal trends in species abundance by introducing purified community gDNA collected at different times into a mixed culture of the ST-R and SA-G sensors (FIG. 3D). Due to the low abundance of target species in the sample, a higher amount of DNA (1 µg/mL) was used for transformation. Consistent with the trends based on 16S rRNA gene sequencing, the number of GFP fluorescent colonies of the SA-G sensor decreased at sequential time points, whereas the number of RFP fluorescent colonies of ST-R sensor were similar at sequential time points (FIGS. 3D-3F). The SA sensor displayed better performance in mirroring the trend from 16S rRNA gene sequencing than the ST sensor, consistent with its higher sensitivity than other sensors (FIGS. 2A, 3E, and 3F). For the community lacking *S. typhimurium* and *S. aureus*, a much smaller number of background colonies was detected than in the 6-member community. This implies that the ST and SA sensors were specific to the target species gDNA and did not generate false positives in the presence of the other constituent community member gDNA (FIGS. 13A and 13B). In sum, our results show that accurate multiplexed DNA detection can be achieved in samples derived from multi-species microbial communities.

Detection of Target Species without DNA Extraction

[0109] Specific bacterial species have been shown to release eDNA in response to environmental stimuli¹¹, suggesting that the DNA sensor could detect species without requiring prior gDNA purification. To test this possibility, we co-cultured individual DNA sensor strains with the corresponding donor species with an initial OD₆₀₀ 0.1 of the target strain (1.22×10⁸ CFU/mL, 1.07×10⁸ CFU/mL, 3.2×10⁸ CFU/mL, and 1.1×10⁷ CFU/mL for *E. coli*, *S. typhimurium*, *S. aureus*, and *C. difficile*, respectively) (FIG. 4A). Since the other species could compete with *B. subtilis*, we introduced specific antibiotics (ABX) to inhibit the growth of the donor cells and enhance donor eDNA release. The DNA sensor strains are resistant to the antibiotics since they harbor the appropriate antibiotic resistance genes. In the presence of 100 µg/mL spectinomycin, the DNA sensors displayed robust detection of *E. coli*, *S. typhimurium*, and *S. aureus* (FIG. 4B). The addition of spectinomycin was not required for *C. difficile* detection since the growth of *C. difficile* is negatively impacted by the presence of oxygen³⁰. To confirm the transformation was mediated by the eDNA released from the target strain, DNase I (1 unit/mL) was added into the co-culture. The number of transformed cells was substantially lower in the presence of DNase I, indicating that DNA detection occurred via natural transformation in the co-cultures (FIG. 4B). Antibiotic resistance is prevalent in microbiomes and may not be used universally as a treatment for the donor cells. Therefore, we tested if heat treatment could be used to efficiently release donor cell DNA. Incubation of *E. coli* at 90° C. for 10 minutes substantially enhanced the EC sensor detection limit (5×10⁶

CFU/mL) compared to the addition of spectinomycin (FIG. 4C). In sum, detection of target DNA sequences directly from crude samples in the absence of DNA purification could enable the deployment of the DNA sensors for different future applications.

[0110] To evaluate the robustness of the DNA sensing function, we characterized the performance of the DNA sensors for multiplexed detection of spike-in bacteria in the presence of cecal contents derived from germ-free mice that were orally gavaged with a defined bacterial consortium (Methods). Mouse ceca contain other bacterial species, host cells and other chemical compounds (e.g. dietary factors), and thus can be used to evaluate the robustness of the DNA sensor. To this end, we introduced varying amounts of *E. coli* and *S. typhimurium* into 10 mg of mouse ceca, incubated these samples at 90° C. for 10 minutes, and then transferred the samples into a mixed culture containing the EC-G and ST-R sensors (FIG. 4D). Our results demonstrated that both sensors can detect *E. coli* and *S. typhimurium* cells in ceca without DNA extraction. In particular, the EC and ST sensors displayed a detection limit of 10⁷ CFU/mL (FIG. 4D). In samples containing a single donor species, high density of *E. coli* or *S. typhimurium* (10⁸ CFU/mL) yielded infrequent false positives for the multiplexed DNA detection. This suggests that further optimization of the DNA sensors may be needed in complex samples containing high donor cell densities that are heat treated (FIGS. 14A and 14B). In sum, we show that the cell-based DNA sensors can robustly perform DNA detection of heat-treated samples that contain the target sequence.

DISCUSSION

[0111] Here we engineered the naturally competent bacterium *B. subtilis* to sense and respond to specific DNA sequences. DNA sensing can be achieved for purified DNA or eDNA released from pre-treated samples containing donor cells harboring the target sequence. We demonstrate that DNA sensing can be sensitive and specific, and multiplexed sensing can be achieved by engineering sensors with orthogonal reporter genes. Detection of species using a living cell-based DNA sensor strain opens avenues for future research for versatile sensing of species and does not rely on chemical or physical signals^{31,32}. Since our circuit design is modular, customized sensors could be constructed in the future for the detection of sequences derived from diverse organisms including viruses, fungi and mammalian cells.

[0112] The DNA detection limit is impacted by the frequency of background mutations and could be improved by reducing the background genetic mutation rate. For example, counter-selectable markers³³ that can achieve lower background mutation rate could be used to optimize the strength of negative growth selection. The mutation rate can also be reduced by deleting endogenous genes in *B. subtilis* that promote mutagenesis such as the transcription conflict factor *mfd*³⁴. The reduction of mutation rates is also critical for long-term implementation of living DNA sensors in the environments³⁵.

[0113] The time required for DNA detection using the cell-based DNA sensors is relatively slow compared to other diagnostic methods³⁶. In particular, the total time required for DNA detection including transformation and selection is approximately one day, which is not suitable for certain applications that require a rapid response. To reduce the detection time, directed evolution or rational design of the

natural competence pathway could be used for enhancing the transformation efficiency of *B. subtilis*³⁷. This in turn would reduce the time required for transformation and detection of GFP in liquid media. In addition, some naturally competent bacteria such as *Streptococcus pneumoniae* can achieve 50% transformation efficiency³⁸. This transformation efficiency is substantially higher than our constructed DNA sensors (10⁻⁵-10⁻⁴). With a suitable chassis with a high transformation ability, GFP expression could be observed more rapidly following transformation, which requires a few hours.

[0114] The living DNA sensors have potential for in vitro DNA detection applications. One unique feature of the cell-based DNA sensor is the long homology within the landing pad region of the circuit, distinct from PCR-based methods that use short recognition sequences. Therefore, the target DNA sequence can be specified at the level of genes or pathways (i.e. biosynthetic gene clusters), and the DNA sensor could be used to mine such sequences from metagenomic DNA³⁹. In addition, the access to NGS sequencing or multiplexed qPCR may not be widely available^{36,40}. By contrast, the cell-based DNA detection is relatively simple and cost-effective. The DNA sensors may be suitable for large-scale screening with limited experimental resources. Further, *B. subtilis* sensors could be stored as spores for easy and long-term storage⁴¹. The metrics used in this study (sequence identity similarity and coverage) should be systematically examined using existing sequencing data and experimental characterization to elucidate sequence design rules for homologous recombination. In addition, tools from machine learning could be used to predict the impact of landing pad sequences on the fitness of *B. subtilis* to minimize any negative effects on growth rate⁴².

[0115] One of the most unique aspects of this system is the potential for in situ DNA detection. *B. subtilis* has been shown to colonize or reside temporarily in diverse environments including soil and the mammalian gastrointestinal tract^{43,44}, enabling in situ DNA monitoring. For example, living DNA sensors could be introduced into gastrointestinal tract or plant-associated environment to monitor microbiome dynamics by sensing and recording in real-time⁴⁵. The sensing mechanisms could be coupled to the release of antimicrobials to target specific pathogens⁴⁶. A recent study demonstrated that the naturally competent bacterium *Acinetobacter baylyi* (*A. baylyi*) can be engineered to detect tumor DNA in the mouse colon⁴⁷, demonstrating a potential application of in situ DNA detection. In their study, the native CRISPR system in *A. baylyi* was exploited to detect a single mutation in the KRAS gene in cancer cells. Similar CRISPR systems could be incorporated into our current circuit design in the future to discriminate between single-nucleotide differences. In sum, we believe that engineering DNA-sensing bacteria could open new avenues for both in vitro and in situ applications in the future.

Materials and Methods

Plasmid and Strain Construction

[0116] All DNA sensor strains were derived from *B. subtilis* PY79. Plasmids constructed in this work are listed in Table 1. The pAX01-comK plasmid was purchased from Bacillus Genetic Stock Center (BGSC ID: ECE222) to introduce P_{xyIA}-comK at the lacA locus in *B. subtilis* PY79 by the selection of MLS (1 µg/mL erythromycin from Sigma-Aldrich and 25 µg/mL lincomycin from Thermo

Fisher Scientific) to enhance the transformation efficiency in LB^{17,48}. Genes of fluorescent protein GFP(Sp), mCherry, and mTagBFP were cloned from plasmid pDR111_GFP(Sp)¹⁰ (BGSC ID: ECE278), plasmid mCherry_Bsu¹¹ (BGSC ID: ECE756), and plasmid mTagBFP_Bsu¹¹ (BGSC ID: ECE745) to construct fluorescent reporter plasmids pOSV00170, pOSV00455 and pOSV00456, respectively. The fluorescent reporter was introduced at the ycgO locus by

the selection of 5 µg/mL chloramphenicol (MilliporeSigma). The null DNA detection plasmid pOSV00157 was composed of Repressor lacI and IPTG-inducible toxin-antitoxin system P_{hyperspank}-txpA-ratA and can be introduced at the amyE locus by the selection of 100 µg/mL spectinomycin (Dot Scientific). The toxin-antitoxin system txpA-ratA was PCR amplified from *B. subtilis* 168 gDNA.

TABLE 1

List of plasmids.		
Plasmid	Description	Genotype
pAX01-comK ¹⁷	Xylose-inducible comK	lacA(up), erm, P _{xytA} -comK, xylR, lacA(down)
pOSV00170	GFP reporter	ycgO(up), cat, P _{hyperspank} -gfp, ycgO(down)
pOSV00455	RFP reporter	ycgO(up), cat, P _{hyperspank} -rfp, ycgO(down)
pOSV00456	BFP reporter	ycgO(up), cat, P _{hyperspank} -bfp, ycgO(down)
pOSV00157	Detection plasmid without target sequence	amyE(up), lacI, P _{hyperspank} -txpA-ratA, spec, amyE(down)
pOSV00169	Detection plasmid with 500 bp EC homology	amyE(up), 0.5 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 0.5 kbp EC(down), spec, amyE(down)
pOSV00205	Detection plasmid with 1000 bp EC homology	amyE(up), 1 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 1 kbp EC(down), spec, amyE(down)
pOSV00206	Detection plasmid with 1500 bp EC homology	amyE(up), 1.5 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 1.5 kbp EC(down), spec, amyE(down)
pOSV00207	Detection plasmid with 2000 bp EC homology	amyE(up), 2 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 2 kbp EC(down), spec, amyE(down)
pOSV00208	Detection plasmid with 2500 bp EC homology	amyE(up), 2.5 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 2.5 kbp EC(down), spec, amyE(down)
pOSV00292	Detection plasmid with 2500 bp ST homology	amyE(up), 2.5 kbp ST(up), lacI, P _{hyperspank} -txpA-ratA, 2.5 kbp ST(down), spec, amyE(down)
pOSV00459	Detection plasmid with 2500 bp SA homology	amyE(up), 2.5 kbp SA(up), lacI, P _{hyperspank} -txpA-ratA, 2.5 kbp SA(down), spec, amyE(down)
pOSV00475	Detection plasmid with 2500 bp CD homology	amyE(up), 2.5 kbp CD(up), lacI, P _{hyperspank} -txpA-ratA, 2.5 kbp CD(down), spec, amyE(down)

[0117] *B. subtilis*, *E. coli*, *S. typhimurium*, *S. aureus* and *S. epidermidis* were all cultured at 37° C. in Lennox LB medium (MilliporeSigma). *C. difficile* and gut bacterial strains *A. caccae*, *B. theta*taoiaomicon, *C. asparagiforme*, *C. hiranonis*, and *B. longum* were cultured at 37° C. in YBHI medium in an anaerobic chamber (Coy Laboratory). YBHI medium is Brain-Heart Infusion Medium (Acumedia Lab) supplemented with 0.5% Bacto Yeast Extract (Thermo Fisher Scientific), 1 mg/mL D-Cellobiose (MilliporeSigma), 1 mg/mL D-maltose (MilliporeSigma), and 0.5 mg/mL L-cysteine (MilliporeSigma). The gDNA of each species was extracted using DNeasy Blood & Tissue Kit (Qiagen). For *S. aureus* gDNA extraction, 0.1 mg/mL Lysostaphin (MilliporeSigma) was added in the pre-treatment step in combination with enzymatic lysis buffer (Qiagen). Bacterial strains are listed in Table 2.

TABLE 2

List of bacterial strains.		
Strain	Description	Genotype
msOSV00487	EC-sensor with 500 bp homology	<i>B. subtilis</i> PY79 amyE::0.5 kbp EC(up), lacI, P _{hyperspank} -txpA-ratA, 0.5 kbp EC(down), spec; ycgO::cat, P _{hyperspank} -gfp; lacA::erm, P _{xytA} -comK, xylR

TABLE 2-continued

List of bacterial strains.		
Strain	Description	Genotype
mOSV00580	EC-sensor with 1000 bp homology	<i>B. subtilis</i> PY79 amyE::1 kbp EC(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 1 kbp EC(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
mOSV00581	EC-sensor with 1500 bp homology	<i>B. subtilis</i> PY79 amyE::1.5 kbp EC(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 1.5 kbp EC(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
mOSV00582	EC-sensor with 2000 bp homology	<i>B. subtilis</i> PY79 amyE::2 kbp EC(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2 kbp EC(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV00495	EC-sensor (EC-G sensor) with 2500 bp homology	<i>B. subtilis</i> PY79 amyE::2.5 kbp EC(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp EC(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV00605	ST sensor	<i>B. subtilis</i> PY79 amyE::2.5 kbp ST(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp ST(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV00906	SA sensor	<i>B. subtilis</i> PY79 amyE::2.5 kbp SA(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp SA(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV01005	CD sensor	<i>B. subtilis</i> PY79 amyE::2.5 kbp CD(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp CD(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -gfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV01009	ST-R sensor	<i>B. subtilis</i> PY79 amyE::2.5 kbp ST(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp ST(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -rfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
msOSV01008	SA-B sensor	<i>B. subtilis</i> PY79 amyE::2.5 kbp SA(up), lacI, <i>P_{hyperspank}</i> -txpA-ratA, 2.5 kbp SA(down), spec; ycgO::cat, <i>P_{hyperspank}</i> -bfp; lacA::erm, <i>P_{xyLA}</i> -comK, xylR
usOSV00264	<i>Escherichia coli</i> MG1655	
usOSV00197	<i>Salmonella enterica</i> serovar Typhimurium LT2	
usOSV00113	ATCC 700720 <i>Staphylococcus aureus</i> DSM 2569	
usOSV00095	<i>Clostridium difficile</i> DSM 27147	
usOSV00165	<i>Staphylococcus epidermidis</i> ATCC 14990	
usOSV00046	<i>Clostridium hiranonis</i> DSM 13275	
usOSV00157	<i>Anaerostipes caccae</i> DSMZ 14662	
usOSV00011	<i>Bacteroides thetaiotaomicron</i> ATCC 29148	
usOSV00041	<i>Clostridium asparagiforme</i> DSM 15981	
usOSV00067	<i>Bifidobacterium longum</i> subs. <i>infantis</i> DSM 20088	

[0118] The target sequences xdhABC were PCR amplified from *E. coli* MG1655 gDNA (NCBI Reference Sequence: NC_000913.3; Location: 3001505-3004004 and 3004005-3006504), sipBCDA from *Salmonella enterica* serovar *Typhimurium* LT2 ATCC 700720 (NCBI Reference Sequence: NC_003197.2; Location: 3025979-3028478 and 3028479-3030978), hemEH from *S. aureus* DSM 2569 (GenBank: LHUS02000002.1; Location: 553-2770 and 2864-5638), and pheST from *C. difficile* DSM 27147 (GenBank: FN545816.1; Location: 770923-773144 and 773157-775686) to construct a set of plasmids (pOSV00169, pOSV00205, pOSV00206, pOSV00207, pOS00208, pOSV00292, pOSV00459 and pOSV00475) using restriction enzymes BamHI-HF (New England Biolabs) and EcoRI-HF (New England Biolabs) or Golden Gate Assembly Mix (New England Biolabs). DNA sequences of genetic parts are listed in Table 3.

TABLE 3
Sequences of genetic parts.

Part	Sequence
P _{hyperspank} -txpA-ratA	ctcgagggtaaatgtgagcactcacaattcattttgcaaaagttgttgactttatctacaaggtgtggcataatgtgtgtaattgtgagcggataacaattaaagcttacataaggaggaactactATGTCGACCTATGAA TCTCTAATGGTCATGATCGGCTTTGCCAATTAATAGCGGGATT ATGACATGGGTAATATCTCTTTTAACATTATTATTCATGCTTAGAA AAAAAGACACTCATCCTATTACATTACTGTAAGGAAAAGTGTC TACACGAGGACCCCTCTATTAAAGGGTAGTTCTTTTTTAAAGCT AGAGTGTCTGCCACACTCTGGCTTTTATATTTTAGCATTTCTCATGA AAGTAACACACATTAAACAAGTGGTAATGTGGTAATGTGGTACCAA CTATAAGCTTACGCCAGTAGTTGCAATACTTTTGCTTGGCACCATT ATAACATGAATATATATTGATTATATAAATTATTGTATCTTTTATT TGTTACTTTTTTTATCTATGAGTTCAAAATGACCTGATCATAGAAG CCTTAACCCTTTTTCTTTTATTAACCCCTCGGATTATGAAAGTG TTATGGTACAATATGGTTTAGTATAAATGAATATTGGCTTTCAAC ATCTCAAGGCGGTCTGGCTCACTCCCTCATGAAAGGGGTGATG CACGTGTCAACATTTCAAGCATTAACTGCTTATGCTTGCTTCGGGT CATTTATAATTGCCCTGTTGACTTATATAAAGAAGAAATAGACCC ACCCCTTGAGCTCGGCAAGTAAAGGGTAA (SEQ ID NO: 5)
gfp (Sp)	ATGGTTTCTAAAGGTGAAGAATTGTTTACAGGTGTTGTTCCAATTT TGGTTGAATTGGATGGTGATGTTAATGGTCATAAATTTCTGTTTC TGGTGAAGGTGAAGGTGATGCTACATACGGTAAATTGACATTGAA ATTTATTTGTACAACCTGGTAAATTGCCAGTTCTTGGCCAACTTG GTTACAACATTGCTTATGGTTTGCAATGTTTGCCTCGTTATCCAG ATCAGATGAACAACATGATTCTTTAAATCTGCTATGCCAGAAG GTTATGTTCAAGAACGTACAATCTTTTCAAGGATGATGGTAATT ATAAGACACGTGCTGAGGTTAAGTTTGAAGGTGATACATTGGTTA ATCGTATCGAATTGAAGGATATCGATTTTAAAGAAGATGGTAATA TCTTGGGTCATAAATTGGAATATAATTATAATTCTCATAATGTTTA TATCATGGCTGATAAACAAGAACGGTATTAAGTTAATTTTAA AATTCGTACATAATATTGAAGATGGTTCTGTCAATTGGCTGATCAT TATCAACAAAATACACCAATTGGTGATGGTCCAGTTTGTGTTGCA GATAATCATTATTTGTCTACACAATCTAAATTGTCTAAAGATCCA AATGAAAACGTGATCACATGGTTTGTGGAATTTGTTACAGCT GCTGGTATTACATGGTATGGATGAATTGTATAAATAA (SEQ ID NO: 6)
rmCherry	ATGGTTAGCAAAGGCGAAGAGGATAATATGGCGATCATCAAAGA ATTTATGCGCTTTAAAGTTCATATGGAAGGCAGCGTTAATGGCCA CGAATTTGAAATTGAAGGCGAAGGTGAAGGCAGACCGTATGAAG GCACACAAACAGCAAACTGAAAGTTACAAAAGCGCGACGCTG CCGTTTGCATGGGATATTCTGTCAACGCAATTTATGTATGGCAGC AAAGCATATGTTAAACATCCGGCAGATATCCCGGATTATCTGAAA CTGTCATTTCCGGAAGGCTTTAAATGGGAACGCGTCATGAATTTT GAAGATGGCGGAGTTGTTACAGTCACACAAGATTCACTACTGCAA GATGGCGAATTTATCTATAAAGTCAAACCTGCGTGGCACGAACCTT CCGTCCAGATGGCCCTGTTATGCGAAAAAAACAATGGGCTGGGA AGCATCAAGCGAAAGAATGTATCCGGAAGATGGTGCCTGAAAG GCGAAATTAAACAACGCTGAAACCTAAAGACGGTGGACATTAT GATGCGGAAGTCAAAACAACGTATAAAGCGAAAAAACCTGTTCA ACTGCTGGCGCATATAACGTTAACATTAACTGGATATCACGAG CCATAACGAAGATTATACAATCGTCGAACAGTATGAAAGAGCAG AAGGACGCCATTCAACAGGCGGAATGGATGAACGTGATAAATAC TAG (SEQ ID NO: 7)
mTagBFP	ATGAGCGAACTGATCAAAGAAAACATGCATATGAACTGTACAT GGAAGGCACAGTCGATAACCATCACTTTAAATGCACATCAGAAG GCGAAGGCAAACCGTATGAAGGCACACAAACAATGAGAATCAAA GTTGTTGAAGGCGACCGCTGCCGTTTGCAATTTGATATTCTGGCA ACATCATTTCTGTATGGCAGCAAAACGTTTATCAATCATACACAA GGCATCCCGGATTTTTTAAACAATCATTTCCGGAAGGCTTTACAT GGGAACGCGTTACAACATATGAAGATGGCGGAGTTCTGACAGCA

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	ACACAAGATACATCATTTGCAAGATGGCTGCCTGATCTATAATGTC AAAATTAGAGGCGTCAACTTTTACAAGCAATGGCCCTGTTATGCAG AAAAAAACACTGGGCTGGGAAGCATTACAGAAACACTGTATCC GGCTGATGGCGGACTGGAAGGCAGAAACGATATGGCACTGAAAC TGGTTGGCGGATCACATCTGATTGCAAAACATCAAAACAACGTACC GCTCAAAAAAACCGCAAAAAATCTGAAAATGCCTGGCGTCTATT ATGTCGATTATAGACTGGAACGCATCAAAGAAGCGAACACGAA ACATATGTCGAACAACATGAAGTTGCAGTTGCGAGATATTGCGAT CTGCCGTCAAAACTGGGCCATAAACTGAATTACTAG (SEQ ID NO: 8)
EC(up) 0.5 kbp	Caattaccatcgaatgcaccattaacgggatgccttttcagcttcacgccgcaccaggcacgccgctctcgg aattactccgcgaacaaggactgctaaagtgtcaacaagggtgctgctgggtgaatgtggtgctgtacggt gttggtcgacggcacagcaatagacagttgcttataccttgcgcctgggtgaaggaaaagagatccgcac ctcggaaggtgaagcgaaaggcggaacatttctcatgttcagcaggcttatgcgaaatccggcgacgtgca gtgcggtttttgtacgctggcctgattatggctaccacggcaatgctggcgaaaccacgcgagaagccatt aaccattacggaaattcgtcgcggactggcggaatctttgtcgtgcacgggtatcagatgattgtaata cagttctggattgcgaaaaacgaagtaa (SEQ ID NO: 9)
EC(down) 0.5 kbp	Ttatgtgtttaacaactcatatttcttaactcttgcgatagagcgtagcaatgccgatgccagttcatcagcaactt gcttcttgctgttatgacgtgaaagcgctcgcggatcatttgcctttccatctcctccagcgccgtgcccgcg catcatcgagtgacaggtgcgcctcactgacctctgttacctactttgctccgttggtgacattatcagcagattt ggcggcaatagcgtgctgtcgataaactcacctgaaggaaaccagtttaaccagatattccatcaaatgtcttaa ctcgcgcaggtttccggggccaacgatgcttacgcaatatttcgacgacatcgggagcaatgccaggataaac cgatcccagacgacgggtatgcagatgtaaaaagtaaatgcaccaatagttcaatatcttctgacgttcacgca gcggtggcagagttatcgggataaacattaagtcggtagaagagatctt (SEQ ID NO: 10)
EC(up) 1 kbp	Atgacgcgaaactggagatccactccccgcggtgttcgtttcgtcccgattaatggctttcacaccgggccc gggcaagtgctctcttgagcatgacgaaatctcgtcgcctttcatttccgccacagccgaaagaacacgcg ggcagcgccgcattttaaatatgccaatgcgcgacgcaatggatattcaacgattgggtgcgcgcgacattgccc gactggataaacggcaatttcagcgaattacgcctggcatttggtgttgccgcgccaacgccgatcgcgtgcca acatgccgaacagactgcacaaaatgcgccattaaacctgcaaacgctgggaagctatcagcgaatctgtcct gcaagatgcgccccgcgttctctcatggcgggcccagtaaaagagtttcgtctgcacatctcatccagacgatgacc aaaaaagtgattagcgaagccgtgcgcgcgggggggaaaaattgcaatgaatcacagcgaaacaattac catcgaatgcaccattaacgggatgccttttcagcttcacgccgcaccaggcacgccgctctcgggaattactc cgcaacaaggactgctaagtgtaacaagggtgctgcgtgggtgaatgtggtgctgtacggtgttggtc gacggcacagcaatagacagttgcttataccttgcgcctgggctgaaggaaaagagatccgcacgctgga aggtaagcgaaaggcggaacatttctcatgttcagcaggcttatgcgaaatccggcgacgtgcagtgcg ggttttgtacgcctggcctgatttggtaccacggcaatgctggcgaaaccacgcgagaagccataaccat tacggaatttcgtcgcggactggcggaatctttgtcgtgcacggggtatcagatgattgtaataacagttc tggtattgcgagaaaacgaagtaaaaggatccggcctgaattcaggccggattcactg (SEQ ID NO: 11)
EC(down) 1 kbp	Aggttatgtgtttaacaactcatatttcttaactcttgcgatagagcgtagcaatgccgatgccagttcatcagca acttgcttcttgctgttatgacgtgaaagcgctcgcggatcatttgcctttccatctcctccagcgccgtgcccgc ccgcatcatcgagtgacaggtgcgcctcactgacctctgttacctactttgctccgttggtgacattatcagca gatttgccggcaatagcgtgctgtcgataaactcacctgaaggaaaccagtttaaccagatattccatcaaatg cttaactcgcgcaggtttccggggccaacgatgcttacgcaatatttcgacgacatcgggagcaatgccaggat aaaccgatcccagacgacgggtatgcagatgtaaaaagtaaatgcaccaatagttcaatatcttctgacgttca cgcagcggtggcagagttatcgggataacattaagtgcgtagaagagatcttcgcggaatttacctcggcaa tgaaactggggccaaattctgattagttgcagaaatgatgcgaatgtcgacttgattgggtgactggcaccatcg gcagaatttcacgtgcctcaatagcgcgcagtaatttagcctgcaacatataatggcatatcacctatttcatcga gaaacagcgtgcccgtatttcgcgcctgaatcaacctgtttacccgtggcagaagcgccagtaaatgcacc tttaacataaacgaaacagttcgtctccagaagctgctccggaatcgcggcacagttgatagcaataaagggt ttattccgtcttccgctcaacttatggattgcacggcgacgaacttcttaccgcgtgcgcgtttcaccaaccacca taacgctggatgggtgggtgcaatacggctaatgagtcgttttaattgcccgcataaacacggcactcgccaac caattgttcaatatgcggttcatcaggtgcattt (SEQ ID NO: 12)
EC(up) 1.5 kbp	cgtaacgcggtgaagatggctaccgggtgtgcaatcaatacactgccgctgacgccccaaacgggttatatgaa gagttccatctggcaggattgatttgaggataaacatcatgtttgatttgccttctaccatcgcgacgcaacccctt gccgatgccatcaacctgctggtgacaacccgcaggccaaactgctcgcgggtggcactgacgtactgattc agctccaccatcacaaatgaccttatcgccatattgttgatattcataatctggcggagctgcgggggaattacgc tgccggaagatggctcgtacgtatgcgtctgcaacgacatttaccagctaataagaagatccataaccata cgtcatctcccggcgttatgtgctgcgcgccagctccattgctggaccgcagatccgtaacgtcgtcactacg gtggaatatattgcaacgggtgccaccagcgcagattctgccacgccaacgctaatttatgacgcgaaactgga gatccactcccgcgcgggtgttgcgtctcccgatataatggctttcacaccggcgccgggcaaatgtctcttg agcatgacgaaatcctcgtcgccttcattttccgccacagccgaaagaacacgcggcgacgcgcatttttaa atatgccatgcgcgacgcaatggatatttcaacgattggctgcgcgcgacatttgcgactggataacggcaat ttcagcgaattacgcctggcatttggtgtgcgcgcgcaacgcgattcgttgccaaatgcccgaacagactg cacaaaatgcgccaattaaactgcgaacgctggaagctatcagcgaatctgtcctgcaagatgtcgcgccgc gttcttcatggcggggcagtaaaagagtttcgtctgcacatctcatccagacgatgacaaaaaagtgattagcgga ccgtgcgcgggggggggaaaaattgcaatgaatcacagcgaaacaattaccatgaatgcaccataaac gggatgccttttcagcttcacgccgcaccaggcacgccgctctcgggaattactccgcgaacaggactgcta

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	agtgtcaaacagggtgctgctgggtgaatgtggtgctgtacgggttgggtcgacggcacagcaatagac agtgtcttataccttgccgctgggtgtaaggaaagagatccgcacgctggaagggtgaagcgaaaggcgg aaaactttctcatgttcagcaggcttatgcgaaatccggcgcagtgacgtgctgggttttgtagcctggcctga ttatggctaccacggcaatgctggcgaaccacgcgagaagccattaacattacggaattcgtcggcgac tggcgggaaatctttgctgctgcacgggtatcagatgattgtaaatacagttctggattgcgagaaaacgaa gtaaaaggatatccggcctgaattcaggcggattcactg (SEQ ID NO: 13)
EC(down) 1.5 kbp	aggttatgtgtttaaacaactcatatttcttaattctgcgatagagcgtagcaatgccgatgccagttcatcagca acttgcttcttgctgttatgacgtgaaagcgctcgcggatcatttgcttttccatctcctccagcgcgtgcccgc ccgcatcatcgagtgcaggtgcgcctcactgacctctgttacatcactttgctccgttgtgccattattcagca gatttggcggcaatagcgtgctgcgataaacttcacctgaaggaaaccacgttaaccagatatccatcaaattg cttaactcgcgcaggtttccgggccaacgatgcttacgcaaatatttcgacgacatcgggagcaatgccaggat aaaccgatcccagacgacgggtatgcagatgtaaaaagtaatgcaccaatagttcaatatcttctgacgttca cgcagcgtggcagagttatcgggataacattaaagtgcgtagaagagatcttcgcggaatttacctcggcga tgaactggggccaaattctgattagttgcagaaatgatgcgaatgcgacttgatattgggtactggcaccatcg gcagaatttcacgtgcctcaatagcgcgcagtaatttagcctgcaacattaatggcatatcacctatttcacga gaaacagcgtgcccgtatttcgcgcctgaatcaacctgtttaccgttggcagaagcgcagtaaatgcacc tttacataaacggaacagttcgtctccagaagctgctccggaatcgcggcacagtttagtagcaataaagggt ttattccgtcttcgcctcaacttatggattgcacggggcagcagacttcttaccctgctgcgctttcaccaaccacca taacgctggatgggtgggtgcaatacggctaatgagtcgttttaattgccgcataaacacggcactcgccaac caattgttcaatatgcgggtcatcaggtgcatttgctacagaaaaactggatgcgatttggtgaaacgccattaa aaataattgtcggcctgaatttatgcaattgaccaatgatttaattcacttttatcgtcccatgaaacaatatgct gcatatgtccatgggtaaaattactctcaaatgttaattggctgaaacgggatagggtttcccaataatattatttgc caaacaccaaggttttttaaggcagtcgtatttaacaaactgaaccgatatttcacatctacaactataacgcccgt atccatattatcgatcatggtcgcaaatattttactgatgttatctcctggcccctgatctccagaagtttcgaaa caaaaaatggtggatataatggcgcaacataatcagaaaaatcgcgtaaatatcactgatatgctcttgttgcctg gtaacggcaatcaacttatcaccaccaacacacagatcctgtaaaatgacaggcgtaccagaaatgcttttt c (SEQ ID NO: 14)
EC(up) 2 kbp	ccttgataaaaaggccatatcgtgctgggtgaacgacccggaagagcgttaatgtcgttaaaagatttggcgat ggacgctttctaccaccctgaacgcggcgggcagctctctgctgaaagctccatcaaaaccaccactaaccc accggcggttggctgtacctttgtgactcgcaggtcgatattgctgctgtgcaaaagtaccatcaaccgcatcct caacgctcatgattcagggcatattcttaactcactgctggcagaaggtcaggtacacggcggaatgggaatg ggcatggctggcgctatttgaagagatgatcagatgctaaaaagcggcgtgggtccgtaaccaccaatctgc tggattacaaaaatgcgacacatgcgggatctgccacaactggaaagcgcgttcgtcgaaatcaatgagccgc aatccgcatacggacataagtcaactgggtgagccaccaataattcctgttgcgctgctattcgttaacggcgtg aagatggctaccggtgttgcaatcaatacactgccgctgacgccccaaacggttatatgaagagttccatctgg caggattgatttgaggataaacatcatgtttgattttgcttcttaccatcgcgcagcaacccttgcgatgccatca acctgctggctgacaaccgcgagggccaaactgctcgcgggtggcactgacgtactgattcagctccaccatc acaatgaccggttatcgccatatttggtgatattcacaatctggcgagctgcggggaattacgctggcggaagat ggctcgctacgtatcggtctgcacacgacatttaccagctaatagaagatcctataactcaacgtcatctccc ggcttatgtgctgcggccacgtccattgctggacgcagatcgtaacgtcgctacctacggtggaaatattt gcaacgggtgccaccagcgagatttctgccacgccaacgctaatttatgacgcaaaactggagatccactccc cgcggggtgttcgttctgcctcgatttaattggctttcacacggggcgggcaaaagtgtctcttgagcatgacgaa atcctcgtcgcctttctatttcgcgcacagcgaaagaacacgcggcgagcgcgcattttaaatatgcctgctg cgacgcaatggatatttcaacgattggctgcgcgcacatttgcgactggataacggcaatttcacgcaattta cgctggcatttggtgttgccgcgcgaacgcgcgatcgcgtgccaaactgcgaacagactgcacaaaaatgccc ccatataaacctgcaaacgctggaagctatcagcgaatctgtcctgcaagatgtcgccccgcgtttctcatggc gggcagtaaaagagtttcgtctgcactctcatccagacgatgacaaaaaagtgtatagcgaagcgtcgcgc cggcgggggggaattgcaatgaatcacagcgaaacaattaccatcgatgacccattaacgggattgcctttt cagcttcacgcgcgaccaggcagcgcgtctcggaattactcgcgcaacaaggactgctaaagtgtcaaaaca aggggtgctgcgtgggtgaatgtggtgctgtacgggttgggtcgacggcacagcaatagacagttgcttatac cttgcgcctgggtgaaagaaaagagatccgcacgctggaagggtgaagcgaaaggcggaataatttctc atggtcagcaggcttatgcgaaatccggcgagtgacgtgctgggttttgtagcgcctggcctgattatgggtacc acggcaatgctggcgaaccacgcgagaagccattaacattacggaattcgtcgcggactggcgggaa atctttgtcgtgcacgggtatcagatgattgtaaatacagttctggattgcgagaaaaacgaagtaaaaggat atccggcctgaattcaggccggattcactg (SEQ ID NO: 15)
EC(down) 2 kbp	aggttatgtgtttaaacaactcatatttcttaattctgcgatagagcgtagcaatgccgatgccagttcatcagca acttgcttcttgctgttatgacgtgaaagcgctcgcggatcatttgcttttccatctcctccagcgcgtgcccgc ccgcatcatcgagtgcaggtgcgcctcactgacctctgttacatcactttgctccgttgtgccattattcagca gatttggcggcaatagcgtgctgcgataaacttcacctgaaggaaaccacgttaaccagatatccatcaaattg cttaactcgcgcaggtttccgggccaacgatgcttacgcaaatatttcgacgacatcgggagcaatgccaggat aaaccgatcccagacgacgggtatgcagatgtaaaaagtaatgcaccaatagttcaatatcttctgacgttca cgcagcgtggcagagttatcgggataacattaaagtgcgtagaagagatcttcgcggaatttacctcggcga tgaactggggccaaattctgattagttgcagaaatgatgcgaatgcgacttgatattgggtactggcaccatcg gcagaatttcacgtgcctcaatagcgcgcagtaatttagcctgcaacattaatggcatatcacctatttcacga gaaacagcgtgcccgtatttcgcgcctgaatcaacctgtttaccgttggcagaagcgcagtaaatgcacc tttacataaacggaacagttcgtctccagaagctgctccggaatcgcggcacagtttagtagcaataaagggt ttattccgtcttcgcctcaacttatggattgcacggggcagcagacttcttaccctgctgcgctttcaccaaccacca taacgctggatgggtgggtgcaatacggctaatgagtcgttttaattgccgcataaacacggcactcgccaac caattgttcaatatgcgggtcatcaggtgcatttgctacagaaaaactggatgcgatttggtgaaacgccattaa aaataattgtcggcctgaatttatgcaattgaccaatgatttaattcacttttatcgtcccatgaaacaatatgct

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	gcatatgtccatgggtaaaattactctcaaatgttaatggctgtaaacggataggtttcccaataatattattttgc acaacaccaagtgttttaaggcagctctgatttaacaaactgaaccgcattttcatcatctacaactaatacgcccgtg atccatattatcgatcatggctcgcaaatattttactgatgttatctcctggccctgatcctccagaagtttcgaaa caaaaatggtagatataatggcgacataatcagaaaatcgcgtaaattatcactgatatgctctgttgcctgtg ggtaacggcaatcaaacttatcccccaacacacagatcctgtaaaatgacaggcgtaccagaaaatgcttttt cgcgccaattttcttactatcgcaaccttcgcaaaaggggatcgaagcgagactgtgtcacaactttttcagtttt cgtttccaggacgtggcgagcaggcgtgagttgcgcgtcaactggcgaccaagaacttcccatacgcgc ccgttcgggcaacgcgcacacagttttcatcaacgatctcaacctcaagctgcaaaacgctggcagcattct ggcaaaacgctgaattgtcggttgaatttgcataactgactgcgtagtagcaagctccatagctttaccttc cagacttacttaaaagtcgatcattgaagacgttgatgggttcacagatcatgatgataaactcaggcgaaatt ggctttgataaaaacataagatttttatcattttctaatgaaattatggaagagatatcacatttctatatcaaat gagaattacggcggtagtttatcaaacgaagagagatagcctgcccccttat (SEQ ID NO: 16)
EC(up) 2.5 kbp	aggagatgctaataccgctcacgggcaaacgtatttacagcgcagggttgcgggagtgctttgaaaaaggcgcg gaaaatctttgaaatgggaaaaacgcgcgtgcagaatgccagaaccagcaaggcaatttgcgcgcggcgttg gcgtcgcctgttttagctacacctctaaccacctggcctgtcggcgtagaaaatagcaggcgcgcgcctcttgat gaatcaggatggaaacctcaacgtgcaaaagcggcgacggaatcggtcagggtgcccagaccgtcttct cgcaaatggtagcagaacacgtgggggttcgggtcagcgacgttcgcgttatttcaactcaagatcaggcgcg ttacgcgcgttcgatcccggcgcatttgcctcacgcgcagagctatgtgcgcgcctgcgtgcgcagtcgcgc cactatttataaaagagaaaatcatcgctcacgcgcagctcatgctacatcagtcagcgatgaattctgacctg ataaaaggccatcatcgctcgtggttgaacgacgcggaagagccgttaattgctcgttaaaagatttggcgatggacg ctttctaccacctgaacgcggggcgagctctctgctgaaagctccatcaaaaccacactaacccacgcg cgtttggctgtacctttgttgatcagcggctcgatattgcgcgtgtgcaaaagtccacatcaaccgcatctcaacgt tcatgattcagggcataattcttaactccactgctggcagaaggtcaggtagacgcgcggaatgggaatggcatt ggctgggcgctatttgaagagatgatcatcgatgctaaaagcggcgtgggtccgtaaccccaatctgctggatt acaaaaatgccgacctgcccgtatctgccacaactggaaagcgcgttcgtcgaaatcaatgagccgcaatcc gcatacggacataagtcactgggtgagccaccaataatctcgttgcgcgtgctattcgtaacgcggtggaagat ggctaccggtgttgcaatcaatacactgcgcgtgacgcgcaaaacggttatatgaagagttccatctggcgagga ttgattttaggataaacatcatgtttgattttgcttctaccatcgcgcagcaaaccttgcgcgatgccatcaacctgc tggctgacaacccgcagcccaaacgtctgcgcgtggcactgacgtactgattcagctccacatcagcgttcg accgttatcgccatattgttgatattcataatctggcgagctgcgggggaattacgctggcggaagatggctcg ctacgtatcggtctgcaacgacatttaccagctaatagaagatcctataactcaacgtcatctccggcggtta tgtctgcgcgcacgtccattgtctgacgcgcagatccgtaacgtgcgtacctacggtggaaatatttgcaacg gtgccaccagcgcagattctgcccacgcaacgctaatttatgacgcgaaactggagatccactccccgcgcg gtgttcgtttcgtcccgattaatggctttcacacccgggcccgggcaagtgctcttgagcatgacgaatcctcg tcgccccttcattttcgcccacagcgaagaacacggggcagcgcgcattttaaatatgcccacgctggcagc aatggatatttcaacgattggctgcgcgcgcacattgcccactggataacggcaattttagcgaattacgcctgg catttgggtgttgcgcgcgcaacgcgcgattcgtctgccaacatgcccgaacagactgcacaaaatgcccattaa acgtgcacacgcgtggaagctatcagcgaatctgctcctgcaagatgtcggccgcggtttctcatggcgggcca gtaaagagtttctgctgcatctcatccagacgatgaccaaaaagtgttagcgaagcgtgcgcgcggcg ggggaaaaatgcaatgaatcacagcgaacaatatacctcgaatgcaccatataccgggatgacctttcagcttc acgcgcgacacgggcagcgcgctctcggaattactcgcggaacaaggactgctaagtgtcaacaagggtgc tgcgtgggtgaattgtgtgcctgtacgggtgttgggtcgacggcacagcaatagacagttgcttataccttgcgc ctgggctgaaggaaaagagatccgcacgtcgaaggtgaagcgaaggcggaaaaactttctcatgttcagc aggcttatcgcaaatccgcgcgcagctgcagtgccgggtttgtacgcctggcctgattatggctcaacgaat gctggcgaaacacgcgcgagaagccattaaccttacggaaattcgtcgcgcgactgggggaaattcttctgc ctgcacggggatcatgagatgattgtaaatcagttctggattgcgcgaaaaaggaatgaaaggatataccggcct gaattcaggccggattcactg (SEQ ID NO: 17)
EC(down) 2.5 kbp	aggttatgtgtttaaacaactcatatttcttaactcttgcgatagagcgtagcaatgccgatgccaggtcatcagca acttgcttcttgcgtgttatgacgtgaaaagcgcctcgcggatcatttgcttttccatctcctccagcgcgcgcgc ccgcatcatcgagtgaacggtgcgcctcactgacctctgttacatcactttgctccgttggccattattcagca gatttggcggaatagcgtgctgcgataacttcacctgaagggaaccagcttaaccagatattccatcaaatg cttaactcgcgcaggtttccgggccaacgatgcttacgcaatatttcgacgacatcgggagcaatgccaggat aaaccgatcccagacgacgggtatgcagatgtaaaaagttaatgcaccaatagttcaatatctcctgacgttca cgcagcgtggcagagttatcgggataacattaaagtcgtagaagagatcttcgcggaatttacctcgggcaa tgaactggggccaaattctgatttgcagaaatgatgcgaatgtcgacttgatttgggtactggcaccacatcg gcagaatttcaacgtgcctcaatagcgcgcagtaatttagcctgcaacatataatggcatataacctatttcatcga gaaacagcgtgcccgtatttcgcgcgcgtgaatcaacctgtttaccgttggcagaagcgcagtaaatgcacc tttaacataaccgaacagttcgtctcgcagaagctgctccggaatcgcggcacagttgtagcaataaaggtt ttattccgtcttccgctcaacttatggattgcacgggcgacgaacttcttaccgctgcccgtttcaccaaccacca taacgctggatgggctgggtgcaatacggctaatgagtcgttttaattgcccataaacacggcactcgccaac cactgttcaaatgcccgttcatcaggtgcatttgcacagaaaaactggtagcgattgggtggaacgaatctaa aaataattgtcggccctgaatttatgcaattgaccaatgatataacttcttctcgtcccatgaaacaatagct gcatatgtccatgggtaaaattactctcaaatgttaatggctgtaaacggataggtttcccaataatattatttgc acaacccaagtgttttaaggcagctctgatttaacaaactgaaccgcattttcatcatctacaactaatcagccctg atccatattatcgatcatggctcgcaaatattttactgatgttatctcctggccctgatcctccagaagtttcgaaa caaaaatggtagatataatggcgacataatcagaaaatcgcgtaaattatcactgatatgctctgttgcctgtg ggtaacggcaatcaaaacttatcccccaacacacagatcctgtaaaatgacaggcgtaccagaaatgctttt cgcgccaattttcttactatcgcaaccttcgcaaaaggggatcgaagcgagactgtgtcacaactttttcagtttt cgtttccaggacgtggcgagcaggcgtgagttgcgcgtcaactggcgaccaagaacttcccatacgcgc ccgttcgggcaacgcgcacacagttttcatcaacgatctcaacctcaagctgcaaaacgctggcagcattct ggcaaaacgctgaatttgcgttgaatttgcataactgactgcgtagtagcaagctccatagctttaccttc

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	cagacttactttaaagtcgatcattgaagacgttgatgggttcacagatcatgatgataattaactcaggcgaatt ggcttttgataaaaaacataagattttttatcatttttctaataaattatggaagagatatacattttctatatcaaat gagaattacggcgggtgagtttatcaaacgaagagagatagcctgcccctttatcttattttctgatacttagcagca aataaataacgcgataaaaaaagccaaacgttttcgtattttacaacaaccagaagctggcatcaatttgtgatc aaccacacacattatccgtcaaaattagtccttttgcagcgcgcggataaattctggcacacttatttgttagtccag gtatagctgtgaaaacaccaatcactttggcaagtacagtgaaaataaaccactttgctgtcattccactaccg ggactttatgatgaaaactgttaatgagctgatgaaggatatacaattcgtgacctctcaccttcacgagaagat ttttgttaacgtgggaacagacgcagatgaactgaaacagtagtgacgttgcgcgagcattaaaagcac tgctgtgctgaaaacatctcaacaaagcttttaatagtggattaggattttccgtattccgcgacaactccacc gtaccgcctctcttatgcttccgcg (SEQ ID NO: 18)
ST(up) 2.5 kbp	acgtagcagcaggggtatcaacgtttgcatttcaaggtgccgggttcccgctcctacgtggtagccctgctctt gcgttaatttttgggtggcacatatacagcgcctcaacagccttcgcgcgcgtttgtcaacaaggtgcgtaaga ttgctgcgggttaacggatctaacgtacagcgaaggtatgttcaatgacgtggcaatatagggtacacctc ctgcataacaagattcgtcgataatttacttaattcacgcgcagtggtatttttgataaatatctaacagctgcttt ccaggttttccagcttcgccttcgcgtttctttgtttctggcagccatggcccaaaagctgactttcttctcaggcca tcttttatgatttgcgcggtatcctcgtcccccactctcatcagtagcgtcttcgcctcaggagaatcactgggtggc gttagcgcgtgaacgaaagagcccgcaaacctcattatcgctttcttaccggcgacatattttgaattggtaaaaa cttcttttaacgcctcagcgttttccgcctttaaacaatgcataccagactcgcctgtttgatcagcgcgggaaa atcttccagttgcgggcttttaatttccctgacagcgtcgtgtggcactttctcgtgactgcggaaaagattcgcg gcaagattcgtggcctgcgttttgatctcgttctgcatacctggcattatgacggggggtgagtccttacactt gtaaccattattaataatcctcttctgttatcttgcaggaagcttttggcggtttccaggctgctacttatcgtact gctcagcacttttaccaggttgctgtacaatgaattggcatttgcataatttttgcgtcagcgtctgtaattgtggtt tcataattttctcctgcgcttttaaacccgactgccaggttgatatttggcggttatccatttcagagttttgagct tttcccgcgcgcttaaacatcaatatcctgaaccattttttgtaattggcgtcagatcaacgggtgacgacataacc ggatccataagatttccaggcagctattcggttaattcaatcactgagccactgtctcgttccgccttcagtggtta cttaacgcgcgtcctgcgtgagctggaaataaaacgggtattactgtttatttgattatattttgactaaactg tttaaatcatttttgagttaggttaacatctagcttaacgggtattaccgtccttacctggtaataaacagcctcccat tttgaaaagaatatcactgaaggcctgataaaaatcggtatagactgcgacaacgttttataaacgccagatagc tgtcactatcgcgcatataatttgggaaaccataatcccaaatccagcatcagaattgggttgcctcgcgcgcg cataggcgaagcgttaataaaggcgcagctcggcgagaaaacgcgctccgcaggttctcattttgttctgc ggataatgacacgcgcgacttcgcgcagcatttcaggtgctggtcaactgctggcgcgccagcgtgcgct cgtcattatctctcagagatcgggtggcgttgactgcagcgtctcgtgtgcgtgggtggatttttagtagccgctg cgataatgaaatgatatactgtaccgcgatttctgtggttagacggtaccacggcagctctcgacgtgctcgtcgc ccgagggagttcgcggcggttcggcaacgatcccggttagaggagaagcggaaataattttgaatatgaagca taatatcccgagttcgcctcaggagcgcgattaaatcacaccatgatggcgatagatgacattcagattaa gcgcgaatattgcctgcgatagcagcagatgcggatgctttcagctggttaatgctctccattgttttcagcattt cctgaatcaggtggtcgatttacctgaactttccagggcttcgtccgattgcggtgctggcaaccgggttattc acctggctaaatttgctgctcggaacgttctcgtgagtagcggcgtactgcgcgagcgccttgcaataaccacgc accgtgaccgagttctcataatcagatcgcccgctcatctgcacttgcgcgcatcgattccgggtcatatccatg gtattctgctcaagcgaatatcggtatcgacagactcaagacgtttcgacagaatagcctgatgttcagggga gatttgtttattactgtcttttaataccagactttccgtggcgctgggttcgggcattagatttaagcgtcgcatcat taagatttttgcgcgcatcggtaccggttttctcatatttaacgatttcagagaatcgacgccttcagcaccaggt ttgacgctattctgcgcgttcagcaggttttaataactgtggctttcagtggtcagtttatcgatcttcgcggcatt atgtttaag (SEQ ID NO: 19)
ST(down) 2.5 kbp	cgcgcctctttcattctgcagcccttatattccagtttggcgcccacgcagtgatcccccaactgaagcgcgc ctcgggaaataactacgggacaacgcattcatccttcgcgcgcatcatggagcttgcgcgtcttttagcgcatacaa aactgactaatgacaaacttaccagacagtttgctatcagcctgggttcaacgtcagcattaaacgtattcgcggca gccaacagcgcgaacggcactggaagacattccgctaatataaaaaactttccgactctcgtcgtcgtcgc gttaactgggtttgcacaacctcattcgtcttagtcgtgacattatttgcagagcatcacaactcgtgattcgtgcg gtattttgaatactggcttttaaaaaggacgtgatcgttccggggggttgcggttaataaccctggcgagggcgcg ctcagtgtaggactcaaccccaggtcactgactttactgctgctaataccaatactattcagaatatctttagcgc taacgggattgcgaagctgtctgtgaactattctcaacagaatgattatttaataaagcggcggggtttattccac attactaattaacataattttctcctcttattttggcagtttttatgcgcgactctggcgagaaataaaacgcgaag catccgcattttgcgtgacgcgagaagacatggctttttgcagttccgcggttaacctctgggttttaccacaaat tctacggattgttttaagccactctgaatctgatccatggcaaaaacggcgagcataaaaatcgacagcgcctcgt ctggcatttttaataaatcgcgcctcggcaacaccacgggtgactggggtcgggtattcgtgacttccatgccc caacgccactttatttagggtattacctaccagctcttacttaaggcattcgtttgcaggcccatcttgctaccca cataccagacgcgtagtaataacgtttgcatccctgggttaagaggtttgctgcggttttgccgcaactgtttc gcacgttaggcaccaactcttaacgtttcgcgccatcatttgctcagcgcgttaccaggtttcgcgcgcgcgc ctttcccgacaactcgcgaccaccacaatgacgcgcaccatggcaatagcggcgacaatcgcaccaacaatg catcgggccaatctcgcgcttttcttatcgacgcctaactcctccagcgttttggttaacgccttgcccaatcagct ccattaacgggttcagcagcatgctccataatcggttttagcgcctgctgaataaacgacactcccgctcgcgcg cttcacaatttcatcggccaccattaccgcaagtcaccacgcagccagcgcagactcgccccaccgggttaa aacagcggccacaacgctgacaatggttagcagcgcgcgaggaacttcccgataacatcccaatcggtt cgtttcctcggctttgcgcgtctcttctggaattcagccgatttctttccatctcgcctgacgcgccttctgca aggcgttgaagaagcgaagatcgttttgcaggctttctccgatttttgcacaaatctcaataaacatggccatg agcatagtgaggcggcgacatttgacagattatcctgctcaccctgggaacactgattctgagaggcgga ttagccgttccctggaatttggtcagaatgttatcgcgtttctcggctttcgccttgcgctctgtgctgctttaac cgtcgcactcgtggccttatctaaggcctcttccgcctctgctcgtcttttccggcctgttctacgcggcgttcag cttgctgcatagcggggtcagcgggtccagcgattgcaatttatttgcgcgtcgcgcagtttttgggtcgagc gtcataaacactcttggcgggtatccgcttttttgatactggcttcatagagatccgtcgcctcctgagcctctccc

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	agagccgtctggaattctttcgatacctgaatccccatctctttttgtgactcaatcatcgctgccataccgcca gacgagactccagttgagacagcgaaacatcgccagtagggcttaacttgccaagcagtaattgtcaattg ccttccgtggagagtttttccgggcggttcgttaggcggttttagaccacccgtatataagcgcctcgc cggactttgttcgggctttaaggctgcggcgtttcggtgccaccacatctttaaagctttatccgcgcttttaa aagtcggtgttcttcaaacgccttcaaacgcgctcagcgaggcgcggttttgggtatatacgcgtacggc taatgctacttgcgtcatttaccataattattcttcttcttgcactgtgctgctcctccgcgcttttagcg cctccagatagaccaacgctttg (SEQ ID NO: 20)
SA(up) 2.5 kbp	ctatcgtgcatgtaattcttgcatcgcattgtgcaacgctgtaaatgtttcgaagccatcctcttctaagaagtgc ctccatcttccacgatttcgaagttccctcttaattgcattcattaaacgctgggtttctttatagaaacgtatttg tcaattttagaactcaatccgtaaaaattgtcaactttcttttaattattatcgtaacatggttacattacttaa atcaatatctaaatctatatcttctgcatcttctttaaagccgctatacaaaaaacgcttcaatcggtgatcaa tcaattcaatatattttaagctgtgattgaacctaaacctatgtgtacaaaatattgatcctttttgcgtacatta atttgtttcgcatagcttcaatccactgatccactgtcttgcgttcaggggattcaaaatataaatggttacgtc atatccttctaagtttaagttatgctccaacctgataccaatgatttctactatttccatgcatagaattgtacaa taattacatctgtcatctcattctctcctttcaacttactacttctttctatttttaaaaaaatgactgattacct ataattgtaaaaataaaacaccttaattagaattgttatatcgcaagtgacatttctaattaaagtgtattgtcat caattcaatattcattcaaaaacagctaaacctttgtctctgcttcaatttcacaaaaataattcccgctgaagtat ctatatttacacattacttccaccattatataaactaaaaatgactatatttcatcaaacattatctaaaggcgctc cacctacaccaacacccatccaacaattacttacaactctgcgattacttcttcagcagcaactttaccttgtgtaa tacaatcaggtagtcacaacgcgacgcgacatggtagctgtggcatacttttcggcaaacgattgacaattgttaattc aggatcacctttaaatgtcatttcaaaagatgcaccggttactctaagtcgtggatattgttgtttaaatgtgctt gaatctgtctaatgttgtgaatatcatttgacttaaatctctacgtacaatcgatactaaattcattatctgtatgat catcaaccacagtatcacctgggttacctacatcgcacgaatcaaaccttaccttctgggtgtgtaaaatggccat tttttcgatgtccaagtacatgcggtaattgtctgtatcactcgttctcgcaatcgcgaagccagttaccatcataagt attttcaatgtcttttcatcaaatgccaatcaaacagttgcaacagtcgtactatccatcggtttaaaagtaattcaa atgtcggatcttgcgcaacgaattcaaaaagacttgatgcgggtgtgtgctactaatcccatcgatcaatctctct tgttgattactgtaaacatttttatattgcttttgagatgtaataatcatccactgacgtattgtagcgtattgt cacacctttattttcacatcttgttctaattgcttcaataaatgagcttaaacatgcttaaatgttgaattgtc cttttgggtgcgcaggatataaattgtctttgtttcagacgcttatttttctcatccttcatacctttatcagacct ccgaatgcctcttcttttctttaaatttaggaaacgtactcatcaacttaatttatcaatatcggtaccataaat acccccattaaaggctcaattaaagttctcaagtaacctcattacctaatcttgcctctgaaaaatgcaccaacagaaa tgtcacctcttgcatttgtataggctttttgattaaatctaatcctgctcttaatttaccagtgggcgatattaat tttgtagtacaacatgggttaatatctgttggaataccataaattgaaccacctggaatcggtataatttattttt cgcaaaaatatatgatgtgcacgtcgtatttgaacaatatcttgttctaatacaatatcttgcgttaattctgtca taatcgtttttctacctaataaagattcaggccctagtccaatcatataaccatctttacgatacgattgaatcttt cccccggacgattcgatgcttcaagatgggtacatcaatattaggatcttgcgtgttttaa
SA(down) 2.5 kbp	gacttgatttcatcaacaattgcacgcgataaataatggatgtgatttcggcatttttggacgatataaattcgcacc aatatcatcgcaaaccaactttacattcataatcatttgtcataaagcactctaaatgctcacatacaaacctactc gcgtatataaaagtttttatactgatgtttttcatataaaatcagtggttaaatcttgcacatctggcccttaaccaagg tgcacctgtattaccttcagattgcacaaccaatcgcgatattgttcaatattagattgttctttaaataaaagcgag atgttctagttcttgggtatggatcatattcttttcgattaaaccttttggcaaacattatgtgccaacacacataat accgtgctttatgttctcttccgggtatttgagctaatgtttcggtgactttattcgccaatattcaataaaatttagg ttgttcataaataatgtttcacatgtgaagtgtgaataccatattttgcagcttcttcatcagcagctttgtcatatgatc ctactgaaaatgaagaataatgtggtgctagtactaccgtgattgcttcagtaataaccatcattgtgcaattgttcaa ccgcatcttgcataaattgggtgaattgttttaactcctaagtatagtttaaatcaacatctgcataatgctttatttaatt gctgaaactagtgcatcagctcgtcatctgttgactgctaatgggtgataaacacacacataaaattcatatctatc tttcaaatcttgaagtcttcttctcagatggacgtttaccatgttcaatatctgtataaatatggctctatgtcactttctt tataaggtgtgcataagccataaactaataacccatttttttagtcattgataataaccttctttaaataaattatctt tcaatgtctcaatgtaataactatgattatctttgtgtatattgtgtgacgaattcgcttactttcaactcgtctctgg ttgcacttctgggaaaacacggtgtcctaataaagatgtgtttaccgttctccataccttgatcataatattggtttc aatctctcttcaatgacattccatgggtgctaataaaatttgatggatctaaattcccttgtaattgttttagtaacgccta attgttgagcctgattaatagacgttctccaattcaggcctaatacatcaatcggttaaatcattccattcatgtatta aatgactggcactacaccgaataaaattacggcacatcatgtttttcttaacctcactgattaatcgaatcata tgtggtttaatgtaacgtctgtaactcctgcacatttaatgcacctaccatgaatcgaaaaattgtaataatcggc acctgcttcgacttgagctgtacatttttaacagatacatcaactaaatgattcattaaagcaaacacggttgctgt catctctatcatcatcgccttttgtaaaaattgtaatttttcgatgggtccgccttcaatcatatagacgctaattgtaaat ggtgccccagtaaatcctattagcggcacatttaactttctctgtttaaagtttaattgtatctaatacatatggta catctcgttcgggtctatttgaaaagttctcaacatcttgattgttttgataggattatgaatcactgaagttttaa taccgatttaatttctacatcgacaccaattggctttaatgggtgcataatatcttgtataaaattgctgcactgt atgataaattcaactggtaaatgtgtacataagcgcacaactccggctgatgtgtaataatcgaatagtgtaatt tttctttcaattttcgatattctgggtgcgaacggccagcttgctgcataaaacaaacagggttgatgtgttcttct caoctttgatcatttttaaaatttgtattgtttttattatgcaccataaaggcctcctaataaaatcattcttattctat attatcatatcgctcattcgttctgattttcaataaaataaattgcataaaactgacatttaattcatagaactatttatt gtaaattttaatttcaaaagtcattatttttgatcattacttcaaatatcgcgaagattcattatagtaatttttaatt caattataatagtggttaattgactagtttatcatcgataataaataaaacataaaggggaccttctcatatgaagaaa ctatatacatcttatggcacttatggatttttcatcaataaaatacaataacccgacccatcaactatccaatttt cagcatcagatacttcagtttattttgaagaaactgatggtagactgttttaaaatcaccttcaatatatgaagttat taagaaaatttggtgaattcagtgacatcatttctattgtgcaatcttcatcttcaacagaagatcatgcatatcaa cttgaaaagaaactgattagtgtagacgataatttcagaaactttgggtgctttaaagctatcgtttgttaagacct gctaaaggtacaacataataaatttattcggatttgctgacgacatgcatacgaagactttaaagcaatctgatg cctttaatgaccatttttcaaaagacgattaaagtatttcttgggttcaagcggacaacattcaagttattttgaaa

TABLE 3-continued

Sequences of genetic parts.	
Part	Sequence
	gatatctataccaataaaaagaatag (SEQ ID NO: 22)
CD(up) 2.5 kbp	atgaagcaatatatagtcattgggtgtgggagatttggaaagttcagttgcgtctactatgcatcttttaggacatc aagtaatggcaatagacaaaatgaagattcagttcaaagtatatctgacaaggtaaccatttcacttatagtg atgttactgatgagcaagcgttaaggtcattaggttaggttaactttgatgtagcagtagttgcaataggttctgat ataagggcatctataatggcgactcttatagccaagaaatgggtgtagagttgataat atgtaaggcgaagg atgaattacaagctaaagtgcctttataaaattggcgagatagagttgtatttccagaaagagatatggggagtaa gagttgcacacaatttagtttcggataatatattagaccataattgaacttgaccagagtagttcaattgttgaatc gtaactccaaatagttgggttgggcaagacacttatagagcttgaat taagagctagatatgagataactgtactt gctataaaaaacaggtaaaaatataaattgttacaccttctccagatgaggaaacttacagcc ggaagtatcctagtt ataactcggtcaaaactactagtataacagcgataacatctggaaataaggggataattagaagaagataatttact atttaatatatatttaattgcaatgaagtaagagtagtatataaattatgaatagttatatgatactatttttattta atcgaaggtagtagtattttttgtaagtagatataagagaagttaaattaaaaagtaaggaggtgtgctaatataaaaa ttataagttattagcatttagataaaaatgattacaaatataaacagtaaggataatgaaaagttaaagtatacaag agcactattaaaaatcaaagaataggaataaagagtcagggttcataatagaaggatacagaatagtaaatgcttg cacttgatgtatggcaaaccttgattatgtattatcaatgaagaatttgaataaagaagaacatgtaaaact attagaagatttggataaaaaaacacaaagatatcaagactactataaaaaactttaaagaattagtgagac agaaaaactcaaggaataataggtgtagtttcatttaagaaaaaaaataaagtgaaggtataataaaaaaaga taaatttgtattgattttatagatagaatacaagaccaggaaatattggggactataataaggactgctgattctgct ggagtagatgccataatagcactaaagggatgtgctgatatatacaacccaaaagtaattaggtctactatggg ttctatttttgatatgaatataattgtagcttccacaagatgaaactgtggacatgcttaaatcattggattttaatatag tttcaagttacttaaatacagaaaaatttttatgacaaaatagattatgggttcaaaagtagcattgggtgtaggaaac gaagcaaatggaataaatgaagaacttgtatcaagctctgatatatttgggttaagataccttatatggttaagcc gagtcgttaaatgctgcatagaagctctgctatactgtagttagtaataaaaaaatacttaatttaattgtattgtaata aatataatgtcatgttataatcttgaattaaatagtagtagtatacaaaatagtcataatataataaaaaaatatta ttaattatataatagtagtatatcatataaaaaatgattgagttgattatgtattagatatataattatataaaaaat ttgagtttaattatgcattagtagtatattatataaaaaaatattgagtttaattatgcattagatatataattatata aataaataaataaattatgtatggatatataataaaaaatagtcacataatttgaatggaaatgatataactactaaat aaacaataataatattgtaaatgcaatgaagaggaaagtagttttgattaaacttagtaagagataaacacctaggct gggagtggttttcaagaggtcatgagaagttccctctggagtaacagagctgaaattttacagtaggctttgacg tcaaaaaacgcgttaagtttggtagaggtgggtttagtagatttttaattgtttaactactagggtggtagccgcgaac tata (SEQ ID NO: 23)
CD(down) 2.5 kbp	tagacagggtatgaggggcttttttatcaaaaaacgaaagggtagatgtgtgcaagaaaaatacttgctt tacgtgaagcagctttggctgaaataaaagaagcacaagcatagaaaggttagaaagtttaaggttaagta cttaggaaaaaaaggtgagataaactgccatacttaagaaatgggttaattatctgctgaagaaagaccagta gttggttaaggttgccaatgaggttaagagaaaaacatgaacttagcataaattctaaaaaagaagaaataaatgc tatgaaaaagaagaaaaatataaagagggaagttagatagttactcaaccaggaaaggttttaaggttggg aagaagcatccaataactcaattatagatgaagtaacagatatattatcggaatgggattctctatagcagaa ggggcagaagttgagactgttgaaaacaaactttgacgcattaaacgctcctaaagaccatccatcaagagata tgagcgatacattctatataatgatgggttatacttagaactcaaacatctccagttcaagtagaactatgag aagtcgaaggttaccaataaaaagtaattgaccagggtagatgttttaggtcagactcgcagatgctacacact accaatgttccatcagatagaagggcttgggttgggaaagatgttactatggcagaatttaagggaactatgg atatcttctgtgaaaaattgttgggtctgatatacaaaactaaagtttagacctcacaacttcccatttacagaacca gtgcagaggttagatgttactgttttcaaatgtgggtggttaaggttgcccaatgtgtaaatatgaaggttgtaga aatattaggttcaggtatgggtcatccaatgtgcttagaaattgtggaatagaccagaagttttagctggattt gcatttggagttgggggttgaaagacttgcaatgcttaaatacgaaatagatgatattagattattatcgaaaatg atatgagattcttaaatcaattttaattaggaggttagtagatgttagtatctttaaattggcttagagactatgttg atatagacatggatgtaaaagagttcgctgataaaatgacaatgacaggaactaaagttgaaacaatagattat tatgggtgaagaaatagaaaaatataatgggttggaagattttagaataaaaacaacatccaatgctgataaggtg gttgtaactaaagtagatatggagataaagttgttcaaatagttacaggagctacaaatatatcagaaggagat tatattccagtagctgtaaatgggtctaaagtacctggaggagttgaaatcaaacagactgatttcagaggtgaat tatcagatggtagatgtgttcagcagctgaactaggtatagatgaacattacattgaggagtagataaaagaggt ggtagatatatatttagaccagaagattcttagaattaggaaaagatataaaagatgttttaggattaaaaagatgc tttaatagattttgaattaaactcaaacagacactgattgtaaatgcattgaggttagctagagaagcagctgca actataggaacaaaagtaaaatctctgaaatcgaagtaaaagaaagtgcgaagagatagatttcaagttg agatagataaatccagattttatgtagaagatatgttgctagaatggttacagatgtaaaaatagaaccttctccata atggttagagcttggccaaccacttcatgcttttgatataaatcaagtagagactggaagaatagtagtaagaat gctaagatggagagaaacttgtaacattatagatgttgagagaaacttagataaaagatatgctagtataaca aatggagaaaaatcacttgggttagctgggtgaatgggtgggtgctaactcagaaataaacttctaatacgaagact gtacttttgaagtgccaaacttcaaacagaaaaacataagaatgacagctaaaaaagttgggtattaggtcagaa gcatcttcaagaaatgaaaaagacttagaccctaatcttgagagatagcagcaaatagagctgcacaacttgt tgaaatgttaggagcaggaaggttttaaaaggtgtgttagatgtatatccaaataaacagaaaccttaaaaaat ggttagtaaatcctcaagaaataaacacctattaggtgttagatgtaccaatggagcagttttaggaaatttaga atcatttagagtttaaatgtaatttggtagctaatgataaattagaatagatgtaccaagctttagaacagatatgg aacaagaagctgatgtatgggaagaaatagctagaatttatggatttgagaata (SEQ ID NO: 24)

[0119] All plasmids were constructed using *E. coli* DH5 α and transformed into *B. subtilis* using MC medium⁴⁹. MC medium is composed of 10.7 g/L potassium phosphate dibasic (Chem-Impex International), 5.2 g/L potassium phosphate monobasic (MilliporeSigma), 20 g/L glucose (MilliporeSigma), 0.88 g/L sodium citrate dihydrate (MilliporeSigma), 0.022 g/L ferric ammonium citrate (MilliporeSigma), 1 g/L Oxoid casein hydrolysate (Thermo Fisher Scientific), 2.2 g/L potassium L-glutamate (MilliporeSigma), and 20 mM magnesium sulfate (MilliporeSigma). Plasmids were extracted from *E. coli* DH5 α using Plasmid Miniprep Kit (Qiagen) for transformation of *B. subtilis*. *B. subtilis* was inoculated into MC medium and incubated at 37° C. for 2 hours. Extracted plasmids were added into *B. subtilis* cell culture and incubated 37° C. for another 4 hours. Transformed *B. subtilis* were selected on the LB agar plate with selective antibiotics. Double cross-over was verified for colonies by the replacement of a different antibiotic resistance gene at the integration locus.

DNA Detection Using Cell-Based Sensors

[0120] DNA sensor strain was inoculated from the -80° C. glycerol stock into LB medium with 100 μ g/mL spectinomycin and incubated at 37° C. with shaking (250 rpm) for 14 hours. On the next day, the OD₆₀₀ of overnight culture was measured by NanoDrop One (Thermo Fisher Scientific) and diluted to OD_{0.1} in 1 mL LB in 14 mL Falcon™ Round-Bottom Tube (Thermo Fisher Scientific) supplemented with 50 mM xylose (Thermo Fisher Scientific) and 100 μ g/mL spectinomycin (Dot Scientific). Xylose was added to induce the competence. Spectinomycin was added to avoid contamination from other bacteria but it was not required. The sample gDNA was quantified by the Quant-iT dsDNA Assay Kit (Thermo Fisher Scientific) and supplemented in sensor culture with known concentration. The DNA sensor culture was incubated at 37° C. with shaking (250 rpm) for 10 hours for transformation. Culture of transformed sensors (5 μ L) was plated onto a 12-well plate (Thermo Fisher Scientific) for selection. In these plates, each well contained 1 mL LB agar supplemented with 2 mM IPTG (Bioline), 5 μ g/mL chloramphenicol (MilliporeSigma), and MLS (1 μ g/mL erythromycin from Sigma-Aldrich and 25 μ g/mL lincomycin from Thermo Fisher Scientific). Antibiotics were used in agar to avoid contamination. GFP-expressing colonies were imaged using Azure Imaging System 300 (Azure Biosystems) using Epi Blue LED Light Imaging with 50 millisecond exposure time. CFU was counted manually. Transformation efficiency is defined as the ratio of CFU on selective plates (transformed *B. subtilis* with GFP expression) to the CFU on non-selective plate (total *B. subtilis*). To count CFU of total *B. subtilis*, cell culture was serially diluted in phosphate-buffered saline (PBS) (Dot Scientific) and plated onto LB agar plate supplemented with 5 μ g/mL chloramphenicol and MLS. Since the total *B. subtilis* CFU was similar for most conditions, CFU of transformed cells was used to indicate detection efficiency.

[0121] To quantify the sensitivity or specificity of sensors, cell culture was transferred to liquid LB medium with a 1:20 dilution after transformation with serially diluted DNA from 1500 ng/mL to 1 ng/mL. To test the specificity towards gDNA from different strains, 100 ng/mL purified DNA was used for transformation. LB medium was supplemented with 2 mM IPTG (Bioline), 5 μ g/mL chloramphenicol (Mil-

liporeSigma), and MLS. Diluted cell culture was transferred to a 96-well black and clear-bottom CELLSTAR® microplate with 100 μ L volume in each well (Greiner Bio-One). Plate was sealed with Breathe-Easy Adhesive Microplate Seals (Thermo Fisher Scientific) and incubated in the SPARK Multimode Microplate Reader (TECAN) at 37° C. with shaking for time-series OD₆₀₀ and GFP measurements. A threshold of GFP fluorescence 400 was used to determine the detection time for different DNA or different concentrations. The threshold was placed in the region of exponential amplification of GFP across all of the conditions where the difference in the detection time between different conditions was not affected much by the choice of threshold. Unpaired t-test was performed to determine if the detection time of specific DNA concentration is different than the background without DNA (N=4). The detection limit is the lowest DNA concentration with a statistical difference. A straight line was fitted to the mean detection time of the four technical replicates with statistical difference versus the logarithmic DNA concentration. DNA mass per mL in detection limit (1 ng, 62.5 ng, 4 ng, and 16 ng) and genome size (4639675 bp, 4857450 bp, 2827820 bp, and 4153430 bp) were converted to chromosome copy number by NEBioCalculator for *E. coli* (2.10×10^5), *S. typhimurium* (1.25×10^7), *S. aureus* (1.38×10^6), and *C. difficile* (3.75×10^6), respectively.

Bioinformatic Analysis of the Specificity of Target DNA

[0122] To analyze if the target sequence is conserved for the target species, we searched the 5000 bp of *E. coli* xdhABC, *S. typhimurium* sipBCDA, *S. aureus* hemEH, and *C. difficile* pheST DNA sequence within the same species in NCBI Nucleotide Collection Database. Nucleotide BLAST was optimized for somewhat similar sequences (blastn). The search was specified to taxid 561 for *E. coli*, taxid 28901 for *S. enterica*, taxid 1280 for *S. aureus*, and taxid 1496 for *C. difficile*. Accession date of data is 2022-08-19. For the same strains with target sequence with more than 95% identity similarity and 95% coverage, the percentage is 99% (N=3462), 94.3% (N=2078), 95.6% (N=1488), and 96.4% (N=139) for *E. coli*, *S. typhimurium*, *S. aureus*, and *C. difficile*, respectively. To analyze if the target sequence is conserved in other species, the same search was performed excluding the same species. Homologs with varying identity similarity and coverage were found in different species for the target sequences *E. coli* xdhABC (N=5000), *S. typhimurium* sipBCDA (N=117), *S. aureus* hemEH (N=2993), and *C. difficile* pheST (N=5000). Dot plot of homology coverage and identity similarity of each hit were shown for each target sequence.

Multiplexed DNA Detection Using Cell-Based Sensors

[0123] Overnight cultures of DNA sensor strains EC-G, ST-R and SA-B were diluted to OD_{0.1}, OD_{0.1}, and OD_{0.01}, respectively, in one single culture containing 1 mL LB supplemented with 50 mM xylose (Thermo Fisher Scientific) and 100 μ g/mL spectinomycin (Dot Scientific). Different combinations of gDNA of *E. coli*, *S. typhimurium*, and *S. aureus* (200 ng/mL each) were added into the mixed culture containing three DNA sensor strains in separate 14 mL Falcon tubes (Thermo Fisher Scientific). The DNA sensor strains were incubated at 37° C. with shaking (250 rpm) for 10 hours and 5 μ L of cell culture was plated onto 12-well plates (Thermo Fisher Scientific). In these plates,

each well contained 1 mL LB agar supplemented with 2 mM IPTG, 5 µg/mL chloramphenicol, and MLS. Plates were incubated at 37° C. overnight for bacterial growth.

[0124] On the next day, each well was imaged using Nikon Eclipse Ti-E Microscope. Brightfield images were collected at 4× magnification using the built-in transilluminator of the microscope. Fluorescence images were collected with the epifluorescence light source X-Cite 120 (Excelitas) and standard band filter cubes including BFP (Excitation: 395/25 nm, Emission: 460/50 nm, Chroma), GFP (Excitation: 470/40 nm, Emission: 525/50 nm, Nikon) and Texas Red (Excitation: 560/40 nm, Emission: 630/70 nm, Nikon) to image BFP, GFP, and RFP, respectively. Pixels were processed with 8×8 binning when taking images. The exposure times for BFP, GFP, RFP were 1.5 ms, 1.5 ms, and 7 ms, respectively. Complete images of each LB agar well were generated from multipoint images after scanning a 24×24 mm area. Once the full images were assembled, each of the four channels were mapped to unity by the minimum and maximum pixel values of all images for that channel using ImageJ. Colonies of different colors were counted manually.

Multiplexed Detection in Complex DNA Samples Using NGS or Cell-Based DNA Sensors

[0125] Bacterial species *A. cacciae*, *B. thetaiotaomicron*, *B. longum*, *C. asparagiforme*, *S. typhimurium*, and *S. aureus* were inoculated from the -80° C. glycerol stock into YBHI medium and incubated at 37° C. anaerobically overnight. On the next day, cell culture of each strain was diluted to OD0.01 in YBHI (Passage 0) and incubated for 24 hours (Passage 1). Cell culture was diluted to OD0.1 into a fresh YBHI and incubated for another 24 hours (Passage 2). At each passage, cell pellet was collected for DNA extraction using DNeasy Blood & Tissue Kit (Qiagen). To extract *S. aureus* gDNA, 0.1 mg/mL Lysostaphin (MilliporeSigma) was added in the pre-treatment step in combination with enzymatic lysis buffer (Qiagen). Purified DNA was stored at -20° C. before further processing for NGS or cell-based detection.

[0126] To use NGS for characterizing microbial community composition, the V3-V4 region of the 16S rRNA gene was PCR amplified from extracted DNA using custom dual-indexed primers on a 96-well PCR plate (detailed method described in Clark et al. Nat. Comm., 2021⁵⁰).

[0127] PCR products from each well were pooled and purified using DNA Clean & Concentrator kit (Zymo). The resulting library was sequenced on an Illumina MiSeq using a MiSeq Reagent Kit v2 (500-cycle) to generate 2×250 paired-end reads. Sequencing data were demultiplexed using Basespace Sequencing Hub's FastQ Generation program. Custom python scripts were used for further data processing and sequences were mapped to the 16S rRNA reference database created using consensus sequences from Sanger sequencing data of monospecies cultures. Relative abundance was calculated as the read count mapped to each species divided by the total number of reads of all species.

[0128] To use cell-based DNA sensors to detect *S. typhimurium* and *S. aureus* gDNA in the community DNA, 1000 ng/mL extracted community DNA was added to the mixture of SA-G (OD0.1) and ST-R sensors (OD0.1) for the multiplexed detection of *S. typhimurium* and *S. aureus*. The DNA sensor strains were incubated at 37° C. with shaking (250 rpm) for 10 hours and 5 µL of cell culture was plated onto 12-well plates (Thermo Fisher Scientific) containing 1 mL

LB agar supplemented with 2 mM IPTG, 5 µg/mL chloramphenicol, and MLS. Plates were incubated at 37° C. overnight for bacterial growth. On the next day, each well was imaged using Nikon Eclipse Ti-E Microscope. Brightfield and fluorescence images of GFP and RFP were collected at 4× magnification using the same procedure described in the previous section for the multiplexed DNA detection. Colony numbers were counted manually. The fold change of mean colony numbers compared to Day 1 was compared with the fold change of relative abundance by NGS results.

Direct Detection of Target Species in the Co-Culture Using Cell-Based Sensors

[0129] *E. coli*, *S. typhimurium*, and *S. aureus* and the corresponding DNA sensor strains were inoculated from the -80° C. glycerol stock into LB medium and incubated at 37° C. with shaking (250 rpm) for 14 hours. *C. difficile* was separately inoculated in YBHI medium and incubated in an anaerobic chamber (Coy Laboratory). On the next day, cell culture of sensor and target strain were diluted to an OD0.1 each in a single culture containing 1 mL LB supplemented with 50 mM xylose with or without 100 µg/mL spectinomycin in 14 mL Falcon tubes (Thermo Fisher Scientific). Overnight culture of target strains (14 hr) was diluted in PBS (Dot Scientific) and plated onto LB agar plates or YBHI agar plates to determine the initial CFU of target bacteria in the co-culture (OD0.1), which was 1.22×10^8 CFU/mL, 1.07×10^8 CFU/mL, 3.2×10^8 CFU/mL, and 1.1×10^7 CFU/mL for *E. coli*, *S. typhimurium*, *S. aureus*, and *C. difficile*, respectively. Sensor and target strains were co-cultured at 37° C. with shaking (250 rpm) for 10 hours, and 5 µL of cell culture was plated onto 12-well plates (Thermo Fisher Scientific). Each well contained 1 mL LB agar supplemented with 2 mM IPTG, 5 µg/mL chloramphenicol, and MLS for the selection of transformed *B. subtilis*. The 12-well plates were incubated overnight at 37° C. On the next day, fluorescent colonies were imaged by Azure Imaging System 300 (Azure Biosystems) using the Epi Blue LED Light Imaging with 50 millisecond exposure time. Colonies with GFP expression were counted manually. To determine if the detection of target bacteria was via transformation, 1 µL (1 unit) DNase I (Thermo Fisher Scientific) was added to the 1 mL co-culture. One unit of DNase I can completely degrade 1 µg of plasmid DNA in 10 min at 37° C. according to the manufacturer's specification.

[0130] To improve the detection efficiency, overnight culture of *E. coli* was incubated at 90° C. in digital dry baths/block heaters (Thermo Fisher Scientific) for 10 min and placed on ice for 3 min before being transferred to the sensor culture containing 1 mL LB, 50 mM xylose, and OD0.1 of sensor strain for detection. Spectinomycin was not used in the sensor culture for heat-treated samples. To test the multiplexed detection of *E. coli* and *S. typhimurium* in mice cecal samples, 10 mg cecal samples were first resuspended with 100 µL LB and 50 mM xylose in 1.7 mL Eppendorf tubes (Dot Scientific). Different amounts of overnight culture of *E. coli* and *S. typhimurium* were spiked into cecal samples. The cecal samples were then incubated at 90° C. for 10 min and sat on ice for 3 min before being transferred to the mixed culture of EC-G and ST-R sensors (1 mL LB, 50 mM xylose, OD0.1 of EC-G, and OD0.1 of ST-R) for multiplexed detection. Cecal samples were collected from germ-free mouse experiments following protocols approved by the University of Wisconsin-Madison

Animal Care and Use Committee. Briefly, 8-week-old C57BL/6 gnotobiotic male mice (wild-type) were inoculated with 8 human gut bacteria—*Dorea formicigenerans*, *Coprococcus comes*, *Anaerostipes caccae*, *Bifidobacterium longum*, *Bifidobacterium adolescentis*, *Bacteroides vulgatus*, *Bacteroides caccae*, and *Bacteroides thetaiotaomicron* via oral gavage. After four weeks of colonization, mice were euthanized for cecal sample collection. Cecal samples were stored at -80°C and thawed for the use in multiplexed detection of spike-in *E. coli* and *S. typhimurium*.

REFERENCES

- [0131] ADDIN Mendeley Bibliography CSL_BIBLIOGRAPHY 1. Brophy, J. A. N. & Voigt, C. A. Principles of genetic circuit design. *Nature Methods* 11, (2014).
- [0132] 2. Levskaya, A. et al. Engineering *Escherichia coli* to see light. *Nature* 438, 441-442 (2005).
- [0133] 3. Bourdeau, R. W. et al. Acoustic reporter genes for noninvasive imaging of microorganisms in mammalian hosts. *Nature* 553, (2018).
- [0134] 4. You, L., Cox, R. S., Weiss, R. & Arnold, F. H. Programmed population control by cell-cell communication and regulated killing. *Nature* 428, (2004).
- [0135] 5. Danino, T., Mondragon-Palomino, O., Tsimring, L. & Hasty, J. A synchronized quorum of genetic clocks. *Nature* 463, (2010).
- [0136] 6. Liao, M. J., Din, M. O., Tsimring, L. & Hasty, J. Rock-paper-scissors: Engineered population dynamics increase genetic stability. *Science* (80-.). 365, (2019).
- [0137] 7. Saeidi, N. et al. Engineering microbes to sense and eradicate *Pseudomonas aeruginosa*, a human pathogen. *Mol. Syst. Biol.* 7, (2011).
- [0138] 8. Borrero, J., Chen, Y., Dunny, G. M. & Kaznessis, Y. N. Modified lactic acid bacteria detect and inhibit multidrug-resistant enterococci. *ACS Synth. Biol.* 4, (2015).
- [0139] 9. Scott, S. R. & Hasty, J. Quorum Sensing Communication Modules for Microbial Consortia. *ACS Synth. Biol.* 5, (2016).
- [0140] 10. Sexton, J. T. & Tabor, J. J. Multiplexing cell-cell communication. *Mol. Syst. Biol.* 16, (2020).
- [0141] 11. Ibáñez de Aldecoa, A. L., Zafra, O. & Gonzalez-Pastor, J. E. Mechanisms and regulation of extracellular DNA release and its biological roles in microbial communities. *Front. Microbiol.* 8, 1390 (2017).
- [0142] 12. Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J.-P. Bacterial transformation: distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.* 12, 181-196 (2014).
- [0143] 13. Carrasco, B., Serrano, E., Martin-Gonzalez, A., Moreno-Herrero, F. & Alonso, J. C. *Bacillus subtilis* MutS modulates RecA-mediated DNA strand exchange between divergent DNA sequences. *Front. Microbiol.* (2019). doi:10.3389/fmicb.2019.00237
- [0144] 14. Popp, P. F., Dotzler, M., Radeck, J., Bartels, J. & Mascher, T. The Bacillus BioBrick Box 2.0: Expanding the genetic toolbox for the standardized work with *Bacillus subtilis*. *Sci. Rep.* 7, (2017).
- [0145] 15. Dubnau, D. Genetic competence in *Bacillus subtilis*. *Microbiological Reviews* (1991).
- [0146] 16. Ji, M. et al. Engineering *Bacillus subtilis* ATCC 6051a for the production of recombinant catalases. *J. Ind. Microbiol. Biotechnol.* 48, (2021).
- [0147] 17. Zhang, X.-Z. & Zhang, Y.-H. P. Simple, fast and high-efficiency transformation system for directed evolution of cellulase in *Bacillus subtilis*. *Microb. Biotechnol.* 4, 98-105 (2011).
- [0148] 18. Silvaggi, J. M., Perkins, J. B. & Losick, R. Small untranslated RNA antitoxin in *Bacillus subtilis*. *J. Bacteriol.* 187, (2005).
- [0149] 19. Maamar, H. & Dubnau, D. Bistability in the *Bacillus subtilis* K-state (competence) system requires a positive feedback loop. *Mol. Microbiol.* (2005). doi:10.1111/j.1365-2958.2005.04592.x
- [0150] 20. Xi, H., Schneider, B. L. & Reitzer, L. Purine catabolism in *Escherichia coli* and function of xanthine dehydrogenase in purine salvage. *J. Bacteriol.* 182, (2000).
- [0151] 21. McClelland, M. et al. Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature* 413, (2001).
- [0152] 22. Stabler, R. A. et al. Comparative genome and phenotypic analysis of *Clostridium difficile* 027 strains provides insight into the evolution of a hypervirulent bacterium. *Genome Biol.* 10, (2009).
- [0153] 23. Soni, I., Chakrapani, H. & Chopra, S. Draft genome sequence of methicillin-sensitive *Staphylococcus aureus* ATCC 29213. *Genome Announc.* 3, (2015).
- [0154] 24. Tucker, S. C. & Galan, J. E. Complex function for SicA, a *Salmonella enterica* serovar typhimurium type III secretion-associated chaperone. *J. Bacteriol.* 182, (2000).
- [0155] 25. Lobo, S. A. L. et al. *Staphylococcus aureus* haem biosynthesis: Characterisation of the enzymes involved in final steps of the pathway. *Mol. Microbiol.* 97, (2015).
- [0156] 26. Beyer, D. et al. New Class of Bacterial Phenylalanyl-tRNA Synthetase Inhibitors with High Potency and Broad-Spectrum Activity. *Antimicrob. Agents Chemother.* 48, (2004).
- [0157] 27. Ibekwe, A. M. & Grieve, C. M. Detection and quantification of *Escherichia coli* O157:H7 in environmental samples by real-time PCR. *J. Appl. Microbiol.* 94, (2003).
- [0158] 28. Jung, B. Y., Suk, C. J. & Chang, H. K. Development of a rapid immunochromatographic strip for detection of *Escherichia coli* O157. *J. Food Prot.* 68, (2005).
- [0159] 29. Serrano, E., Ramos, C., Alonso, J. C. & Ayora, S. Recombination proteins differently control the acquisition of homeologous DNA during *Bacillus subtilis* natural chromosomal transformation. *Environ. Microbiol.* 23, (2021).
- [0160] 30. Giordano, N. et al. Cysteine desulfurase IscS2 plays a role in oxygen resistance in *Clostridium difficile*. *Infect. Immun.* 86, (2018).
- [0161] 31. Papagiannopoulou, C., Parchen, R., Rubbens, P. & Waegeman, W. Fast Pathogen Identification Using Single-Cell Matrix-Assisted Laser Desorption/Ionization-Aerosol Time-of-Flight Mass Spectrometry Data and Deep Learning Methods. *Anal. Chem.* 92, (2020).
- [0162] 32. Nisßler, R. et al. Remote near infrared identification of pathogens with multiplexed nanosensors. *Nat. Commun.* 11, (2020).
- [0163] 33. Dong, H. & Zhang, D. Current development in genetic engineering strategies of *Bacillus* species. *Microb. Cell Fact.* 13, (2014).

- [0164] 34. Ragheb, M. N., Merrikh, C., Browning, K. & Merrikh, H. Mfd regulates RNA polymerase association with hard-to-transcribe regions in vivo, especially those with structured RNAs. *Proc. Natl. Acad. Sci. U.S.A* 118, (2020).
- [0165] 35. Renda, B. A., Hammerling, M. J. & Barrick, J. E. Engineering reduced evolutionary potential for synthetic biology. *Molecular BioSystems* 10, (2014).
- [0166] 36. Law, J. W. F., Mutalib, N. S. A., Chan, K. G. & Lee, L. H. Rapid methods for the detection of foodborne bacterial pathogens: Principles, applications, advantages and limitations. *Front. Microbiol.* 5, (2014).
- [0167] 37. Rahmer, R., Heravi, K. M. & Altenbuchner, J. Construction of a super-competent *Bacillus subtilis* 168 using the Pmt1A-comKS inducible cassette. *Front. Microbiol.* 6, 1431 (2015).
- [0168] 38. Kurushima, J. et al. Unbiased homeologous recombination during pneumococcal transformation allows for multiple chromosomal integration events. *Elife* 9, (2020).
- [0169] 39. Burian, J. et al. High-throughput retrieval of target sequences from complex clone libraries using CRISPRi. *Nat. Biotechnol.* 2022 1-5 (2022). doi:10.1038/s41587-022-01531-8
- [0170] 40. Gu, W., Miller, S. & Chiu, C. Y. Clinical Metagenomic Next-Generation Sequencing for Pathogen Detection. *Annu. Rev. Pathol. Mech. Dis.* 14, (2019).
- [0171] 41. Liu, F., Li, J., Zhang, T., Chen, J. & Ho, C. L. Engineered Spore-Forming *Bacillus* as a Microbial Vessel for Long-Term DNA Data Storage. *ACS Synth. Biol.* (2022). doi:10.1021/ACSSYNBIO.2C00291/SUPPL_FILE/SB2C00291_SI_001.PDF
- [0172] 42. Gomes, A. L. C. et al. Genome and sequence determinants governing the expression of horizontally acquired DNA in bacteria. *ISME J.* 14, (2020).
- [0173] 43. Beauregard, P. B., Chai, Y., Vlamakis, H., Losick, R. & Kolter, R. *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proc. Natl. Acad. Sci. U.S.A* 110, (2013).
- [0174] 44. Tam, N. K. M. et al. The intestinal life cycle of *Bacillus subtilis* and close relatives. *J. Bacteriol.* (2006). doi: 10.1128/JB.0.188.7.2692-2700.2006
- [0175] 45. Sheth, R. U. & Wang, H. H. DNA-based memory devices for recording cellular events. *Nature Reviews Genetics* (2018). doi:10.1038/s41576-018-0052-8
- [0176] 46. Gonzalez, L. M., Mukhitov, N. & Voigt, C. A. Resilient living materials built by printing bacterial spores. *Nat. Chem. Biol.* 16, (2020).
- [0177] 47. Cooper, R. M. et al. Engineered bacteria detect tumor DNA in vivo. *bioRxiv* 2021.09.10.459858 (2021). doi:10.1101/2021.09.10.459858
- [0178] 48. Cheng, Y.-Y., Papadopoulos, J. M., Falbel, T., Burton, B. M. & Venturelli, O. S. Efficient plasmid transfer via natural competence in a synthetic microbial community. *bioRxiv* 2020.10.19.342733v2 (2020). doi:10.1101/2020.10.19.342733
- [0179] 49. Konkol, M. A., Blair, K. M. & Kearns, D. B. Plasmid-encoded comI inhibits competence in the ancestral 3610 strain of *Bacillus subtilis*. *J. Bacteriol.* 195, (2013).
- [0180] 50. Clark, R. L. et al. Design of synthetic human gut microbiome assembly and butyrate production. *Nat. Commun.* 12, (2021).
- [0181] 51. Overkamp, W. et al. Benchmarking various green fluorescent protein variants in *Bacillus subtilis*, *Streptococcus pneumoniae*, and *Lactococcus lactis* for live cell imaging. *Appl. Environ. Microbiol.* 79, 6481-6490 (2013).
- [0182] 52. Tack, D. S. et al. The genotype-phenotype landscape of an allosteric protein. *Mol. Syst. Biol.* 17, (2021).
- [0183] 53. Prindle, A. et al. Ion channels enable electrical communication in bacterial communities. *Nature* 527, (2015).
- [0184] 54. Miller, M. B. & Bassler, B. L. Quorum sensing in bacteria. *Annual Review of Microbiology* 55, (2001).
- [0185] 55. Du, P. et al. De novo design of an intercellular signaling toolbox for multi-channel cell-cell communication and biological computation. *Nat. Commun.* 11, (2020).
- [0186] 56. Marchand, N. & Collins, C. H. Peptide-based communication system enables *Escherichia coli* to *Bacillus megaterium* interspecies signaling. *Biotechnol. Bioeng.* 110, (2013).
- [0187] 57. Morens, D. M. & Fauci, A. S. Emerging Pandemic Diseases: How We Got to COVID-19. *Cell* 182, (2020).
- [0188] 58. Iqbal, S. S. et al. A review of molecular recognition technologies for detection of biological threat agents. *Biosens. Bioelectron.* 15, (2000).
- [0189] 59. Chiu, C. Y. & Miller, S. A. Clinical metagenomics. *Nature Reviews Genetics* 20, (2019).
- [0190] 60. East-Seletsky, A. et al. Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature* 538, (2016).
- [0191] 61. Gootenberg J S, Abudayyeh O O, Lee J W, Essletzbichler P, Dy A J, Joung J, Verdine V, Donghia N, Daringer N M, Freije C A, Myhrvold C, Bhattacharyya R P, Livny J, Regev A, Koonin E V, Hung D T, Sabeti P C, Collins J J, Zhang F. Nucleic acid detection with CRISPR-Cas13a/C2c2. *Science*. 2017 Apr. 28; 356(6336):438-442.

SEQUENCE LISTING

Sequence total quantity: 24
 SEQ ID NO: 1 moltype = DNA length = 107
 FEATURE Location/Qualifiers
 source 1..107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 1
 tgtaaataca gttctggatt gcgagaaaac gaagtaaaag gatataccgc ctgaattcag 60
 gccggattca ctggtagcga ccggcgctca ggatcctaac tcacatt 107

-continued

SEQ ID NO: 2 moltype = DNA length = 107
 FEATURE Location/Qualifiers
 source 1..107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 2
 tgccccaagc tagagagtcg aattcctgca gccctggcag gttatgtgtt taacaactca 60
 tatttccttaa tcttgcgata gaggcgtagca atgccgatgc ccagttc 107

SEQ ID NO: 3 moltype = DNA length = 107
 FEATURE Location/Qualifiers
 source 1..107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 3
 aactaccctt taataggagg gtccctcgtgt agacactttt cctttacagt aatgtaaata 60
 ggatgagtggt ctttttttct aagcatgaat aataatgtta aaagaga 107

SEQ ID NO: 4 moltype = DNA length = 108
 FEATURE Location/Qualifiers
 source 1..108
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 4
 ctgaatttga ttgcgagtga gatatttatg ccagccagcc agacgcagac gcgccgagac 60
 agaacttaat gggcccgccta acagcgcgat ttgctgggtga cccaatgc 108

SEQ ID NO: 5 moltype = DNA length = 809
 FEATURE Location/Qualifiers
 source 1..809
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 5
 ctcgagggtta aatgtgagca ctcacaattc attttgc aaa agttgttgac tttatctaca 60
 aggtgtggca taatgtgtgt aattgtgagc ggataacaat taagcttaca taaggaggaa 120
 ctactatgtc gacctatgaa tctctaattg tcatgatcgg ctttgccaat ttaataggcg 180
 ggattatgac atggggaata tctcttttaa cattattatt catgcttaga aaaaaagaca 240
 ctcacccat ttacattact gtaaaaggaaa agtgtctaca cgaggaccct cctattaaag 300
 ggtagtttct tttttaaaag cttagagtgt gccacactct ggcttttata ttttagcatt 360
 tctcatgaaa gtaacacaca ttaacaagtg gtaatgtggt aatgtggtac caactataag 420
 cttacgccag tagttgcaat acttttgctt ggcaccatta taacatgaat atattattgat 480
 tatataatta tttgtatctt ttatttggtta ctttttttat ctatgagttc aaaatgacct 540
 gatcatagaa gccttaaccc tttttctttt attaaaaacc ctcggattat gaaagtgtta 600
 tggtaacaata tggtttagta taaatgaata ttggctttca acatctcaag ggcggtctgg 660
 ctcactccct catgaaaagg ggtgatgcac gtgtcaacat ttcaagcatt aatgcttatg 720
 cttgtcttcg ggtcatttat aattgccctg ttgacttata taaagaagaa atagaccac 780
 cccttgagct cggcaaagta aaagggttaa 809

SEQ ID NO: 6 moltype = DNA length = 720
 FEATURE Location/Qualifiers
 source 1..720
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 6
 atggtttcta aaggtgaaga attgtttaca ggtgtgtgtc caattttggt tgaattggat 60
 ggtgatgtta atggtcataa attttctgtt tctggtgaag gtgaagggtga tgctacatac 120
 ggtaaatgga cattgaaatt tatttgtaca actggtaaat tgcagttcc ttggccaaca 180
 ttggttacaa catttgctta tggtttgcaa tgttttgctc gttatccaga tcacatgaaa 240
 caacatgatt tctttaaatc tgctatgcc aagggttatg ttcaagaacg tacaatcttt 300
 ttcaaggatg atggttaata taagacacgt gctgagggtta agtttgaagg tgatacattg 360
 gttaatcgta tcgaattgaa ggggtatcgat tttaagaag atggtaatat cttgggtcat 420
 aaattggaat ataattataa ttctcataat gtttatatca tggctgataa acaaaagaac 480
 ggtattaaag ttaattttta aattcgtcac aatattgaag atggttctgt tcaattggct 540
 gatcattatc aacaaaatac accaattggg gatggtccag ttttgggtcc agataatcat 600
 tatttgtcta cacaatctaa attgtctaaa gatccaaatg aaaaacgtga tcacatggtt 660
 ttgttggaat ttgttacagc tgtcgtgtatt acacatggta tggatgaatt gtataataa 720

SEQ ID NO: 7 moltype = DNA length = 714
 FEATURE Location/Qualifiers
 source 1..714
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 7
 atggttagca aaggcgaaga ggataaatg gcgatcatca aagaatttat gcgctttaaa 60
 gttcatatgg aaggcagcgt taatggccac gaatttgaaa ttgaaggcga aggtgaaggc 120

-continued

```

agaccgtatg aaggcacaca aacagcaaaa ctgaaagtta caaaaggcgg accgctgccg 180
tttgcattgg atattctgtc accgcaattt atgtatggca gcaaagcata tgttaaacat 240
ccggcagata tcccggaata tctgaaactg tcatttccgg aaggctttaa atgggaacgc 300
gtcatgaatt ttgaagatgg cggagttgtt acagtcacac aagattcatc actgcaagat 360
ggcgaattta tctataaagt caaactgcgt ggcacgaact ttccgtcaga tggccctgtt 420
atgcagaaaa aaacaatggg ctgggaagca tcaagcgaaa gaatgtatcc ggaagatggt 480
gcactgaaag gcgaaattaa acaacgctg aaacttaaag acggtggaca ttatgatgcg 540
gaagtcaaaa caacgtataa agcgaaaaaa cctgttcaac tgccctggcg atataacgtt 600
aacattaaac tggatatcac gagccataac gaagattata caatcgtcga acagtatgaa 660
agagcagaag gacgccattc aacaggcgga atggatgaac tgtataaata ctag 714

```

```

SEQ ID NO: 8      moltype = DNA length = 705
FEATURE          Location/Qualifiers
source           1..705
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 8
atgagcgaac tgatcaaaaga aaacatgcat atgaaactgt acatggaagg cacagtogat 60
aaccatcact ttaaatgcac atcagaaggc gaaggcaaac cgtatgaagg cacacaaaca 120
atgagaatca aagttgttga aggcggaccg ctgccgtttg catttgatat tctggcaaca 180
tcatttctgt atggcagcaa aacgtttatc aatcatacac aaggcatccc ggattttttt 240
aaacaatcat ttccggaagg ctttacatgg gaacgcgtta caacatatga agatggcgga 300
gtttctgacg caacacaaga tacatcattg caagatggct gcctgatcta taatgtcaaa 360
attagaggcg tcaactttac aagcaatggc cctgttatgc agaaaaaac actgggcttg 420
gaagcattta cagaaacact gtatccggct gatggcggac tggaaaggcag aaacgatatg 480
gcactgaaac tggttggcgg atcacatctg attgcaaaac tcaaaacaac gtaccgtca 540
aaaaaacccg caaaaaatct gaaaatgcct ggcgtctatt atgtcgatta tagactggaa 600
cgcattcaag aagcgaacaa cgaacacatg gtcgaacaac atgaagttgc agttgcgaga 660
tattgcgacg tgcggtcaaa actggggccat aaactgaatt actag 705

```

```

SEQ ID NO: 9      moltype = DNA length = 464
FEATURE          Location/Qualifiers
source           1..464
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 9
caattaccat cgaatgcacc attaacggga tgccttttca gcttcacgcc gcaccaggca 60
cgccgctctc ggaattactc cgcgaacaag gactgctaag tgtcaaaaca ggggtgctgcg 120
tgggtgaatg tggtgccctg acggtgttgg tgcacggcac agcaatagac agttgcttat 180
accttgccgc tgggttgtaa ggaaaagaga tccgcacgct ggaaggtgaa gcgaaaggcg 240
gaaaactttc tcatgttcag caggcttatg cgaaatccgg cgcagtgcag tgcgggtttt 300
tgacgcctgg cctgattatg gctaccacgg caatgctggc gaaaccacgc gagaagccat 360
taaccattac ggaattctgt cgcggactgg cgggaaatct ttgtcgctgc acggggtatc 420
agatgattgt aaatacagtt ctggattgcg agaaaacgaa gtaa 464

```

```

SEQ ID NO: 10     moltype = DNA length = 497
FEATURE          Location/Qualifiers
source           1..497
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 10
ttatgtgttt aacaactcat atttcttaat ctgctgatag agcgtagcaa tgccgatgcc 60
cagttcatca gcaacttgct tcttctgtgt atgacgtgaa agcgccctgc ggatcatttg 120
cttttccatc tctccacagc ccgtgcccgc cgcacatcgc agtgacaggt gcgcctcact 180
gacctctgtt acatcaactt gtcccgttgt gccattatc agcagatttg gcggcaatag 240
cgtgctgtcg ataacttcac ctgaagggaac cacgttaaac agatattcca tcaaatgct 300
taactcgcgc aggtttccgg gccaacgatg cttacgcaat atttcgacga catcgggagc 360
aatgccagga taaaccgatc ccagacgacg ggtatgcaga tgtaaaaagt aatgcacca 420
tagttcaata tcttctgacg gttcacgcag cgggtggcaga gttatcgga taacattaag 480
tcggtagaag agatctt 497

```

```

SEQ ID NO: 11     moltype = DNA length = 1000
FEATURE          Location/Qualifiers
source           1..1000
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 11
atgacgcgaa actggagatc cactccccgc gcggtgttcg tttcgtcccg attaatggct 60
ttcacaccgg gccgggcaaa gtgtctcttg agcatgacga aatcctcgtc gcctttcatt 120
ttccgccaca gccgaaagaa cacgcgggca gcgcgcatct taaatatgcc atgcgcgacg 180
caatggatat ttcaacgatt ggctgcgcgc cacattgccg actggataac ggcaatttca 240
gcgaattacg cctggcattt ggtgttgccg gcgcaacgcc gattcgtgc caacatgccg 300
aacagactgc acaaaatgct ccattaaacc tgcaaacgct ggaagctatc agcgaatctg 360
tcctgcaaga tgtcgcctcg cgttcttcat ggcggggccg taaagagttt cgtctgcac 420
tcacccagac gatgaccaaa aaagtgatta gcgaagccgt cgcgcggcgg gggggaaaat 480
tgcaatgaat cacagcgaaa caattaccat cgaatgcacc attaacggga tgccttttca 540

```

-continued

```

gcttcacgcc gcaccaggca cgcgcgtctc ggaattactc cgcgaacaag gactgctaag 600
tgtcaaacaa ggggtgctgcg tgggtgaatg tgggtcctgt acggtgttgg tcgacggcac 660
agcaatagac agttgcttat accttgccgc ctgggctgaa ggaaaagaga tccgcacgct 720
ggaaggtgaa gcgaaaggcg gaaaactttc tcatgttcag caggcttatg cgaaatccgg 780
cgcagtgcag tgcgggtttt gtacgcctgg cctgattatg gctaccacgg caatgctggc 840
gaaaccacgc gagaaagccat taaccattac ggaattcgt cgcggactgg cgggaaatct 900
ttgtcgctgc acgggggtac agatgattgt aaatacagtt ctggattgcg agaaaacgaa 960
gtaaaaggat atccggcctg aattcaggcc ggattcactg 1000

```

```

SEQ ID NO: 12      moltype = DNA length = 1000
FEATURE            Location/Qualifiers
source              1..1000
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 12
aggttatgtg tttacaact catatttctt aatcttgcca tagagcgtag caatgccgat 60
gccaggttca tcagcaactt gcttcttctt gttatgacgt gaaagcgccct cgcggatcat 120
ttgcttttcc atctcctcca gcgcgctgcc gcccgcatca tcgagtgcac ggtgcgcctc 180
actgacctct gttacatcac tttgtccgtg tgtgccatta ttcagcagat ttggcggcaa 240
tagcgtgctg tcgataactt cacctgaagg aaccacgtta accagatatt ccatcaaatt 300
gcttaactcg cgcagggttc cgggccaaac atgcttacgc aatatttcga cgacatcggg 360
agcaatgcca ggataaacgg atccagacg acgggtatgc agatgtaaaa agtaatgcac 420
caatagttca atatcttctt gacgttcacg cagcgggtggc agagtatatc ggataacatt 480
aagtcggtag aagagatctt cgcggaattt accctcggca atgaactggg ccaaattctg 540
attagtgcga gaaatgatgc gaatgtcgac ttgtattggg ctactggcac caatcggcag 600
aatttcacgt gcctcaaatg cgcgcagtaa tttagcctgc aacattaatg gcataacacc 660
tatttcacgt agaaacagcg tgcgcgtatt cgcgcgctga atcaaccctg ttttaccggt 720
ggcagaagcg ccagttaaat cacctttaac ataaccgaac agttcgctct ccagaagctg 780
ctccggaatc cgcgcacagt tgatagcaat aaagggttta ttcgctcttc cgtcacaact 840
atggattgca cgggcgacga cttctttacc cgtgcgcgtt tcaccaacca ccataacgct 900
ggatgggctg ggtgcaatgc ggctaagtag tcgttttaac tgccgcataa caccggcactc 960
gccaaaccaat tgttcaatat gcgggttcac aggtgcattt 1000

```

```

SEQ ID NO: 13      moltype = DNA length = 1500
FEATURE            Location/Qualifiers
source              1..1500
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 13
cgtaacgcgg tgaagatggc tacccgtgtt gcaatcaata cactgccgct gacgcacaaa 60
cgggttatatg aagagttcca tctggcagga ttgatttgag gataacatca tgtttgattt 120
tgcttcttac catcgcgacg caacccttgc cgtatgccatc aacctgctgg ctgacaaccc 180
gcaggccaaa ctgctcgccg gtggcactga cgtactgatt cagctccacc atcacaatga 240
ccgttatcgc catattgttg atattcataa tctggcggag ctgcccggaa ttacgctggc 300
ggaagatggc tcgctacgta tcggctctgc aacgacattt acccagctaa tagaagatcc 360
tataactcaa cgtcatctcc cgcgcttatg tgcgtcgccc acgtccattg ctggacgcga 420
gatccgtaac gtcgctacct acgggtggaa tatttgcaac ggtgccacca gcgcagattc 480
tgccacgcca acgctaattt atgacgcgaa actggagatc cactccccgc gcggtgttctg 540
tttcgtcccc attaatggct ttcacaccgg gccgggcaaa gtgtctcttg agcatgacga 600
aatcctcgtc gcctttcatt ttcgccaca gccgaaagaa caccgcccga gcgcgcattt 660
taaatatgcc atgcgcgacg caatggatat ttcacacgatt ggctgcgcgc cacattgccc 720
actggataac ggcaatttac gcgaattacg cctggcattt ggtgttgccc gcaccaagcc 780
gattcgctgc caacatgcgc aacagactgc acaaaatgcg ccattaaacc tgcaaacgct 840
ggaagctatc agcgaatctg tcctgcaaga tgtcgcctcg cgttcttcat ggcgggccag 900
taaagagttt cgtctgcac tcatccagac gatgacaaaa aaagtgatta gcgaagccgt 960
cgccgcggcg ggggggaaat tgcaatgaat cacagcgaaa caattaccat cgaatgcacc 1020
attaacggga tgcttttcca gcttcacgcc gcaccaggca cgccgctctc ggaattactc 1080
cgcgaaacaag gactgctaag tgtcaaacaa ggggtgctgcg tgggtgaatg ttgtgcctgt 1140
acgggtgttg tcgacggcac agcaatagac agttgcttat acctgcccgc ctgggctgaa 1200
ggaaaagaga tcgcacacgt ggaaggtgaa gcgaaaggcg gaaaactttc tcatgttcag 1260
caggcttatg cgaatcccg cgcagtgcag tcggggtttt gtacgcctgg cctgattatg 1320
gtaccacacg caatgctggc gaaaccacgc gagaagccat taaccattac ggaaattcgt 1380
cgccgactgg cgggaaatct ttgtcgctgc acgggggtac agatgattgt aaatacagtt 1440
ctggattgcg agaaaacgaa gtaaaaggat atccggcctg aattcaggcc ggattcactg 1500

```

```

SEQ ID NO: 14      moltype = DNA length = 1500
FEATURE            Location/Qualifiers
source              1..1500
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 14
aggttatgtg tttacaact catatttctt aatcttgcca tagagcgtag caatgccgat 60
gccaggttca tcagcaactt gcttcttctt gttatgacgt gaaagcgccct cgcggatcat 120
ttgcttttcc atctcctcca gcgcgctgcc gcccgcatca tcgagtgcac ggtgcgcctc 180
actgacctct gttacatcac tttgtccgtg tgtgccatta ttcagcagat ttggcggcaa 240
tagcgtgctg tcgataactt cacctgaagg aaccacgtta accagatatt ccatcaaatt 300

```

-continued

```

gcttaactcg cgcagggttc cgggccaacg atgcttacgc aatatttcga cgacatcggg 360
agcaatgccg ggataaacccg atcccagacg acgggtatgc agatgtaaaa agtaatgcac 420
caatagtcca atattcttct gacgttcacg cagcggtggc agagtattcg ggataacatt 480
aagtcggtag aagagatctt cgcggaattt accttcggca atgaactggg ccaaattctg 540
attagtgcga gaaatgatgc gaatgacgac ttgtattggg ctactggcac caatcggcag 600
aatttcacgt gcctcaatag cgcgcagtaa tttagctgcg aacattaatg gcataccacc 660
tatttcacgt agaaacagcg tgcccgtatt cgcgcctga atcaaccctg tttaccggtt 720
ggcagaagcg ccagtaaatg cacctttaac ataaccgaac agttcgctct ccagaagctg 780
ctccggaatc ggcgcacagt tgatagcaat aaagggttta ttccgtcttc cgctcaactt 840
atggattgca cgggcgacga cttctttacc cgtgcgcgtt tcaccaacca ccataacgct 900
ggatgggctg ggtgcaatag ggctaagtgc tcgttttaac tgccgcataa cacggcactc 960
gccaaccaat tgttcaatat gcggttcacg aggtgcattt gctacagaaa aactggtatg 1020
cgattggtga aacgccatta aaaataattg tcggccctga atgttatgca attgaccaat 1080
gattaattca cttttatcgt cccatgaaac aatattgctg atattgctcc gggtaaaatt 1140
actctcaaat gtttaattgtc tgaacaggat aggtttccca ataataattt ttgacacaac 1200
accaagtgtt ttaaggcgag tctgattaac aaactgaacc cgattttcat catctacaac 1260
taatacgcgc tgatccatat tatcgatcat ggtcgcaaat attttactga tgttatcttc 1320
tgggccctga cctccagaaa gtttcgaaac aaaaatggtg gatataatgg gaacataatc 1380
agaaaattcg tcgtcaattat cactgatatg ctcttggtgc tcgtgggtaa cggcaatcaa 1440
acttatcacc ccaacacaaac gatcctgtga aatgacaggc gtaccagaaa atgctttttc 1500

```

```

SEQ ID NO: 15      moltype = DNA length = 2000
FEATURE            Location/Qualifiers
source              1..2000
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 15
cctgataaaa aggccatatac gtgctgggtg aacgaccgga agagccgtta atgtcgttaa 60
aagatttggc gatggagcgt ttctaccacc ctgaacgcgg cgggcagctc tctgctgaaa 120
gtcccatcaa aaccaccact aaccaccagg cgtttggctg tacctttggt gatctgacgg 180
tcgatattgc tctgtgcaaa gtcacacatca accgcaccc ccaacgttcat gattcagggc 240
atattcttaa tccactgctg gcagaaggtc aggtacacgg cggaaatggga atgggcattg 300
gtcggggcgt atttgaagag atgatcatcg atgctaaaa cggcggtggc cgttaacccca 360
atctgctgga ttacaaaaatg ccgacatgc cggatctgcc acaactggaa agcgcgttcg 420
tcgaatacaa tgagccgcaa tccgcatacg gacataagtc actgggtgag ccaccaataa 480
ttcctgttgc cgctgctatt cgtaacgcgg tgaagatggc taccggtgtt gcaatcaata 540
cactgccgct gacgccaaaa cggttatatg aagagttcca tctggcagga ttgatttgag 600
gataaacatca tgtttgattt tgettcttac catcgcgag caacccttgc cgtatgccatc 660
aacctgctgg ctgacacccc gcaggccaaa ctgctcgccg gtggcactga cgtactgatt 720
cagctccacc atcacaatga ccgttatcgc catattgttg atattcataa tctggcggag 780
ctgcggggaa ttacgctggc ggaagatggc tcgctacgta tcggctctgc aacgacattt 840
accagctaa tagaagatcc tataactcaa cgtcatctcc cggcggttatg tgcctgggcc 900
acgtccattg ctggaccgca gatccgtaac gtgcctacct acggtggaaa tatttgcaac 960
ggtgccacca cgcagatctc tgccacgcca acgctaattt atgacgcgaa actggagatc 1020
cactccccgc cgggtgttcg ttctgtcccg attaatggct ttacacccgg gccgggcaaa 1080
gtgtctcttg agcatgacga aatcctcgtc gcctttcatt ttccgcccaca gccgaaagaa 1140
cacgcgggca cgcgcgcat taaatatgcc atgcgcgacg caatggatat ttcaacgatt 1200
ggctgcgcgc cacattgcgc actggataac ggcaatttca gcgaattacg cctggcattt 1260
cgtgttgccg tgcacaacgc gattcgtcgc caacatgccc aacagactgc acaaaatgcg 1320
ccattaaacc tgcaaacgct ggaagctatc agcgaatctg tcctgcaaga tgtcgccccg 1380
cgttcttcat gggggccagc taaagagttt cgtctgcac tcatccagac gatgacaaa 1440
aaagtgatta cgaagccgct cgcgcggcg gggggaaaat tgcaatgaat cacagcgaaa 1500
caattaccat cgaatgcacc attaacggga tgccctttta gcttcacgcc gcaccaggca 1560
cgccgctctc ggaattactc cgcgaacaag gactgctaag tgtcaacaa ggggtgctgcg 1620
tgggtgaatg tgggtgctgt acgggtgttg tcgacggcac agcaatagac agttgcttat 1680
accttgccgc ctgggctgaa ggaagagaga tccgcacgct ggaaggtgaa gcgaaaggcg 1740
gaaaacttct tcatgttcag caggcttatg cgaaatccgg cgcagtgacg tcggggtttt 1800
gtacgcctgg cctgattatg gctaccacgg caatgctggc gaaaccacgc gagaagccat 1860
taaccattac ggaattcgt cgcggactgg cgggaaatct ttgtcgtgc acggggtatc 1920
agatgattgt aaatacagtt ctggattgcg agaaaaagaa gtaaaaggat atccggcctg 1980
aattcaggcc ggattcactg                                     2000

```

```

SEQ ID NO: 16      moltype = DNA length = 2000
FEATURE            Location/Qualifiers
source              1..2000
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 16
aggttatgtg ttaacaact catatttctt aatcttgca tagagcgtag caatgccgat 60
gcccagttca tcagcaactt gcttcttctg gttatgacgt gaaagccct cgcggatcat 120
ttgcttttcc atctctcca gcgcggtgcc gcccgcatca tcgagtgaac ggtgcgctc 180
actgacctct gttacatcac ttgctccgt tgtgccatta ttcagcagat ttggcgccaa 240
tagcgtgctg tcgataactt cacctgaagg aaccacgcta accagatatt ccatcaaatt 300
gcttaactcg cgcagggttc cgggccaacg atgcttacgc aatatttcga cgacatcggg 360
agcaatgccg ggataaacccg atcccagacg acgggtatgc agatgtaaaa agtaatgcac 420
caatagtcca atattcttct gacgttcacg cagcggtggc agagtattcg ggataacatt 480

```

-continued

```

aagtcggtag aagagatctt cgcggaattt accttcggca atgaactggg ccaaattctg 540
attagttgca gaaatgatgc gaatgtcgac ttgtattggg ctactggcac caatcggcag 600
aatttcacgt gcctcaatag cgcgcagtaa tttagcctgc aacattaatg gcataacacc 660
tatttcacgt agaaacagcg tgcgcgtatt cgcgcctga atcaaccctg tttaccggt 720
ggcagaagcg ccagtaaatg cacccttaac ataaccgaac agttcgctct ccagaagctg 780
ctccggatc cgcgcacagt tgatagcaat aaaggggtta ttccgtcttc cgctcaactt 840
atggattgca cgggcgacga cttctttacc cgtgcgcgtt tcaccaacca ccataacgct 900
ggatgggctg ggtgcaatc ggcataatgag tcgttttaac tgccgcataa cacggcactc 960
gccaaccaat tgttcaatat gcggttcac aggtgcattt gctacagaaa aactggtatg 1020
cgattggtga aacgccatta aaaataattg tcggccctga atgttatgca attgaccaat 1080
gattaattca cttttatcgt cccatgaaac aatatgctgc atatgtccat gggtaaaatt 1140
actctcaaat gtttaatggtc tgaacgggat aggtttccca ataataattt tttgcacaac 1200
accaagtgtt ttttaaggcag tctgattaac aaactgaacc cgattttcat catctacaac 1260
taatacgcgc tgatccatat tatcgatcat ggtcgcaaat attttactga tgttatcttc 1320
tgcccccctg tctccagaaa gtttcgaaac aaaaatggtg gatatatggc gaacataatc 1380
agaaaattcg cgtaaatatt catgatgatg cttctgttgc tcgtgggtta cggcaatcaa 1440
acttatcacc ccaacacaaac gatcctgtaa aatgacagcg gtaccagaaa atgctttttc 1500
gcggcaattt tctttactat cgcacacctc gcaaggggga tcgaagcgag actgtgtcac 1560
aactttttca gttttcgttt ccaggacgtg cgggagcagg cgtgagttgc cgctcaactg 1620
gcgaccaaga aacttcccat acgcgcctgt tccggcaacg cgacacaagt tttcatcaac 1680
gatctcaacc tcaagctgca aaacgctggc aagcattctg gcaaaacgct gaattgtcgg 1740
ttgaatttgc atcaatagc actgcgtagt agcaagctcc atagctttac ctccagact 1800
tacttaaaag tcgatacttg aagacgttga tgggtcacag atcatgatga tattaactca 1860
ggcgaatttg gctttgataa aaacataaga tttttatcat tttctaataa aattatggaa 1920
gagatatcac attctatat caatatgaga attacggcgg tgagtttatc aaactgaaga 1980
gagatagcct gcccttttat

```

```

SEQ ID NO: 17      moltype = DNA length = 2500
FEATURE            Location/Qualifiers
source              1..2500
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 17
aggagatgct aatcgcgtca cgggcaaacg tattcacagc gcagggttgc cggagtgtct 60
tgaataaggc cggaaaatct ttgaatggga aaacgcctgc gcagaatgcc agaaccagca 120
aggcaatttg cgcgcggcgc ttggcgctgc ctgttttagc tacacctcta acacctggcc 180
tgtcggcgta gaaatagcag cgcgcgcgct tctgatgaat caggatggaa ccatcaacgt 240
gcaaaagcgg cgcagcgaag tcggtcaggg tgccgacacc gtcttctcgc aaatggtggc 300
agaaacgttg ggggttccgg tcagcgacgt tcgcgttatt tcaactcaag ataccgacgt 360
tacgccgttc cgtccggcgc catttgccct acgcagagac tatgttgccg cgctcgcgct 420
gcgcagtgcg gcactattat taaaagagaa aatcatcgct cagccgcgag tcatgctaca 480
tcagtcagcg atgaatctga ccttgataaa aggcacatcc gtgtgtgttg aacgaccgga 540
agagccgtta atgtcgttaa aagatttggc gatggacgct ttctaccacc ctgaacgcgg 600
cgggcagctc tctgctgaaa gctccatcaa aaccaccact aaccaccagg cgtttggctg 660
tacctttggt gatctgacgg tcgatattgc gctgtgcaaa gtcaccatca accgcaccc 720
caacgttcat gatcagggc atattcttaa tccactgctg gcagaaggct aggtacacgg 780
cggaatggga atggggcatt gctgggcgct atttgaagag atgatcatcg atgctaaaag 840
cggcggtggt cgtaacccca atctgctgga ttacaaaatg ccgaccatgc cggatctgcc 900
acaaactggaa ccacgcttcg tcgaaatcaa tgagccgcaa tccgcatacg gacataagtc 960
actgggtgag ccaccaataa ttctctgttc cgtctctatt cgtaacgcgg tgaagatggc 1020
taccggtggt gcaatcaata cactgcccgt gacgcaaaaa cggttatatg aagagttcca 1080
tctggcagga ttgatttgag gataacatca tgtttgattt tgcttcttac catcgcgag 1140
caaccttgcg cgtatgcac cactgctggt ctgacaaccc gcaggccaaa ctgctcgccg 1200
gtggcactga cgtactgatt cagctccacc atcacatga ccgttatcgc catattgttg 1260
atattcataa tctggcggag ctcgggggaa ttacgctggc ggaagatggc tcgctacgta 1320
tcggctctgc aacgacattt acccagctaa tagaagatcc tataactcaa cgtcatctcc 1380
cggcggtatg tgctgcggcc acgtccattg ctggaccgca gatccgtaac gtctgtacct 1440
acgggtgaaa tatttgcaac ggtgccacca gcgcagattc tgccacgcca acgctaattt 1500
atgacgcgaa actggagatc cactccccgc gcgggtgtcg tttcgtcccc attaatggct 1560
ttcacaccgg gccgggcaaa gtgtctcttg agcatgacga aatcctcgct gcctttcatt 1620
ttccgccaca gccgaagaaa caccgggcca gcgcgcattt taaatatgcc atgcgcgagc 1680
caatggatat ttcaacgatt ggtcgcgcgc cacattgccc actggataac ggcaatttca 1740
gcaattacg cctggcattt ggtgttgccg ccgcaacgcc gatcgcgtgc caacatgcgc 1800
aacagactgc acaaaatgcg ccattaaacc tgcaaacgct ggaagctatc agcgaatctg 1860
tctgcgaaga tgtcgccccg cgttcttcat ggccggccag taaagagttt cgtctgcatc 1920
tcatccagac gatgacaaaa aaagtgatta gcgaagccgt gcgcgcggcg gggggaaaat 1980
tgcaatgaat cacagcgaag caattaccat cgaatgcacc attaacggga tgctttttca 2040
gcttcacgcc gcaccaggca cgcgcctctc ggaattactc cgcgaacaag gactgctaag 2100
tgtcaaacaa ggggtgctgc tgggtgaatg tgggtgctgt acggtgttgg tcgacggcac 2160
agcaatagac agttgcttat acctgcccgc ctgggctgaa ggaaaagaga tccgcacgct 2220
ggaaggtgaa gcgaaggcgc gaaaactttc tcatgttcag caggttatg cgaaatccgg 2280
cgagtgacg tgccgggttt gtacgccttg cctgattatg gctaccacgg caatgctggc 2340
gaaaccacgc gagaagccat taaccattac ggaattctgt cgcggactgg cgggaaatct 2400
ttgtcgtcgc acggggatc agatgattgt aaatacagtt ctggattgcg agaaaacgaa 2460
gtaaaaggat atccggcctg aattcaggcc ggattcactg

```

-continued

```

SEQ ID NO: 18      moltype = DNA length = 2500
FEATURE           Location/Qualifiers
source            1..2500
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 18
aggttatgtg ttaacaact catatttctt aatcttgcca tagagcgtag caatgccgat 60
gccaggttca tcagcaactt gcttcttgct gttatgacgt gaaagcgccct cgcggatcat 120
ttgcttttcc atctctccca gcgcggtgcc gcccgcatca tcgagtgaca ggtgcgcctc 180
actgacctct gttacatcac tttgtccgtg tgtgccatta ttcagcagat ttggcggcaa 240
tagcgtgctg tcgataactt cacctgaagg aaccacgtta accagatatt ccatcaaat 300
gcttaactcg cgcaggtttc cgggccaacg atgcttacgc aatatttcga cgacatcggg 360
agcaatgcca ggataaaccc atcccagacg acgggtatgc agatgtaaaa agtaatgcac 420
caatagttca atatctctct gacgttcacg cagcgggtggc agagttatcg ggataacatt 480
aagtcggtag aagagatctt cgcggaattt accttcggca atgaactggg ccaaattctg 540
attagttgca gaaatgatgc gaatgtcgac ttgtattggg ctactggcac caatcggcag 600
aatttcacgt gccccaatg cgcgcagtaa tttagcctgc aacattaatg gcataacacc 660
tatttcacgt agaaacagcg tgcctgtatt cgcgcctgca atcaaccctg ttttaccggt 720
ggcagaagcg ccagtaaatg cactttaac ataacggaac agttcgctct ccagaagctg 780
ctccggaatc cgggcacagt tgatagcaat aaagggttta ttcgctcttc cgtcacaact 840
atggattgca cgggcgacga cttctttacc cgtgcgcgtt tcaccaacca ccataacgct 900
ggatgggctg ggtgcaatag ggctaagtga tcgttttaac tgccgcataa cacggcactc 960
gccaaccaat ggtccaatat gcggttcacg aggtgcattt gctacagaaa aactggtatg 1020
cgattggtga aacgccatta aaaataattg tcggccctga atgttatgca attgaccaat 1080
gattaattca cttttatcgt cccatgaaac aatatgctgc atatgtccat gggtaaaatt 1140
actctcaaat gttaatggtc tgaacgggat aggtttccca ataataattt ttgacacaac 1200
accaagtgtt ttaaggcgag tctgattaac aaactgaacc cgattttcat catctacaac 1260
taatacgccc tgatccatat tctcgatcat ggtcgcaaat atttactga tgttatctcc 1320
tggccctcga tctccagaaa gtttcgaaac aaaaatgggt gatatatggc gaacataatc 1380
agaaaattcg cgtaaatatt cactgatatg ctctgttgcc tcgtgggtga cggcaatcaa 1440
acttatcacc ccaacacaac gcactgttaa aatgacagcg gtaccagaaa atgctttttc 1500
gcggcaattt tctttactat gcacaacctc gcaagggtga tcgaagcgag actgtgtcac 1560
aactttttca gttttcgttt ccaggacgtg cgggagcagg cgtgagttgc cgtcacaactg 1620
gcgaccaaga aacttcccat acgcgccctg tccggcaacg cgacacaagt tttcatcaac 1680
gatctcaacc tcaagctgca aaacgctggc aagcattctg gcaaaacgct gaattgtcgg 1740
tgaatttgc atcaactatg aactgcgtat agcaagctcc atagctttac cttccagact 1800
tacttaaaag tcgatccatt agacgttga tgggttcacag atcatgatga tattaactca 1860
ggcgaaattg gctttgataa aaacataaga tttttatcat tttctaataa aattatggaa 1920
gagatatcac attctatat caatatgaga attacggcgg tgagtttatc aaactgaaga 1980
gagatagcct gcccttttat cttatttctg atacttagca gcaataaaat aacgcgataa 2040
aaaaagccaa acgttttctg attttacaaa caaccagaaag ctggcatcaa tttgtgatca 2100
acccacacac ttaaccgtca aattagtctt ttgcagccgc gcggataatt ctggcacact 2160
tattgttagt accaggttat gctgtgaaaa caccaatcac tttggcaagt cacagtgaag 2220
taaaaccact tgctgtcat tccactaccg ggactttatg atgaaaactg ttaatgagct 2280
gattaaggat atcaattcgc tgacctctca ccttcacgaa aaagattttt tgttaacgtg 2340
ggaaacagac cgagatgaac tgaaacaagt actggacgtt gccgcagcat taaaagcact 2400
gcgtgctgaa aacatctcaa ccaaaagtct taatagtggg ttaggtattt ccgtattccg 2460
cgacaactcc acccgtaacc gcttctctta tgcctccgcg 2500

```

```

SEQ ID NO: 19      moltype = DNA length = 2500
FEATURE           Location/Qualifiers
source            1..2500
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 19
acgtagcagc aggggtatca acgtttgcat ttcaaggtgc cgggcttccc gtctacgct 60
ggtaccctgc tcttgcggtt atttttggtg gcacatatca agcgccctcaa cagcctctgc 120
cgccgctttg tcaacaagggt gcgtaagatt gctgcgggtt aacggatcta acgtacagcc 180
aaagttagtg tcaatgcagc tggcaatata gggcatcacc tctgcataa caagattcgt 240
cgataaattt cttaattcac cgcacgtggt atttttgata atatctaaca gctgcttttc 300
caggttttcc agcttcgcct cgcctttctt tgtttctggc agccatggcc caaaagctga 360
cttttctttc agggcatctt ttatgatttg cgcggtatag tctgccccca ccttcatcag 420
tagcgtcttc gccctcaggag aatcactggt ggcgttgagc gctgaacgaa agagcccgcc 480
aaactccatt atcgctttct taccggcgac attatttgaa ttggtaaaaa cttcttttaa 540
cgctcagcg tctttccgcg atttaacaaa tgcattccaga ctgcgctggt tgatcagcgc 600
gggaaaaatc tccagttgcg gccctttaat ttccctgag agcgtcgctg tggcactttc 660
tctgactgcg gaaagattcg ccgcaagatt cgtggcctgc gttttgatct cggctctgcat 720
acctggcatt atgacggggg gctgagtcct tacacttgta accattatta atacctctt 780
ctgttatcct tgcaggaagc ttttggcggg ttccaggctg ctacttatcg tactgctcag 840
cacttttacc aggttgcgtg acaatgaatt ggcattgcta tatttttgcg tcagcgtctg 900
taatgtggtt ttcataattt ctctctgcgc tttaaaaccc gactgccagg cttgatattt 960
ggcggttatc atttcagatt ttgagtcctt tcccgccgcg cctaaaccat caataacctg 1020
aaccattttt tgtaatggcg tcagatcaac ggtgacgaca taaccggatc cataagattt 1080
caggcagcta ttccgtaaat tcaattcact gagccactgt ctgccttccg cttcagtggc 1140
tactttaacg ccgctgcctg actgcgctgg aaataaaacg gtattactgt ttatttgatt 1200
atattttatt actaaactgt ttaaatcatt tttgagttag gtaacatcta gcttaacggt 1260

```

-continued

```

attaccgtcc ttacctggta ataaccagcc tcccattttg gaaagaatat cactgaaggc 1320
ctgataaaaa tcggtataga ctgcgacaac gttttcataa acgccagat agctgtcacc 1380
tatccgccgat atattttggg aaaccatata ccaaatctca gcatcagaaa tggttgttct 1440
cggtcgccgc ataggcgaag cgctaaataa ggccgacgtc ggccagaaa acgcgtccg 1500
caggttctca tttgtttctg cggataatga cagcgcgac ttcgccagcg cattcaggct 1560
gctggtcaac tgctggcgcg ccagcgtgcg ctgcctatta ttctcttcag agatcggtgg 1620
cgttgactgc agcgtctgct gtgcctggtg gatttttagta gccgctgctg ataatgaaat 1680
gatattctga ccgcgatggt ctgtggtaga cggtaaccag gcagtctcga cgtgctcgct 1740
cgccgagggg gtctgcggcc gttcggcaac gatccccga tgaggagaag cggaataatt 1800
ttgaatatta agcataatat cccagtttcg ccatacaggag cgcgattaaa tcacacccat 1860
gatggcggtat agatgacctt tcagattaag cgcgaatatt gcctgcgata gcagcgagtg 1920
cggatgcttt cgactggtta atgctctcca ttgttttcag catttctcga atcagcgctg 1980
tcgatttacg tgaactttca cgggcttcgt ccgatgcggt gctggcaacc cggttattca 2040
cctgctaata ttgctgctcg gaacgttcct gagtagcggc gtactgcccg gacgcccctg 2100
caataccacc gacgtgaccc gagttcttca taatcagatc gcccgctcat tgcattctgc 2160
gcgcatacat tcgggtcaca tccatgggtat tctgctcaag acgaatatcg gattcgacag 2220
actcaagacg tttcgacaga atagcctgat gttcagggga gatttggtta ttactgtctt 2280
taataccacc actttccgtg gcgctgggtt cggcattaga ttttagcgct gcatcattaa 2340
gatttttctg cgcactggta ccggttttct tcataattaa cgatttcaga gaatcgacgc 2400
cttcagcacc gagtttgacg ctattctgcc cgttcagcac gtttttaata ctgtggcttt 2460
cagtggtcag tttatcgatc ttccggcgat tatgtttaag 2500

```

```

SEQ ID NO: 20      moltype = DNA length = 2500
FEATURE           Location/Qualifiers
source            1..2500
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 20
cgcgcctctt tcattctgca gcccttata ttccagtttg gcgcccacgc cagtgtatcc 60
caactgaagc gcgctctggg aaatactacc ggacaacgca ttcactccct cgcgcatcat 120
ggagcttgcc gtgcgttttag ctgcatacaa actgactaat gacaacttac cagacagttt 180
gctatcagcg tggttcaacg tcagcattaa cgtattcgcg gcagccaaca gcgcaacggc 240
actggaagac attcgcgtaa tctcaaaaaa ctttccgact tctgcctgct gctcgcgtaa 300
ctgggttttg acaacctcat tcgctttagt cgtgacatta tttgccagag cattcaaatc 360
ctgattcatg tcggtatttt gaatactggc ttttaaaaag gacgtgatcg ttccgggggt 420
ttgcgttaat accctgggct caggcgcgct cagtgtagga ctcaacccca ggctcactgac 480
tttactgctg ctaataccaa tactattcag aatatcttta gcgctaaccg attgcgaagc 540
tgtctgtgaa ctattctcaa cagaatgatt atttaataaa gcggcgggat ttattccac 600
attactaatt aacataattt ttccctttta ttttggcagt ttttatgcgc gactctggcg 660
cagaataaaa cgcgaagcat ccgcattttg ctgtaccgca gaagacatgg cttttgcag 720
ttccgcgctt accctctggt ttccaccaaa tatctctacg gattgtttaa gccactgctg 780
aatctgatcc atggcaaaaac gggcgagcat aaaatcagca agcgccctcg tggcattttt 840
aataaatacg cctcggcaca caccaccggc tgactgggct gcggtattcg tgactccat 900
gcccacgcgc actttattta ggggtattacc taccagctct ttacttaagg cattcgtttg 960
caggcccatc ttgctaccca cattaccacg accgctagta atacgttgca tcccctgggt 1020
aaagagtttg ctgcgctttt gcgccaactg tttcagcacg ttaggcacca acttcttaat 1080
cgtttcgccc atcattttgc tcagcgcgctt acccagtttc gccgcgcgc ctttcccgac 1140
aactgcgacc accacaatga ccgccaccat ggcaatagcg gcgacaatcg caccacaat 1200
gctgcgggcc atctctgcg ttttcttatc gacgcctaata ccttcacgcg ctttggtaat 1260
cgcttgcaca atcagctcca ttaacggctt cagcacatgc tccataatcg ggtttagcgc 1320
ctgctgaata aacgacactc ccgtcgccgc cttcacaatt tcacggcca ccattaccgc 1380
aagtccacc gcagccagcg ccagactcgc cccaccggta aaaacagcgc ccacaacgct 1440
gacaatgggt agcagcgcgc cgaggacttt cccgatacat cccataatgc ggttcggttc 1500
ctcgcttttg cgcgtctctt cctggaattc agccgatttc ttttccatct ccgcctgacg 1560
cccttctgca aaggcggtga aaagcgcaag atcgttttgc aggcctttct cgtattttt 1620
gcccacaatc tcaataaaca tggccatgag catagttagg cgggcgacat ttgacagatt 1680
atcctgctca cctgggaaa cctgattctg agaggcgga ttagcggttc cctggaattt 1740
ggtcagaatg ttatccgctt cctcgctttt cgctttggcg tctgtgctcg ctttaaccgt 1800
gcctccgctg gccttatcta aggcctcttt cgcctctgtc gcttcttttc cggcctgttc 1860
taccgcggct tcagctttgt catagccggg gtcagccggg tccagcgatt gcaatttatt 1920
ttgcgcctgc tgcagttttt tggtcgcagc gtcataaaca ctcttgccg taccgctctt 1980
tttgatactg gcttcataga gatccgtcgc ctctgagcc tctccagag cgtctggaa 2040
ttctttcgat acctgaatcc ccattctctt ttgtgactca atcatcgct gccataccgc 2100
cagacgagac tccagttgag acagcgaaac atcgcccagt agggctatta acttgccaag 2160
cagtaatgct aattgcctt cgtctgagag tttttccgg gcggcgctcg tagggcgctt 2220
tagacccacc gtattaatag cgctctcgcc ggactttgtt ccgcttttaa ggtcgccgc 2280
tttcggtgcc accacatctt taaaagcttt atccgcgct tttaaaaagt ccgtgttctt 2340
acgaacgcct tcaaaagcgc cctcagcgag gcgcggtatt tgggtatatt cgctacggct 2400
aatgctactt gcgtcattta ccataattat tctctttctt gttcactgtg ctgctctgtc 2460
tccgcgcttt ttagcgcttc cagatagacc aacgcttttg 2500

```

```

SEQ ID NO: 21      moltype = DNA length = 2218
FEATURE           Location/Qualifiers
source            1..2218
                  mol_type = other DNA
                  organism = synthetic construct

```

-continued

SEQUENCE: 21

```

ctatcgtgtc atgtaatctt gcatccgata ttgcaacgct gtaaatgttt cgaagccatc 60
ctcttctaag aagtgccctc catcttccac gattcgcgaag ttccctccta atgcattcat 120
taaacgctgg gttcttttat atgaaacgta ttgtgcattt ttagaactca atccgtaaaa 180
attgtcaact ttctttttta tattatcgta atcaatgggt acattactta aatcaatatc 240
taaatctata tttctctgat ctcttttaaa gcccgctata ctaaaaaagc cttcaatcgg 300
ctgatcaatc atttcaatat attttaaagc tgtgattgaa cctaaccatc gtgttacaaa 360
atatgtatcc tttttgctga cattaatttg ttctgcata gcttcaatcc actgatccac 420
tgtcttcgct tcaggggatt caaaattaaa taatgttacg tcatatcctt ctaaagttaa 480
gttatgtccc aaccactgat accaatgatt tctactattt ccatgcatag aatgtacaat 540
aattacatct gtcactcatc tctctcctt caacttacta ctctttttct atttttaaaa 600
aaatgactga ttacctataa ttgtaaaata aaaacacctt aattagaatc gttatatcgc 660
aaagtgcacat ttctaattaa agtgtattgt catcatttca atatcattca aaaacagcta 720
aacctttgtc tctgtctcaa ttccacaaaa ataattcccg ctgaaagtat ctatatattac 780
acattacttc caccattata taacttaaaa atgactatata ttcatcaaac attatctaaa 840
ggcgctgcac ctacaccaac accatccaac aattaactta caactctgcg attacttctt 900
cagcagcaac tttaccttgt gtaatacaat caggtagtcc aaccgcttca aaagatgcac 960
cagttactct aagtcgtgga tatgtttgtt taatatgtgc ttgaatctgt ctaatttgtt 1020
gaatatgacc gacatggata tgtggcatac ttttcggcaa acgattgaca attgtaaat 1080
caggatcacc tttaaatgtc atcatttgac ttaaactctc acgtacaatc gataactaatt 1140
cattatctgt atgatcatca accacagtat cacctgggtt acctacatac gcacgaatca 1200
aaaccttacc ttctgggtga gtaaatggcc attttttcga tgtccaagta catgcggtaa 1260
tgtctgtatc actcgtcttc gcaatcacga agccagtagc atcataagta ttttcaatgt 1320
ctttttcatc aaatgccaat acaacagttg caacagtcgt actatccatc gttttaaagt 1380
aatcaaatgc tggatcttgt ccgaaccaat tcaaaaagac ttgatgcggt gttgtcacta 1440
atacccatc ggaatcacat tcttgttgat tactgtaaac aatttttatat tgcttttgag 1500
atgtaataat atcatccact gacgtattgt agcgtattgt cacaccttta tttttcacat 1560
ctgtttctaa tgcttcaata atgagcttta aaccatgctt aaattgtttg aattgtcctt 1620
ttgggtgcgc aggatataat tgtctttgtt tcagacgctt attttttcct tcttccatac 1680
cttttatcag acttccgaat gccctctctt tttctttaa ataggaaac gtactcatca 1740
aaacttaatt atcaatatcg gtaccataaa taccacccat taaaggctca attaatgtct 1800
caagtacctc attacctaatt ctgtctctga aaaatgcacc aacagaaatg tccaccatct 1860
gcatttgtat aggttttttg attaaatcta atcctgctct taattttacca agtggcgata 1920
ttaattttgt gtaacaaat ggtttaatat ctggtggaat acccataatt gaaccacctg 1980
gaatcggata taattttatt ttcgcaaaaa tatatgattg tccagtcgta tttgtacaaa 2040
tatcttgttc taatccaata tctttcgcta attctgtcat aatcggtttt ctacctaaat 2100
aagattcagg ccctagttca atcatataac catctttacg atacgattga atctttcccc 2160
cggacgatt cgatgcttca aagatgggta catcaatatt aggatcttgc tgttttaa 2218

```

SEQ ID NO: 22

moltype = DNA length = 2775

FEATURE

Location/Qualifiers

source

1..2775

mol_type = other DNA

organism = synthetic construct

SEQUENCE: 22

```

gacttgattt catcaacaat tgcaccgata aataatggat gtgtattcgg catttttggg 60
cgataataat tcgcaccaat atcatcgcaa acaactttac attcataatc attgtcataa 120
agcacctcta aatgctcaca taaaaaacct actggcgat atataaagtt tttatactga 180
tgtttttcat aataatcacg ttttaaatct tgtacatctg gccctaacca aggtgtacct 240
gtattacctt cagattgcc aaccaatcgc atatgttcaa tattagattg ttctttaatt 300
aaaagcgcag tatgttctag ttctgttgga tatggatcat tattcttttc gattaaacct 360
tttgccaac tatgtgcgga acaactaat accgtgtctt tatgttcttc ttccggattt 420
tgagctaatt tttcgttgac tttattcgtc caatattcaa taaatttagg ttgttcataa 480
taattgttca catgtgtaag ttgaatacca tattttgag ctctctcatc agcacgtttg 540
tcatatgato ctactgaaaa tgaagaataa tgtgttgcta gtactaccgt gattgcttca 600
gtaataccat catttgtcat ttgttcaacc gcatcttcca taaatgggtg aatgtgtttt 660
aatcctaagt atagtttaaa ttcaacatct gcatatgctt tatttaatgc tgaacctagt 720
gcatcagctt ggtcatctgt tgtacctgt aatgggtgata aaccacctat aaattcatat 780
ctatctttca aatcttgaag ttcttcttca gatggacgtt taccatgtct aatatctgta 840
taatatggct ctatgtcact ttctttataa ggtgtgccat aagccataac taataacccc 900
atttttttag ctattgataa taccctcctt taaatgaatt atctttcatg tgcttcaatg 960
taatactatg attatctttg tgtatatgtg tgtacgaatt cgcttacttt acgtaacgtc 1020
tctggttgca ctctgggaa aacaccgtgt cctaaattaa agatgtgttt accgttctcc 1080
ataccttgat ctaaatattg ttccaatctc tcttcaatga cattccatgg tgctaataaa 1140
attgatggat ctaaatctcc ttgtaattgt tttagtaacgc ctaattgtttg agcctgatta 1200
atagacgttc tccaatctag gccataataa tcaatcggtg aatcattcca ttcatgattt 1260
aaatgactgg caccatcac gaataaaatt accggcacat catgtttttc tttaacctca 1320
ctgatttaac gaatcatatg tggtttaatg taacgtctgt aatcctcgac atttaatgca 1380
cctaccatg aatcgaaaat ttgaatcaat tcggcacctg ctctgacttg agctgttaca 1440
tatttaacag atacatcaac taaatgattc attaaagcaa accatgttgc ttcatctcta 1500
tacatcatcg ctcttgtaaa atttgtaatt ttcatggttc cgccttcaat catatatgac 1560
gctaattgaa atgggtgccc agttaaactc attagcggca catttaactt ttcttctgtt 1620
aaaagtttaa ttgatctaa tacatatggt acatctcgtt cggggctctat ttgagaaagt 1680
ttctcaacat ctgaatttgt ttgatagga ttatgaatca ctggaccaat acccgattta 1740
atttctacat cgacaccaat tggctttaat ggtgtcataa tatctttgta taaaattgct 1800
gcatctgtat gataattatc aactggtaaa tgtgttacat aagcgcacaa ctccggctga 1860

```

-continued

tgtgtaatat	cgaatagtga	atatttttct	ttcaattttc	gatattctgg	ttgcgaacgg	1920
ccagcttgct	gcataaaacca	aacaggtgta	tgtgatgttt	cttcaccttt	gatcattttt	1980
aaaattgtat	tgttttttatt	atgcaccata	aaggcctcct	aaattaaaat	cattctttatc	2040
tatattatca	tatcgctcat	tcggttcgtat	tttcaataaa	taaatgtcat	aaaactgaca	2100
tttaatacata	gaactattta	ttgtaaattt	aaattctaaa	gtccattatt	ttgtatcatt	2160
actttctaaat	atctcgcaag	attcattata	gtaattttta	tcaattatta	atagtggtaa	2220
tgactagttt	atcatcgat	aataaataaa	aacataaggg	ggacctttca	tatgaagaaa	2280
ctatatacat	cttatggcac	ttatggattt	ttacatcaaa	taaaaatcaa	taacccgacc	2340
catcaactat	tccaattttc	agcatcagat	acttcagtta	tttttgaaga	aactgatggt	2400
gagactgttt	taaaatcacc	ttcaatatat	gaagtattta	aagaaattgg	tgaattcagt	2460
gaacatcatt	tctattgtgc	aatcttcatt	ccttcaacag	aagatcatgc	atatcaactt	2520
gaaaagaaac	tgatttagtg	agacgataat	ttcagaaact	ttggtggctt	taaaagctat	2580
cgtttgttaa	gacctgctaa	aggtacaaca	tataaaattt	atttcggatt	tgctgatcga	2640
catgcatacg	aagactttta	gcactctgat	gcctttaatg	accatttttc	aaaagacgca	2700
ttaagtcatt	actttgtgtc	aagcggacaa	cattcaagtt	attttgaag	atatctatac	2760
ccaataaaag	aatag					2775

SEQ ID NO: 23 moltype = DNA length = 2222
 FEATURE Location/Qualifiers
 source 1..2222
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 23

atgaagcaat	atatagtcac	tgggtgtggg	agatttggaa	gttcagttgc	gtctactatg	60
catcttttag	gacatacaag	aatggccaata	gacaaaaatg	aagattcagt	tcaaagtata	120
tctgacaagg	taacccattc	acttatagtg	gatgttactg	atgagcaagc	gttaagggtca	180
ttaggtttag	gtaactttga	tgtagcagta	gttgcaatag	gttctgatat	aagggcatct	240
ataatggcga	ctcttatagc	caaagaaatg	ggtgtagagt	tgataaatatg	taaggcaaaag	300
gatgaattac	aagctaaagt	gctttataaa	attggcgagc	atagagtgtg	atttccagaa	360
agagatatgg	gagtaagagt	tgacacacaat	ttagtttcgg	ataatatatt	agaccatatt	420
gaacttgacc	cagagtagtc	aattgttgaa	atcgtaactc	caaatagttg	ggttggcaag	480
acacttatag	agcttgaatt	aagagctaga	tatgagataa	ctgtacttgc	tataaaaaaca	540
ggtaaaaata	taaatgtttac	accttctcca	gatgaggaaac	ttacagccgg	aagtatccta	600
gttataatcg	gtcaaaaatac	tagtataaca	gcgataacat	ctggaaaataa	ggggataaatt	660
agaagaagat	aattttactat	ttaatatata	tttaattgca	atgaaagtta	agagtatcat	720
ataattatga	atagtttatat	gatactattt	ttatttaaac	gaaggtagta	tttatttgta	780
agattagata	aagagaagtt	aaattaaaag	taaggaggct	gtgctaatat	aaaaatttat	840
aagttattag	catttagata	aaatgattac	aaatataaac	agtaaggata	atgaaaagtt	900
aaagtataca	agagcactat	taaaatcaaa	gaataggaat	aaagagtcac	agttcataat	960
agaaggatag	agaatagtaa	tgcttgcaact	tgaatgtatg	gcaaaccttg	attatgtatt	1020
tatcaatgaa	gaatttgaaa	ataagaaaaga	acatgtaaaa	ctattagaag	atttggataa	1080
aaaaaacaca	aagatatata	agactactaa	taaaaacttt	aaagaattag	tggtacacaga	1140
aaataactca	ggaataatag	gtgtagtgttc	atttaagaaa	aaaaaattaa	gtgaaagtat	1200
aaataaaaaa	gataaatttg	tattgatatt	agatagaata	caagaccag	gaaatatggg	1260
gactataata	aggactgctg	attctgctgg	agtagatgcc	ataatagcac	taaaaggatg	1320
tgctgatata	tacaaccaca	aagtaattag	gtctactatg	ggttctattt	ttgatatgaa	1380
tataattgat	gcttcacaag	atgaaactgt	ggacatgctt	aatcatttgg	attttaatat	1440
agtttcaagt	tacttaataa	cagaaaattt	ttatgacaaa	atagattatg	gttcaaaagt	1500
agcattgggtg	ataggaacac	aagcaaatgg	aataaatgaa	gaacttgtat	caaagtctga	1560
tattttgggt	aagataacct	tatatggtaa	agccgagtcg	ttaaatgctg	cgataagctc	1620
tgctatactg	atgtatgaaa	taaaaaata	cttaatttaa	tgtattgtaa	taaatataat	1680
gtcatgttat	aatcttgta	taaatagtag	agtatcaaaa	atagtcataa	tatatataaa	1740
aaataaattaa	attaattata	tatttagatg	atatcatata	aaaaatgatt	gagttgatta	1800
tgtattagat	atatattata	taaaaaata	ttgagttaat	tatgcattag	atgtatatta	1860
tataaaaaat	aattgagtta	attatgcatt	agatatatat	tataaaaaaa	ataattaaat	1920
taattatgta	ttggatatat	attaaaaata	gtcacataat	ttgaatggaa	atgatataat	1980
actaaaaata	acaatataat	attgtaaatg	caatgaaaga	ggaaagtatt	ttgattaaac	2040
ttagtaaaaga	gataaaacac	taggctggga	gtgttttcta	agaggctcat	agaagttccc	2100
tctggagtaa	cagagctgaa	attttacagt	agggcttgac	gtcaaaaacg	cgtttaagtt	2160
gttagagggtg	gtttgatgat	ttttaaattg	ttaaactact	aggggtgtac	cgcgaaacta	2220
ta						2222

SEQ ID NO: 24 moltype = DNA length = 2530
 FEATURE Location/Qualifiers
 source 1..2530
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 24

tagacagggg	ttgaggggct	ttttttatag	aaaaaaacga	aaggggtgata	tgtgtgcaag	60
aaaaattact	tgctttacgt	gaagcagctt	tggtgcaaat	aaaagaagca	caaagcatag	120
aaagtgtaga	aaagtttaaga	tcttaagtact	taggaaaaaa	aggtgagata	actgccatac	180
ttaaagaaat	gggtaaaatta	tctgctgaag	aaagaccagt	agttgggtaag	gttgccaatg	240
aggtaagaga	aaacattgaa	cttagcataa	attctaaaaa	agaagaaata	aatgctattg	300
aaaaagaaag	aaaattaaaa	gaggaagtga	tagatgttac	tcaaccagga	aaagttttta	360
aggttgggaa	gaagcatcca	ataactcaaa	ttatagatga	agtaacagat	atatttatcg	420
gaatgggatt	ctctatagca	gaagggccag	aagttgagac	tgttgaaaac	aactttgacg	480

-continued

cattaacgc	tcctaaagac	catccatcaa	gagatatgag	cgatacatc	tatatcaatg	540
atgggggtatt	acttagaact	caaacatctc	cagttcaagt	aagaactatg	agaagtcaag	600
agttaccaat	aaaagtaatt	gcaccaggta	gatgttttag	gtcagactcg	ccagatgcta	660
cacactcacc	aatgttccat	cagatagaag	ggcttgtgtg	tggaaaagat	gttactatgg	720
cagaatttaa	aggaactatg	gatatcttcg	ttgaaaaatt	gtttgggtct	gatatcaaaa	780
ctaagtttag	acctcacac	ttcccattta	cagaaccaag	tgcagaggtt	gatgttactt	840
gtttcaaatg	tggtggtaaa	gggtgcccac	tgtgtaaata	tgaaggttgg	atagaaatat	900
taggttcagg	tatggttcat	ccaaatgtgc	ttagaaattg	tggaaatagac	ccagaagttt	960
acagtgatt	tgcatttggg	gttgggggtg	aaagacttgc	aatgcttaaa	tacgaaatag	1020
atgatattag	attattattc	gaaaatgata	tgagattctt	aaatcaattt	taattaggag	1080
ggtagttaga	tgttagtatt	tttaaatggg	cttagagact	atgttgatat	agacatggat	1140
gtaaaagagt	tcgctgataa	aatgacaatg	acaggaaacta	aagttgaaac	aatagattat	1200
tatggtgaag	aaatagaaaa	tatatgtggt	ggaaagattt	tagaaataaa	acaacatcca	1260
aatgctgata	agttgggtgt	aactaaagta	gatattggag	ataaagttgt	tcaaatagtt	1320
acaggagcta	caaataatc	agaaggagat	tatatccag	tagctgtaaa	tggttctaag	1380
ttacctggag	gagttgaaat	caaacagact	gatttcagag	gtgaattatc	agatggtatg	1440
atgtgttcag	cagctgaact	aggtatagat	gaacattaca	ttgaggagta	taaaagaggt	1500
ggtagatata	tttttagacca	cgaagattct	tatgaattag	gaaaagatat	aaaagatgtt	1560
ttaggattaa	aagatgcttt	aatagatttt	gaattaactt	caaacagacc	tgattgtaaa	1620
tgcatgatgg	gtatagctag	agaagcagct	gcaactatag	gaacaaaagt	aaaatatcct	1680
gaaatcgaag	taaaagaaag	tgacgaagag	atagatttca	aagttgagat	agataatcca	1740
tatttatgta	gaagatatgt	tgctagaatg	gttacagatg	taaaaataga	accttctcca	1800
tattggatgc	aaagaagact	tacagaagca	ggagtaagac	ctataagtaa	catagtcgat	1860
ataacaaact	tcgtaattgt	agagcttggt	caaccacttc	atgcttttga	tataaatcaa	1920
gtagagactg	gaagaatagt	agtaagaaat	gctaagatg	gagagaaact	tgtaacatta	1980
gatgatgttg	agagaacatt	agataaagat	atgctagtta	taacaaatgg	agaaaaatca	2040
cttggtttag	ctggtgtaat	gggtggtgct	aactcagaaa	taacttctaa	tacgaagact	2100
gtactttttg	aaagtgcaca	tttcaaacca	gaaaacataa	gaatgcacgc	taaaaaagtt	2160
ggtagttagt	cagaagcatc	ttcaagaaat	gaaaagact	tagaccctaa	tcttgacagag	2220
atagcagcaa	atagagctgc	acaacttgtt	gaaatgttag	gagcaggaaa	agtttttaaaa	2280
gggtgtgtag	atgtatatcc	aaataaacca	gaacctaaaa	aattggtagt	aaatcctcaa	2340
agaatttaacc	acctattagg	tgtagatgta	ccaatggagc	agtttgtagg	aatttttagaa	2400
tcattagagt	ttaaatgtaa	tttggttagct	aatgataaat	tagaaataga	tgtaccaagc	2460
tttagaacag	atatggaaca	agaagctgat	gtatgggaag	aaatagctag	aatttatgga	2520
tttgagaata						2530

1. A cell-based DNA sensor comprising a competent cell comprising a genetic circuit, wherein the genetic circuit comprises:

a pair of homology arms in a DNA strand, wherein the homology arms comprise a first homology arm and a second homology arm, the first homology arm is homologous to a first portion of a target DNA, the second homology arm is homologous to a second portion of the target DNA, and the first homology arm and the second homology arm are separated within the DNA strand by an interstitial region of the DNA strand; and

at least one of a reporter switch and a kill switch, wherein:

the reporter switch comprises a reporter gene and a negative regulator of the reporter gene, wherein the reporter gene comprises a promoter and a coding sequence that are not comprised within the interstitial region of the DNA strand, and wherein the negative regulator of the reporter gene is comprised within the interstitial region of the DNA strand; and the kill switch comprises one or more genetic elements configured for inhibiting growth of the competent cell, wherein at least one of the one or more genetic elements is comprised within the interstitial region of the DNA strand.

2. The DNA sensor of claim 1, wherein the negative regulator of the reporter gene comprises a repressor gene, wherein the repressor gene expresses a repressor that represses expression of the reporter gene.

3. The DNA sensor of claim 1, wherein the reporter gene is a fluorescent or luminescent protein gene.

4. The DNA sensor of claim 1, wherein the interstitial region comprises at least one of a growth inhibitor gene, a positive regulator of a growth inhibitor gene, and a negative regulator of a selectable marker gene.

5-7. (canceled)

8. The DNA sensor of claim 1, wherein the kill switch comprises a toxin gene and a repressor gene, wherein the repressor gene expresses an inducible repressor of the toxin gene, and wherein the toxin gene is comprised within the interstitial region.

9. The DNA sensor of claim 1, wherein the kill switch comprises a toxin gene and a positive regulator of the toxin gene, wherein one or both of the toxin gene and the positive regulator of the toxin gene is comprised within the interstitial region.

10. (canceled)

11. The DNA sensor of claim 1, wherein the kill switch comprises a counter-selectable marker gene comprised within the interstitial region.

12. The DNA sensor of claim 1, wherein the kill switch comprises a selectable marker gene and a negative regulator of the selectable marker gene, wherein the selectable marker gene is not comprised within the interstitial region of the DNA strand, and wherein the negative regulator of the selectable marker gene is comprised within the interstitial region of the DNA strand.

13. The DNA sensor of claim 12, wherein the negative regulator of the selectable marker gene is also the negative regulator of the reporter gene.

14. The DNA sensor of claim 1, wherein the genetic circuit comprises both the reporter switch and the kill switch.

- 15-16.** (canceled)
- 17.** The DNA sensor of claim **1**, wherein the target DNA is native DNA.
- 18-19.** (canceled)
- 20.** A composition comprising two or more cell-based DNA sensors, wherein each of the two or more cell-based DNA sensors is a cell-based DNA sensor as recited in claim **1**, wherein:
- the pairs of homology arms in the two or more cell-based sensors are each homologous to different target DNA sequences;
 - each of the two or more cell-based DNA sensors comprises the reporter switch; and
 - the reporter genes in the two or more cell-based DNA sensors express reporters that are each detectably different from each other.
- 21.** (canceled)
- 22.** A method of detecting target DNA with the cell-based DNA sensor of claim **1**, the method comprising:
- culturing the DNA sensor in a culture medium comprising the target DNA for a time effective to transform the DNA sensor with the target DNA; and
 - detecting the transformed DNA sensor.
- 23.** The method of claim **22**, wherein the target DNA is non-isolated cellular DNA.
- 24.** The method of claim **22**, wherein the target DNA is non-isolated bacterial DNA.
- 25-26.** (canceled)
- 27.** The method of claim **22**, wherein the cell-based DNA sensor comprises two or more cell-based DNA sensors, wherein:

- the pairs of homology arms in the two or more cell-based sensors are each homologous to different target DNA sequences;
 - each of the two or more cell-based DNA sensors comprises the reporter switch; and
 - the reporter genes in the two or more cell-based DNA sensors express reporters that are each detectably different from each other.
- 28.** (canceled)
- 29.** A method of detecting a target cell comprising target DNA with the cell-based DNA sensor of claim **1**, the method comprising:
- culturing the DNA sensor in a culture medium with the target cell for a time effective to transform the DNA sensor with the target DNA; and
 - detecting the transformed DNA sensor.
- 30.** The method of claim **29**, wherein the culturing is performed without lysing the target cell.
- 31.** The method of claim **29**, wherein the target cell comprises a bacterium.
- 32-33.** (canceled)
- 34.** The method of claim **29**, wherein the cell-based DNA sensor comprises two or more cell-based DNA sensors, wherein:
- the pairs of homology arms the two or more cell-based sensors are each homologous to different target DNA sequences from different target cells;
 - each of the two or more cell-based DNA sensors comprises the reporter switch; and
 - the reporter genes in the two or more cell-based DNA sensors express reporters that are each detectably different from each other.
- 35.** (canceled)

* * * * *